

LA-UR-21-21791

Approved for public release; distribution is unlimited.

Title: Quantum Monte Carlo calculations of nuclei with local chiral interactions

Author(s): Lonardoni, Diego

Intended for: One-Page Publication Highlights

Issued: 2021-02-23


Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Quantum Monte Carlo calculations of nuclei with local chiral interactions

Objectives

- We use the Auxiliary Field Diffusion Monte Carlo (AFDMC) method to calculate ground state properties of nuclei up to $A=16$.
- We employ local chiral interactions up to next-to-next-to-leading order ($N^2\text{LO}$), including consistent three-nucleon forces, fit to few-body observables.



Energies and charge radii of nuclei with $3 \leq A \leq 16$ employing local chiral interactions up to $N^2\text{LO}$ for different parametrizations of the three-body force ($E\tau$, $E1$) and coordinate-space cutoff $R_0=1.0\text{fm}$.

Impact

- The AFDMC method is successfully applied to the study of both closed- and open-shell nuclei employing the full two- plus three-body Hamiltonian in the imaginary time propagation.
- The employed local chiral interactions provide a good description of ground state properties (binding energies, charge radii, charge form factors) of nuclei up to oxygen, while simultaneously predicting properties of neutron matter compatible with neutron star observations.
- In addition to the Monte Carlo statistical uncertainties, we examine the theoretical uncertainties associated with the chiral expansion and the cutoff in the theory, as well as the associated operator choices in the three-nucleon forces. Harder interactions show independence of such a choice up to $A=16$.

Accomplishments

- Publications:
 - D. Lonardoni, J. Carlson, S. Gandolfi, J. E. Lynn, K. E. Schmidt, A. Schwenk, and X. B. Wang, [Phys. Rev Lett. 120, 122502 \(2018\)](#)
 - D. Lonardoni, S. Gandolfi, J. E. Lynn, C. Petrie, J. Carlson, K. E. Schmidt, and A. Schwenk, [Phys. Rev. C 97, 044318 \(2018\)](#)