

Initialized Earth System prediction from subseasonal to decadal timescales

Gerald A. Meehl^{1†}, Jadwiga H. Richter¹, Haiyan Teng², Antonietta Capotondi^{3,4}, Kim Cobb⁵, Francisco Doblas-Reyes^{6,7}, Markus G. Donat⁶, Matthew H. England⁸, John C. Fyfe⁹, Weiqing Han¹⁰, Hyemi Kim¹¹, Ben P. Kirtman¹², Yochanan Kushnir¹³, Nicole S. Lovenduski^{10,14}, Michael E. Mann^{15,16}, William J. Merryfield⁹, Veronica Nieves¹⁷, Kathy Pegion¹⁸, Nan Rosenbloom¹, Sara C. Sanchez¹⁹, Adam A. Scaife^{20,21}, Doug Smith²⁰, Aneesh C. Subramanian¹⁰, Lantao Sun²², Diane Thompson²³, Caroline C. Ummenhofer^{24,25}, Shang-Ping Xie²⁶

1. Climate and Global Dynamics Laboratory, National Center for Atmospheric Research, Boulder, CO, USA
2. Pacific Northwest National Lab, Richland, WA, USA
3. Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA
4. NOAA Physical Sciences Laboratory, Boulder, CO, USA
5. School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA
6. Barcelona Supercomputing Center and ICREA, Barcelona, Spain
7. Institució Catalana de Recerca i Estudis Avançats, Barcelona, Spain
8. Climate Change Research Centre, University of New South Wales, Sydney, NSW, Australia
9. Canadian Centre for Climate Modeling and Analysis, Environment and Climate Change Canada, Victoria, BC, Canada
10. Department of Atmospheric and Oceanic Sciences, University of Colorado, Boulder, CO, USA
11. School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, NY, USA
12. Rosenstiel School for Marine and Atmospheric Science, University of Miami, Miami, FL, USA
13. Lamont Doherty Earth Observatory, Columbia University, Palisades, NY, USA
14. Institute of Arctic and Alpine Research, University of Colorado, Boulder, CO, USA
15. Dept. of Meteorology & Atmospheric Science, Pennsylvania State University, State College, PA, USA
16. Earth & Environmental Systems Institute, Pennsylvania State University, State College, PA, USA
17. Image Processing Laboratory, University of Valencia, Valencia, Spain
18. Department of Atmospheric Oceanic and Earth Sciences, George Mason University, Fairfax, VA, USA
19. Joint Institute for the Study of the Atmosphere and Ocean, University of Washington, Seattle, WA, USA
20. Hadley Centre, Exeter, UK
21. College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, UK
22. Department of Atmospheric Science, Colorado State University, Ft. Collins, CO, USA
23. Department of Geosciences, University of Arizona, Tucson, AZ, USA

24. Department of Physical Oceanography, Woods Hole Oceanographic Institution, Woods Hole, MA, USA
25. ARC Centre of Excellence for Climate Extremes, Sydney, NSW, Australia
26. Scripps Institution of Oceanography, La Jolla, CA USA

[†]email: meehl@ucar.edu

February 22, 2021

Abstract

Initialized Earth System predictions are made by starting a numerical prediction model in a state as consistent as possible to observations and running it forward in time for up to ten years. Skillful predictions at time slices from subseasonal to seasonal (S2S), seasonal to interannual (S2I) and seasonal to decadal (S2D) offer information useful for various stakeholders, from agriculture to water resource management, and human and infrastructure safety. In this Review, we examine the processes influencing predictability, and discuss estimates of skill across S2S, S2I and S2D timescales. There are encouraging signs that skillful predictions can be made: at S2S timescales, there has been some skill in predicting the Madden-Julian Oscillation and North Atlantic Oscillation; at S2I in predicting the El Niño-Southern Oscillation; and at S2D, in predicting variability in North Atlantic sea surface temperatures. However, challenges remain, and future work must prioritize reducing model error, more effectively communicating forecasts to users, and increasing process and mechanistic understanding that could increase predictive skill and, in turn, confidence. As numerical models progress towards Earth System models, initialized predictions are expanding to include prediction of sea-ice, air pollution, terrestrial and ocean biochemistry which can bring clear benefit to society and various stakeholders.

[H1] Introduction

In recent decades there has been an increasing desire for climatic information on timescales from weeks, to months, to seasons and years. Such information offers clear benefits to society and various stakeholders alike. For instance, prediction of the hydroclimate could allow for better water resource management and improved agricultural maintenance, while temperature and wind predictions could provide critical information for infrastructure planning and expected energy consumption. To obtain this climatic information, initialized predictions on various near-term timescales must be used.

Initialized Earth System prediction describes a suite of climate model simulations wherein the starting conditions are set as close to observations as possible and the model run forward for up to 10 years (1). Internally-generated, naturally-occurring variability is therefore considered a key aspect of these time-evolving climate predictions (2). They differ from uninitialized simulations – or climate change projections – where internal variability is removed through ensemble averaging and focus is instead given to quantifying the effects of external forcing such as anthropogenic greenhouse gases (3,4).

Given the duration of simulations, initialized predictions span various timescales (**Fig. 1a**): subseasonal-to-seasonal (S2S; ~2 weeks to 2 months) (5, 6); seasonal-to-interannual (S2I; 2 to 12 months) (7); and seasonal-to-decadal (S2D; 3 months to 10 years) (1,2). In each case, efforts have focused on climate phenomena that also operate on similar timescales. For example, S2S research has concentrated on the Madden-Julian Oscillation (MJO) and sudden stratospheric warmings (SSW); S2I on the El Niño-Southern Oscillation (ENSO), North Atlantic Oscillation (NAO), Indian Ocean Dipole (IOD), Southern Annular Mode (SAM) and Quasi-Biennial

Oscillation (QBO); and S2D on slowly evolving oceanic processes such as Pacific Decadal Variability (PDV) and Atlantic Multidecadal Variability (AMV).

Distinct communities have therefore formed to coordinate research and perform initialized predictions at each timescale. Efforts such as the S2S Prediction Project and Database (5) and the Subseasonal Experiment (SubX (6)) emerged for S2S; the North American Multi-Model Ensemble [PR, hyperlink to: <https://www.cpc.ncep.noaa.gov/products/NMME/>] (NMME (7) and the Copernicus Climate Change Service [PR, hyperlink to: <https://climate.copernicus.eu/>]) (C3S) for S2I; and sets of hindcasts and predictions as part of the Coupled Model Intercomparison Project phase 5 (1,2) (CMIP5) and CMIP6 (8) for S2D.

While these communities are often separate, however, all rely on similar methodologies (**Table 1, Supplementary Tables 1-3**). Thus, there is potential for “seamless prediction” (9), whereby one framework can be used to address prediction across all timescales, with skill increasingly associated with external forcing as simulations progress (10) (**Fig. 1b**). Yet in practice, community differences with regards to initialization frequency, for example, make seamless prediction challenging (1,2).

In this Review, we bring together research on initialized predictions on timescales of weeks to years. We begin by outlining current methodologies for initialized predictions, incorporating discussion of the process, ensemble size, verification and prediction skill. We subsequently outline prediction at S2S, S2I and S2D timescales, before discussing priorities for future research that will increase the feasibility for seamless prediction.

[H1] Making Predictions

S2I research using initialized prediction has been taking place since the late 1980s (11). In contrast, it was not until 20 years later that initialized S2D climate predictions began, in turn, initiating a rapid acceleration of research from which operational systems are now routinely produced (12). We begin by describing the process of initialized prediction, focusing on the methodological aspects involving forecast verification and measures of prediction skill (the level of agreement between an initialized prediction and the observed state it is meant to predict).

[H2] Process of initialized prediction

Predictions for S2S, S2I and S2D timescales, ranging from weeks to years, use numerical models with components of (at least) atmosphere, ocean, land and sea ice that are started from a particular observed state. The process of bringing the model components into close correspondence to that observed state is termed initialization, and predictions that are started from such observed states initialized predictions. There are currently many activities taking place in the S2S, S2I and S2D communities with regards to initialized prediction, with key differences amongst centers regarding how models are used (**Table 1, Supplementary Tables 1-3**).

One key difference between the subseasonal and longer timescale systems is the origin of the model. Many S2S (and some S2I) prediction systems originate in the numerical weather prediction (NWP) community. As such, they tend to have the highest horizontal resolution in the atmosphere, largely $\sim 0.25\text{-}0.5^\circ$ (**Table 1**). Atmospheric initialization in these NWP-derived models uses data assimilation, such as 3D variational assimilation (as in the CMA model). Moreover, to produce the initial perturbations for ensemble generation, they sometimes use data

assimilation with an Ensemble Kalman Filter (14) (as in the ECCC model) or singular vectors (15) (as in the JMA model). In comparison, most S2I, and all but one S2D, prediction systems are based on climate or Earth System Models (ESMs) previously used for IPCC climate projections. In these cases, the majority of models have a horizontal resolution of $\sim 0.5\text{--}1^\circ$ (**Table 1**).

In addition to differences in the models and their resolution across prediction timescales, contrasts are also evident in the components that are initialized and the degree of coupling between Earth System components. In S2S predictions, for example, coupling between the atmosphere, ocean, land and sea ice is not considered crucial (**Fig. 1a**). As such, only a small number of models initialize the ocean and employ atmosphere-ocean coupling, but the majority initialize land surface conditions (**Supplementary Table 1**). For S2D predictions, however, oceanic processes are vital, and as a result, all models initialize the ocean and have at least partial coupling with the atmosphere and sea ice; only a fraction initialize the atmosphere and land surface (**Supplementary Table 3**). As S2I falls in the time window where predictability comes from all Earth System components (**Fig. 1a**), care is typically taken to initialize each of them. Atmospheric initialization is often achieved by interpolating an existing analysis to the model grid and generating an ensemble spread using the random field perturbation method (16) (as in CESM1 for S2S), the lagged ensemble method (17) (as in CCSM3), or nudging to reanalyses in coupled mode (19) (as in the CCCma model). A variety of approaches have also been used to initialize the ocean state, including a hindcast spin-up in an ocean forced by observed atmospheric conditions (20), nudging the ocean model to some observed ocean state (21), or using full ocean data assimilation (22). Land variables are initialized either by assimilation of land observations (23) or by running an offline land-only model that is forced with observed

atmospheric conditions (24). The initialization strategy also differs between the shorter and longer-term prediction models. All S2S and S2I prediction models use full fields (such as sea surface temperature, SST). By contrast, about half of the S2D modes use anomaly initialization, meaning an initial condition is constructed by adding observed (or reanalysis) anomalies to the model's climatology in order to minimize initialization shock and model drift (25, 26, 27).

As individual model components are often initialized in different ways, there is frequently no coupling between initial conditions for various parts of the Earth System, thereby creating an imbalance in the initial state of the model. New methodologies, such as weakly coupled and strongly coupled data assimilation, offer promising approaches to reduce initialization shock and imbalance in the model (28). In the weakly coupled approach, the assimilation is applied to each of the components of the coupled model independently, whereas interaction between the components is provided by the coupled forecasting system (28). In the strongly coupled method, however, assimilation is applied to the full Earth System state simultaneously, treating the coupled system as one single integrated system (28).

There are currently very few modeling centers that have been able to apply seamless prediction owing to numerous practical aspects (including initialization method, initialization frequency, number of ensemble members, among others). The most seamless system is currently operated by the UK Met Office which is providing S2S, S2I, and S2D forecasts operationally, using almost identical configurations of the model for all prediction systems (29). NCAR, although not an operational center, is also using the same models, CESM1 and CESM2, to generate S2S, S2I and S2D hindcasts (and predictions for research purposes) using the same modeling framework, although at this time initialization details vary among the three prediction systems.

[H2] Ensemble size

Ensemble size is an important aspect determining predictive skill and reliability. In most prediction systems, ensemble sizes typically range between 10 and 50 (**Table 1**). There is potential of increasing the number of ensembles by combining those from multiple systems (30) or time-lagged ensembles (31), or using other techniques such as subsampling (32, 33) to improve the ensemble properties. Typically, the more ensemble members, the higher the anomaly correlation coefficient (ACC), a measure of prediction skill. For example, at S2S timescales, ACC of global surface air temperature over land is ~0.29 when using only 4 CESM1 hindcast ensemble members (34), increasing to ~0.33 for 8 members, and ~0.36 for 16 members (**Fig. 2a**).

Very large ensembles are also advantageous for improving seasonal prediction skill of the NAO (35), including at S2D timescales (36, 33). For example, ACC values are ~ 0.6 for an average of years 2 to 8 when using 40 ensemble members (**Fig. 2b**) (37). Further increases in multi-year NAO skill with ACC of 0.8 are possible with a lagged ensemble of 676 members (33) as a result of the modelled signal to noise ratio being too small.

Yet, there are consequences in terms of computing costs when using more ensemble members. For instance, an S2S reforecast could run 16 years (SubX) * 4 members * 2 months long * weekly start dates for ~600 model years; an S2I example could run 30 years *9 members*1 year long* 4 start dates per year for ~1000 model years; and an S2D example (DCPP) could run 60 years*10 members *10 years long for ~6000 model years.

[H2] Verification using observations

A key element of initialized prediction is having a solid understanding of the climate phenomena that are being predicted. Analyses of observations in comparison to the model simulations are thus required. On S2S and S2I timescales, the observational record provides a good source of data to verify initialized hindcasts. For example, observations cover roughly 30 ENSO events and as many as 300 MJO cycles. However, these data have their limitations. For instance, 3D observations of the atmosphere and ocean are desired for prediction verification, for understanding of processes and mechanisms, and for initialization of the predictions in the first place (38). Yet such 3D gridded data are limited to the period of the satellite record (dating from the late-1970s) and to reanalyses that assimilate all available observations. Moreover, while several ENSO (and similar timescale) events have been observed, these can exhibit different expressions (39) and undergo large decadal-to-millennial variations (40, 41, 42), requiring a long observational record to perform robust analyses.

Researchers in the field of initialized Earth System prediction on S2D timescales often cite the short observational record as a factor inhibiting understanding. For example, with reliable observations limited to the latter half of the 20th century (43), only \sim 3 PDV or AMV transitions have occurred by which to compare to predictions. While some observations are available earlier in the 20th century, these are sparse and reanalyses are highly uncertain, making consistent comparisons of prediction skill between the pre- and post- satellite era difficult. Added to that, subsurface ocean observations and critical state atmospheric variables (such as surface winds) are crucial to understanding slow variations in the climate system (44), but such observations also have a very short duration. Moreover, it is also difficult to objectively separate forced (natural and anthropogenic) and internal decadal-to-multidecadal climate variability, adding

further challenges for S2D prediction verification and triggering debate of best practices for signal separation (45, 46, 47, 48).

Nevertheless, efforts are underway to improve methodological approaches and data provisions for prediction verification. The crucial need for better observations of the full depth of the ocean have started to be addressed by Argo floats, first for the upper 2000m (49) but with plans to be expanded to the full ocean depth (50).

Proxy-based reconstructions are also increasingly available, shedding light on processes associated with interannual and decadal timescales of variability (51) beyond that possible by instrumental observations. Indeed, the particular limitations of instrumental data length and coverage for verification of S2D predictions have pointed to paleoclimate reconstructions -- using trees, corals and speleothems -- to extend observations and provide further realizations of decadal variability (52; 40; 53; 54; 55; 42; 56) (**Fig. 3**). Additionally, such records can provide insights into the physical mechanisms associated with that variability, including westerly wind anomalies (51), upwelling, gyre circulation (57) and links among major modes of variability (58). Together with further advances in paleoclimate research – including paleoclimate synthesis (59, 60, 61, 62), paleo data assimilation techniques (63, 64, 65), and development and expansion of proxy system models and toolboxes (66, 67) – paleoclimate data will not only help with the verification of climate model simulations, particularly on the S2D timescale, but also will provide context for initialized predictions by providing insights into the timescales of variability beyond the instrumental record.

[H2] Bias correction and prediction skill

To account for model drifts and biases, the skill of initialized predictions is typically evaluated in terms of forecast time-dependent anomalies that are departures from some measure of mean climate. However, a prediction will drift rapidly from the initial observed state towards its own climatology owing to model error. These drifts start almost immediately in a prediction, and by lead year 1, are already considerable (**Fig. 4**).

The calculation of anomalies and correction of model biases are addressed together, typically by calculating and removing the model climatology. For S2S predictions, the common methodology is to calculate a lead time dependent model climatology from a set of hindcasts and to compute anomalies from this climatology. However, such a procedure is complicated owing to the inhomogeneous nature of current subseasonal prediction systems (6, 68). The climatology for S2I predictions is similarly accomplished by averaging over all years of the hindcast for a particular start time and lead or target time (68), thereby assuming stationarity of biases and drifts in the predictions.

For S2D predictions, model drift is acute and is addressed by multiple approaches for computing anomalies (**Fig. 4**). One method is to calculate the model climatology of drifts from hindcasts over a prediction period of interest (for example, the average of lead years 3 to 7), and subtract that climatology from each 3 to 7 year prediction (69); this approach works well for short timescale predictions where externally-forced trends are less of a factor, but can be problematic for longer timescales. An alternative method is to compute a mean time-evolving drift from a set of hindcasts, subtract that mean drift from a prediction, and compute anomalies as differences from the drift-adjusted prediction and time period (such as the previous 15 year average)

immediately prior to the prediction (70). This alternative approach better reduces the effects of an externally-forced trend, but raises the issue of how big a role the recent observed period should play in prediction verification. When long-term trends in the hindcasts differ from observations, a further method is to correct biases in the trends in addition to those in the mean model climatology over the hindcast period (71), though such an approach can yield an overestimation of the skill of the system.

Models can also underestimate the magnitude of predictable signals relative to unpredictable internal variability, especially on seasonal and longer timescales in the extra-tropical north Atlantic sector (33). This underestimation leads to the counterintuitive implication that models are better at predicting the real climate variability than they are at predicting themselves, a phenomenon termed the “signal to noise paradox”, when observed signal to noise ratios are larger than in models (72). Given that such features also occur in uninitialized climate simulations of the historical period (73; 74), and potentially in modelled responses to volcanoes and solar variations (72), they are not believed to arise from initialization itself. As a result of the signal to noise paradox, it is necessary to take the mean of a very large ensemble to extract the predictable signal and then adjust its variance (33).

Although discrepancies between signal to noise measures in models and observations highlight an important model deficiency, it also implies an optimistic potential to use adjusted climate model outputs to predict the observed system (36, 33). Additionally, there has been a growing interest in the influence of decadal variability on the predictability and skill of seasonal forecasts (75). Sometimes the impact of this variability can obscure the gradual skill improvements that are found from advancing the science and modelling (76).

Clearly a major challenge for initialized prediction on any timescale is the mean drift of the model away from its initialized state to its preferred systematic error state (**Fig. 4**). All the efforts at bias adjustment and drift correction arise from this fundamental characteristic of model error, but improvements in initialized prediction require increased understanding of the processes and mechanisms at work in the climate system in order to reduce model error.

[H1] S2S initialized predictions

All initialized predictions start with a particular observed state that could contribute to some combination of externally forced and internally generated variability. However, owing to the relatively short timescales, subseasonal (S2S) predictability is largely an initial value problem in which the atmosphere, ocean, land and sea-ice contribute to prediction skill through their memory of the initial state, and not external forcing (**Fig. 1**)). Considerable resources are therefore allocated to initialization of atmosphere and land, including generation of ensemble spread. Ocean initialization and coupling are additionally important, especially in tropical regions where sources of predictability can come from modes of variability such as the MJO (77; 6), as well as the stratosphere, both of which are now discussed.

[H2] Modes of variability

The MJO is recognized as one of the leading sources of S2S predictability (78) owing to the strong interaction between the tropics and extratropics on subseasonal timescales (79). For example, forecast models involved in SubX and the Subseasonal-to-Seasonal Project can predict

the MJO skillfully up to 4 weeks (5, 80, 81). Furthermore, skill has been shown in predicting the MJO in a multi-model framework consisting of six SubX models for week three predictions averaged over days 15-21 (6) (Fig. 5), whereby most reproduce the eastward propagation of outgoing longwave radiation anomalies. Some models, however, have difficulty in simulating the propagation of the MJO across the Maritime Continent (eastward of 120°E), the so-called Maritime Continent “barrier” (78). MJO-related Rossby wave propagation into the extratropics also provides predictability for extreme events such as storm tracks (82), atmospheric rivers (83) and tornadoes (84).

S2S predictability is also influenced by the NAO (itself influenced by ENSO (85)), sea-ice and the stratosphere (86), which has bearing on extremes in large regions of Europe and North America. Using the NCEP Climate Forecast System version 2 (CFSv2) and the Met Office Global Seasonal forecast System 5 (GloSea5), it has been suggested that the NAO exhibits predictability out to at least several months ahead (87, 88, 35). Indeed, all SubX models demonstrate significant NAO skill at week 3, specifically an ACC of ~ 0.27 to 0.5 (ref 6).

Similarly, the SAM is a source of predictability and prediction skill of rainfall, temperature and heat extremes over Australia (89, 90). Although SAM predictability is typically low beyond \sim two weeks, there is the potential to make seasonal predictions (91) because of its association with ENSO (92) and the influence of the stratosphere (81, 93).

Consideration of these modes offer ‘windows of opportunity’ in S2S prediction, where in certain situations, there could be better predictability owing to active periods of the MJO or certain large-scale atmospheric regimes, for example (94).

[H2] Initial state

Given that the land surface varies more slowly than the atmosphere, it provides a source of predictability for temperature and precipitation on S2S timescales, the greatest contribution coming from soil moisture (95). This predictability is most pronounced during boreal spring and summer when synoptic systems have a smaller influence on soil moisture variability. The contribution of soil moisture anomalies to subseasonal predictability also varies regionally, with the largest contribution in areas of strong land-atmosphere interactions (96). As such, the land-surface is initialized in most current operational subseasonal prediction systems and all research subseasonal systems (**Supplementary Table 1, 2**). In doing so, improved skill for S2S predictions of temperature and precipitation have been observed, although model errors impact the full realization of this skill (97, 98, 95).

The coupling of the atmosphere to the ocean and sea-ice are further thought to be important for predictability at lead times longer than two weeks, and accordingly, ocean-sea ice-atmosphere coupled models are routinely used in operational S2S initialized predictions. For Arctic sea ice, there is rising demand for reliable projections up to months ahead owing to increased human activities. Currently the best subseasonal models show skillful forecasts of more than 1.5 months ahead (99). Yet, many current operational forecast models lack skill even on timescales of a week (100). Hence, there is more work to be done to improve the S2S forecast skill of Arctic sea-ice variables, though many systems are capable of predicting sea ice extent on seasonal time scales, at least in some regions and seasons (101, 102, 103, 104).

Sea-ice conditions (such as the location of the sea-ice edge) can have significant feedbacks with the atmosphere and thus impact the forecast of the coupled system in initialized predictions

(105). For example, the largest midlatitude forecast skill improvements have occurred owing to improved Arctic predictions over eastern Europe, northern Asia and North America relating to sea ice reductions and anomalous anticyclonic circulation (106).

[H2] Stratosphere

The largest recognized influence of the stratosphere on the troposphere comes from extreme states of the stratospheric polar vortex, particularly SSWs. SSWs are followed by tropospheric circulation anomalies that can last up to 60 days and resemble the negative phase of the NAO (107, 108). S2S forecasts initialized near the onset of an SSW thus show increased skill for mid-to high-latitude surface climate (109), and seasonal predictability of the NAO is dependent on the presence of SSWs in ensemble predictions (110). While SSWs are not as common in the Southern Hemisphere, weakening and warming of the stratospheric polar vortex is predictable a season in advance, and through connections with a negative SAM, can offer some predictability of hot and dry extremes over Australia (81, 93).

The QBO can further influence the troposphere on S2S timescales. Specifically, phase changes in the QBO modify the strength of the stratospheric polar vortex (111), in turn affecting the subtropical jet and storm tracks (112, 113), and strength of the MJO (114, 115). For example, the phase of the QBO in the initial state influences the prediction skill of the MJO, with higher skill during easterly-QBO boreal winters compared to westerly-QBO winters and improved skill for lead times of 1-10 days (116). The prediction skill of the QBO itself is very high on the S2S timescales with ACC of 0.85 to 1.0 on a one month timescale (93).

[H1] S2I initialized predictions

S2I initialized predictions are relatively mature compared to S2S and S2D, as evidenced by the number of national operational meteorological services that maintain state-of-the-art initialized S2I prediction systems (7; 117). Primary sources and mechanisms of S2I predictability consist of slowly evolving boundary conditions of SST, land-surface conditions (moisture, snow cover), sea-ice variations (118) and stratospheric state. Additional predictability might be gained from atmospheric composition, not typically represented in S2I models. Each of these factors are now discussed.

[H2] ENSO

The largest source of S2I predictability is associated with ENSO. ENSO provides skill in predicting rainfall across the tropics (119) and surface climate across the globe given their teleconnections (120). This predictability skill is primarily derived from subsurface ocean processes (121). Specifically, given that winds and SST in the deep tropical Pacific are largely in equilibrium, and the sub-surface temperature or thermocline variations are in dis-equilibrium, capturing the latter in the initial state of ESMs offers predictability (121).

However, ENSO events exhibit a large diversity in spatial patterns, with the location of maximum SST anomalies ranging from the central Pacific to the far-eastern Pacific (39; 122). ENSO diversity raises predictability issues in terms of precursor mechanisms such as Pacific

Meridional Modes (123; 124; 125; 126; 127), forecast skill (128, 129), teleconnections (130), multi-year events (131) and interpretation in the paleo-record (132)--many of which remain unresolved.

Overall, current state-of-the-art prediction systems are able to predict SSTs in the eastern Pacific up to 6-9 months in advance with modest skill, especially for forecasts initialized in June and verifying in the following boreal winter. Yet, current prediction systems consistently struggle to predict through the boreal spring season, that is, the so-called spring prediction barrier. The rapid onset or initiation of canonical, eastern Pacific, ENSO events also remains a challenge to predict, largely because onset often requires stochastic triggers such as westerly wind bursts (133, 134). Indeed, inclusion of westerly wind bursts (or other triggers) as stochastic parameterizations has been found to improve model simulations of ENSO (135) and forecast skill (136). Prediction of different ENSO types appears to be limited to about one month (137), and owing to the models' systematic tendency to produce more warming in the east, strong eastern Pacific events are generally better predicted (that is, exhibit better forecast skill) than central Pacific events (7).

[H2] Other modes of variability

Tropical Atlantic SST anomalies are also predictable on S2I time-scales. SST anomaly variability in this region is broadly categorized into two spatial patterns. The first is often referred to as the “Atlantic Niño” and involves many of the feedback mechanisms noted for ENSO (138), but is shorter lived and weaker. In comparison to ENSO, however, Atlantic Niño are less studied and also less predictable (139;140). The second pattern of variability is referred

to the Atlantic Meridional Mode (87). It is estimated that the Atlantic Meridional Mode is predictable one to two seasons in advance, with the mechanisms for predictability largely stemming from near surface air-sea interactions (thermocline variability is of secondary importance). However, even with some indications of successful predictions in certain circumstances including interactions with the tropical Pacific (138), as with all timescales of initialized predictions, persistent regional systematic errors with current initialized Earth prediction systems continue to be a factor in limiting predictive abilities of tropical Atlantic S2I variability (141; 142).

Much like the Atlantic, Indian Ocean SST anomaly variability is weaker and less predictable than the Pacific, but is important for regional teleconnections and impacts. Indian Ocean SST variability has three distinct patterns of interest: the IOD, that can be triggered by ENSO but can also emerge independently (58; 143); a basin-wide pattern that is an ENSO teleconnection (144); and a meridional mode pattern that depends on near surface air-sea interactions similar to that in the Atlantic (145). Earth System prediction models typically struggle to predict the connection between ENSO and the IOD, the northward propagation of the meridional mode, and the persistence of the IOD, except in large amplitude cases (146). The IOD also can affect processes on the S2S timescale (147), including the MJO. There are also other possible sources of S2I predictive skill involving the NAO (148) and Atlantic Ocean state which appears to drive aspects of summer European rainfall (149).

[H2] Land Surface Processes

Slowly varying S2I soil moisture anomalies influence prediction skill of precipitation and temperature (150). Currently, the memory resulting from large soil moisture anomalies in the initial conditions is believed to last ~2-3 months (151), but there are case-by-case examples where predictability can be considerably longer under conditions where soil moisture anomalies persist for more than one season, particularly for surface temperature. Indeed, some seasonal temperature predictability has been confirmed to arise from soil moisture, but the realization of skill is severely hampered by model biases (152; 153). Thus, reducing model error in the land surface components could considerably improve forecast skill, as seen in a large sample of initialized Earth System prediction experiments (17).

[H2] Stratosphere

Improved surface prediction resulting from stratosphere-related processes has been demonstrated on the seasonal timescale: having a higher vertical resolution in the stratosphere in a GCM captures SSWs earlier compared to the standard model configuration and has a positive influence on the simulations of European surface climate (154). Southern Hemisphere SSWs also affect predictions of Australian extremes (81; 93). The QBO, discussed earlier with respect to S2S predictability, has also been shown to lead to enhanced predictability on seasonal timescales (155; 156), is predictable out to several years ahead (157), and can also involve the MJO (116).

[H2] Atmospheric composition and other possible sources of predictive skill

There are additional sources and mechanisms for S2I predictability that are not particularly well modeled in S2I prediction. For example, slowly evolving greenhouse gases such as carbon dioxide and methane are known to be a source of forecast skill owing to their role as external forcing agents (158). However, an approximate time-history of carbon dioxide, methane and chlorofluorocarbons is typically specified and not predicted, thus limiting the potential to capture S2I variability or regional effects. Moreover, dust and aerosol concentrations are known to affect human health, but these changes in atmospheric composition are usually not included in prediction systems.

[H1] S2D initialized predictions

There is a high level of interest in, and expectations of, initialized Earth System predictions on timescales beyond S2S and S2I. For example, even with their limitations, there is evidence of skill in predicting surface temperature over and above that of simple persistence (Fig. 6a,b), and also precipitation and sea level pressure when using large multi-model ensembles, albeit with less skill (36) . These skillful multi-year predictions of precipitation over land indicate potential benefit to communities, as demonstrated with summer drought indicators in major European agricultural regions being predictable on multi-year timescales (159). Here we review the evidence for processes and mechanisms acting on the S2D timescale that could contribute to the skill of initialized predictions (12; 36).

[H2] Modes of decadal SST variability

Processes and mechanisms have been identified that could provide skill for fundamental quantities like SST in initialized predictions. Attention has been focused on AMV (160), but predictions of PDV (160; 161) -- which are often described in terms of the Interdecadal Pacific Oscillation (IPO) (162) over the Pacific basin and the Pacific Decadal Oscillation (163; 164) over the North Pacific -- are also of interest. Other modes of variability associated with decadal timescales include the Meridional Modes (165) and the North Pacific Gyre Oscillation (166).

Basin-wide warming and cooling patterns of SSTs and upper ocean heat content (0-400 m averaged temperature) have also been shown to characterize decadal-timescale variability in the Indian Ocean (167, 168; 169), as have decadal variations of the IOD (56, 170). Decadal variability in the Indian Ocean could influence warming events near the Australian west coast (171; 172). Furthermore, a rapid rise in Indian Ocean subsurface heat content in the 2000s in observations and model simulations is associated with a redistribution of heat from the Pacific to the Indian Ocean and has been suggested to account for a large portion of the global ocean heat gain during that period (173, 174). IPO variability could thus be affecting Indian Ocean variability, transmitted through both the atmospheric and oceanic bridges (175). These low-frequency connections have been implicated in modulating interannual variability associated with the IOD on decadal timescales (176, 172).

One issue that remains to be resolved for S2D related to prediction skill is whether there are well-defined timescales of variability that are distinct from the background of climatic noise; that is, if there are modes of large-scale variability that might display a statistically significant spectral peak in the decadal-to-multidecadal range and that could be predicted. Such signals could offer the best prospect for long-term predictability, but on this timescale, there is more of a broad-band spectral peak. For example, CMIP5 control simulations showed patterns and multi-

decadal timescales of variability in the Pacific associated with the IPO that resemble observations but with lower amplitude (177). Moreover, analysis of three generations of climate models (CMIP3, CMIP5 and CMIP6) shows progressive improvement of climate models' simulations of PDV (178). However, there was no convincing evidence across these state-of-the-art coupled models for distinct oscillatory signals, other than on the interannual (3-7 year) ENSO timescales (179). These observations suggest, as noted previously, that low frequency variability on interdecadal timescales is characterized by broadband rather than oscillatory behavior.

[H2] Global temperatures

The idealized “rising staircase” (**Fig. 6c**) of global mean surface temperature (GMST) trends represents actual epochs of larger or smaller amplitude positive GMST trends (**Fig. 6d**) in a world with steadily increasing positive radiative forcing from increasing greenhouse gases (180). This increase in radiative forcing means that the entire Earth System warms continuously, but the manifestation of that warming at the Earth’s surface on decadal timescales depends on how heat is redistributed in the climate system: if more heat remains near the ocean surface, the GMST rate of warming will be larger, but if more heat is distributed into the deeper ocean, then the GMST trend will be reduced (44, 181).

It is recognized that the slowdown in the rate of GMST warming in the early 2000s was likely a combination of internal variability from the negative phase of the IPO (182, 183, 184, 185, 186) and/or variations in the strength of the Atlantic meridional overturning circulation (187), both of

which acted to re-distribute heat into the subsurface ocean. However, there is disagreement on whether the heat is primarily stored in the tropics (174) or at high-latitudes (181). External forcing from a collection of moderate sized volcanic eruptions (188) and from anthropogenic aerosols (189), might have also played a role in the slowdown, though their contribution is not entirely settled (190).

Initialized predictions have been shown to successfully predict the onset of the GMST warming slowdown, linked to increased ocean heat uptake in the tropical Pacific and Atlantic oceans (191; 183). Spatial patterns of predicted 20-year surface air temperature trends have been shown to depend on the initial state of the Pacific Ocean (192), with initialized model predictions exhibiting a large spread in projected multi-decadal global warming unless the initial state of the Pacific Ocean is known and well represented in the model. Apart from its connection to the recent global warming slowdown, the negative phase of the IPO has also been linked to regional climate changes at higher latitudes, including the rate of Arctic sea ice decrease in the early 2000s (193) and Antarctic sea ice expansion during that same period (194, 195).

Statistical methods (47) and initialized predictions (196, 197) foretold a transition of the IPO in the tropical Pacific from negative to positive in the 2014-2015 time frame, with a resumption of more rapid rates of global warming thereafter. There is observational evidence that this IPO transition also contributed to initiating rapid Antarctic sea ice retreat (198).

There is a chronic shortage of observed data in the ocean to document heat redistribution. In models, this redistribution has been shown to involve the subtropical cells in the Pacific, Antarctic Bottom Water formation and the AMOC in the Atlantic (44; 2), as well as changes in the zonal slope of the equatorial thermocline (182; 199) associated with changes in tropical

winds. However, deciphering decadal timescale variability in the observed climate system, and interpreting such variability in the context of initialized predictions, is complicated by the presence of external forcings (such as anthropogenic and volcanic aerosols and solar forcing) that can produce decadal variability in the Pacific (189) or Atlantic (200; 201) with similar patterns to presumptive internally generated decadal climate variability (180; 202, 203)

[H2] Interactions between ocean basins

Interactions between various ocean basins is one of the most compelling science questions that has arisen regarding the origins and nature of decadal climate variability, with implications for initialized prediction skill (160, 204, 205). For instance, if a skillful prediction of climate in one basin is achieved, then skillful simulations in the other basins could follow (if the models capture these connections realistically), thus improving the skill of initialized S2D predictions.

SST variability in one ocean basin can affect the others through the tropical large-scale east-west atmospheric Walker Circulation, though the direction of those influences differs (205, 206). For example, model simulations have indicated that decadal timescale variability in the Atlantic could produce decadal timescale variability in the Pacific (61; 207; 208; 209). Pacific decadal variability can also affect the Atlantic (210; 211; 194) and control a large fraction of decadal variability in the Indian Ocean (58, 172, 212, 213, 214). Similarly, the Indian Ocean could influence decadal variability in the Pacific (168; 204; 215). There also could be staggered responses based on decadal timescales, with the tropical Pacific driving the tropical Atlantic on interannual timescales, with the Atlantic then affecting the Indian Ocean and subsequently the Pacific on decadal timescales (216; 217). It has further been postulated that the tropical Atlantic

and Pacific Oceans are mutually interactive on decadal timescales, with each alternately affecting the other (206), and that the tropical Pacific could be driving the extra-tropical Pacific (218).

External forcing, particularly from time-evolving anthropogenic aerosols, is another factor that could produce decadal climate variability and inter-basin connections (200; 189; 219). Such fundamental interactions all currently fall under the heading of a compelling research frontier that, with increased understanding, will certainly advance the science of initialized prediction.

[H1] Summary and future perspectives

Numerical models initialized with observations for specific time periods and integrated forward in time provide a continuum of predictions on different timescales from S2S, S2I and S2D.

Results so far demonstrate initialized prediction skill for variables such as surface temperature and key modes of atmospheric and ocean variability. Such skill has been demonstrated, for example, for the MJO on S2S timescales, for ENSO on S2I timescales, and for surface temperatures in most ocean regions on S2D timescales. Yet despite progress in predictions and processes, there are still many challenges and priorities for future research.

[H2] Model error

Almost every science-related aspect of subseasonal to decadal climate variability has considerable uncertainty associated with it. Therefore, apart from fundamental scientific

understanding, perhaps the key obstacle to progress is model error, particularly resolving biases and drifts and drifts and errors in the signal to noise ratio. Progress thus requires model improvement, developments of which are difficult but not impossible. In recent years for instance, model development work has been undertaken in the coupled space, improving simulation of atmosphere-ocean phenomena that give rise to predictability (such as the MJO and ENSO), and therefore minimizing the exacerbation of drift when developed in isolation. Model improvements depend critically on our understanding of processes and mechanisms and how they work in the climate system since it is difficult to model what is not understood. Therefore, enhanced observational and analysis projects must continue to provide the knowledge base from which to make improvements to the model simulations.

Model error remains a significant obstacle against which future progress will be measured, with profound implications for possible applications to stakeholder communities. Such applications could include energy supply (wind, solar) and demand (220), agriculture (drought, freezing), transport (221) and numerous others spanning a range of timescales. Notably, S2S prediction could inform preparedness for specific large-scale extreme events weeks ahead (5), and S2I and S2D initialized predictions are beginning to inform planning at ranges between the seasonal to multi-decadal climate change time scales (222).

In addition to coupled model development, increased model resolution has also shown ability to improve model bias and signal to noise ratio. Consequently, the benefit of increased model resolution is one of the research frontiers of initialized prediction. However, such increased resolution must also be accompanied by comparable increases in the quality of the physical parameterizations such as cloud feedbacks and cloud-aerosol interactions (198). Though we are still very likely decades away from having global coupled models (and suitable machines)

capable of explicitly resolving processes that would improve model bias (such as atmospheric convection and ocean eddies), approaches have been developed to reduce computational cost and bias. These approaches include flux correction techniques (223); parameter estimation (224); reducing the precision of some variables (225); and stochastic modelling (226). Additionally, machine learning techniques are providing indications of improving predictive skill. For example, a deep-learning approach using a statistical forecast model has been shown to produce skillful ENSO forecasts for lead times of up to one and a half years (227). Utilization of GPU-based computer architectures could become useful and open the way to better parametrizations that depend on intensive calculations that can be addressed with GPU architectures.

[H2] Initialization

Integrating the vast amount of observed information into an Earth System model is central to the S2D prediction. Traditionally the most advanced data assimilation techniques were implemented in the atmospheric component. In the last decade, however, there have been growing interests in how to fully utilize relevant satellite and in situ observations to improve S2S and S2I predictions. Coupled ocean-atmosphere data assimilation (28, 228, 229) shows promising evidence that coupling can reduce “initialization shock” and improve forecast performance on time scales of weeks to decades (230). The advancement has led to coupled reanalysis products for both ocean and atmosphere (CFSR by NCEP, (231) and CERA by ECMWF, (232)) and is expected to substantially improve S2S and S2I predictions.

Compared to S2S and S2I predictions, there remain critical obstacles as to how to initialize decadal predictions. First, there is a lack of observations. S2D models need to be initialized in the 1960s and 1970s in order to calibrate the decadal prediction systems and achieve the potential to capture the evolution of low-frequency modes of variability (such as PDV and AMV). Reconstruction of global ocean subsurface temperature and salinity prior to the advent of Argo floats remain a large problem. Currently most modeling centers performing decadal predictions don't carry out their own assimilation exercise, rather they simply nudge some reanalysis products in the ocean and atmosphere (Supplementary Table 3). It has not been carefully investigated how to best initialize the ocean without reliable subsurface observations, and how the inhomogeneity of the observations can impact the model performance.

Building ensembles is another key obstacle to decadal prediction, as the common practice in the community is to use an ensemble of 10 members following the CMIP5 and CMIP6 experimental designs. A large ensemble consisting of 40 members can provide better opportunities for skillful predictions of low-frequency climate variability over land in selected regions (20). However, compared to the atmosphere, there is very limited understanding of the mechanisms and uncertainty associated with the low-frequency internal variability in the ocean owing to the lack of long-term observations of the subsurface ocean, and thus lack of guidance as to how to build the ensemble.

Machine learning methods could help address this problem, though lack of long-term subsurface ocean observations will always be a factor for the S2D timescale.

Finally, a major constraint is computational capability, both for initialization and for running adequate numbers of ensembles to improve skill (33). The future of initialized prediction will depend on computational resources balanced with factors involving

increased resolution, artificial intelligence, use of new high performance computing architectures, and developments in exascale computing.

[H2] Predictability of internal variability

There are considerable future challenges for understanding internal variability in the context of initialized prediction. These include the need to have a better understanding and better estimates of predictability. Additionally, research is needed regarding why models appear to underestimate the magnitude of predictable signals compared to unpredictable variability, and this involves the response to external forcing as well (233).

One issue that remains to be resolved for S2D initialized predictions is whether there are well-defined processes and mechanisms that, if initialized properly, could provide predictable signals distinct from the background of climatic noise. Signals from PDV and AMV offer the best prospect for long-term predictability. Strong low-frequency variability in paleoclimate “proxy” records, which is not captured by most climate models, suggests that either models do indeed underestimate low-frequency modes of variability, that proxy observations contain significant residual non-climatic sources of variation, or some combination thereof (234; 235, 236, 237). Even if there is no distinct low-frequency (oscillating) phenomenon, predictability on decadal timescales could also come from memory and slowly varying components of the Earth System such as the slow propagation of oceanic planetary waves (238; 239) or natural volcanic forcing (47), and initialization could be expected to contribute to skill in such cases.

[H2] Expanding predicted variables

There is interest, and corresponding applications, for expanding beyond the prediction of surface temperature, precipitation and SST. There have been efforts at predicting soil moisture with implications for drought prediction (240) and ecosystem respiration (241, as well as snowpack with ramifications for water resources (242; 243) and marine heat waves (244). There is also a great societal need for prediction of sea-ice on S2I and S2D timescales. Some S2I models show some skill in predicting sea-ice edge in the Arctic (245), while S2S models show a very wide range of skill in predicting the sea-ice edge in the Arctic, with the most skillful models producing useful forecasts up to 45 days (99). While the potential for skillful initialized predictions of Arctic sea-ice on S2S timescales has improved in the last decade, there is still a lot more to be explored and improved (101). We still need to understand what are the key processes driving sub-seasonal variations of sea-ice and improve the representation of these processes in the S2S models. Improved coupled data assimilation of the ocean, sea-ice and the atmospheric coupled system can help improve initial conditions for coupled forecasts and concomitantly the forecast skill of features that are sensitive to the initial state (14, 246; 247).

Other important aspects of the cryosphere relevant to initialized prediction on S2D timescales are ice sheets. As new interactive ice sheet simulations and spin-up procedures come increasingly online (248), this will provide an additional opportunity for initialized S2D predictions.

Air pollution and air quality are other very society-relevant applications which have been largely unexplored owing to the lack of inclusion of interactive tropospheric chemistry in most S2S, S2I and S2D models. However, new comprehensive ESMs, such as the Community Earth System

Model with the Whole Atmosphere Community Climate Model as its atmospheric component (CESM2-WACCM, 249) will be able to explore this research area.

In the broader Earth System, there is growing interest in predicting the biosphere and biogeochemical state variables and fluxes that could inform management decisions. Skillful initialized predictions of SST on S2S timescales can engender predictability of fish yields in the California Current System (250) and other Large Marine Ecosystems (251). S2S initialized predictions of heat stress and coral bleaching risk have also demonstrated considerable skill and have provided critical advanced warning for coral reef scientists, managers, and stakeholders (252). SST anomalies in the western tropical Pacific and northern subtropics, often associated with ENSO events, appear to be skillful precursors for variations in temperature and related biological productivity along the U.S. West Coast at S2I timescales (253).

Emerging literature on S2D predictions of biogeochemistry in the terrestrial biosphere and ocean suggests that slowly evolving state variables could enable prediction of biogeochemically relevant quantities with greater skill than physical state variables such as temperature and precipitation. For example, predictions of marine net primary production by photosynthesizing phytoplankton (including algae, eukaryotes and cyanobacteria) might foretell future potential fisheries catch, predict harmful algal blooms (254), and aid with fisheries management strategies (255; 254; 256; 257), as would skillful predictions of ocean oxygen content or acidity (258; 259). Reliable forecasts of the changing global carbon budget, including the rate of ocean carbon absorption (217; 260; 261; 262) or the rate of terrestrial biosphere-atmosphere net ecosystem exchange (260; 241) could help to generate forecasts of atmospheric CO₂ growth rate and contribute to CO₂ emissions management strategies. Additionally, there has been demonstrated S2I skill at predicting net primary production related to fire risk (263).

Recently reported skillful predictions of chlorophyll concentrations over the global oceans at seasonal to multi-annual timescales have been related to the successful simulation of the chlorophyll response to ENSO, and to the winter re-emergence of subsurface nutrient anomalies in the extra-tropics (256). Chlorophyll not only responds to ENSO, but can also constitute a potentially useful ENSO precursor (264).

In the ocean biogeochemical system, variables of interest for prediction are rarely directly observed at the spatial and temporal scales needed for forecast verification, regardless of the timescale of the prediction (265; 266). Thus, most of the literature is focused on the potential to make predictions of these quantities, rather than on skill as measured by historical observations (255, 260; 261, 257), with exceptions (258; 259; 217). On the global scale, verification is limited to variables measured or derived from satellite observations, such as ocean chlorophyll (256), marine primary productivity (19), or interpolated estimates of the surface ocean partial pressure of CO₂ (262). Nevertheless, there is promising potential to make ocean biogeochemical initialized predictions across multiple timescales.

For S2S, S2I, and S2D initialized predictions to be useful, they must be shown to be not only skillful but reliable (267), and this is a considerable challenge that the community is only starting to attempt to address (5; 21). The ultimate challenge in this emerging area of research, and one that is igniting excitement and interest in the scientific community, is to provide predictions with maximum skill that take into account all relevant processes across subseasonal to decadal timescales (268, 269). Toward that end, initialized prediction is already put to task and being applied in various sectors even as improvements in understanding and prediction capability are being improved, thus driving rapid advances in this burgeoning field.

References

1. Meehl, G. A., et al. Decadal Prediction. *Bull. Am. Meteorol. Soc.*, 90, 1467–1486, doi:10.1175/2009BAMS2778.1. <https://doi.org/10.1175/2009BAMS2778.1> (2009).
2. Meehl, G.A., Hu, A., Arblaster, J.M., Fasullo, J., & Trenberth, K.E. Externally forced and internally generated decadal climate variability associated with the Interdecadal Pacific Oscillation, *J. Climate*, 26, 7298-7310, doi: <http://dx.doi.org/10.1175/JCLI-D-12-00548.1> (2013).
3. Hawkins, E., & Sutton, R. (2009). The Potential to Narrow Uncertainty in Regional Climate Predictions, *Bulletin of the American Meteorological Society*, 90(8), 1095-1108.
4. Lehner, F., Deser, C., Maher, N., Marotzke, J., Fischer, E. M., Brunner, L., Knutti, R., and Hawkins, E.: Partitioning climate projection uncertainty with multiple large ensembles and CMIP5/6, *Earth Syst. Dynam.*, 11, 491–508, <https://doi.org/10.5194/esd-11-491-2020>, 2020.
5. Vitart, F., and A.W. Robertson, The sub-seasonal to seasonal prediction project (S2S) and the prediction of extreme events. *Npj Clim. Atmos. Sci.*, 1, 3, doi:10.1038/s41612-018-0013-0 (2018).
6. Pegion, K. et al., The Subseasonal Experiment (SubX), *Bull. Amer. Meteorol. Soc.*, 100, 2043—2060, doi:10.1175/BAMS-D-18-0270.1 (2019).
7. Kirtman, B.P., Min, D., & J.M., et al. The North American Multimodel Ensemble: Phase-1 seasonal to interannual prediction; Phase-2 toward developing intraseasonal prediction. *Bull. Amer. Meteorol. Soc.*, <https://doi.org/10.1175/BAMS-D-12-00050.1> (2014).

8. Boer, G. J. et al. The Decadal Climate Prediction Project (DCPP) contribution to CMIP6. *Geosci. Model Dev.* **9**, 3751–3777 (2016).
9. Palmer, T.N., F.J. Doblas-Reyes, A. Weisheimer, and J.J. Rodwell, Toward seamless prediction: Calibration of climate change projections using seasonal forecasts. *Bull. Amer. Meteorol. Soc.*, **89**, 459—470 (2008).
10. Branstator, G., & Teng, H. Potential impact of initialization on decadal predictions as assessed for CMIP5 models. *Geophy Res Lett*, **39**, doi: 10.1029/2012GL051974 (2012).
11. Barnett, T., Graham, N., Cane, M., Zebiak, S., Dolan, S., O'Brien, J. and Legler, D., On the prediction of the *El Niño* of 1986-1987. *Science*, **241**, 192-196 (1988).
12. Kushnir, Y., et al. Towards operational predictions of the near-term climate. *Nat. Clim. Chang.*, **9**, doi:10.1038/s41558-018-0359-7 (2019).
13. Lean, P., Hólm, E.V., Bonavita, M., Bormann, N., McNally, A.P. and Järvinen, H. Continuous data assimilation for global numerical weather prediction. *Quarterly Journal of the Royal Meteorological Society*. <https://doi.org/10.1002/qj.3917> (2020).
14. Sandery, PA, O’Kane TJ, Kitsios V, and Sakov P. Climate model state estimation using variants of EnKF coupled data assimilation, *Mon. Wea. Rev.* **148**, 2411-2431, <https://doi.org/10.1175/MWR-D-18-0443.1> (2020).
15. Johnson, C., Hoskins, B.J. & Nichols, N.K. A singular vector perspective of 4D-Var: Filtering and interpolation, *Q. J. R. Meteorol. Soc.* **131**, 1–19, doi: 10.1256/qj.03.231 .(2005).
16. Magnusson, L., Nycander, J. & Kallen, E. Flow dependent versus flow independent initial perturbations for ensemble prediction. *Tellus A*, **61**, 194-209. doi:10.1111/j.1600-0870.2008.00385.x, (2009).

17. Infant, J. M., and B. P. Kirtman Prediction and predictability of land and atmosphere initialized CCSM4 climate forecasts over North America, *J. Geophys. Res. Atmos.*, **121**, 12,690–12,701, doi:[10.1002/2016JD024932](https://doi.org/10.1002/2016JD024932) (2016b).

18. Trenary, L., T. DelSole, M. K. Tippett, and K. Pegion A new method for determining the optimal lagged ensemble, *J. Adv. Model. Earth Syst.*, **9**, 291–306, doi:[10.1002/2016MS000838](https://doi.org/10.1002/2016MS000838) (2017).

19. Kirtman, B. P., & Min, D. Multi-model ensemble ENSO prediction with CCSM and CFS. *Mon. Wea. Rev.*, DOI: [10.1175/2009MWR2672.1](https://doi.org/10.1175/2009MWR2672.1) (2009).

20. Yeager, S. G., and Coauthors, Predicting Near-Term Changes in the Earth System: A Large Ensemble of Initialized Decadal Prediction Simulations Using the Community Earth System Model. *Bull. Am. Meteorol. Soc.*, **99**, 1867–1886, doi:[10.1175/BAMS-D-17-0098.1](https://doi.org/10.1175/BAMS-D-17-0098.1) (2018).

21. Smith, D. M. et al. Real-time multi-model decadal climate predictions. *Clim. Dyn.* **41**, 2875–2888 (2013a).

22. MacLachlan C., A. Arribas, K.A. Peterson, A. Maidens, D. Fereday, A.A. Scaife, M. Gordon, M. Vellinga, A. Williams, R. E. Comer, J. Camp and P. Xavier. Global Seasonal forecast system version 5 (GloSea5): a high resolution seasonal forecast system. *Q. J. R. Met. Soc.*, DOI: [10.1002/qj.2396](https://doi.org/10.1002/qj.2396), (2014)

23. Muñoz-Sabater et al. Assimilation of SMOS brightness temperatures in the ECMWF Integrated Forecasting System. *Q. J. R. Met. Soc.*, DOI: [10.1002/qj.3577](https://doi.org/10.1002/qj.3577) (2019)

24. Drewitt, G., A.A. Berg, W.J. Merryfield & W.-S. Lee Effect of Realistic Soil Moisture Initialization on the Canadian CanCM3 Seasonal Forecast Model, *Atmosphere-Ocean*, 50:4, 466-474, DOI: 10.1080/07055900.2012.722910 (2012)

25. Polkova, I., Köhl, A. & Stammer, Climate-mode initialization for decadal climate predictions, *Clim Dyn*, 53: 7097-7111, <https://doi.org/10.1007/s00382-019-04975-y> (2019)

26. Smith, D. M., R. Eade and H. Pohlmann, A comparison of full-field and anomaly initialization for seasonal to decadal climate prediction, *Climate Dynamics*, 41, 3325-3338, DOI 10.1007/s00382-013-1683-2 (2013b)

27. Volpi, D.; Guemas, V.; Doblas-Reyes, F. J. Comparison of full field and anomaly initialisation for decadal climate prediction: towards an optimal consistency between the ocean and sea-ice anomaly initialisation state. *Climate Dynamics*, 49, 1181-1195 (2017)

28. Penny, S.G., S. Akella, O. Alves, C. Bishop, M. Buehner, M. Chevallier, F. Counillon, C. Draper, S. Frolov, Y. Fujii, A. Karspeck, A. Kumar, P. Laloyaux, J.-F. Mahfouf, M. Martin, M. Pena, P. De Rosnay, A. Subramanian, R. Tardif, Y. Wang, and X. Wu. Coupled Data Assimilation for Integrated Earth System Analysis and Prediction: Goals, Challenges and Recommendations. Technical report, World Meteorological Organisation, (2017)

29. Williams, K. D., et al. The Met Office Global Coupled model 2.0 (GC2) configuration, *Geosci. Model Dev.*, 8, 1509–1524, <https://doi.org/10.5194/gmd-8-1509-2015>, (2015)

30. Becker, E., & Van Den Dool, H. Probabilistic seasonal forecasts in the North American Multimodel Ensemble: A baseline skill assessment. *J. Climate*, 29, 3015–3026. <https://doi.org/10.1175/JCLI-D-14-00862.1> (2016)

31. Kadow, C., et al. Decadal climate predictions improved by ocean ensemble dispersion filtering. *Journal of Advances in Modeling Earth Systems*, 9.2, 1138-1149 (2017)

32. Dobrynin, M., Domeisen, D. I., Müller, W. A., Bell, L., Brune, S., Bunzel, F., et al. Improved teleconnection based dynamical seasonal predictions of boreal winter. *Geophysical Research Letters*, 45, 3605– 3614. <https://doi.org/10.1002/2018GL077209> (2018)

33. Smith, D. M., and co-authors, North Atlantic climate far more predictable than models imply, *Nature*, 583, 796-800, <https://doi.org/10.1038/s41586-020-2525-0> (2020)

34. Richter, J. H. and co-authors. Subseasonal prediction with and without a well-represented stratosphere in CESM1, *Weather and Forecasting*, <https://journals.ametsoc.org/view/journals/wefo/aop/WAF-D-20-0029.1/WAF-D-20-0029.1.xml> (2020)

35. Scaife, A. A., Arribas, A., Blockley, E., Brookshaw, A., Clark, R., Dunstone, R. N., et al. Skillful long-range prediction of European and North American winters. *Geophysical Research Letters*, 41, 2514–2519. <https://doi.org/10.1002/2014GL059637> (2014a)

36. Smith, D.M., and co-authors, Robust skill of decadal climate predictions. *Npj Clim. Atmos. Sci.* 2, 13, <https://doi.org/10.1038/s41612-019-0071-y> (2019)

37. Athanasiadis, P.J., Yeager, S., Kwon, Y.-O., Bellucci, A., Smith, D.W., & Tibaldi, S. Decadal predictability of North Atlantic blocking and the NAO. *NPJ Climate and Atmos. Sci.* **3**, 20 <https://doi.org/10.1038/s41612-020-0120-6> (2020)

38. Nie Y., et al. Stratospheric initial conditions provide seasonal predictability of the North Atlantic and Arctic Oscillations. *Env. Res. Lett.*, **14**, 3, (2019)

39. Capotondi, A., et al. Understanding ENSO diversity. *Bull Amer. Meteor. Soc.*, **96**, 921-938, doi:10.1175/BAMS-D-13-00117.1 (2015)

40. Cobb, K.M., Westphal, N., Sayani, H.R., Watson, J.T., Di Lorenzo, E., Cheng, H., Edwards, R.L. & Charles, C.D.. Highly variable El Niño–Southern Oscillation throughout the Holocene. *Science*, **339**(6115), pp.67-70, (2013)

41. Capotondi, A., & Sardeshmukh, P.D. Is El Niño really changing? *Geophys. Res. Lett.*, **44**, doi:10.1002/2017GL074515. (2017)

42. Grothe, P. R., Cobb, K. M., Liguori, G., Di Lorenzo, E., Capotondi, A., Lu, Y., ... & Deocampo, D. M. Enhanced El Niño–Southern Oscillation variability in recent decades. *Geophysical Research Letters* (2019)

43. Deser, C., Phillips, A.S., & Alexander, M.A., 2010. Twentieth century tropical sea surface temperature trends revisited. *Geophys. Res. Lett.*, **37**(10) (2010)

44. Meehl, G.A., Arblaster, J.M., Fasullo, J., Hu, A., & Trenberth, K.E. Model-based evidence of deep ocean heat uptake during surface temperature hiatus periods. *Nature Climate Change*, **1**, 360—364, doi:10.1038/NCLIMATE1229 (2011)

45. Mann, M.E., Emanuel, K.A. Atlantic Hurricane Trends linked to Climate Change, *Eos*, **87**, 24, p 233, 238, 241 (2006)

46. Mann, M.E., Steinman, B.A., Miller, S.K., On Forced Temperature Changes, Internal Variability and the AMO, *Geophys. Res. Lett.* (“Frontier” article), 41, 3211-3219, doi:10.1002/2014GL059233, (2014)

47. Mann, M. E., Steinman, B. A., Miller, S. K., Frankcombe, L. M., England, M. H., & Cheung, A. H. Predictability of the recent slowdown and subsequent recovery of large-scale surface warming using statistical methods. *Geophysical Research Letters* 43, 3459–3467, doi:10.1002/2016GL068159 (2016)

48. Steinman, B.A. Frankcombe, L.M., Mann, M.E., Miller, S.K., England, M.H. Response to Comment on “Atlantic and Pacific multidecadal oscillations and Northern Hemisphere temperatures”, *Science*, 350, 1326, (2015)

49. Roemmich, D. & Gilson, J. The 2004-2008 mean and annual cycle of temperature, salinity, and steric height in the global ocean from the Argo Program. *Progress in Oceanography*, 82, 81-100, (2009)

50. Roemmich, D. et al. On the Future of Argo: A Global, Full-Depth, Multi-Disciplinary Array. *Front. Mar. Sci.* <https://doi.org/10.3389/fmars.2019.00439>, (2019)

51. Thompson, D. M., J. E. Cole, G. T. Shen, A. W. Tudhope, and G. A. Meehl. Early twentieth-century warming linked to tropical Pacific wind strength. *Nature Geoscience* 8(2), (2015)

52. Cook, E. R. et al. Megadroughts in North America: placing IPCC projections of hydroclimatic change in a long-term palaeoclimate context. *J Quaternary Sci* 25, 48-61 (2010)

53. Emile-Geay, J., Cobb, K. M., Mann, M. E., & Wittenberg, A. T. Estimating Central Equatorial Pacific SST Variability over the Past Millennium. Part II: Reconstructions and Implications. *Journal of Climate*, 26(7), 2329–2352
<http://doi.org/10.1175/JCLI-D-11-00511.1> (2013)

54. Linsley, B. K., Wu, H. C., Dassié, E. P., & Schrag, D. P. Decadal changes in South Pacific sea surface temperatures and the relationship to the Pacific decadal oscillation and upper ocean heat content. *Geophysical Research Letters*, 42(7), 2358-2366 (2015)

55. Buckley, B. M., Ummenhofer, C. C., D'Arrigo, R. D., Hansen, K. G., Truong, L. H., Le, C. N., & Stahle, D. K. Interdecadal Pacific Oscillation reconstructed from trans-Pacific tree rings: 1350–2004 CE. *Climate Dynamics*, 1-16 (2019)

56. Abram, N. J., Hargreaves, J. A., Wright, N. M., Thirumalai, K., Ummenhofer, C. C., & England, M. H. Palaeoclimate perspectives on the Indian Ocean Dipole. *Quaternary Science Reviews*, 237, doi: 10.1016/j.quascirev.2020.106302 (2020a)

57. Sanchez, S. C., Charles, C. D., Carriquiry, J. D., and Villaescusa, J. A. Two centuries of coherent decadal climate variability across the Pacific North American region, *Geophys. Res. Lett.*, 43, 9208– 9216, doi:10.1002/2016GL069037 (2016)

58. Abram, N.J., Wright, N.M., Ellis, B., Dixon, B.C., Wurtzel, J.B., England, M.H., Ummenhofer, C.C., Philibosian, B., Cahyarini, S.Y., Yu, T.L. & Shen, C.C.. Coupling of Indo-Pacific climate variability over the last millennium. *Nature*, 579(7799), pp.385-392 (2020b)

59. Konecky, B., Dee, S. G., & Noone, D. WaxPSM: A forward model of leaf wax hydrogen isotope ratios to bridge proxy and model estimates of past climate. *Journal of Geophysical Research: Biogeosciences* (2019)

60. Neukom, R., Barboza, L.A., Erb, M.P., Shi, F., Emile-Geay, J., Evans, M.N., Franke, J., Kaufman, D.S., Lücke, L., Rehfeld, K. & Schurer, A.. Consistent multi-decadal variability in global temperature reconstructions and simulations over the common era. *Nature Geoscience*, 12(8), p.643 (2019)

61. McGregor, H.V., Evans, M.N., Goosse, H., Leduc, G., Martrat, B., Addison, J.A., Mortyn, P.G., Oppo, D.W., Seidenkrantz, M.S., Sicre, M.A. & Phipps, S.J.. Robust global ocean cooling trend for the pre-industrial Common Era. *Nature Geoscience*, 8(9), pp.671-677, (2015)

62. Tierney, J.E., Abram, N.J., Anchukaitis, K.J., Evans, M.N., Giry, C., Kilbourne, K.H., Saenger, C.P., Wu, H.C. & Zinke, J.. Tropical sea surface temperatures for the past four centuries reconstructed from coral archives. *Paleoceanography*, 30(3), pp.226-252, (2015)

63. Goosse, H., Crespin, E., de Montety, A., Mann, M. E., Ressen, H., & Timmermann, A. Reconstructing surface temperature changes over the past 600 years using climate model simulations with data assimilation. *Journal of Geophysical Research*, 115, D09108 (2010)

64. Hakim, G. J., Emile Geay, J., Steig, E. J., Noone, D., Anderson, D. M., Tardif, R., Steiger, N., and Perkins, W. A. The last millennium climate reanalysis project: Framework and first results, *J. Geophys. Res. Atmos.*, 121, 6745– 6764, doi:[10.1002/2016JD024751](https://doi.org/10.1002/2016JD024751) (2016)

65. Steiger, Nathan J., Jason E. Smerdon, Edward R. Cook, and Benjamin I. Cook. A reconstruction of global hydroclimate and dynamical variables over the Common Era. *Scientific Data* 5 180086 doi: 10.1086/sdata.2018.86 (2018)

66. Evans, M. N., Tolwinski-Ward, S. E., Thompson, D. M., & Anchukaitis, K. J. Applications of proxy system modeling in high resolution paleoclimatology. *Quaternary science reviews*, 76, 16-28 (2013)

67. Dee, S., Emile Geay, J., Evans, M. N., Allam, A., Steig, E. J., & Thompson, D. M. PRYSM: An open source framework for PRoXy System Modeling, with applications to oxygen isotope systems. *Journal of Advances in Modeling Earth Systems*, 7(3), 1220-1247 (2015)

68. Becker, E., Dool, den, H. V., & Zhang, Q. Predictability and Forecast Skill in NMME. *J. Climate*, 27, 5891–5906. <http://doi.org/10.1175/JCLI-D-13-00597.1> (2014)

69. Doblas-Reyes, F. J. et al. Initialized near-term regional climate change prediction. *Nat. Commun.* 4, 1715 (2013)

70. Meehl, G.A., Hu, A., & Teng, H. Initialized decadal prediction for transition to positive phase of the Interdecadal Pacific Oscillation. *Nature Communications.*, 7, doi:10.1038/NCOMMS11718 (2016b)

71. Kharin, V. V., Boer, G. J., Merryfield, W. J., Scinocca, J. F. & Lee, W.-S. Statistical adjustment of decadal predictions in a changing climate. *Geophys. Res. Lett.* 39, L19705 <https://doi.org/10.1029/2012GL052647> (2012)

72. Scaife, A. A., & Smith, D. A signal-to-noise paradox in climate science. *npj Climate and Atmospheric Science*, 1, 28. (2018)

73. Sévellec, F., & Drijfhout, S. S. The signal-to-noise paradox for interannual surface atmospheric temperature predictions. *Geophysical Research Letters*, 46, 9031–9041. DOI: 10.1029/2019GL083855. (2019)

74. Zhang W. and B. Kirtman, Estimates of decadal climate predictability from an interactive ensemble model. *Geophy Res Letts*, 46, 3387-3397. (2019)

75. Weisheimer, A., Decremer,D., MacLeod, D., O'Reilly, C., Stokdale, T.N., Johnson, S., & Palmer, T.N. How confident are predictability estimates of the winter North Atlantic Oscillation? *Quarterly Journal of the Royal Meteorological Society*, 1–20. <https://doi.org/10.1002/qj.3446> (2019)

76. Barnston, A. G., Tippett, M. K., L'Heureux, M. L., Li, S., & DeWitt, D. G. Skill of Real-Time Seasonal ENSO Model Predictions during 2002–11: Is Our Capability Increasing? *Bulletin of the American Meteorological Society*, 93(5), 631–651. <https://doi.org/10.1175/BAMS-D-11-00111.1>, (2012)

77. Robertson, A.W. and F. Vitart (eds.), Sub-seasonal to seasonal prediction. Elsevier, ISBN:9780128117156, 585pp. (2018)

78. Kim, H., F. Vitart, and D.E. Waliser Prediction of the Madden–Julian Oscillation: A Review. *J. Climate*, 31, 9425–9443, <https://doi.org/10.1175/JCLI-D-18-0210.1> (2018)

79. Stan, C., Straus, D. M., Frederiksen, J. S., Lin, H., Maloney, E. D., & Schumacher, C. Review of Tropical-Extratropical Teleconnections on Intraseasonal Time Scales. *Reviews of Geophysics*, 55(4), 902–937. <http://doi.org/10.1002/2016RG000538> (2017)

80. Kim, H., Richter, J. H., & Martin, Z. Insignificant QBO–MJO prediction skill relationship in the SubX and S2S subseasonal reforecasts. *Journal of Geophysical Research: Atmospheres*, 124. <https://doi.org/10.1029/2019JD031416> (2019a)

81. Lim, E.-P., Hendon, H. H., & Thompson, D. W. J. Seasonal evolution of stratosphere-troposphere coupling in the Southern Hemisphere and implications for the predictability of surface climate. *J. Geophys. Res. Atmos.*, 123, 1–15, doi:10.1029/2018JD029321 (2018)

82. Zheng, C., E.K.M. Chang, H. Kim, M. Zhang, and W. Wang, Subseasonal to seasonal prediction of wintertime Northern Hemisphere extratropical cyclone activity by S2S and NMME models. *J. Geophys. Res. Atmos.*, Doi: 10.1029/2019JD031252 (2019)

83. DeFlorio, M.J., Waliser, D.E., Guan, B. *et al.* Global evaluation of atmospheric river subseasonal prediction skill. *Clim Dyn* 52, 3039–3060. <https://doi.org/10.1007/s00382-018-4309-x>. (2019)

84. Baggett, C. *et al.*, Skillful subseasonal forecasts of weekly tornado and hail activity using the Madden-Julian Oscillation. *J. Geophys Res Atmos*, 123, 12,661–12,675. (2018)

85. Broennimann, S. Impact of El Niño–Southern Oscillation on European climate, *Rev. Geophys.*, 45, RG3003, doi:10.1029/2006RG000199 (2007)

86. Ambaum, P., & Hoskins, B.J. The NAO Troposphere–Stratosphere Connection, *J. Climate*, 15, 1969–1978 (2002)

87. Kushnir, Y., Robinson, W. A., Chang, P., & Robertson, A. W. The physical basis for predicting Atlantic sector seasonal-to-interannual climate variability. *Journal of Climate*, **19**, 5949-5970 (2006)

88. Riddle, E. E., A. H. Butler, J. C. Furtado, J. L. Cohen, and A. Kumar CFSv2 ensemble prediction of the wintertime Arctic Oscillation. *Climate Dyn.*, **41**, 1099–1116, doi:10.1007/s00382-013-1850-5 (2013)

89. Hendon, H. H., Thompson, D. W. J. & Wheeler, M. C. Australian rainfall and surface temperature variations associated with the Southern Hemisphere annular mode. *J. Climate*, **20**, 2452–2467 (2007)

90. Marshall, A.G., Hudson, D., Wheeler, M., Alves,O., Hendon, H.H., Pook, M.J. & Risbey, J.S. Intra-seasonal drivers of extreme heat over Australia in observations and POAMA-2. *Climate Dynamics*. **43**, 1915-1937, (2014)

91. Seviour, W. J. M., S. C. Hardiman, L. J. Gray, N. Butchart, C. MacLachlan, and A. A. Scaife, 2014: Skillful Seasonal Prediction of the Southern Annular Mode and Antarctic Ozone. *J. Climate*, **27**, 7462–7474, DOI: 10.1175/JCLI-D-14-00264.1. (2014)

92. Lim, E.-P., Hendon, H.H., & Rashid, H.A. Seasonal predictability of the Southern Annular Mode due to its association with ENSO. *Journal of Climate*, **26**, 8037-8054 (2013)

93. Lim, E., Hendon, H.H., Boschat, G., Hudson, D., Thompson, D.W.J, Dowdy, A., & Arblaster, J.M. Australian hot and dry extremes induced by weakenings of the stratospheric polar vortex. *Nature Geoscience* **12**, 896–901 (2019)

94. Mariotti, A., et al. Windows of opportunity for skillful forecasts subseasonal to seasonal and beyond. *Bull. Amer. Meteorol. Soc.*, <https://doi.org/10.1175/BAMS-D-18-0326.1> (2020)

95. Dirmeyer, P. A., Halder, S., & Bombardi, R. On the Harvest of Predictability From Land States in a Global Forecast Model. *Journal of Geophysical Research: Atmospheres*, 123(23), 145–17. <http://doi.org/10.1029/2018JD029103> (2018)

96. Koster, R. D. et al., Regions of strong coupling between soil moisture and precipitation. *Science*, 305, 1138-1140 (2004)

97. Koster, R. D., and Coauthors, The Second Phase of the Global Land–Atmosphere Coupling Experiment: Soil Moisture Contributions to Subseasonal Forecast Skill. *J. Hydrometeorol.*, 12, 805–822, <https://doi.org/10.1175/2011JHM1365.1> (2011)

98. Seo E., M.-I. Lee, J.-H. Jeong, R.D. Koster, S.D. Schubert, H.-M. Kim, D. Kim, H.-S. Kang, H.-K. Kim, C. MacLachlan and A.A. Scaife. Impact of soil moisture initialization on boreal summer subseasonal forecasts: mid-latitude surface air temperature and heat wave events. *Clim. Dyn.*, doi:10.1007/s00382-018-4221-4, (2018)

99. Zampieri, L., Goessling, H. F., & Jung, T. Bright prospects for Arctic sea ice prediction on subseasonal time scales. *Geophysical Research Letters*, 45, 9731–9738. <https://doi.org/10.1029/2018GL079394> (2018)

100. Zampieri, L., Goessling, H. F., & Jung, T. Predictability of Antarctic sea ice edge on subseasonal time scales, *Geophysical Research Letters*, 46, 9719–9727. (2019)

101. Bushuk, M., et al. A mechanism for the Arctic sea ice spring predictability barrier. *Geophysical Research Letters*, <https://doi.org/10.1029/2020GL088335>, (2020)

102. Kimmritz, M., et al. Impact of Ocean and Sea Ice Initialisation On Seasonal Prediction Skill in the Arctic, *JAMES*, <https://doi.org/10.1029/2019MS001825> (2019)

103. Ono, J., Yoshiki Komuro, Y., & Tatebe, H. Impact of sea-ice thickness initialized in April on Arctic sea-ice extent predictability with the MIROC climate model. *Annals of Glaciology*: 1-9, (2020)

104. Liu, J., Chen, Z., Hu, Y., Zhang, Y., Ding, Y., & co authors. Towards reliable Arctic sea ice prediction using multivariate data assimilation. *Science Bulletin*, 64(1), 63–72. doi: 10.1016/j.scib.2018.11.018 (2019)

105. Jung, T., Gordon, N. D., Bauer, P., Bromwich, D. H., Chevallier, M., Day, J. J., Dawson, J., Doblas Reyes, F., Fairall, C., Goessling, H. F., Holland, M., Inoue, J., Iversen, T., Klebe, S., Lemke, P., Losch, M., Makshtas, A., Mills, B., Nurmi, P., Perovich, D., Reid, P., Renfrew, I. A., Smith, G., Svensson, G., Tolstykh, M., & Yang, Q. Advancing polar prediction capabilities on daily to seasonal time scales. *Bulletin of the American Meteorological Society*, 97(9), 1631– 1647. <https://journals.ametsoc.org/doi/full/10.1175/BAMS-D-14-00246.1> (2016)

106. Jung, T., Kasper, M. A., Semmler, T., and Serrar, S. Arctic influence on subseasonal midlatitude prediction, *Geophys. Res. Lett.*, 41, 3676– 3680, doi:10.1002/2014GL059961 (2014)

107. Baldwin, M. P., Stephenson, D. B., Thompson, D. W. J., Dunkerton, T. J., Charlton, A. J., & O'Neill, A. Stratospheric memory and skill of extended- range weather forecasts. *Science*, 301, 636–640. (2003)

108. Butler, A.H., Polvani, L.M., & Deser, C. Separating the stratospheric and tropospheric pathways of El Niño-Southern Oscillation teleconnections, *Environmental Res. Lett.*, , 024014, doi: 10.1088/1748-9326/9/2/024014 (2014)

109. Sigmond, M., Scinocca, J., Kharin, V. et al. Enhanced seasonal forecast skill following stratospheric sudden warmings. *Nature Geosci* 6, 98–102 doi:10.1038/ngeo1698 (2013)

110. Scaife A.A., A.-Yu. Karpechko, M.P. Baldwin, A. Brookshaw, A.H. Butler, R. Eade, M. Gordon, C. MacLachlan, N. Martin, N. Dunstone and D. Smith. Seasonal winter forecasts and the stratosphere. *Atm. Sci. Lett.*, DOI: 10.1002/asl.598, (2016)

111. Anstey, J. A. & T. G. Shepherd, Review Article High-latitude influence of the quasi-biennial oscillation. *Quart. J. Roy. Meteor. Soc.*, 140, 1–21 (2014)

112. Garfinkel, C. I., &and Hartmann, D. L. Influence of the quasi- biennial oscillation on the North Pacific and El Niño teleconnections, *J. Geophys. Res.*, 115, D20116, doi:10.1029/2010JD014181 (2010)

113. Wang, J., Kim, H. M., & Chang, E. K. M. Interannual modulation of Northern Hemisphere winter storm tracks by the QBO. *Geophysical Research Letters*, 45, 2786– 2794. <https://doi.org/10.1002/2017GL076929> (2018)

114. Yoo, C., and S.-W. Son, Modulation of the boreal wintertime Madden-Julian oscillation by the stratospheric quasi-biennial oscillation, *Geophys. Res. Lett.*, 43, 1392–1398, doi:10.1002/2016GL067762 (2016)

115. Son, S.-W., Y. Lim, C. Yoo, H. H. Hendon, and J. Kim, Stratospheric control of Madden–Julian oscillation. *J. Climate*, 30, 1909–1922, doi:10.1175/JCLI-D-16-0620.1 (2017)

116. Lim, Y., Son, SW., Marshall, A.G. et al. Influence of the QBO on MJO prediction skill in the subseasonal-to-seasonal prediction models. *Climate Dynamics* <https://doi.org/10.1007/s00382-019-04719-y>, (2019)

117. Tompkins, A. M., and Coauthors, The climate-system historical forecast project: Providing open access to seasonal forecast ensembles from centers around the globe. *Bulletin of the American Meteorological Society*, 98 (11), 2293–2301 (2017)

118. Acosta Navarro, J. C., Ortega, P., Batté, L., Smith, D., Bretonnière, P. A., Guemas, V., ... & Doblas Reyes, F. J. Link between autumnal Arctic Sea ice and Northern Hemisphere winter forecast skill. *Geophysical Research Letters*, 47(5), (2020)

119. Scaife A.A. et al. Skill of tropical rainfall predictions in multiple seasonal forecast systems, *Int. J. Climatol.*, 1-15, doi:10.1002/joc.5855, (2018)

120. Hu, Z., Kumar, A., Jha, B. et al. How much of monthly mean precipitation variability over global land is associated with SST anomalies?. *Clim Dyn* **54**, 701–712 (2020)

121. Kirtman, B. P., D. Anderson, G. Brunet, I.-S. Kang, A. A. Scaife and D. Smith, Prediction from weeks to decades, Climate Science for Serving Society: Research, Modelling and Prediction Priorities. G. R. Asrar and J. W. Hurrell, Eds. Springer, DOI 10.1007/978-94-007-6692-1_8 (2013)

122. Capotondi, A., Wittenberg, A.T., Kug, J.-S., Takahashi, K. & McPhaden, M.J., ENSO Diversity. *AGU Monograph* “El Niño Southern Oscillation in a changing

climate”, M. McPhaden, A. Santoso, and W. Cai, Editors, DOI: 10.1002/9781119548164.ch4. (2020)

123. Vimont, D. J., M. A. Alexander, and M. Newman, Optimal growth of central and east Pacific ENSO events. *Geophysical Research Letters*, 41 (11), 4027–4034, doi:10.1002/2014GL059997 (2014)

124. Zhang, H., A. Clement, & DiNezio. The South Pacific Meridional Mode: A mechanism for ENSO-like variability. *J. Climate*, 27, 769-783 (2014)

125. Larson, S., and B. P. Kirtman, The Pacific meridional mode as a trigger for ENSO in a high-resolution coupled model. *Geophys. Res. Lett.*, DOI: 10.1002/grl.50571 (2013)

126. Capotondi, A., & Sardeshmukh, P.D. Optimal precursors of different types of ENSO events. *Geophys. Res. Lett.*, 42, 9952-9960, doi:10.1002/2015GL066171 (2015)

127. Amaya, D. The Pacific meridional mode and ENSO: a review. *Current Clim. Change. Reps.*, <https://doi.org/10.1007/s40641-019-00142-x> (2019)

128. Larson, S. M., and B. P. Kirtman, Assessing Pacific Meridional Mode forecasts and its role as an ENSO precursor and predictor in the North American Multi-Model Ensemble. *J. Climate* 27, 7018-7032 (2014)

129. Ren H., F-F. Jin, B. Tian, A.A. Scaife. Distinct persistence barriers in two types of ENSO. *Geophys. Res. Lett.*, 43, 10,973-10,979, doi:10.1002/2016GL071015, (2016)

130. Infant, J. M., and B. P. Kirtman, North American rainfall and temperature prediction response to the diversity of ENSO, *Climate Dynamics* doi: 10.1007/s00382-015-2749-0 (2016a)

131. DiNezio, P., Deser, C., Karspeck, A., Yeager, S., Okumura, Y., Danabasoglu, G., Rosenbloom, N., Caron, J., & Meehl, G.A. A two-year forecast for a 60-80% chance of La Niña in 2017-2018. *Geophys. Res. Lett.*, DOI: 10.1002/2017GL074904 (2017)

132. Freund, M.B., Henley, B.J., Karoly, D.J., McGregor, H.V., Abram, N.J. & Dommenget, D.. Higher frequency of Central Pacific El Niño events in recent decades relative to past centuries. *Nature Geoscience*, 12(6), 450-455, (2019)

133. McPhaden MJ Genesis and evolution of the 1997–98 El Niño. *Science* 283:950–954 (1999)

134. Capotondi, A., Sardeshmukh, P. D., & Ricciardulli, L.. The nature of the stochastic wind forcing of ENSO. *Journal of Climate*, 31(19), 8081-8099 (2018)

135. Tan, X., Tang, Y., Lian, T. *et al.* A study of the effects of westerly wind bursts on ENSO based on CESM. *Clim Dyn* 54, 885–899 <https://doi.org/10.1007/s00382-019-05034-2> (2020)

136. Lopez, H., and B. P. Kirtman WWBs, ENSO predictability, the spring barrier and extreme events, *J. Geophys. Res. Atmos.*, 119, 10,114–10,138, doi:[10.1002/2014JD021908](https://doi.org/10.1002/2014JD021908) (2014)

137. Ren, H.L., A.A. Scaife, N. Dunstone, B. Tian, Y. Liu, S. Ineson, et al. Seasonal predictability of winter ENSO types in operational dynamical model predictions. *Climate Dynamics*, 52, 3869-3890, (2019)

138. Chang, P., et al., Climate fluctuations of tropical coupled systems: the role of ocean dynamics. *J. Clim.* 19, 5122–5174. (2006)

139. Lübbecke, J.F., & McPhaden, M.J. Symmetry of the Atlantic Niño mode. *Geophysical Research Letters* 44.2, 965-973 (2017)

140. Richter, I., et al. On the link between mean state biases and prediction skill in the tropics: an atmospheric perspective. *Climate Dynamics* 50.9-10, 3355-3374 (2018)

141. Stockdale, T. N., Balmaseda, M. A., & Vidard, A. Tropical atlantic sst prediction with coupled ocean– atmosphere GCMs. *Journal of Climate* 19 (23), 6047–6061 (2006)

142. Ding, H., et al. The impact of mean state errors on equatorial Atlantic interannual variability in a climate model. *Journal of Geophysical Research: Oceans* 120.2 1133-1151 (2015)

143. Saji NH, Goswami BN, Vinayachandran PN, Yamagata T A dipole mode in the tropical Indian Ocean. *Nature* 401:360–363 (1999)

144. Krishnamurthy V, Kirtman BP Variability of the Indian Ocean: relation to monsoon and ENSO. *Q J R Meteorol Soc* 129:1623–1646 (2003)

145. Wu R, Kirtman BP, Krishnamurthy V An asymmetric mode of tropical Indian Ocean rainfall variability in boreal spring. *J Geophys Res-Atmos* 113 (2008)

146. Lu B., H.-L. Ren, A.A. Scaife, J. Wu, N. Dunstone, D. Smith, J. Wan, R. Eade, C. MacLachlan and M. Gordon. An extreme negative Indian Ocean dipole event in 2016: dynamics and predictability. *Clim. Dyn.*, doi:10.1007/s00382-017-3908-2, (2017)

147. Shinoda T. and W. Han, Influence of Indian Ocean dipole on atmospheric subseasonal variability. *J. Climate*, 18, 3891-3909 (2005)

148. Dunstone, N., Smith, D., Scaife, A. et al. Skilful predictions of the winter North Atlantic Oscillation one year ahead. *Nature Geosci* 9, 809–814 doi:10.1038/ngeo2824 (2016)

149. Dunstone, N., Smith, D., Scaife, A., Hermanson, L., Fereday, D., O'Reilly, C., ... & Woollings, T. Skilful seasonal predictions of summer European rainfall. *Geophysical Research Letters*, 45(7), 3246-3254, (2018)

150. Paolino, D. A., J. L. Kinter, B. P. Kirtman, D. Min, and D. M. Straus The impact of land surface and atmospheric initialization on seasonal forecasts with CCSM, *J. Clim.*, 25(3), 1007– 1021, doi:10.1175/2011JCLI3934.1 (2011)

151. Dirmeyer, P. A. The role of the land surface background state in climate predictability, *J. Hydrometeorol.*, 4(3), 599– 610, doi:[10.1175/1525-7541\(2003\)004<0599:TROTLS>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)004<0599:TROTLS>2.0.CO;2) (2003)

152. Prodhomme, C., Doblas-Reyes, F., Bellprat, O., & Dutra, E. Impact of land-surface initialization on sub-seasonal to seasonal forecasts over Europe. *Climate Dynamics*, 47(3-4), 919-935 (2016)

153. Ardilouze, C., Batté, L., Decharme, B., & Déqué, M. On the link between summer dry bias over the US Great Plains and seasonal temperature prediction skill in a dynamical forecast system. *Weather and Forecasting*, 34(4), 1161-1172 (2019)

154. Marshall, A. G., & Scaife, A. A. Improved predictability of stratospheric sudden warming events in an atmospheric general circulation model with enhanced

stratospheric resolution, *J. Geophys. Res.*, 115, D16114,

doi:10.1029/2009JD012643 (2010)

155. Boer, G. J., & Hamilton, K. QBO influence on extratropical predictive skill, *Clim. Dyn.*, 31, 987–1000 (2008)

156. Marshall A.G., & Scaife A.A. Impact of the QBO on surface winter climate. *J. Geophys. Res.* 114, D18110, DOI: 10.1029/2009JD011737 (2009)

157. Scaife, A.A., M. Athanassiadou, M. Andrews, A. Arribas, M. Baldwin, N. Dunstone, J. Knight, C. MacLachlan, E. Manzini, W.A. Muller, H. Pohlmann, D. Smith, T. Stockdale and A. Williams, Predictability of the quasi-biennial oscillation and its northern winter teleconnection on seasonal to decadal timescales, *Geophys. Res. Letts.*, 41, 1752-1758, 10.1002/2013GL059160 (2014b)

158. Doblas-Reyes, F.J., R. Hagedorn, T. N. Palmer, and J.-J. Morcrette, Impact of increasing greenhouse gas concentrations in seasonal ensemble forecasts. *Geophys. Res. Lett.*, 33, L07708, doi:10.1029/2005GL025061 (2006)

159. Solaraju-Murali, B., Caron, L.-P. , González-Reviriego, N., Doblas-Reyes, F.J., Multi-year prediction of European summer drought conditions for the agricultural sector. *Environmental Research Letters*, doi:10.1088/1748-9326/ab5043 (2019)

160. Cassou, C., Y. Kushnir, Y., E., A., F., I.-S., & N. Decadal Climate Variability and Predictability: Challenges and opportunities. *Bull. Am. Meteorol. Soc.*, 479–490, doi:10.1175/bams-d-16-0286.1 (2018)

161. Liu, Z., Di Lorenzo, E. Mechanisms and Predictability of Pacific Decadal Variability. *Curr. Clim. Chang. Reports*, 4, 128–144, doi:10.1007/s40641-018-0090-5 (2018)

162. Power, S., T. Casey, C. Folland, A. Colman, and V. Mehta, Inter-decadal modulation of the impact of ENSO on Australia. *Clim. Dyn.*, 15, 319–324, doi:10.1007/s003820050284 (1999)

163. Mantua, Nathan J.; Hare, Steven R.; Zhang, Yuan; Wallace, John M.; Francis, Robert C. A Pacific interdecadal climate oscillation with impacts on salmon production. *Bulletin of the American Meteorological Society*. 78 (6): 1069–79. (1997)

164. Newman, M., and Coauthors, The Pacific decadal oscillation, revisited. *J. Clim.*, 29, 4399–4427, doi:10.1175/JCLI-D-15-0508.1 (2016)

165. Chiang, J.C.H. & Vimont, D.J. Analogous Pacific and Atlantic meridional modes of tropical atmosphere-ocean variability. *J. Climate*, 17, 4143-4158 (2004)

166. Di Lorenzo, E., et al., North Pacific Gyre Oscillation links ocean climate and ecosystem change. *GRL*, vol. 35, L08607, doi:10.1029/2007GL032838. (2008)

167. Han W., J. Vialard, M.J. McPhaden, T. Lee, Y. Masumoto, M. Feng, and W. de Ruijter, Indian Ocean Decadal Variability: A Review. *Bull. Amer. Meteor. Soc.*, 95, 1679-1703 (2014a)

168. Han W., G.A. Meehl, A. Hu, M. Alexander, T. Yamagata, D. Yuan, M. Ishii, P. Pegion, J. Zheng, B. Hamlington, X.-W. Quan, and R. Leben, Intensification of decadal and multi-decadal sea level variability in the western tropical Pacific during recent decades. *Climate Dynamics*, 43:1357-1379 (2014b)

169. Li, Y., Han, W., Wang, F., Zhang, L, & Duan, J. Vertical structure of the upper-Indian Ocean thermal variability. *J. Clim.*, 33, 7233–7253, <https://doi.org/10.1175/JCLI-D-19-0851.1> (2020)

170. Tozuka, T., J. Luo, S. Masson, and T. Yamagata, Decadal modulations of the Indian Ocean dipole in the SINTEX-F1 coupled GCM. *J. Climate*, 20, 2881–2894, doi:10.1175/JCLI4168.1 (2007)

171. Feng, M., H.H. Hendon, S.-P. Xie, A. G. Marshall, A. Schiller, Y. Kosaka, N. Caputi, and A. Pearce, Decadal increase in Ningaloo Niño since the late 1990s, *Geophys. Res. Lett.*, 42, 104–112, doi:10.1002/2014GL062509 (2015)

172. Ummenhofer CC, Biastoch A, and Böning CW. Multi-decadal Indian Ocean variability linked to the Pacific and implications for preconditioning Indian Ocean Dipole events. *Journal of Climate*, 30, 1739-1751. (2017)

173. Lee, S.-K., W. Park, M. O. Baringer, A. L. Gordon, B. Huber, and Y. Liu. Pacific origin of the abrupt increase in Indian Ocean heat content during the warming hiatus. *Nat. Geo.*, 8, 445–450 (2015)

174. Nieves, V., J. K. Willis, and W. C. Patzert. Recent hiatus caused by decadal shift in Indo-Pacific heating. *Science* 349(6247):532–535, doi:10.1126/science. Aaa4521 (2015)

175. Jin, X., Kwon, Y.-O., Ummenhofer, C. C., Seo, H., Kosaka, Y., & Wright, J. S. Distinct mechanisms of decadal subsurface heat content variations in the eastern and western Indian Ocean modulated by tropical Pacific SST. *Journal of Climate*, 31, 7751-7769 (2018)

176. Annamalai, H., J. Potemra, R. Murtugudde, and J. P. McCreary. Effect of preconditioning on the extreme climate events in the tropical Indian Ocean. *Journal of Climate*, 18, 3450–3469 (2005)

177. Henley, B.J., et al. Spatial and temporal agreement in climate model simulations of the Interdecadal Pacific Oscillation, *Env. Res. Lett.*, 12, 044011 (2017)

178. Fasullo, J. T., Phillips, A.S., & Deser C. Evaluation of leading modes of climate variability in the CMIP Archives. *J. Clim.* doi:10.1175/JCLI-D-19-1024.1 (2020)

179. Mann, M.E., Steinman, B.A., Miller, S.K., Absence of internal multidecadal and interdecadal oscillations in climate model simulations, *Nature Communications*, <https://doi.org/10.1038/s41467-019-13823-w> (2020)

180. Kosaka, Y., & and Xie, S.-P. The tropical Pacific as a key pacemaker of the variable rates of global warming. *Nature. Geo.*, DOI: 10.1038/NGEO2770 (2016)

181. Tung, K.-K., & X. Chen, X. Understanding the recent global surface warming slowdown: A review. *Climate*, 6, 82, doi:10.3390/cli6040082 (2018)

182. England, M. H., S. McGregor, P. Spence, et al. Recent intensification of wind-driven circulation in the Pacific and the ongoing warming hiatus, *Nat. Clim. Change*, 4(3), 222–227. (2014)

183. Meehl, G.A., Teng, H., & Arblaster, J.M. Climate model simulations of the observed early-2000s hiatus of global warming. *Nature Climate Change*, 4, 898—902, DOI: 10.1038/NCLIMATE2357 (2014)

184. Fyfe, J. C., Meehl, G. A., M. H., M. E., B. D., G. M., Hawkins, E., Gillett, N. P., S.-P., Kosaka, Y., &, N.C. Making sense of the early-2000s warming slowdown. *Nature Climate Change* 6(3), (2016)

185. Xie, S.-P., and Y. Kosaka, What caused the global surface warming hiatus of 1998-2013? *Curr. Clim. Change Rep.*, 3, 128-140, doi: 10.1007/s40641-017-0063-0 (2017)

186. Seager, R., Cane, M., Henderson, N. et al. Strengthening tropical Pacific zonal sea surface temperature gradient consistent with rising greenhouse gases. *Nat. Clim. Change.* **9**, 517–522 <https://doi.org/10.1038/s41558-019-0505-x> (2019)

187. Chen, X. & Tung, K.-K. Varying planetary heat sink led to global-warming slowdown and acceleration, *Science*, **345**, 897-903 (2014)

188. Santer, B. D., et al. Observed multivariable signals of late 20th and early 21st century volcanic activity, *Geophys. Res. Lett.*, **42**, 500–509, doi:10.1002/2014GL062366 (2015)

189. Smith, D. M., and Coauthors, Role of volcanic and anthropogenic aerosols in the recent global surface warming slowdown. *Nat. Clim. Chang.*, **6**, 936. <https://doi.org/10.1038/nclimate3058> (2016)

190. Oudar, T., Kushner, P. J., Fyfe, J., & Sigmond, M. No impact of anthropogenic aerosols on early 21st century global temperature trends in a large initial-condition ensemble. *Geophysical Research Letters*, **45**, 9245–9252. <https://doi.org/10.1029/2018GL078841> (2018)

191. Guemas, V., Doblas-Reyes, F. J., Andreu-Burillo, I., Asif, M. Retrospective prediction of the global warming slowdown in the past decade. *Nature Climate Change*, **3**, 649-653, doi:10.1038/nclimate1863. (2013)

192. Bordbar, M. H., M. H. England, A. Sen Gupta, A. Santoso, A. S. Taschetto, T. Martin, W. Park and M. Latif, Uncertainty in near-term global surface warming linked to tropical Pacific climate variability, *Nature Communications*, **10**:1990, DOI: 10.1038/s41467-019-09761-2 (2019)

193. Meehl, G.A., Chung, C.T.Y., Arblaster, J.M., Holland, M.M., & Bitz, C.M. Tropical decadal variability and the rate of Arctic sea ice retreat, *Geophys. Res. Lett.*, 10.1029/2018GL079989 (2018)

194. Meehl, G.A., Arblaster, J.M., Bitz, C., Chung, C.T.Y., & Teng, H. Antarctic sea ice expansion between 2000-2014 driven by tropical Pacific decadal climate variability. *Nature Geoscience*, DOI: 10.1038/NGEO2751 (2016a)

195. Purich, A., M. H. England, W. Cai, Y. Chikamoto, A. Timmermann, J. C. Fyfe, L. Frankcombe, G. A. Meehl and J. M. Arblaster. Tropical Pacific SST drivers of recent Antarctic sea ice trends, *Journal of Climate*, 29, 8931-8948. (2016)

196. Meehl, G.A., Hu, A., & Teng, H. Initialized decadal prediction for transition to positive phase of the Interdecadal Pacific Oscillation. *Nature Communications.*, 7, doi:10.1038/NCOMMS11718 (2016b)

197. Thoma, M., Greatbatch, R.J., Kadow, C., & Gerdes, R. Decadal hindcasts initialized using observed surface wind stress: evaluation and prediction out to 2024. *Geophys. Res. Lett.*, 42, 6454–6461, <https://doi.org/10.1002/2015GL064833> (2015)

198. Meehl, G.A., Arblaster, J.M., Chung, C.T.Y., Holland, M. M., DuVivier, A., Thompson, L., Yang, D., & Bitz, C.M. Recent sudden Antarctic sea ice retreat caused by connections to the tropics and sustained ocean changes around Antarctica, *Nature Comms.*, 10:14, <https://doi.org/10.1038/s41467-018-07865-9> (2019)

199. Yin, J., Overpeck, J., Peyser, C., & Stouffer, R. Big jump of record warm global mean surface temperature in 2014–2016 related to unusually large oceanic heat

releases. *Geophysical Research Letters*, **45**, 1069–1078.
<https://doi.org/10.1002/2017GL076500> (2018)

200. Booth, B. B. B., Dunstone, N. J., P. R., Andrews, T., & N. Aerosols implicated as a prime driver of twentieth-century North Atlantic climate variability. *Nature*, **484**, 228. <https://doi.org/10.1038/nature10946> (2012)

201. Watanabe, M., & Tatebe, H. Reconciling roles of sulphate aerosol forcing and internal variability in Atlantic multidecadal climate changes. *Clim. Dyn.*, **53**, 4651–4665, <https://doi.org/10.1007/s00382-019-04811-3>. (2019)

202. Hermanson, L., Bilbao, R., Dunstone, N., Ménégoz, M., Ortega, P., Pohlmann, H., ... & Yeager, S. Robust multiyear climate impacts of volcanic eruptions in decadal prediction systems. *Journal of Geophysical Research: Atmospheres*, e2019JD031739, <https://doi.org/10.1029/2019JD031739>, (2020)

203. Menary, M.B. & Scaife, A.A. Forced multidecadal variability of the Atlantic meridional overturning circulation. *Climate Dynamics* **42** (2014)

204. Cai, W., Wu, L., Lengaigne, M., Li, T., McGregor, S., et al. Pantropical climate interactions. *Science*, **363**, eaav4236, doi:10.1126/science.aav4236. (2019)

205. Mechoso, R. ed. *Interacting Climates of Ocean Basins: Observations, Mechanisms, Predictability, and Impacts*, Cambridge University Press, ISBN: 9781108492706 (2020)

206. Meehl, G.A., A. Hu, F. Castruccio, M.H. England, S.C. Bates, G. Donabasoglu, S. McGregor, J.M. Arblaster, S.-P. Xie, and N. Rosenbloom, Atlantic and Pacific tropics connected by mutually interactive decadal-timescale processes , *Nature Geo.*, doi:10.1038/s41561-020-00669-x (2020)

207. Chikamoto, Y. et al. Skillful multi-year predictions of tropical trans-basin climate variability. *Nature Commun.* 6, 6869 (2015)

208. Ruprich-Robert, Y., and co-authors, Assessing the climate impacts of the observed Atlantic Multidecadal Variability using the GFDL CM2.1 and NCAR CESM1 global coupled models, *J. Clim.*, 30, 2785—2810, DOI: 10.1175/JCLI-D-16-0127.1 (2017)

209. Levine, A. F. Z., McPhaden, M. J., & Frierson, D. M. W. The impact of the AMV on multidecadal ENSO variability, *Geophys. Res. Lett.*, 44, 3877–3886, doi:10.1002/2017GL072524 (2017)

210. Kumar, A., Bhaskar, J., & Wang, H. Attribution of SST variability in global oceans and the role of ENSO. *Clim Dyn.*, 43, 209–220, doi:10.1007/s00382-013-1865-y (2014)

211. Taschetto, A.S., R.R. Rodrigues, G. A. Meehl, S. McGregor, and M. H. England, How sensitive are the Pacific-North Atlantic teleconnections to the position and intensity of El Niño-related warming. *Clim. Dyn.*, DOI:10.1007/s00382-015-2679-x (2015)

212. Han W., G.A. Meehl, A. Hu, J. Zheng, J. Kenigson, J. Vialard, B. Rajagopalan, and Yanto, Decadal variability of Indian and Pacific Walker Cells: Do they co-vary on decadal timescales? *J. Climate*, 30, 8447-8468, DOI: 10.1175/JCLI-D-16-0783.1 (2017)

213. Han W., et al.. Multi-decadal trend and decadal variability of the regional sea level over the Indian Ocean since the 1960s: Roles of climate modes and external forcing, *Climate*, 6(2), 51; <https://doi.org/10.3390/cli6020051> (2018)

214. Deepa JS, Gnanaseelan C, Sandeep Mohapatra JS, Chowdary AK, Kakatkar R, Parekh A The tropical Indian Ocean decadal sea level response to the Pacific Decadal Oscillation forcing. *Clim Dyn* 52:5045. <https://doi.org/10.1007/s00382-018-4431-9> (2019)

215. Zhang, R., R. Sutton, G. Danabasoglu, Y.-O. Kwon, R. Marsh, S. G. Yeager, D. E. Amrhein, and C. M. Little, A review of the role of the Atlantic meridional overturning circulation in Atlantic multidecadal variability and associated climate impacts. *Rev. Geophys.*, 57, 316-375, doi: 10.1029/2019RG000644 (2019)

216. Li, X., Xie, S.-P., Gille, S.T., & Yoo, C. Atlantic-induced pan-tropical climate change over the past three decades. *Nature Clim. Change*, 6, DOI: 10.1038/NCLIMATE2840 (2015)

217. Li, H., Ilyina, T., Müller, W.A., & Seinz, F. Decadal prediction of the North Atlantic CO₂ uptake, *Nature Communications*, 7, 11076, doi:10.1038/ncomms11076 (2016)

218. Jin, D., and B. P. Kirtman How the annual cycle affects the extratropical response to ENSO, *J. Geophys. Res.*, 115, D06102, doi:10.1029/2009JD012660 (2010)

219. Zhang L., W. Han, and F. Sienz, Unraveling causes for the changing behavior of tropical Indian Ocean in the past few decades. *J. Clim.*, 31, 2377-2388, doi: 10.1175/JCLI-D-17-0445.1 (2018)

220. Thornton H. et al. Skillful seasonal prediction of winter gas demand. *Env. Res. Lett.*, 14, 2, 024009, (2019)

221. Palin, E. J., A. A. Scaife, E. Wallace, E. C. D. Pope, A. Arribas, and A. Brookshaw, Skillful seasonal forecasts of winter disruption to the U.K. transport

system. *J. Appl. Meteor. Climatol.*, 55, 325–344,
doi:<https://doi.org/10.1175/JAMC-D-15-0102.1> (2016)

222. Towler E, Paimazumder, D., and Done, J., Toward application of decadal climate predictions. *J. Applied Meteorology and climatology*, 57, 555-568. (2018)

223. Vecchi, G. A. et al. On the seasonal forecasting of regional tropical cyclone activity. *J. Clim.* 27, 7994–8016 (2014)

224. Annan, J. D., et al. Parameter estimation in an atmospheric GCM using the Ensemble Kalman Filter. *Nonlinear Processes in Geophysics* 12(3), DOI: 10.5194/npg-12-363-2005 (2005)

225. Düben, P.D., Hugh McNamara, H. & Palmer, T.N. The use of imprecise processing to improve accuracy in weather & climate prediction. *Journal of Computational Physics* 271, 2-18 (2014)

226. Palmer, T.N., Peter Düben, P., & McNamara, H. Stochastic modelling and energy-efficient computing for weather and climate prediction. *Phil. Trans. Roy. Soc. A*, 20140118, <https://doi.org/10.1098/rsta.2014.0118>, (2014)

227. Ham, Y.-G., Kim, J.-H. & Luo, J.-J. Deep learning for multi-year ENSO forecasts, *Nature*, 573 <https://doi.org/10.1038/s41586-019-1559-7> (2019)

228. Zhang , S. and coauthors. Coupled data assimilation and parameter estimation in coupled ocean-atmosphere models: a review. *Clim Dyn* 54, 5127-5144 (2020)

229. Karspeck A.R. and coauthors, A global coupled ensemble data assimilation system using the Community Earth System Model and the Data Assimilation Research Testbed. *Q J R Meteorol Soc* 144, 2404-2430 (2018)

230. Mulholland, D., Laloyaux, P., Haines, K. and Balmaseda, M. Origin and impact of initialization shocks in coupled atmosphere–ocean forecasts. *Monthly Weather Review*, **143**, 4631–4644 (2015)

231. Saha S and coauthors. The NCEP climate forecast system reanalysis. *Bull Am Meteorol Soc*, **91**, 1015-1057 (2010)

232. Laloyaux P et al. A coupled data assimilation system for climate reanalysis. *Q J R Meteorol Soc*, **142**, 65-78 (2016)

233. Herman, R.J., Giannini, A., Biasutti, M. *et al.* The effects of anthropogenic and volcanic aerosols and greenhouse gases on twentieth century Sahel precipitation. *Sci Rep* **10**, 12203 <https://doi.org/10.1038/s41598-020-68356-w> (2020)

234. Schurer, A., Hegerl, G., Mann, M.E., Tett, S.F.B., Separating forced from chaotic climate variability over the past millennium, *J. Climate*, **26**, 6954-6973, (2013)

235. Ault, T. R., Cole, J. E., Overpeck, J. T., Pederson, G. T., St. George, S., Otto-Bliesner, B., Woodhouse, C. A., & Deser, C. The continuum of hydroclimate variability in western North America during the last millennium. *J. Climate*, **26**, 5863–5878, doi:<https://doi.org/10.1175/JCLI-D-11-00732.1> (2013)

236. Laepple, T., and P. Huybers. Global and regional variability in marine surface temperatures. *Geophys. Res. Lett.*, **41**, 2528–2534, doi:<https://doi.org/10.1002/2014GL059345>. (2014)

237. Loope, G., Thompson, D.M., Cole, J.E., & Overpeck, J. Is there a low-frequency bias in multiproxy reconstructions of Pacific SST variability? *Quaternary Science Reviews* **246**, 106530, <https://doi.org/10.1016/j.quascirev.2020> (2020)

238. Frankignoul, C., Muller, P. & Zorita, E. A simple model of the decadal response of the ocean to stochastic wind forcing. *J. Phys. Oceanogr.*, **27**, 1533-1546 (1997)

239. Capotondi, A., Alexander, M.A. & Deser, C. Why are there Rossby wave maxima in the Pacific at 10S and 13N? *J. Phys. Oceanogr.*, **33**, 1549-1563 (2003)

240. Chikamoto, Y., Timmermann, A., Widlansky, M. J., M. A., & L. Multi-year predictability of climate, drought, and wildfire in southwestern North America. *Sci. Rep.* <https://www.ncbi.nlm.nih.gov/pubmed/28747719> (2017)

241. Lovenduski, N.S., Bonan, G.B., Yeager, S.G., Lindsay, K., Lombardozzi, D.L. High predictability of terrestrial carbon fluxes from an initialized decadal prediction system, *Environmental Research Letters*, **14**, 124074, doi:10.1088/1748-9326/ab5c55 (2019a)

242. Sospedra-Alfonso, R., W. J. Merryfield, and V. V. Kharin, Representation of snow in the Canadian Seasonal to Interannual Prediction System: Part II. Potential predictability and hindcast skill. *J. Hydrometeor.*, **17**, 2511–2535, <https://doi.org/10.1175/JHM-D-16-0027.1> (2016)

243. Kapnick, S. B., and Coauthors. Potential for western US seasonal snowpack prediction. *Proc. Natl. Acad. Sci. USA*, **115**, 1180–1185, <https://doi.org/10.1073/pnas.1716760115> (2018)

244. Holbrook, N.J et al. Keeping pace with marine heatwaves. *Nat. Rev. Earth Environ.* **1**, 482-493 (2020)

245. Batté, L., Valisuo, I., Chevallier, M., Acosta Navarro, J. C., Ortega, P., & Smith, D. Summer predictions of Arctic sea ice edge in multi-model seasonal reforecasts. *Climate Dynamics*, **54**, 5013-5029, DOI:10.1007/s00382-020-05273-8 (2020)

246. Subramanian, A., Juricke, S., Dueben, P., Palmer, T., A Stochastic Representation of Subgrid Uncertainty for Dynamical Core Development. *Bulletin of the American Meteorological Society* 100 (6), 1091-1101. (2019)

247. Penny, S.G., et al Observational needs for improving ocean and coupled reanalysis, S2S prediction, and decadal prediction, *Front. Mar. Sci.*, 11, <https://doi.org/10.3389/fmars.2019.00391>, (2019)

248. Lofverstrom, et al. An efficient ice sheet/Earth system model spin up procedure for CESM2.1 and CISM2.1: description, evaluation, and broader applicability, *JAMES*, <https://doi.org/10.1029/2019MS001984> (2020)

249. Gettelman, A., Mills, M. J., Kinnison, D. E., Garcia, R. R., Smith, A. K., Marsh, D. R., et al. The Whole Atmosphere Community Climate Model Version 6 (WACCM6). *J. Geophys. Res. Atmos.*, 124. <https://doi.org/10.1029/2019JD030943> (2019)

250. Tommasi, D. C., and coauthors Managing living marine resources in a dynamic environment: The role of seasonal to decadal climate forecasts, *Progress in Oceanography*, 152, 15-49, doi:10.1016/j.pocean.2016.12.011 (2017)

251. Stock, C. A., and coauthors, Seasonal sea surface temperature anomaly prediction for coastal ecosystems, *Progress in Oceanography*, 137, 219-236, doi:10.1016/j.pocean.2015.06.007 (2015)

252. Liu, G. et al. Predicting heat stress to inform reef management: NOAA Coral Reef Watch's 4-month coral bleaching outlook. *Front. Mar. Sci.*, <https://doi.org/10.3389/fmars.2018.00057> (2018)

253. Capotondi, A., Sardeshmukh, P. D., Di Lorenzo, E., Subramanian, A., & Miller, A.J. Predictability of US West Coast ocean temperatures is not solely due to ENSO. *Sci. Rep.* 9:10993, <https://doi.org/10.1038/s41598-019-47400-4>, (2019a)

254. Wells, M.L. et al. Harmful algal blooms and climate change: Learning from the past and present to forecast the future. *Harmful Algae*, 49, 68-93 (2015)

255. Séférian, R., and coauthors, Multiyear predictability of tropical marine productivity, *Proceedings of the National Academy of Sciences*, 111, 11646-11651, doi:10.1073/pnas.1315855111 (2014)

256. Park, J.-Y., C. A. Stock, J. P. Dunne, X. Yang, and A. Rosati Seasonal to multiannual marine ecosystem prediction with a global Earth system model, *Science* 365(6450), 284-288, doi:10.1126/science.aav6634 (2019)

257. Krumhardt, K. M., Lovenduski, N. S., Long, M. C., Luo, J. Y., Lindsay, K., Yeager, S. G., & Harrison, C. Potential predictability of net primary production in the ocean, *Global Biogeochemical Cycles*, 34, e2020GB006531, doi:10.1029/2020GB006531 (2020)

258. Siedlecki, S. A., and coauthors, Experiments with seasonal forecasts of ocean conditions for the northern region of the California Current upwelling system, (2016)

259. Brady, R. X., Lovenduski, N. S., Yeager, S. G., Long, M. C., & Lindsay, K. Skillful multiyear predictions of ocean acidification in the California Current System, *Nature Communications*, 11: 2166, doi:10.1038/s41467-020-15722-x (2020)

260. Séférian, R., S. Berthet, and M. Chevallier, Assessing the decadal predictability of land and ocean carbon uptake, *Geophysical Research Letters*, 45(5), 2455-2466, doi:10.1002/2017GL076092 (2018)

261. Lovenduski, N.S., Yeager, S.G., Lindsay, K., & Long, M.C. Predicting near-term variability in ocean carbon uptake, *Earth System Dynamics*, 10, 45-57, doi:10.5194/esd-10-45-2019 (2019b)

262. Li, H, Ilyina, T., Müller, W.A., & Landschützer, P. Predicting the variable ocean carbon sink, *Science Advances*, 5, doi:10.1126/sciadv.aav6471 (2019)

263. Bett, P. E., Williams, K. E., Burton, C., Scaife, A. A., Wiltshire, A. J., & Gilham, R. Skillful seasonal prediction of key carbon cycle components: NPP and fire risk. *Environmental Research Communications*, 2(5), 055002, (2020)

264. Park, J.-Y., Dunne, J. P., & Stock, C. A. Ocean chlorophyll as a precursor of ENSO: An earth system modeling study. *Geophys. Res. Lett.*, <https://doi.org/10.1002/2017GL076077> (2018)

265. Capotondi, A., et al. Observational needs supporting marine ecosystem modeling and forecasting: From the global ocean to regional and coastal systems. *Frontiers in Marine Science*, 6:623, doi:10.3389/fmars.2019.00623, (2019b)

266. Fennel K., et al. Advancing Marine Biogeochemical and Ecosystem Reanalyses and Forecasts as Tools for Monitoring and Managing Ecosystem Health. *Front. Mar. Sci.* 6:89. (2019)

267. Weisheimer, A., Palmer, T. N. On the reliability of seasonal climate forecasts *J. R. Soc. Interface*, <http://doi.org/10.1098/rsif.2013.1162> (2014)

268. National Academies of Sciences, Engineering and Medicine. Next Generation Earth System Prediction: Strategies for Subseasonal to Seasonal Forecasts (pp. 1–351). Washington, DC: National Academies Press (2017).

269. National Research Council. Assessment of Intraseasonal to Interannual Climate Prediction and Predictability (pp. 1–193). Washington, D.C.: National Academies Press. <http://doi.org/10.17226/12878> (2010).

Acknowledgements

The foundations of this paper emerged from a workshop held by National Academies of Sciences, Engineering and Medicine in 2015 at Woods Hole, MA, and the authors gratefully acknowledge support from Amanda Purcell and Nancy Huddleston. Portions of this study were supported by the Regional and Global Model Analysis (RGMA) component of the Earth and Environmental System Modeling Program of the U.S. Department of Energy's Office of Biological & Environmental Research (BER) via National Science Foundation IA 1844590. This work also was supported by the National Center for Atmospheric Research, which is a major facility sponsored by the National Science Foundation under Cooperative Agreement No. 1852977. M.E.M. was supported by a grant from the NSF Paleoclimate Program #1748097. F.J.D.R. and M.G.D. were supported by the H2020 EUCLIP project under Grant agreement no. 776613, M.G.D also by the Ramón y Cajal 2017 grant reference RYC-2017-22964. A. C. was supported by the NOAA Climate Program Office Climate Variability and Predictability (CVP), and Modeling, Analysis, Predictions and Projections (MAPP) Programs.

A.C.S. acknowledges support from the NOAA Climate Variability and Predictability Program (Award NA18OAR4310405) and the National Oceanic and Atmospheric Administration (NOAA MAPP; NA17OAR4310106) for support. N.S.L. is grateful for support from the NSF (OCE-1752724). D.M.T. acknowledges support from NCAR Advanced Study Program and NSF (OCE-1931242). S.C.S was supported by the Joint Institute for the Study of the Atmosphere and Ocean (JISAO) Postdoctoral Fellowship. A.A.S. and D.M.S were supported by the Met Office Hadley Centre Climate Programme funded by BEIS and Defra and by the European Commission Horizon 2020 EUCLIP project (GA 776613).

Author contributions

H.T. suggested the original concept. G.A.M. led the overall conceptual design, and coordinated the writing. J.H.R. and H.T. made major contributions to the conceptual design and organization. J.H.R. generated Fig. 1a. H.T. generated Fig. 4. All authors discussed the concepts presented and contributed to the writing.

Competing Interests

The authors declare no competing interests.

Peer review information

Nature Reviews Earth & Environment thanks Debra Hudson, Constantin Ardilouze, Terrence O’Kane and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations

Supplementary information

Supplementary information is available for this paper at <https://doi.org/10.1038/s415XX-XXX-XXXX-X>

Related links

Copernicus Climate Change Service: <https://www.copernicus.eu/en>

APCC climate predictions: <https://www.apcc21.org/>

North American Multi-model Ensemble: <https://www.cpc.ncep.noaa.gov/products/NMME/>

Key points

- Initialization methods vary greatly across different prediction timescales creating difficulties for seamless prediction.
- Model error and drift limit predictability across all timescales. Although higher resolution models show promise in reducing these errors, improvement in physical parameterizations are needed to improve predictability.


- The effects of land processes, interactions across various ocean basins and the role of stratospheric processes in predictability are not well understood.
- Predictability on S2D timescales is largely associated with predictability of the major modes of variability in the atmosphere and the ocean.
- Evolution of Earth System models will lead to predictability of more societal-relevant variables spanning multiple parts of the Earth System

Table 1. General characteristics of models used for S2S, S2I and S2D initialized predictions*.

Timescale	Number of models	Atmospheric resolution & levels	Ocean resolution & levels	Components initialized	Initialization	Number of ensembles	Prediction length
S2S	18	25—200 km 17—91 levels	8—200 km 25—75 levels	Most initialize atmosphere, ocean, land and sea ice	Full field	4—51	31—62 days
S2I	13	36—200 km 24—95 levels	25—200 km 24—74 levels	All initialize atmosphere, ocean, land and sea ice	Full field	10—51	6—12 months
S2D	14	50—20 0km 26—95 levels	25—100 km 30—75 levels	Models range from initializing only ocean, to initializing atmosphere, ocean, land and sea ice	Full field, anomaly	10—40	5—10 years

*A full and more complete accounting of model features is given in Supplementary Table 1, 2 and 3 for S2S, S2I and S2D models.

a Predictability sources and timescales



b Predictability skill from initial state vs. external forcing




Figure 1.

Timescales and processes involved with initialized predictions. **a** Timescales and sources of predictability for S2S, S2I, and S2D. Lighter green shading indicates larger uncertainty. MJO: Madden-Julian Oscillation; NAO: North Atlantic Oscillation; QBO: Quasi-Biennial Oscillation; SSWs: Sudden Stratospheric Warmings; ENSO: El Niño-Southern Oscillation; PDV: Pacific Decadal Variability; AMV: Atlantic Multi-decadal variability; GMST: Global Mean Surface Temperature; GHG: Greenhouse Gas. **b** skill in predicting the upper 300m of the Atlantic Ocean temperature, as measured by relative entropy, in initialized models (blue) and those forced by RCP4.5 (red). Skill is high for initialized predictions at S2S and S2I timescales (<2 years), but decreases toward S2D (year 3-9), after which time skill from external forcing increases. Panel b adapted, with permission, from ref 10.




Figure 2. Influence of ensemble size and lead year ranges on predictive skill. **a|** Skill (as measured by anomaly correlation coefficient) in predicting S2S globally averaged NDJFM surface air temperature (excluding the Antarctic) from CESM initialized hindcasts of various ensemble size (grey line). Shading denotes the 5% and 95% significance levels. Blue and red whiskers illustrate predictive skill for NCEP CFSv2 and ECMWF subseasonal hindcasts, respectively. **b|** Skill (as measured by the anomaly correlation coefficient) in predicting S2D wintertime NAO using ensembles of different sizes from the Decadal Prediction Large Ensemble, DPLE (20). Each line depicts a different lead year range, with those that are colored corresponding to statistically significant correlations; the darker the shading, the greater the statistical significance. The dashed-dotted line shows the skill of the sub-ensemble mean against a single member of the ensemble (averaged for all possible combinations). Both panels illustrate that the more ensemble members, the higher the skill for longer lead year ranges. Panel b adapted, with permission, from ref 37.




Figure 3. Extending proxy observations of S2D variability back in time derived from corals. **a** | Global mean surface temperature anomalies , **b** | 30 year running means of the coral-based Indian Ocean Dipole (IOD) (blue) and El Niño-Southern Oscillation (ENSO) (red); **c** | scatter plot of coral-based IOD and ENSO; **d** | equatorial Pacific west-east SST gradient, shading represents uncertainty ; **e** | central and eastern Pacific El Niño derived from teleconnected climate patterns. **f** | An indication of reconstructions considered robust in panel e. Collectively, the figures illustrate a strengthening of IOD-ENSO decadal variability after ~ 1590 . Figure adapted, with permission, from ref 58.




Figure 4. Impact of model drift on initialized predictions. Globally averaged surface temperature predictions from the Decadal Prediction Large Ensemble (20) as a function of simulation year. Initial state predictions (blue dots) compare well to observations (black line), but drift (progression of blue dots to red dots) toward the model's systematic error state represented by the uninitialized state (dark gray line; gray shading is range of uninitialized projections).




Figure 5. Initialized S2S predictions of the MJO. **a** | observed outgoing longwave radiation (OLR) anomalies averaged over 5°S to 5°N as a function of the stage of the Madden-Julian Oscillation (MJO). **b-g** | as in **a**, but for various initialized predictions, with OLR anomalies taken as the average of simulations days 15-21. MJO events are identified based on the Real-time Multivariate MJO (RMM) index amplitude ≥ 1 . The eastward propagation of MJO-related OLR anomalies is well captured by all six models. Figure adapted, with permission, from ref 6.




Figure 6. S2D predictions and aspects of time-evolving globally averaged temperature.

a| Prediction skill, measured as the anomaly correlation coefficient, of sea surface temperature (SST) averaged over lead years 5-9 from the decadal prediction large ensemble (20); darker red indicates higher skill. **b|** improvement in prediction skill from initialized predictions in a over and above a persistence prediction; darker red indicates better skill in the initialized predictions, thus showing the value-added of initialized predictions **c|** Schematic of the “rising staircase”, illustrating how natural decadal-scale temperature fluctuations (blue) are tilted upwards owing to anthropogenic greenhouse gas emissions (red), producing accelerated warming in some decades, and reduced warming in others. **d|** time series of observed global mean surface temperature anomalies showing characteristics of the rising staircase: accelerated warming over 1980-2000 and 2014-present, and a slow-down in the rate of warming over 2000-2014. Panels a and b adapted, with permission, from ref 20. Panel c adapted, with permission, from ref 268 Panel d adapted, with permission, from NASA.

TOC summary

Initialized climate predictions offer distinct benefits for multiple stakeholders. This Review discusses initialized prediction at subseasonal-to-seasonal (S2S), seasonal-to-interannual (S2I) and seasonal-to-decadal (S2D) timescales, highlighting potential for skillful predictions in the years to come.