

## LA-UR-21-31229

Approved for public release; distribution is unlimited.

Title: Analytical solutions of the Arrhenius-Semenov problem for constant volume burn

Author(s): Craven, Galen Thomas

Intended for: Report

Issued: 2021-11-10

---

**Disclaimer:**

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

# Analytical solutions of the Arrhenius-Semenov problem for constant volume burn

Galen T. Craven

*Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico, USA*

Analytical solutions to the Semenov thermal ignition problem for constant volume burn governed by Arrhenius reaction kinetics are derived. Specifically, an approximate analytical solution technique for the Arrhenius-Semenov differential equation is derived for reaction orders  $n \in \mathbb{R}_{>0}$  and exact solutions are also constructed for reaction orders  $n \in \mathbb{N} : n \leq 3$ . The approximation technique relies on expansion of the respective nondominant terms in the differential equation at the lower and upper bounds of the reaction progress variable in order to create a pair of integrable series. The two integrated series are then connected to create a single continuous analytical solution. Excellent agreement is observed between the analytical approximation and solutions obtained numerically. The presented approximation constitutes a simple and robust strategy for solving the Arrhenius-Semenov problem analytically.

## I. INTRODUCTION

Theories describing equilibrium<sup>1–3</sup> and nonequilibrium<sup>4–10</sup> chemical kinetics play a fundamental role in computational treatments of shock-induced chemical reactions and high explosive detonations<sup>11–18</sup>. The principal motivations for developing these theories are to (a) determine chemical reaction rates, (b) to predict reaction products, and (c) to elucidate complex molecular behavior that arises due to energy and charge redistribution that occurs as the reaction proceeds. While various advanced computational methods<sup>19,20</sup> can provide high-level insight into a system’s properties<sup>21–25</sup>, it is often advantageous to describe molecular processes using analytical methods, for example, to develop physical intuition about the system or for computational efficiency. In complex systems, however, nonlinearities typically preclude exact analytical solutions to a system’s dynamical equations and, therefore, approximation methods must be employed. In this report, we develop an analytical approximation method to solve the kinetic equation describing a constant volume burn governed

by Arrhenius reaction kinetics.

Constant volume burn occurs when a thermal explosion takes place in a rigid reaction vessel over a fast enough time scale such that no heat is exchanged across the vessel boundary and the chemical and thermodynamic properties of the system are spatially uniform<sup>26,27</sup>. Following Menikoff's mathematical exposition for constant volume burn from Ref. 26, the ordinary differential equation (ODE) that describes the temperature evolution of a constant volume burn is<sup>26</sup>

$$\frac{dT}{dt} = \frac{\mathcal{Q}}{C_v(\lambda, T)} \mathcal{R}(\lambda, T), \quad (1)$$

where  $\lambda \in [0, 1]$  is a reaction progress variable,  $\mathcal{R}$  is the reaction rate,  $\mathcal{Q}$  is the specific energy released, and  $C_v$  is the specific heat. When  $\lambda = 0$  the system consists of completely unburned reactant and when  $\lambda = 1$  the reactant is completely burned. The initial ( $\lambda = 0$ ) system temperature is  $T_0$  and the final ( $\lambda = 1$ ) system temperature at the end of the reaction is  $T_1$ . Equation (1) is coupled with an equation that describes the time evolution of the reaction progress variable

$$\frac{d\lambda}{dt} = \mathcal{R}(\lambda, T). \quad (2)$$

Here, the reaction rate is assumed to take the Arrhenius form

$$\mathcal{R}(\lambda, T) = (1 - \lambda)^n k \exp \left[ -\frac{T_a}{T(\lambda)} \right], \quad (3)$$

where  $n$  is the reaction order,  $k$  is a pre-exponential frequency factor, and  $T_a$  is the activation temperature. Assuming that  $\mathcal{Q}$  and  $C_v$  are constant, the temperature of the system as a function of reaction progress is

$$T(\lambda) = T_0 + (\mathcal{Q}/C_v)\lambda = T_0 + (T_1 - T_0)\lambda, \quad (4)$$

with  $T_1 = T_0 + \mathcal{Q}/C_v$ . The time-dependence of  $T$  in Eq. (4) arises through the reaction progress variable  $\lambda \rightarrow \lambda(t)$ . Under these conditions, the time evolution of the reaction can be described by the single ODE

$$\frac{d\lambda}{dt} = (1 - \lambda)^n k \exp \left[ -\frac{T_a}{T_0 + (\mathcal{Q}/C_v)\lambda} \right] = (1 - \lambda)^n k \exp \left[ -\frac{T_a}{T_0 + (T_1 - T_0)\lambda} \right]. \quad (5)$$

Our aim is to develop analytical techniques that can be used to solve Eq. (5) over a broad class of reaction orders. Note that this ODE is separable,

$$\int k dt = \int (1 - \lambda)^{-n} \exp \left[ \frac{T_a}{T_0 + (T_1 - T_0)\lambda} \right] d\lambda, \quad (6)$$

and, therefore, the Semenov ignition problem in the presence of Arrhenius rate kinetics can in essence be solved by evaluating the integral on the RHS of Eq. (6):

$$I_n(\lambda) \equiv \int (1 - \lambda)^{-n} \exp \left[ \frac{T_a}{T_0 + (T_1 - T_0)\lambda} \right] d\lambda. \quad (7)$$

Below, in Sec. II we illustrate an analytical technique for approximating  $I_n$  and, therefore, for solving the Semenov problem. In Sec. III we derive exact solutions for reaction orders  $n \in \mathbb{N} : n \leq 3$ . Section IV contains a comparison between the developed approximation and numerical solutions for various reaction conditions.

## II. APPROXIMATE SOLUTION FOR $n \in \mathbb{R}_{>0}$

An exact solution of  $I_n$  is not known for  $n \in \mathbb{R}^+ \setminus \mathbb{N}$ . Below we give an approximate analytical solution for  $n \in \mathbb{R}_{>0}$  which relies on series expansion of different terms in the integrand at different ends of the  $\lambda \in [0, 1]$  interval.

As  $\lambda \rightarrow 1$ , the term  $(1 - \lambda)^{-n}$  dominates the integrand for  $n > 0$ :

$$\lim_{\lambda \rightarrow 1} \frac{\exp [T_a/T(\lambda)]}{(1 - \lambda)^{-n}} = 0. \quad (8)$$

In this fully-burned limit, we therefore expand  $\exp [T_a/T(\lambda)]$  about  $\lambda = 1$  to obtain the approximation

$$\int (1 - \lambda)^{-n} \exp \left[ \frac{T_a}{T(\lambda)} \right] d\lambda \approx \int \sum_{m=0}^{\alpha} (1 - \lambda)^{m-n} \frac{D_{\lambda}^m \exp \left[ \frac{T_a}{T(\lambda)} \right] \Big|_{\lambda=1}}{m!} d\lambda, \quad (9)$$

where  $\alpha$  is the series expansion order and  $D_{\lambda}^m$  is  $m$ -th order derivative operator with respect

to  $\lambda$ . The integrals in the series approximation up to third order ( $\alpha = 3$ ) are

$$\begin{aligned}
I_n^+(\lambda) \approx & \underbrace{\int (1-\lambda)^{-n} \exp\left[\frac{T_a}{T_1}\right] d\lambda}_{I_n^{(0+)}} \\
& + \underbrace{\int (1-\lambda)^{1-n} \exp\left[\frac{T_a}{T_1}\right] \left(\frac{\mathcal{Q}}{C_v}\right) \frac{T_a}{T_1^2} d\lambda}_{I_n^{(1+)}} \\
& + \underbrace{\int (1-\lambda)^{2-n} \exp\left[\frac{T_a}{T_1}\right] \left(\frac{\mathcal{Q}}{C_v}\right)^2 \frac{T_a(T_a + 2T_1)}{2T_1^4} d\lambda}_{I_n^{(2+)}} \\
& + \underbrace{\int (1-\lambda)^{3-n} \exp\left[\frac{T_a}{T_1}\right] \left(\frac{\mathcal{Q}}{C_v}\right)^3 \frac{T_a(T_a^2 + 6T_1^2 + 6T_a T_1)}{6T_1^6} d\lambda}_{I_n^{(3+)}} ,
\end{aligned} \tag{10}$$

where the superscript “+” in  $I_n^+$  denotes that the approximate solution is approached from the upper bound of  $\lambda$  and  $I_n^{(j+)}$  denotes the  $j$ th order term in the upper-bound expansion.

Evaluating each integral in the series yields

$$I_n^{(0+)} = \frac{(1-\lambda)^{1-n}}{n-1} \exp\left[\frac{T_a}{T_1}\right], \tag{11}$$

$$I_n^{(1+)} = \frac{(1-\lambda)^{2-n}}{n-2} \exp\left[\frac{T_a}{T_1}\right] \left(\frac{\mathcal{Q}}{C_v}\right) \frac{T_a}{T_1^2}, \tag{12}$$

$$I_n^{(2+)} = \frac{(1-\lambda)^{3-n}}{n-3} \exp\left[\frac{T_a}{T_1}\right] \left(\frac{\mathcal{Q}}{C_v}\right)^2 \frac{T_a(T_a + 2T_1)}{2T_1^4}, \tag{13}$$

$$I_n^{(3+)} = \frac{(1-\lambda)^{4-n}}{n-4} \exp\left[\frac{T_a}{T_1}\right] \left(\frac{\mathcal{Q}}{C_v}\right)^3 \frac{T_a(T_a^2 + 6T_1^2 + 6T_a T_1)}{6T_1^6}. \tag{14}$$

Note that each of the  $I_n^{(j+)}$  functions above is vertically asymptotic for  $n = j + 1$ . However, as illustrated below in Sec. II A, after applying the initial value to the full solution, there is no asymptote with respect to variation of  $n$ .

In the limit  $\lambda \rightarrow 0$ , the ratio between the two functions in the integrand of  $I_n$  is

$$\lim_{\lambda \rightarrow 0} \frac{\exp[T_a/T(\lambda)]}{(1-\lambda)^{-n}} = \exp\left[\frac{T_a}{T_0}\right]. \tag{15}$$

In this limit, we approximate the integral by expanding the  $(1-\lambda)^{-n}$  term about  $\lambda = 0$ :

$$\int (1-\lambda)^{-n} \exp\left[\frac{T_a}{T(\lambda)}\right] d\lambda \approx \int \sum_{m=0}^{\alpha} \exp\left[\frac{T_a}{T(\lambda)}\right] \lambda^m \frac{D_{\lambda}^m (1-\lambda)^{-n} \Big|_{\lambda=0}}{m!} d\lambda. \tag{16}$$

Expanding the RHS of Eq. (16) up to third order gives the following approximation for  $I_n$ :

$$\begin{aligned}
I_n^-(\lambda) \approx & \underbrace{\int \exp \left[ \frac{T_a}{T(\lambda)} \right] d\lambda}_{I_n^{(0-)}} \\
& + \underbrace{\int \exp \left[ \frac{T_a}{T(\lambda)} \right] n\lambda d\lambda}_{I_n^{(1-)}} \\
& + \underbrace{\int \exp \left[ \frac{T_a}{T(\lambda)} \right] \frac{(n+n^2)}{2} \lambda^2 d\lambda}_{I_n^{(2-)}} \\
& + \underbrace{\int \exp \left[ \frac{T_a}{T(\lambda)} \right] \frac{(2n+3n^2+n^3)}{6} \lambda^3 d\lambda}_{I_n^{(3-)}},
\end{aligned} \tag{17}$$

where the superscript “ $-$ ” denotes that the approximate solution is approached from the lower bound of  $\lambda$  and  $I_n^{(j-)}$  denotes the  $j$ th-order term in the lower-bound expansion.

Evaluating these integrals gives

$$I_n^{(0-)} = \left( \frac{C_v}{Q} \right) \left[ T(\lambda) \exp [h(\lambda)] - T_a \text{Ei} [h(\lambda)] \right], \tag{18}$$

$$I_n^{(1-)} = \frac{n}{2} \left( \frac{C_v}{Q} \right)^2 \left[ T(\lambda) \left( \frac{Q}{C_v} \lambda - T_0 + T_a \right) \exp [h(\lambda)] + T_a (2T_0 - T_a) \text{Ei} [h(\lambda)] \right], \tag{19}$$

$$\begin{aligned}
I_n^{(2-)} = & \frac{n+n^2}{12} \left( \frac{C_v}{Q} \right)^3 \\
& \times \left[ T(\lambda) \left( 2 \left( \frac{Q}{C_v} \right)^2 \lambda^2 + \left( \frac{Q}{C_v} T_a - 2T_0 T_1 + 2T_0^2 \right) \lambda + (T_a^2 - 5T_a T_0 + 2T_0^2) \right) \right. \\
& \left. \times \exp [h(\lambda)] - T_a (T_a^2 - 6T_a T_0 + 6T_0^2) \text{Ei} [h(\lambda)] \right],
\end{aligned} \tag{20}$$

$$\begin{aligned}
I_n^{(3-)} = & \frac{(2n+3n^2+n^3)}{144} \left( \frac{C_v}{Q} \right)^4 \left[ \left( 6 \left( \frac{Q}{C_v} \right)^4 \lambda^4 + 2T_a \left( \frac{Q}{C_v} \right)^3 \lambda^3 - \left( \frac{Q}{C_v} \right)^2 T_a (6T_0 - T_a) \lambda^2 \right. \right. \\
& + \left( \frac{Q}{C_v} \right) T_a (T_a^2 - 10T_a T_0 + 18T_0^2) \lambda - T_0 (3T_0 - T_a) (T_a^2 - 8T_a T_0 + 2T_0^2) \left. \right) \exp [h(\lambda)] \\
& \left. - T_a (T_a^3 - 12T_a^2 T_0 + 36T_a T_0^2 - 24T_0^3) \text{Ei} [h(\lambda)] \right].
\end{aligned} \tag{21}$$

where

$$h(\lambda) = \frac{T_a}{T(\lambda)}. \quad (22)$$

We now enforce the condition that the complete solution must be continuous in  $t$ . First, we apply the initial value  $\lambda(0) = 0$  to both solution branches leading to the relations

$$kt^+(\lambda) = I_n^+(\lambda) - I_n^+(0), \quad (23)$$

for the upper-bound approximation and

$$kt^-(\lambda) = I_n^-(\lambda) - I_n^-(0), \quad (24)$$

for the lower-bound approximation, where  $t^\pm(\lambda)$  denotes time as a function of  $\lambda$  given by the respective “ $\pm$ ” solution. We want to enforce the condition  $t^+(\lambda^*) = t^-(\lambda^*)$  at some point  $\lambda^*$  in the interval  $[0, 1]$  and then to connect the two series using a continuous piecewise function that shifts between branches at  $\lambda^*$ . The method we apply to construct this continuous function is to shift the upper-bound approximation  $I_n^+(\lambda) - I_n^+(0)$  by a factor  $S$  such that the lower-bound approximation  $I_n^-(\lambda) - I_n^-(0)$  and the *shifted* upper-bound approximation  $I_n^+(\lambda) - I_n^+(0) + S$  are equal at  $\lambda^*$ . The shift is obtained by solving the equation

$$I_n^+(\lambda^*) - I_n^+(0) + S = I_n^-(\lambda^*) - I_n^-(0), \quad (25)$$

for  $S$ . An empirically-motivated and simple choice for  $\lambda^*$  is to take  $\lambda^* = 1/2$ , i.e., to connect the “+” and “−” approximations at the point where the reaction is half burned. With  $\lambda^* = 1/2$ , a complete solution to the Arrhenius-Semenov ODE that is continuous in  $\lambda$  is given by

$$kt = I_n(\lambda) - I_n(0) \approx \begin{cases} I_n^-(\lambda) - I_n^-(0), & \lambda < 1/2 \\ I_n^+(\lambda) - I_n^+(1/2) + I_n^-(1/2) - I_n^-(0), & \lambda \geq 1/2 \end{cases}. \quad (26)$$

Equation (26) is the primary result in this report.

It is important to note that the analytical approximation in Eq. (26) is not a smooth function and contains a derivative discontinuity at  $t(\lambda = 1/2)$ . This lack of smoothness will not be significant in situations in which  $\lambda(t)$  is the desired quantity, but it may limit the applicability of the present approach when  $\frac{d\lambda}{dt}$  is explicitly needed. We have found that, under typically physical conditions, the derivative discontinuity is small in comparison to the magnitude of the derivative itself and, as expected, the discontinuity magnitude decreases with increasing series expansion order.

### A. Approximate Solutions for $n \in \mathbb{N}$

In the limit  $n \rightarrow j + 1$  with  $j \in \mathbb{N}$  (where  $n$  is the reaction order), the functions  $I_n^{(j+)}(\lambda)$  given in Eqs. (11)-(14) are vertically asymptotic, and therefore, before applying an initial value  $\lambda(t = 0) = 0$ , Eq. (10) is divergent for  $n \in \mathbb{N}$  if  $n \leq \alpha$ . This can be seen in the general expression for these functions:

$$I_n^{(j+)}(\lambda) = \frac{(1 - \lambda)^{j+1-n}}{n - j + 1} C_n^{(j+)}, \quad (27)$$

where  $C_n^{(j+)}$  is a nonzero constant. However, when the  $j$ th order term in the full solution  $f_n^{(j+)}(\lambda)$  is constructed that includes the initial value:

$$f_n^{(j+)}(\lambda) = I_n^{(j+)}(\lambda) - I_n^{(j+)}(0), \quad (28)$$

we find that

$$\lim_{n \rightarrow j+1} f_n^{(j+)}(\lambda) = -C_n^{(j+)} \ln(1 - \lambda). \quad (29)$$

This shows that the approximate solution does not have a vertical asymptote with respect to variation of  $n$ .

### III. EXACT SOLUTIONS FOR $n \in \mathbb{N}$

In this section, we derive exact analytical solutions for the RHS of Eq. (6) for  $n \leq 3$  :  $n \in \mathbb{N}$ . In all cases the initial value is  $\lambda(t = 0) = 0$ .

#### A. $n = 0$

For  $n = 0$ , the problem reduces to evaluating the integral

$$I_0 = \int \exp \left[ \frac{T_a}{T_0 + \mathcal{Q}/C_v \lambda} \right] d\lambda. \quad (30)$$

With  $u = T_0 + \mathcal{Q}/C_v \lambda$ , the integral takes the form

$$I_0 = \frac{C_v}{\mathcal{Q}} \int \exp \left[ \frac{T_a}{u} \right] du. \quad (31)$$

Now, let  $v = u/T_a = 1/h(\lambda)$ , and

$$I_0 = \frac{C_v T_a}{\mathcal{Q}} \int \exp \left[ \frac{1}{v} \right] dv, \quad (32)$$

which is a known integral form resulting in

$$I_0 = \frac{C_v T_a}{\mathcal{Q}} \left[ v \exp \left[ \frac{1}{v} \right] - \text{Ei} \left[ \frac{1}{v} \right] \right] \quad (33)$$

where Ei is the exponential integral. After substitution for  $v$  we obtain

$$I_0 = \frac{C_v}{\mathcal{Q}} \left[ T(\lambda) \exp [h(\lambda)] - T_a \text{Ei} [h(\lambda)] \right]. \quad (34)$$

The final solution for  $n = 0$  is

$$kt = \left( \frac{C_v}{\mathcal{Q}} \right) \left[ T(\lambda) \exp [h(\lambda)] - T_a \text{Ei} [h(\lambda)] - T_0 \exp \left[ \frac{T_a}{T_0} \right] + T_a \text{Ei} \left[ \frac{T_a}{T_0} \right] \right], \quad (35)$$

with  $h(\lambda)$  defined in Eq. (22) above.

### B. $n = 1$

An exact analytical solution for the  $n = 1$  case is known (see Menikoff's derivation in Section 3 of Ref. 26). Solving  $I_1$  gives

$$kt = \text{Ei} [h(\lambda)] - \exp \left[ \frac{T_a}{T_1} \right] \text{Ei} [g(\lambda)] - \text{Ei} \left[ \frac{T_a}{T_0} \right] + \exp \left[ \frac{T_a}{T_1} \right] \text{Ei} [g(0)], \quad (36)$$

with

$$g(\lambda) = h(\lambda) - \frac{T_a}{T_1}. \quad (37)$$

### C. $n = 2$

For  $n = 2$ , the integral to evaluate in order to construct the exact solution is

$$I_2 = \int (1 - \lambda)^{-2} \exp \left[ \frac{T_a}{T_0 + \mathcal{Q}/C_v \lambda} \right] d\lambda. \quad (38)$$

Let  $u = \frac{T_0 + \mathcal{Q}/C_v \lambda}{T_a} = 1/h(\lambda)$ . The integral now takes the form

$$I_2 = \frac{C_v T_a}{\mathcal{Q}} \int \exp \left[ \frac{1}{u} \right] \left( 1 + \frac{C_v T_0}{\mathcal{Q}} - \frac{C_v T_a}{\mathcal{Q}} u \right)^{-2} du. \quad (39)$$

Now, let  $v = 1/u = h(\lambda)$ , then,

$$I_2 = -\frac{C_v T_a}{\mathcal{Q}} \int \exp [v] \left( \frac{C_v T_1}{\mathcal{Q}} v - \frac{C_v T_a}{\mathcal{Q}} \right)^{-2} dv. \quad (40)$$

Integration by parts and algebraic manipulation gives

$$I_2 = \frac{\mathcal{Q}T_a}{C_v} \left[ \frac{\exp[v]}{(T_1^2 v - T_1 T_a)} - \int \frac{\exp[v]}{(T_1^2 v - T_1 T_a)} dv \right]. \quad (41)$$

Finally, let  $w = v - T_a/T_1 = g(\lambda)$  in the integral in previous equation, leading to

$$I_2 = \frac{\mathcal{Q}T_a}{C_v} \left[ \frac{\exp[v]}{(T_1^2 v - T_1 T_a)} - \frac{1}{T_1^2} \exp\left[\frac{T_a}{T_1}\right] \int \frac{\exp[w]}{w} dw \right]. \quad (42)$$

The integral in the second term is the exponential integral:

$$I_2 = \frac{\mathcal{Q}T_a}{C_v} \left[ \frac{\exp[v]}{(T_1^2 v - T_1 T_a)} - \frac{1}{T_1^2} \exp\left[\frac{T_a}{T_1}\right] \text{Ei}[w] \right]. \quad (43)$$

After substitution for  $w$  and  $v$  we obtain

$$I_2 = \frac{T(\lambda)}{(1-\lambda)T_1} \exp[h(\lambda)] - \frac{\mathcal{Q}T_a}{C_v T_1^2} \exp\left[\frac{T_a}{T_1}\right] \text{Ei}[g(\lambda)]. \quad (44)$$

The final solution for  $n = 2$  is

$$\begin{aligned} kt = & \frac{T(\lambda)}{T_1(1-\lambda)} \exp[h(\lambda)] - \frac{\mathcal{Q}T_a}{C_v T_1^2} \exp\left[\frac{T_a}{T_1}\right] \text{Ei}[g(\lambda)] \\ & - \frac{T_0}{T_1} \exp\left[\frac{T_a}{T_0}\right] + \frac{\mathcal{Q}T_a}{C_v T_1^2} \exp\left[\frac{T_a}{T_1}\right] \text{Ei}[g(0)]. \end{aligned} \quad (45)$$

#### D. $n = 3$

For  $n = 3$ , the solution is derived using similar techniques to the cases described above.

After some algebraic manipulation, the solution can be expressed as

$$\begin{aligned} kt = & \frac{\mathcal{Q}^2 T_a}{2C_v^2 T_1^3 g(\lambda)} \left( 2 + \frac{T_a}{T_1} + \frac{T_a}{T_1 g(\lambda)} \right) \exp[h(\lambda)] - \frac{\mathcal{Q}^2 T_a (T_a + 2T_1)}{2C_v^2 T_1^4} \exp\left[\frac{T_a}{T_1}\right] \text{Ei}[g(\lambda)] \\ & - \frac{\mathcal{Q}^2 T_a}{2C_v^2 T_1^3 g(0)} \left( 2 + \frac{T_a}{T_1} + \frac{T_a}{T_1 g(0)} \right) \exp\left[\frac{T_a}{T_0}\right] + \frac{\mathcal{Q}^2 T_a (T_a + 2T_1)}{2C_v^2 T_1^4} \exp\left[\frac{T_a}{T_1}\right] \text{Ei}[g(0)]. \end{aligned} \quad (46)$$

## IV. RESULTS

Figure 1 illustrates a comparison between the solution given by the approximation in Eq. (26) and the corresponding solution generated numerically for various reaction orders. In all cases we define time  $t$  in units of the adiabatic induction time<sup>26</sup>

$$t_{\text{adb}} = \frac{C_v T_0^2}{k \mathcal{Q} T_a} \exp\left[\frac{T_a}{T_0}\right], \quad (47)$$

and, therefore, the frequency factor  $k$  does not enter into the presented results. All numerical solutions are obtained using the **Mathematica** program **NDSolve** function. The different color curves correspond to different reaction orders. The system is characterized by three temperatures:  $T_0$ ,  $T_1$ , and  $T_a$ . In this Figure, the value of these parameters are  $T_0 = 800$  K,  $T_1 = 4000$  K, and  $T_a = 6000$  K which gives rise to the three stages of ignition observed in a system with large activation temperature  $T_a \gg \frac{C_v T_0^2}{Q}$ : induction, thermal runaway, and burn-out. Over each stage of the burn process, the analytical approximation agrees with the numerical solutions. The accuracy of approximation decreases as  $\lambda$  approaches  $1/2$ . The cause of this accuracy decrease is that we expand the upper-bound solution about  $\lambda = 1$  and the lower-bound solution about  $\lambda = 0$ , and then connect those solutions at  $\lambda = 1/2$ . This means that in the complete piecewise solution,  $\lambda = 1/2$  is the point furthest from the respective expansion point for each solution branch, and therefore each series generates the greatest error at this point. In general, the accuracy of the analytical approximation increases as  $n \rightarrow 0$  and decreases as  $n$  is increased. In order to illustrate the effectiveness of present approach for relatively small expansion order, we have used  $\alpha = 3$  in all analytical calculations. For  $n \gg 1$ , the analytical approximation can also lead to accurate solutions by increasing the expansion order as illustrated later.




FIG. 1. Reaction progress variable  $\lambda$  as a function of time  $t$  (units of  $t_{adb}$ ) for varying reaction orders  $n$  shown in the legend. The system parameters are  $T_0 = 800$  K,  $T_1 = 4000$  K, and  $T_a = 6000$  K. The dashed black curves are numerical solutions and the solid curves are the results given by the analytical approximation in Eq. (26) with expansion order  $\alpha = 3$ .




FIG. 2. Reaction progress variable  $\lambda$  as a function of time  $t$  (units of  $t_{\text{adb}}$ ) for varying reaction orders  $n$  shown in the legend. The system parameters are  $T_0 = 3500$  K,  $T_1 = 4000$  K, and  $T_a = 8000$  K. The dashed black curves are numerical solutions and the solid curves are the results given by the analytical approximation in Eq. (26) with expansion order  $\alpha = 3$ .

A comparison between numerical and analytical solutions for a constant volume burn with small energy release  $\mathcal{Q} \ll \frac{C_v T_0^2}{T_a}$  is shown in Fig. 2 for various reaction orders. The system parameters in this Figure are  $T_0 = 3500$  K,  $T_1 = 4000$  K, and  $T_a = 8000$  K. As before, the numerical and analytical results are in agreement, illustrating that the presented approximation is robust with respect to system parameters.

For simplicity, we have only explicitly shown the algebra for analytical expansions up to third order. Greater solution accuracy, however, can be obtained by increasing  $\alpha$ . Shown in Fig. 3 is a comparison between analytical and numerical solutions for the same system parameters used in Fig. 1 with expansion order increased to  $\alpha = 6$ . The increased value of  $\alpha$  results in excellent agreement between solutions for all values of  $n$  shown. In fact, for small  $n$ , the analytical result is essentially indistinguishable from the numerical solution at the shown level of visual fidelity. This illustrates that the analytical method developed here approaches the exact solution in a relatively small number of expansion terms. Note that we have added a curve for  $n = 2.5$  in this plot (cf. Fig. 1) in order to illustrate that accurate solutions for relatively large reaction orders can be constructed by increasing  $\alpha$ .




FIG. 3. Progress variable as a function of time for the same system parameters as Fig. 1 with the analytical series expansion order increased to  $\alpha = 6$ . The different color curves correspond to different reaction orders shown in the legend.

## V. ACKNOWLEDGMENTS

We thank Josh Coe for insightful discussions. This work was supported by Los Alamos National Laboratory (LANL) Directed Research and Development funds (LDRD).

- 
- <sup>1</sup> T. Uzer, C. Jaffé, J. Palacián, P. Yanguas, and S. Wiggins, “The geometry of reaction dynamics,” *Nonlinearity* **15**, 957–992 (2002), doi:10.1088/0951-7715/15/4/301.
  - <sup>2</sup> E. Pollak and P. Talkner, “Reaction rate theory: What it was, where is it today, and where is it going?” *Chaos* **15**, 026116 (2005), 10.1063/1.1858782.
  - <sup>3</sup> T. Bartsch, R. Hernandez, and T. Uzer, “Transition state in a noisy environment,” *Phys. Rev. Lett.* **95**, 058301(1)–058301(4) (2005), doi:10.1103/PhysRevLett.95.058301.
  - <sup>4</sup> M. Z. Bazant, “Theory of chemical kinetics and charge transfer based on nonequilibrium thermodynamics,” *Acc. Chem. Res.* **46**, 1144–1160 (2013).
  - <sup>5</sup> G. T. Craven and R. Hernandez, “Lagrangian descriptors of thermalized transition states on time-varying energy surfaces,” *Phys. Rev. Lett.* **115**, 148301 (2015), doi:10.1103/PhysRevLett.115.148301.

- <sup>6</sup> G. T. Craven and A. Nitzan, “Electron transfer across a thermal gradient,” *Proc. Natl. Acad. Sci.* **113**, 9421–9429 (2016), doi:10.1073/pnas.1609141113.
- <sup>7</sup> D. V. Matyushov, “Activated kinetics in a nonequilibrium thermal bath,” *Proc. Natl. Acad. Sci.* **113**, 9401–9403 (2016), doi:10.1073/pnas.1610542113.
- <sup>8</sup> G. T. Craven and R. Hernandez, “Deconstructing field-induced ketene isomerization through Lagrangian descriptors,” *Phys. Chem. Chem. Phys.* **18**, 4008–4018 (2016), doi:10.1039/C5CP06624G.
- <sup>9</sup> G. T. Craven, T. Bartsch, and R. Hernandez, “Communication: Transition state trajectory stability determines barrier crossing rates in chemical reactions induced by time-dependent oscillating fields,” *J. Chem. Phys.* **141**, 041106 (2014), doi:10.1063/1.489147.
- <sup>10</sup> I. Santamaría-Holek and A. Pérez-Madrid, “Eyring equation and fluctuation–dissipation far away from equilibrium,” *J. Chem. Phys.* **153**, 244116 (2020), doi:10.1063/5.0032634.
- <sup>11</sup> C. M. Tarver, J. W. Forbes, and P. A. Urtiew, “Nonequilibrium Zeldovich-von Neumann-Doring theory and reactive flow modeling of detonation,” *Russ. J. Phys. Chem. B* **1**, 39–45 (2007).
- <sup>12</sup> R. Menikoff, “On beyond the standard model for high explosives: Challenges & obstacles to surmount,” *AIP Conference Proceedings* **1195**, 18–25 (2009), doi:10.1063/1.3295100.
- <sup>13</sup> O. Kunova, E. Nagnibeda, and I. Sharafutdinov, “Non-equilibrium reaction rates in air flows behind shock waves. State-to-state and three-temperature description,” *AIP Conference Proceedings* **1786**, 150005 (2016), doi:10.1063/1.4967646.
- <sup>14</sup> T. D. Aslam, “Shock temperature dependent rate law for plastic bonded explosives,” *J. Appl. Phys.* **123**, 145901 (2018), doi:10.1063/1.5020172.
- <sup>15</sup> R. Menikoff, *Detonation wave profile*, Tech. Rep. (Los Alamos National Laboratory, 2015).
- <sup>16</sup> R. Menikoff, *Introduction to high explosives*, Tech. Rep. (Los Alamos National Laboratory, 2019).
- <sup>17</sup> J. H. Peterson and J. D. Coe, *Beyond the Arrhenius rate law in simulating the shock-driven decomposition of polyimide*, Tech. Rep. (Los Alamos National Laboratory, 2020).
- <sup>18</sup> R. C. Huber, J. Peterson, J. D. Coe, D. M. Dattelbaum, L. L. Gibson, R. L. Gustavsen, J. M. Lang, and S. A. Sheffield, “Polysulfone shock compressed above the decomposition threshold: Velocimetry and modeling of two-wave structures,” *J. Appl. Phys.* **127**, 105902 (2020), doi:10.1063/1.5124252.

- <sup>19</sup> M. Ceriotti, C. Clementi, and O. Anatole von Lilienfeld, “Machine learning meets chemical physics,” *J. Chem. Phys.* **154**, 160401 (2021), doi:10.1063/5.0051418.
- <sup>20</sup> K. T. Butler, D. W. Davies, H. Cartwright, O. Isayev, and A. Walsh, “Machine learning for molecular and materials science,” *Nature* **559**, 547 (2018), doi:10.1038/s41586-018-0337-2.
- <sup>21</sup> P. L. Houston, A. Nandi, and J. M. Bowman, “A machine learning approach for prediction of rate constants,” *J. Phys. Chem. Lett.* **10**, 5250–5258 (2019), doi:10.1021/acs.jpcclett.9b01810.
- <sup>22</sup> E. Komp and S. Valleau, “Machine learning quantum reaction rate constants,” *J. Phys. Chem. A* **124**, 8607–8613 (2020), doi:10.1021/acs.jpca.0c05992.
- <sup>23</sup> T. L. Greaves, K. S. Schaffarczyk McHale, R. F. Burkart-Radke, J. B. Harper, and T. C. Le, “Machine learning approaches to understand and predict rate constants for organic processes in mixtures containing ionic liquids,” *Phys. Chem. Chem. Phys.* **23**, 2742–2752 (2021), doi:10.1039/D0CP04227G.
- <sup>24</sup> Y. Liu, W. Hong, and B. Cao, “Machine learning for predicting thermodynamic properties of pure fluids and their mixtures,” *Energy* **188**, 116091 (2019), doi:10.1016/j.energy.2019.116091.
- <sup>25</sup> G. T. Craven, N. Lubbers, K. Barros, and S. Tretiak, “Machine learning approaches for structural and thermodynamic properties of a Lennard-Jones fluid,” *J. Chem. Phys.* **153**, 104502 (2020), doi:10.1063/5.0017894.
- <sup>26</sup> R. Menikoff, *Arrhenius rate: Constant volume burn*, Tech. Rep. (Los Alamos National Laboratory, 2017).
- <sup>27</sup> T. Boddington, P. Gray, and G. Wake, “Criteria for thermal explosions with and without reactant consumption,” *Proc. R. Soc. Lond.* **357**, 403–422 (1977).