

# Engineering and optimization of lignin catabolic pathways in *Rhodosporidium toruloides*


Valentina E. Garcia,<sup>1,2 \*</sup> Peter B. Otopal,<sup>1,2</sup> (PeterOtopal@lbl.gov) Gina M. Geiselman,<sup>1,2</sup> Joonhoon Kim,<sup>1,3</sup> Hyun Gyu Lim,<sup>5</sup> Ramu Kakumanu,<sup>1,4</sup> Jon K. Magnuson<sup>1,3</sup> Edward Baidoo,<sup>1,4</sup> Adam M. Feist,<sup>5</sup> Jay D. Keasling,<sup>1,4</sup> Blake A. Simmons,<sup>1,4</sup> & John M. Gladden<sup>1,2</sup>

Deconstruction Division – Fungal Biotechnology  
Joint BioEnergy Institute, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

## Abstract

*Rhodosporidium toruloides* is an ideal chassis for valorizing lignocellulosic biomass, with substantial malonyl-CoA pools and the ability to co-utilize multiple carbon sources<sup>1</sup>. Here, we engineer the yeast to better utilize the lignin portion of biomass by manipulating its native *p*-Coumarate consumption pathway. We explore production of six *p*-Coumarate-derived compounds and obtain significant titers for many (6.7 g/L PCA, 4.6 g/L 4HBA, and 0.4 g/L PDC). Finally, we employ Tolerance Adaptive Laboratory Evolution (TALE) to enable robust growth of *R. toruloides* in 20 g/L *p*-Coumarate.


## Lignin Valorization in *R. toruloides*



Native *p*-Coumarate degradation in *R. toruloides* (in black) and engineered pathways (in color). Completed gene deletions are highlighted in red.

## Deleting Key Steps in Pathway

Employing CRISPR-Cas9<sup>2</sup> to delete 12623, 12923 enables conversion of *p*-Coumarate into  $6.7 \pm 2.2$  g/L PCA or  $4.6 \pm 0.1$  4HBA respectively in minimal media




Successful KOs give brown phenotypes.


Growth of  $\Delta 12923$  in Ensiled Sorghum Hydrolysate produces  $1.3 \pm 0.1$  g/L 4HBA

## Production of PDC

LigABC pathway for producing PDC from PCA integrated in Rhodo, grown with PCA




PCA hindered growth only of strains harboring LigAB. Best explained by LigAB conversion of PCA to CHMS, an aldehyde (& potentially toxic) PDC precursor



Observed peaks for PDC, CHMS on GC-MS. Follow-ups failed, due to bad storage

Integrated LigC @ 12623 to prevent PCA consumption. Stacking with LigAB



## Author Affiliations:

<sup>1</sup>DOE Joint BioEnergy Institute, USA


<sup>2</sup>Biomass Science and Conversion Technologies, Sandia National Laboratories, USA

<sup>3</sup>Chemical and Biological Processing Group, Pacific Northwest National Laboratory, USA

<sup>4</sup>Biological Systems and Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, California, USA


<sup>5</sup>Department of Bioengineering, University of California, San Diego, CA, USA

## *p*-Coumarate Adaptation



Six independent cultures (A-F) adapted over  $\sim 10^{12}$  cumulative cell divisions

Adapted strains grow much better than WT, easily tolerating 20 g/L *p*-Coumarate




Growth is robust in *p*-Coumarate; Strain "F" even grows directly in 20g/L

## Resveratrol Degradation Products

Integration of Resveratrol STS pathway resulted in dark brown color. Two unique peaks observed in HPLC spectra



*Rhodo*'s cousin, *U. maydis*, has a known Resveratrol cleavage enzyme Rco1<sup>3</sup>, with homology to two *Rhodo* genes



*Rhodo* could be producing, & subsequently cleaving, resveratrol

## Enhancing Substrate Availability

In general, *p*-Coumaroyl-CoA based pathways failed, possibly due to localization in the peroxisome

Moving first step in *p*-Coumarate out of peroxisome by deleting PTS's

12555: ...SVYEEKPR**AKL**\* ...LYTKAEGKAR  
**AKL**\*

## Conclusions & Future Work

- High titers of PCA (6.7 g/L), 4HBA (4.6 g/L) were recovered from *R. toruloides*.
- Strong signs of CHMS production, and of potential PDC production (0.4 g/L detected).
- R. toruloides* adapted to high *p*-Coumarate levels, growing directly in 20g/L. Incorporate product pathways into these evolved strains
- Resveratrol degradation products appear from STS expressing *R. toruloides*. Peroxisome localization may hinder product formation. Delete *rco1* and PTS's to fix this.

## References

1. Yaegashi, "Rhodosporidium toruloides: a new platform organism for conversion of lignocellulose into terpene biofuels and bioproducts". Biotech. Biofuels, 2017
2. Otopal, "Multiplexed CRISPR-Cas9-based genome editing of *Rhodosporidium toruloides*". mSphere, 2019
3. Brefort, "Cleavage of resveratrol in fungi: Characterization of the enzyme Rco1 from *Ustilago maydis*". Fungal Genetics and Biology, 2011

## Acknowledgments

This work conducted by the Joint BioEnergy Institute was supported by the Office of Science, Office of Biological and Environmental Research, of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231

**JBEI 5-Year Goal:** Develop fungal strains that deconstruct biomass and generate 10g/L of targeted bioproducts



U.S. DEPARTMENT OF  
**ENERGY**

Office of  
Science