

DISCLAIMER

CONF-950660-10

RECEIVED

FEB 05 1996

OSTI

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PROTON RADIOACTIVITY STUDIES AT THE FMA *

H. Penttilä^{1,2,7}, C. N. Davids¹, P. J. Woods³, J. C. Batchelder⁴, C. R. Bingham⁵, D. J. Blumenthal¹, L. T. Brown^{1,10}, B. C. Busse^{1,6}, L. F. Conticchio^{1,2}, S. J. Freeman⁸, M. Freer⁹, D. J. Henderson¹, R. D. Page^{3,12}, A. V. Ramayya¹⁰, K. S. Toth¹¹, W.B. Walters², A. H. Wuosmaa¹, B. E. Zimmerman⁵

1 Argonne National Laboratory, Argonne, IL, USA

2 University of Maryland, College Park, MD, USA

3 University of Edinburgh, Edinburgh, UK

4 Louisiana State University, Baton Rouge, LA, USA

5 University of Tennessee, Knoxville, TN, USA

6 Oregon State University, Corvallis, OR, USA

7 Present address: University of Jyväskylä, PL 35, 40351 Jyväskylä, Finland

8 Manchester University, UK

9 University of Birmingham, UK

10 Vanderbilt University, Nashville, TN, USA

11 Oak Ridge National Laboratory, Oak Ridge, TN, USA

12 Present address: University of Liverpool, UK

* Work supported by U.S. Department of Energy, Nuclear Physics Division, contract W-31-109-ENG-38

A double sided silicon strip (DSSD) setup is installed at the Argonne National Laboratory recoil separator FMA. Ground state proton emitters ^{146,147}Tm, ¹⁶⁰Re, ¹⁵⁶Ta, ^{150,151}Lu were produced in a series of test experiments. Improved T_{1/2} values were obtained for ¹⁴⁷Tm, ¹⁶⁰Re, and ¹⁵⁶Ta. Improved alpha T_{1/2} value of (13.2 ± 1.1) ms was obtained for ¹⁶¹Re. In a search of new proton radioactivities a (5 ± 2) % proton decay branch with E_p = (1345 ± 13) keV for ¹⁶¹Re was found.

A direct proton decay is a decay mode where a non-excited nucleus emits a proton [1]. The first proton emitter ¹⁵¹Lu was found in 1981 [1], but only 12 proton radioactive nuclei were known prior to the present experiments. A DSSD [2] setup was installed at the the Fragment Mass Analyzer (FMA) [3] to provide an efficient detection system of proton activities. The operation of both the FMA and the DSSD are described more in detail in the references [2,3] and in the contribution of C. N. Davids elsewhere in these Proceedings.

The aim of the experiments was to reinvestigate the proton emitters $^{146,147}\text{Tm}$ [4], ^{160}Re and ^{156}Ta [5], and to search for a proton decay branch in ^{161}Re . The studied decays provided a test case for experimental setup before advancing to searches for new proton emitters like ^{166}Ir and ^{170}Au . The reactions that were used, the observed decay energies and the half lives are given in Table I. In general, the present experiment resulted improved statistics compared with the previous experiments. A previously unknown 50 % β -decay branch was found for ^{156}Ta . To search for the proton decay branch of ^{161}Re , the decays following the implantation of recoils with mass $A = 161$ were gated by the ^{161}Re proton decay daughter ^{160}W and the granddaughter ^{156}Hf α -decays in subsequent generations. In the resulting energy spectrum a peak appears at 1345 ± 13 keV. The proton decay half-life $T_{1/2} = 15^{+8}_{-4}$ ms agrees with the α -decay half-life of ^{161}Re . The proton decay branch 5 ± 2 % results in a partial proton decay half-life of 260^{+180}_{-80} ms. This corresponds to emission of a proton from $h_{11/2}$ orbital.

Table I

Nuclide	E_p [keV]	E_α [keV]	$T_{1/2}$	Production/Comments
^{147}Tm	1051 *)		559 ± 26 ms	^{58}Ni (261MeV) + ^{92}Mo , p2n
^{147m}Tm	1115 ± 8 *)		390^{+93}_{-71} μs	^{58}Ni (261MeV) + ^{92}Mo , p2n
^{146m}Tm	1118 ± 7 *) 1120 ± 4 †)		206 ± 25 ms	^{58}Ni (290MeV) + ^{92}Mo , p3n ^{58}Ni (279MeV) + ^{92}Mo , p3n
^{146}Tm	1189 ± 10 *)		62^{+19}_{-14} ms	^{58}Ni (290MeV) + ^{92}Mo , p3n
^{160}Re	1258 ± 4 †)	6545 ± 14	637^{+90}_{-70} μs 739^{+220}_{-138} μs	^{58}Ni (300MeV) + ^{106}Cd , p3n
^{156}Ta	1011 ± 7 †)		105^{+58}_{-28} ms	^{160}Re α decay
^{161}Re	1345 ± 13 †)	6244 ± 5	15^{+8}_{-4} ms 13.2 ± 1.1 ms	^{58}Ni (300MeV) + ^{106}Cd , p2n

†) Calibrated with respect to ^{147}Tm (1051.1 ± 2.6) keV and ^{151}Lu (1232.9 ± 2.0) keV [1]

*) Calibrated with respect to ^{147}Tm (1051.1 ± 2.6) keV [1]

References:

- [1] S. Hofmann, in *Particle Emission From Nuclei*, ed. D. N. Poenaru and M. Ivascu (CRC Press, Boca Raton, FL, 1989), Vol 2, Chap. 2
- [2] P. J. Sellin *et al*, Nucl. Inst. Meth., **A311** (1992) 217
- [3] C. N. Davids *et al*, Nucl. Inst. Meth., **B70** (1992) 358
- [4] K. Livingstone *et al*, Phys. Lett, **B 312** (1993) 46, and references therein.
- [5] R. D. Page *et al*, Phys. Rev. Lett., **68** (1992) 1287