

First Measurement of the Neutron-Emission Probability with a Surrogate Reaction in Inverse Kinematics at a Heavy-Ion Storage Ring

Sguazzin, M; Jurado, B; Pibernat, J; Swartz, J. A.; Grieser, M; Glorius, J.; Litvinov, Y. A.; Adamczewski-Musch, J; Alfaurt, P; Ascher, P; Audouin, L.; Berthelot, C; Blank, B; Blaum, K.; Brueckner, B; Dellmann, S. F.; Dillmann, I.; Domingo-Pardo, C.; Dupuis, Marc; Erbacher, P; Flayol, M; et al.

Provided by the author(s) and the Los Alamos National Laboratory (1930-01-01).

To be published in: Physical Review Letters

DOI to publisher's version: 10.1103/PhysRevLett.134.072501

Permalink to record:

<https://permalink.lanl.gov/object/view?what=info:lanl-repo/lareport/LA-UR-24-20011>



Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

First Measurement of the Neutron-Emission Probability with a Surrogate Reaction in Inverse Kinematics at a Heavy-Ion Storage Ring

M. Sguazzin^{1,†}, B. Jurado^{1,*}, J. Pibernat,¹ J. A. Swartz^{1,‡}, M. Grieser^{1,§}, J. Glorius^{1,§}, Yu. A. Litvinov^{1,§}, J. Adamczewski-Musch^{1,§}, P. Alfaut,¹ P. Ascher^{1,§}, L. Audouin^{1,§}, C. Berthelot^{1,§}, B. Blank^{1,§}, K. Blaum^{1,§}, B. Brückner,⁵, S. Dellmann^{1,§}, I. Dillmann^{1,§}, C. Domingo-Pardo^{1,§}, M. Dupuis^{1,§,10}, P. Erbacher,⁵, M. Flayol,¹ O. Forstner^{1,§}, D. Freire-Fernández^{1,11}, M. Gerbaux^{1,§}, J. Giovinazzo^{1,§}, S. Grévy^{1,§}, C. J. Griffin^{1,§}, A. Gumberidze^{1,§}, S. Heil⁵, A. Heinz^{1,§}, R. Hess³, D. Kurtulgil⁵, N. Kurz^{1,§}, G. Leckenby^{1,§,13}, S. Litvinov^{1,§}, B. Lorentz^{1,§}, V. Méot^{1,§,10}, J. Michaud^{1,§,14}, S. Pérard,¹ N. Petridis^{1,§}, U. Popp,³, D. Ramos^{1,§,14}, R. Reifarth^{1,§}, M. Roche,¹ M. S. Sanjari^{1,§,16}, R. S. Sidhu^{1,§,17,3,2,§}, U. Spillmann^{1,§}, M. Steck,³, Th. Stöhlker,^{3,18,19}, B. Thomas,¹ L. Thulliez,²⁰, M. Versteegen,¹, and B. Włoch^{1,§}

¹Université de Bordeaux, CNRS, LP2I Bordeaux, 33170 Gradignan, France

²Max-Planck-Institut für Kernphysik, 69117 Heidelberg, Germany

³GSI Helmholtzzentrum für Schwerionenforschung, 64291 Darmstadt, Germany

⁴Université Paris-Saclay, CNRS, IJCLab, 91405 Orsay, France

⁵Goethe University of Frankfurt, 60438 Frankfurt, Germany

⁶TRIUMF, Vancouver, British Columbia, V6T 2A3, Canada

⁷Department of Physics and Astronomy, University of Victoria, Victoria, British Columbia, V8P 5C2, Canada

⁸IFIC, CSIC-Universidad de Valencia, 46980 Valencia, Spain

⁹CEA, DAM, DIF, 91297 Arpajon, France

¹⁰Université Paris-Saclay, CEA, LMCE, 91680 Bruyères-Le-Châtel, France

¹¹Ruprecht-Karls-Universität Heidelberg, 69117 Heidelberg, Germany

¹²Chalmers University of Technology, 41296 Gothenburg, Sweden

¹³Department of Physics and Astronomy, University of British Columbia, Vancouver, British Columbia, V6T 1Z1, Canada

¹⁴GANIL, CRNS/IN2P3-CEA/DRF, 14000 Caen, France

¹⁵Los Alamos National Laboratory, Los Alamos, New Mexico 87544, USA

¹⁶Aachen University of Applied Sciences, Aachen, Germany

¹⁷School of Physics and Astronomy, University of Edinburgh, EH9 3FD Edinburgh, United Kingdom

¹⁸Helmholtz Institute Jena, Fröbelstieg 3, 07743 Jena, Germany

¹⁹Institut für Optik und Quantenelektronik, Friedrich-Schiller-Universität Jena, Max-Wien-Platz 1, 07743 Jena, Germany

²⁰IRFU, CEA, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

(Received 22 December 2023; revised 24 July 2024; accepted 18 November 2024; published 18 February 2025)

Neutron-induced reaction cross sections of short-lived nuclei are imperative to understand the origin of heavy elements in stellar nucleosynthesis and for societal applications, but their measurement is extremely complicated due to the radioactivity of the targets involved. One way of overcoming this issue is to combine surrogate reactions with the unique possibilities offered by heavy-ion storage rings. In this work, we describe the first surrogate-reaction experiment in inverse kinematics, which we successfully conducted at the Experimental Storage Ring (ESR) of the GSI/FAIR facility, using the $^{208}\text{Pb}(p, p')$ reaction as a surrogate for neutron capture on ^{207}Pb . Thanks to the outstanding detection efficiencies possible at the ESR, we were able to measure for the first time the neutron-emission probability as a function of the excitation energy of ^{208}Pb . We have used this probability to select different descriptions of the γ -ray strength function and nuclear level density, and provide reliable results for the neutron-induced radiative capture cross section of ^{207}Pb at energies for which no experimental data exist.

DOI: 10.1103/PhysRevLett.134.072501

*Contact author: jurado@cenbg.in2p3.fr

†Present address: IJCLab, 91405 Orsay, France.

‡Present address: FRIB, MSU, East Lansing, Michigan 48824, USA.

§Present address: School of Mathematics and Physics, University of Surrey, Guildford, GU2 7XH, United Kingdom.

Knowledge of neutron-induced reaction cross sections of short-lived nuclei is pivotal to our understanding of the synthesis of elements via the astrophysical slow (*s*) and rapid (*r*) neutron capture processes, about which there are still many uncertainties and open questions [1]. It is also of interest for applications such as nuclear waste management and innovative fuel cycles [2]. In traditional experiments, the direct measurement of neutron-induced cross sections of short-lived nuclei is very challenging because of the difficulties to produce and handle radioactive targets. Performing the same reaction in inverse kinematics, with the heavy, radioactive nucleus impinging upon a target of neutrons, is not possible either, since free neutron targets are currently not available. For these reasons, when the target nuclei are highly radioactive, experimental data are scarce and most of the neutron-induced reaction cross sections rely on theoretical model predictions. However, these predictions often have large uncertainties due to difficulties in describing the deexcitation process of the nucleus formed after the capture of the neutron. This process is ruled by fundamental properties (γ -ray strength functions, nuclear level densities, fission barriers, etc.) for which the existing nuclear models give very different predictions. This can lead to discrepancies between the calculated cross sections as large as 2 orders of magnitude or more [3,4].

Indirect methods have been developed to infer neutron-induced reaction cross sections of short-lived nuclei [5–9]. Here we use the surrogate reaction method [7]. In this method, the excited nucleus produced in the neutron-induced reaction of interest is formed through an alternative and experimentally feasible binary reaction, typically an inelastic scattering or a transfer reaction. The measurement of the probabilities of the different decay channels of the excited nucleus (γ -ray emission, neutron emission, fission, etc.) as a function of its excitation energy provides the information which is required to constrain the models of the above-mentioned nuclear properties. This significantly improves the predictions of the cross sections of the neutron-induced reactions of interest. To date, the surrogate-reaction method has been used and successfully benchmarked in direct kinematics, see, e.g., Refs. [7,10–12].

The probability P_χ that a nucleus with excitation energy E^* formed with a surrogate reaction $X(a, b)$ decays via channel χ is given by the expression

$$P_\chi(E^*) = \frac{N_{c,\chi}(E^*)}{N_s(E^*)\epsilon_\chi(E^*)}, \quad (1)$$

where N_s is the number of light ejectiles b measured, the so-called single events. $N_{c,\chi}$ is the number of products of decay channel χ measured in coincidence with the ejectiles b and ϵ_χ is the efficiency for detecting the products of decay χ for the reactions in which the outgoing ejectile b is detected. The excitation energy E^* is obtained by measuring the kinetic energies of the projectile beam and of the ejectile b , and the angle θ between them.

Surrogate-reaction experiments in direct kinematics (where the light nucleus a is the projectile and the heavy nucleus X is at rest) have significant limitations. When the nuclei of interest are far from stability, the targets required for the surrogate reaction are also unavailable. Additionally, competing reactions in target contaminants (such as oxygen) and backings produce a large background, which is very complicated or even impossible to remove [13]. Furthermore, the heavy products of the decay of the excited nucleus are stopped in the target and cannot be detected. Therefore, the measurement of γ - and neutron-emission probabilities requires detecting the emitted γ rays and neutrons. However, the γ -ray-cascade detection efficiencies in surrogate-reaction experiments are limited to about 20% [14]. The measurement of the neutron-emission probability is extremely challenging and to our knowledge has never been accomplished.

Some of the latter limitations can be solved by using the surrogate-reaction method in inverse kinematics, which enables the study of short-lived nuclei by using a radioactive ion beam and the detection of the heavy, beamlike residues produced after the emission of γ rays and neutrons. However, the decay probabilities change very rapidly with excitation energy at the neutron-emission and at the fission thresholds; see, e.g., Ref. [12]. The excitation-energy resolution required to scan this rapid evolution is a few 100 keV (FWHM), which is quite difficult to achieve for heavy nuclei in inverse kinematics, due to so-far unresolved target issues. Indeed, the required large target density and thickness result in significant energy loss and straggling effects that translate into a large uncertainty in the energies of the projectile and the targetlike residue, and in the emission angle θ at the interaction point. In addition, the presence of target windows and impurities induces background.

Here we address these target issues by investigating for the first time surrogate reactions at a heavy-ion storage ring [15]. A key component of storage rings is the electron cooler, which significantly reduces the size, angular divergence and energy spread of the revolving ion beam. If a gas target is present in the ring, the electron cooler compensates for the energy loss and for the energy and angular straggling of the beam taking place during each passage of the beam through the target. As a result, the ion beam always reaches the target with the same energy and the same outstanding quality, making energy loss and straggling effects in the target negligible. Additionally, the frequent passing of the target zone (about a million times per second at a few tens of MeV/nucleon) makes possible the use of pure gas targets with ultralow density ($\approx 10^{13}$ atoms/cm²) and no windows are necessary. The very low gas target density makes the probability of two consecutive reactions occurring in the target, a nuclear reaction followed by an atomic reaction and vice versa, extremely low ($\approx 10^{-20}$). The beamlike residues resulting from the nuclear reaction will therefore possess the same charge state as the beam.

Heavy-ion storage rings have to be operated in ultrahigh vacuum (UHV) conditions (10^{-10} to 10^{-11} mbar), which pose severe constraints on in-ring detection systems. UHV-compatible silicon detectors have only started to be used for a few years in pioneering in-ring nuclear reaction experiments [16–18] at the Experimental Storage Ring (ESR) [19] and the CRYRING storage ring [20] of the GSI/FAIR facility.

We have conducted the first surrogate-reaction experiment at the ESR with the aim to use the $^{208}\text{Pb}(p, p')$ surrogate reaction to assess theoretical models and provide predictions for the neutron-induced radiative capture cross section (n, γ) of ^{207}Pb at neutron energies above 800 keV, where no experimental data are available. These data are important for the design of lead-cooled fast reactors [21]. In our experiment, $^{208}\text{Pb}^{82+}$ projectiles at 30.77 MeV/nucleon were excited by inelastic scattering reactions with a gas-jet target of hydrogen. We had on average 5×10^7 cooled and decelerated, bare $^{208}\text{Pb}^{82+}$ ions per measurement cycle, revolving at a frequency of 0.695 MHz. The average target thickness was 6×10^{13} atoms/cm². We measured the inelastically scattered protons with a Si $\Delta E-E$ telescope and the beamlike residues produced after the deexcitation of $^{208}\text{Pb}^*$ via γ -ray and neutron emission with a position-sensitive Si-strip detector placed behind the dipole magnet downstream from the target (denoted beamlike residue detector in Fig. 1). The dipole separated the unreacted beam, the $^{208}\text{Pb}^{82+}$ residues produced after γ -ray emission and the $^{207}\text{Pb}^{82+}$ residues produced after neutron emission, see Fig. 1.

To prevent detector components from degrading the UHV of the ring, the telescope and the beamlike residue detector were housed in pockets behind 25 μm thick stainless-steel windows through which the scattered protons and the heavy beam residues could pass. The telescope was placed at 60° with respect to the beam axis, at a distance of 10.13 cm from the target. The ΔE detector of the telescope consisted of a 530 μm -thick double-sided silicon-strip detector (DSSD) of $20 \times 20 \text{ mm}^2$ with 16 vertical and 16 horizontal strips, which enabled the measurement of the angle θ within the angular range from 54.8 to 64.6° . The angular resolution was estimated to be 0.2° (rms), assuming isotropic emission of the target residues from the center of the target. The ΔE detector was followed by a stack of six single area Si detectors for full energy measurements. Each of the latter E detectors had an active area of $20 \times 20 \text{ mm}^2$ and a thickness of 1.51 mm. In inverse kinematics it is possible to have two kinematic solutions leading to two groups of ejectiles having different kinetic energies, but the same angle θ [22]. In our experiment, scattered protons from the first kinematic solution with kinetic energies above 9.2 MeV passed through the ΔE detector, while scattered protons from the second kinematic solution with kinetic energies between 2.5 and 9.2 MeV were stopped in the ΔE detector, see [22]. The beamlike residue detector was a DSSD with a thickness of




FIG. 1. The lower part shows a schematic view of the ESR. The upper part shows the portion of the ring where our detectors have been installed. The trajectories of the scattered protons, the beam, the $^{208}\text{Pb}^{82+}$ residues produced after γ emission and the $^{207}\text{Pb}^{82+}$ residues formed after neutron emission are represented by the solid pink, black, blue, and green arrows, respectively.

500 μm , an active area of $122 \times 40 \text{ mm}^2$, 122 vertical strips, and 40 horizontal strips. It was positioned 15.0 ± 0.1 mm from the beam axis. With this distance we ensured that the rate of elastic scattered beam ions over the whole detector was well within the radiation-damage tolerance range of the detector, which remained operational throughout the experiment.

Figure 2 shows the position spectrum of beamlike residues detected in coincidence with scattered protons detected in the telescope. In panel (a) we see the heavy residues measured in coincidence with protons from the first kinematic solution. We can clearly distinguish two peaks; the left peak contains the $^{208}\text{Pb}^{82+}$ nuclei formed after γ emission and the right peak the $^{207}\text{Pb}^{82+}$ nuclei produced after neutron emission. In panel (b) are shown the heavy residues detected in coincidence with protons from the second kinematic solution for $E^* = 6.5\text{--}9.1$ MeV and $\theta = 56.1\text{--}60.40^\circ$. In this case, the beamlike residues have larger kinetic energies and their trajectories after the dipole magnet are closer to the beam axis. The $^{208}\text{Pb}^{82+}$ residues formed after γ emission cannot be detected, but all the trajectories of the $^{207}\text{Pb}^{82+}$ residues formed after neutron emission impinge on the beamlike detector, leading to the observed peak in the position spectrum. We emphasize that in this experiment the efficiency ϵ_n for the neutron emission channel is 100% [22]. The largest loss of efficiency comes from electron capture of the $^{207}\text{Pb}^{82+}$ residues in the residual gas between the target and the beamlike detector. The probability for this event has been calculated to account only to $\approx 10^{-10}$, so it can be neglected.




FIG. 2. Position of beamlike residues measured in coincidence with detected scattered protons from the first (a) and the second kinematic solution (b), see text for details.

In this work, we only consider the results obtained with the second kinematic solution, the results of the first kinematic solution are discussed in [22]. We obtained the singles spectrum $N_s(E^*)$ by representing the number of detected protons as a function of the E^* of ^{208}Pb . The coincidence spectrum $N_{c,n}(E^*)$ was inferred by representing the number of protons detected in coincidence with the beamlike residues located within the peak of Fig. 2(b). The bin size of these two histograms was 200 keV. By computing the ratio of $N_{c,n}(E^*)$ over $N_s(E^*)$ and using $\epsilon_n = 1$ [see Eq. (1)], we were able to measure for the first time the neutron-emission probability $P_n(E^*)$, as illustrated in Fig. 3. The displayed error bars include the covariance between $N_{c,n}(E^*)$ and $N_s(E^*)$ [22]. Thanks to the 100% detection efficiency for the heavy residues, it has been possible to achieve relative uncertainties of less than 6%, despite the small total number of 1581 single events measured. The experimental data show an onset of P_n below the neutron separation energy S_n . As discussed below, this is due to the excitation energy resolution ΔE^* , which is ≈ 240 keV (rms). We estimated ΔE^* with a simulation, which was benchmarked with the well-separated ground-state peak of ^{208}Pb at $E^* = 0$ MeV, see [22]. In this experiment, ΔE^* is dominated by the uncertainty in the proton scattering angle θ induced by the target radius of 2.5 mm.

To compare our results with theory, we have calculated $P_n(E^*)$ with the statistical model using the expression:

$$P_n(E^*) = \sum_{J^\pi} F(E^*, J^\pi) G_n(E^*, J^\pi), \quad (2)$$

where $F(E^*, J^\pi)$ is the probability to form the excited nucleus in a state of spin J and parity π at an excitation energy E^* by the $^{208}\text{Pb}(p, p')$ reaction, and $G_n(E^*, J^\pi)$ is the probability that the nucleus decays from that state via neutron emission. The J^π distributions given by F were calculated with the microscopic description developed in [23,24]. The theoretical formalism and the results for $F(J^\pi)$ at $E^* = 8$ and 9 MeV are presented in the Supplemental Material [25]. To determine G_n we used the statistical Hauser-Feshbach model of TALYS 1.96 [39]. Among all the quantities needed to describe the deexcitation of ^{208}Pb , the γ -ray strength function (GSF) and the nuclear level density (NLD) are the most uncertain ones. We considered different models for these two quantities with adjusted parameters for ^{208}Pb , which we obtained from literature. For the GSF, we utilized three models: the model of Kopecky and Uhl [40], the simple modified Lorentzian model (SMLO) [41] and the results of Hartree-Fock-Bogolyubov (HFB) and quasiparticle random phase approximation (QRPA) calculations based on the Gogny D1M nuclear interaction [42], which we will denote as D1M + QRPA. Regarding the NLD, we employed six distinct descriptions, three of these were based on the constant-temperature (CT) model [43] with different adjusted parameters, they are denoted CT1, CT2, and CT3. One description was based on the back-shifted Fermi-gas (BSFG) model [44]. The two others were the microscopic NLDs by Goriely *et al.* [45] and Hilaire *et al.* [46]. Further details on the models and used parameters can be found in [25].

We combined the three GSF descriptions with the six NLD models leading to 18 different TALYS calculations. We expect to observe significant differences between the calculations and our data at S_n due to the excitation energy resolution ΔE^* . To account for ΔE^* we convoluted the calculations with a Gaussian function with a standard deviation of 240 keV. We have evaluated the deviations between the calculations and our data by computing the reduced χ^2 before and after the convolution, see Table II in [25]. The deviations decrease drastically after the convolution for all the calculations. The calculations that use the NLD CT3 and the calculation utilising the SMLO and the BSFG models have a reduced χ^2 exceeding 1.63 and can be excluded by our data with a confidence level of 93%. These calculations are also excluded by the data from the first kinematic solution, see [22]. We obtained the best agreement (lowest residuals and reduced χ^2 , see [25]) with the convoluted calculation using the D1M+QRPA GSF model and the NLD by Goriely *et al.* The latter calculation is compared with our data before and after convolution in




FIG. 3. Neutron-emission probability as a function of the excitation energy E^* of ^{208}Pb measured for the $^{208}\text{Pb}(p, p')$ reaction in comparison with TALYS calculations. The arrows indicate the E^* at which the three first excited states of ^{207}Pb become accessible. The spin and parity of the states are also given. The neutron separation energy S_n of ^{208}Pb at $E^* = 7.37$ MeV is indicated by the vertical dotted line, see text for details.

Fig. 3. As shown by the red dashed-dotted line, the convoluted result exhibits a significantly improved agreement with the experimental data below S_n . The calculation obtained with the SMLO and the CT3 models yields the largest reduced χ^2 . Between S_n and 8.5 MeV, this calculation is systematically below our data, the best TALYS calculation and the blue shaded area, which includes all the calculations except those that are excluded by our data, see Fig. 3. The calculations show an increase at $E^* \approx 8$ and 9 MeV. These increases occur when the E^* of the ^{207}Pb residue formed after neutron emission is high enough to populate the 1st and 3rd excited states of ^{207}Pb , see the arrows in Fig. 3. The population of the 3rd excited state is particularly favoured because its spin $J = 13/2$ is closer to the spins populated in the $^{208}\text{Pb}(p, p')$ reaction (average spin $\bar{J} \approx 5.4$, see [25]).

We have used the GSF and NLD models that are not excluded by our data to calculate the $^{207}\text{Pb}(n, \gamma)$ cross section above neutron energies of 800 keV, see blue line and shaded area in Fig. 4. Above 800 keV, several evaluations (i.e., databases of recommended values for the cross sections [47]) based on the Hauser-Feshbach formalism are available to which we can compare our results. We expect that the calculation obtained with the SMLO and the CT3 models will result in a larger cross section, as this calculation leads to lower values of $P_n(E^*)$




FIG. 4. $^{207}\text{Pb}(n, \gamma)$ cross section as a function of neutron energy E_n . The results of our TALYS calculations are compared with the JENDL-5.0 [49], TENDL-2021 [50], CENDL-3.2 [48], and ENDF/B-VIII.0 [51] evaluations, see text for details.

and thus higher values of the γ -emission probability, since in the covered E^* range γ and neutron emission are the only open decay channels. The green dashed line in Fig. 4 shows that this is indeed the case, this TALYS calculation is well above all the other TALYS calculations. This demonstrates the strong connection between $P_n(E^*)$ and the (n, γ) cross section, and the usefulness of employing the $P_n(E^*)$ from a surrogate reaction for constraining predictions for radiative capture cross sections. As shown in Fig. 4, our calculations encompass all the evaluations except CENDL-3.2 [48], which shows a very different shape and is above our results.

In conclusion, we have measured for the first time the neutron emission probability as a function of the excitation energy, $P_n(E^*)$, of ^{208}Pb . Our measurement benefited from the unrivaled advantages of the ESR heavy-ion storage ring, which allowed us to detect the beamlike residues formed after neutron emission with an efficiency of 100%. We employed our results for $P_n(E^*)$ to select various combinations of models for the γ -ray strength function and the nuclear level density of ^{208}Pb available in the literature. The selected models were used to infer the $^{207}\text{Pb}(n, \gamma)$ cross section at neutron energies for which no experimental data are available. This demonstrates the advantage of using the $P_n(E^*)$ obtained through surrogate-reaction experiments to constrain predictions for (n, γ) cross sections. Our results are in good agreement with the JENDL-5.0, TENDL-2021, and ENDF/B-VIII.0 evaluations, but disagree with the CENDL-3.2 evaluation. In the future, we will complete our setup with fission detectors to measure also the fission probabilities, increase the solid angle of the telescope and use a target with a smaller radius, which will allow us to improve the excitation energy resolution. With these improvements we will be able to conduct next-generation experiments with radioactive stored beams, where we will measure simultaneously and with high precision the probabilities for all the deexcitation channels (fission, γ ,

neutron, and two-neutron emission) of many short-lived nuclei for which the neutron-induced cross sections are considered impossible to measure.

Acknowledgments—This work is supported by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (ERC-Advanced grant NECTAR, Grant Agreement No. 884715). The results presented here are based on the experiment E146, which was performed at the ESR storage ring of the GSI Helmholtzzentrum fuer Schwerionenforschung, Darmstadt (Germany) in the context of FAIR Phase-0. We thank the Prime 80 program from the CNRS for funding the Ph.D. thesis of M. S. and the GSI/IN2P3 Collaboration 19-80. J. G., Yu. A. L., R. R., and Th. S. acknowledge support by the State of Hesse within the Cluster Project ELEMENTS (Project ID 500/10.006). A. H. is grateful for funding from the Knut and Alice Wallenberg Foundation under KAW 2020.0076.

- [20] C. G. Bruno *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **1048**, 168007 (2023).
- [21] C. Domingo-Pardo *et al.*, *Phys. Rev. C* **74**, 055802 (2006).
- [22] M. Sguazzin *et al.*, companion paper, *Phys. Rev. C* **111**, 024614 (2025).
- [23] M. Dupuis, S. Karataglidis, E. Bauge, J. P. Delaroche, and D. Gogny, *Phys. Rev. C* **73**, 014605 (2006).
- [24] M. Dupuis, G. Haouat, J. P. Delaroche, E. Bauge, and J. Lachkar, *Phys. Rev. C* **100**, 044607 (2019).
- [25] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.134.072501> for more details on the calculations of the spin-parity distributions, on the γ -ray strength functions and level densities used, and on the comparison of the calculations and the experimental data, which includes Refs. [26–38].
- [26] J. Berger, M. Girod, and D. Gogny, *Nucl. Phys.* **A502**, 85 (1989).
- [27] M. Dupuis *et al.* (to be published).
- [28] M. Dupuis, PhD thesis, University of Bordeaux, 2006, <https://theses.hal.science/tel-00412169>.
- [29] M. Dupuis, *Eur. Phys. J. A* **53**, 111 (2017).
- [30] M. Kerveno *et al.*, *Phys. Rev. C* **104**, 044605 (2021).
- [31] E. Party, PhD thesis, University of Strasbourg, 2019, <https://theses.hal.science/tel-02496853>.
- [32] G. Henning *et al.*, *Phys. Rev. C* (to be published).
- [33] A. Koning and J. Delaroche, *Nucl. Phys.* **A713**, 231 (2003).
- [34] B. Morillon and P. Romain, *Phys. Rev. C* **74**, 014601 (2006).
- [35] B. Morillon and P. Romain, *Phys. Rev. C* **76**, 044601 (2007).
- [36] S. Bassauer, P. von Neumann-Cosel, and A. Tamii, *Phys. Rev. C* **94**, 054313 (2016).
- [37] A. J. Koning *et al.*, *Nucl. Phys.* **A810**, 13 (2008).
- [38] O. Iwamoto, *J. Nucl. Sci. Technol.* **44**, 687 (2007).
- [39] A. Koning, S. Hilaire, and S. Goriely, *Eur. Phys. J. A* **59**, 131 (2023).
- [40] J. Kopecky and M. Uhl, *Phys. Rev. C* **41**, 1941 (1990).
- [41] S. Goriely and V. Plujko, *Phys. Rev. C* **99**, 014303 (2019).
- [42] S. Goriely, S. Hilaire, S. Péru, and K. Sieja, *Phys. Rev. C* **98**, 014327 (2018).
- [43] A. Gilbert and A. Cameron, *Can. J. Phys.* **43**, 1446 (1965).
- [44] W. Dilg, W. Schantl, H. Vonach, and M. Uhl, *Nucl. Phys. A217*, 269 (1973).
- [45] S. Goriely, S. Hilaire, and A. J. Koning, *Phys. Rev. C* **78**, 064307 (2008).
- [46] S. Hilaire, M. Girod, S. Goriely, and A. J. Koning, *Phys. Rev. C* **86**, 064317 (2012).
- [47] L. A. Bernstein, D. A. Brown, A. J. Koning, B. T. Rearden, C. E. Romano, A. A. Sonzogni, A. S. Volden, and W. Younes, *Annu. Rev. Nucl. Part. Sci.* **69**, 109 (2019).
- [48] Z. Ge *et al.*, *EPJ Web Conf.* **239**, 09001 (2020).
- [49] O. Iwamoto *et al.*, *J. Nucl. Sci. Technol.* **60**, 1 (2023).
- [50] A. Koning, D. Rochman, J. Sublet, N. Dzysiuk, M. Fleming, and S. van der Marck, *Nucl. Data Sheets* **155**, 1 (2019).
- [51] D. A. Brown *et al.*, *Nucl. Data Sheets* **148**, 1 (2018).