

CONF-960888--1

UCRL-JC-122871  
PREPRINT

# A Sensitivity Study of the Mesoscale Characteristics of Squall-Line Systems to Environmental Conditions: Implication of Anvil Cirrus Parameterization


Hung-Neng S. Chin

RECEIVED  
JUN 27 1996  
OSTI

This paper was prepared for submittal to the  
*12th International Conference on Clouds and Precipitation*  
*Zurich, Switzerland*  
*August 19-23, 1996*

# MASTER

April 1996



This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

#### DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

# A Sensitivity Study of the Mesoscale Characteristics of Squall-Line Systems to Environmental Conditions: Implication of Anvil Cirrus Parameterization

Hung-Neng S. Chin

Atmospheric Sciences Division  
Lawrence Livermore National Laboratory

## 1. INTRODUCTION

Cloud-radiation feedback has been identified as the single most important effect limiting general circulation models (GCMs) to further progress in climate change research, and regarded as major uncertainties in estimating the impact of increasing concentrations of greenhouse gases on climate simulations. Therefore, it is crucially important to further understand the physical processes involved in order to improve the representation of cloud processes in GCMs. To this end, a cloud resolving model with enhanced model physics (Chin 1994; Chin *et al.* 1995) was used to study the impact of microphysics, long- and shortwave radiation on midlatitude and tropical mesoscale convective systems (MCSs).

These earlier studies indicated that a strong coupling exists in the MCSs between deep convection and its related anvil cloud through the interaction among dynamical, thermodynamical and radiative processes. They also showed that the tilting structure of MCSs (sub-GCM-grid feature) has an important contribution to the water budget of anvil clouds, particularly the tropical anvil due to the jetlike wind profile. However, most earlier GCMs did not include a direct and physically consistent representation of this coupling. To this end, Randall *et al.* (1989) suggested a more realistic anvil parameterization by adding prognostic cloud water (or ice) variables to account for the formation of anvil clouds from cumulus detrainment. In addition to this effort, our recent studies further suggest the need to parameterize the tilting structure of MCSs in GCMs.

The objective of this work is to parameterize the large-scale effects of an important sub-GCM-grid-scale process associated with the tilting structure of MCSs. To this end, our primary interest focuses on MCSs in an environment with substantial wind shear, such as squall-line systems, since they have longer lifetimes and wider coverage to affect the earth-atmosphere radiation budget and climate. Using varied convective available potential energy (CAPE), wind shear intensity, shear depth, and shear profile pattern (i.e., jetlike or non-jetlike wind profile) over a wide range of bulk Richardson number ( $R_i$ ; Weisman and Klemp 1984) suitable for multicellular convection, a sensitivity study is performed in a cloud resolving

model to link its resulting mesoscale ascent / descent with GCM-resolvable variables. The ultimate goal of this research is to develop an anvil cirrus parameterization (ACP), that will couple with cumulus parameterizations in GCMs to improve the cloud-radiation feedback on large-scale climate.

## 2. MODEL AND INITIALIZATION




Fig. 1. The modified atmospheric composite sounding for the midlatitude broken-line squall systems. (a) temperature (T) and dewpoint temperature ( $T_d$ ) profiles. moist adiabats with increasing surface mixing ratio ( $g \text{ kg}^{-1}$ ) of moisture represent low, medium, and high CAPE, respectively. (b) - (c) the non-jetlike and jetlike wind profiles.  $P_0$  is set at 750 (500) mb for shallow (deep) shear layer.  $U_0$  ranges from 10 to 25  $\text{m s}^{-1}$ .

Corresponding author address: Dr. H.-N. S. Chin, P. O. Box 808 (L-103), Livermore, CA 94551.

The model used is an extension of Chin and Ogura's (1989) two-dimensional (2-D) cloud model, which is non-hydrostatic and fully compressible. The major improvements include ice microphysics and radiation transfer schemes for long- (LW) and short-wave (SW). The modified parameterizations of ice microphysics and radiation can simulate midlatitude and tropical squall-line systems with prominent anvils and realistic mesoscale structures. The radiation schemes used can also distinguish the impacts of hydrometeor phase, size, and shape on cloud optical properties. Refer to Chin (1994) and Chin et al. (1995) for the details of model physics.

Due to the computational constraint for a large number of simulations, radiation is not considered in this sensitivity study. The pre-storm conditions of this study are shown in Fig. 1, where the bulk Richardson number is chosen between 35 and 240 for multicellular convection (Weisman and Klemp 1984; Fovell and Ogura 1989). The detail of these 2-D experiments is listed in Table 1, that contains a total of 36 experiments. The model is initialized by a warm, moist bubble and a horizontally homogeneous sounding.

(a) Non-Jetlike Wind Profile																		
Experiment	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
CAPE	L	L	L	L	L	L	M	M	M	M	M	H	H	H	H	H	H	
Shear Depth ( $R_0$ )	S	S	S	D	D	S	S	D	D	S	S	D	D	S	D	D	D	
Shear Intensity ( $U_0$ )	17.5	15.0	10.0	20.0	15.0	12.5	20.0	5.0	10.0	25.0	20.0	15.0	20.0	15.0	10.0	25.0	20.0	
Bulk Ri	36.9	50.2	113.	72.7	129.	186.	37.1	65.8	148.	61.0	95.0	170.	46.7	83.0	187.	76.9	120.	214.

(b) Jetlike Wind Profile																		
Experiment	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
CAPE	L	L	L	L	L	M	M	M	M	M	H	H	H	H	H	H	H	
Shear Depth ( $R_0$ )	S	S	D	D	S	S	D	D	S	S	D	D	S	S	D	D	D	
Shear Intensity ( $U_0$ )	17.5	15.0	12.5	20.0	15.0	12.5	20.0	15.0	12.5	25.0	20.0	15.0	20.0	15.0	12.5	25.0	20.0	15.0
Bulk Ri	50.0	11.9	112.	72.9	129.	186.	18.3	94.5	147.	61.0	93.0	170.	60.7	119.	185.	76.9	120.	214.

In addition, a 3-D simulation of the GATE 4 September 1974 squall line was performed in this research (Chin and Wilhelmson 1996). The comparison of this simulation with its 2-D counterpart

is used to assess the representative of 2D-based ACP into 3-D applications to GCMs.

### 3. RESULTS

#### a. 2-D sensitivity experiments

These sensitivity experiments are used to study the dependence of mesoscale characteristics of squall-line systems on the convective instability, wind shear intensity, shear layer depth, and the pattern of shear profile. Our results indicate that under constant (medium) CAPE and constant (shallow) shear layer depth of a non-jetlike wind profile, the convective strength of the simulated storm is in positive correlation with the shear intensity, while an opposite relation is found in the stratiform region (see exp7, 8, and 9 in Fig 2). With given (medium) CAPE and shear intensity, the deeper shear layer of the non-jetlike wind profile results in stronger convective activity, while it weakens the stratiform region (see exp9 and 11 in Fig. 2). Further, under constant (medium) CAPE and constant velocity difference ( $U_0$ ) of the shear layer for the non-jetlike wind profile, the deeper shear layer (i.e., weaker shear) leads to stronger convective updraft, stratiform descent, and weaker stratiform ascent (see exp7 and 11 in Fig. 2). Similar findings are also seen in each comparison for low and high CAPE, deep shear depth, and the jetlike wind profile, respectively.




Fig. 2. Vertical profiles of horizontally-averaged vertical velocity for experiment 7, 8, 9, and 11. The averaging domain is selected from the leading edge to the upshear side for 200 km wide. (a) in the convective region. (b) in the stratiform region.

The scatter diagram of horizontally-averaged maximum stratiform ascent and  $Ri$  (Fig. 3) clearly exhibits a close correlation between maximum stratiform ascent and  $Ri$  for any given CAPE, shear depth, and shear profile. Another interesting feature of this scatter diagram is attributed to the prominent separation of two different regimes, which are related to the low and deep shear depth cases. In addition, the stratiform ascent is intensified by the jetlike shear profile and the increasing CAPE; however, these two impacts seem to be weaker than the one caused by the shear depth. All of the features aforementioned related

to the stratiform ascent are also seen in its descent counterpart (not shown).




Fig. 3. Horizontally-averaged maximum stratiform ascent strength ( $(W_m)_{\max}$ ) versus the bulk Richardson number ( $Ri$ ) for varied convective available potential energy (CAPE), wind shear intensity, shear depth, and shear profile pattern for multicellular storms. The numbers beside the dots represent the experiments, listed in Table 1.


In general, our results indicate that the stratiform (convective) ascent / descent is strengthened (weakened) with the increasing bulk Richardson number, except the cases involved in varied shear depths. This suggests that the shallow shear depth case should be treated differently in the ACP from its deep depth counterpart. As compared to the tropical MCS environment, the deep shear depth is a typical representative of most midlatitude cases. As a result of the secondary impact of CAPE and shear profile pattern on the stratiform ascent / descent, the upper regime of Fig. 3 may represent the general mesoscale characteristics of tropical MCSs, and the lower one fits the midlatitude cases. However, this suggestion needs more validations for the tropical case before we can generalize the large-scale effects of the sub-GCM-grid process of concern.

#### b. 3-D simulation and its comparison with 2-D results

To calibrate our 2D-based ACP into 3-D applications, we performed a 3-D simulation of the GATE 4 September 1974 squall line (Chin and Wilhelmson 1996). This 3-D simulation replicates many observed features (Houze 1977), such as the arc-shaped rainband structure and its orientation normal to the principal wind shear (Fig. 4). The comparison of 3-D and 2-D simulations in the multicellular portion of the modeled GATE storm exhibits strong similarity at the dynamical structure, except the difference at the magnitude (Fig. 5). More case studies of 3-D simulations for midlatitude MCSs are also being undertaken to establish the relationship between 3-D

and 2-D simulation as the physical basis for the 3-D applications of the ACP.

#### (a) Observation



#### (b) Simulation




Fig. 4. (a) Radar image taken from the ship Oceanographer at 1015 UTC September 4 1974. The radius of the radar scope is 200 km. (b) 3-D depiction of the isosurface at  $0.5 \text{ g kg}^{-1}$  for the total water mixing ratio of the control run at 4 hour of simulation time. The arrow denotes the principal wind shear of environmental winds below 4 km.

#### 4. SUMMARY AND DISCUSSION

Our results suggest that the bulk Richardson number is a valuable index to categorize the mesoscale characteristics of MCSs. Therefore, it might be feasible to parameterize the sub-GCM-grid process associated with the tilting structure of MCSs.

Due to the computational constraint, we are developing our ACP, based on 2-D simulations. Nonetheless, the strong similarity of resolved mesoscale structure of MCSs between 3-D and 2-D

models suggests a promising clue to calibrate the 2D-based ACP into 3-D applications to GCM.

**ACKNOWLEDGMENTS.** This work was supported by the Department of Energy (DOE) ARM (Atmospheric Radiation Measurement) program and conducted under the auspices of the U. S. DOE by the Lawrence Livermore National Laboratory under contract No. W-7405-ENG-48.

#### REFERENCES:

Chin, H.-N. S., and Y. Ogura, 1989: Supplementary modeling study of a tropical convective band. *J. Atmos. Sci.*, **46**, 1440-1447.

Chin, H.-N. S., 1994: The impact of the ice phase and radiation on a mid-latitude squall line. *J. Atmos. Sci.*, **51**, 3320-3343.

Chin, H.-N. S., Q. Fu, M. M. Bradley, and C. R. Molenkamp, 1995: Modeling of a tropical squall line in two dimensions: Sensitivity to radiation and comparison with a midlatitude case. *J. Atmos. Sci.*, **52**, 3172-3193.

Chin, H.-N. S., and R. B. Wilhelmson, 1996: Modeling of a tropical squall line in three dimensions: The arc-shaped rainband structure sustained by a supercell-like storm. *J. Atmos. Sci.*, (to be submitted).

Fovell, R. G., and Y. Ogura, 1989: Numerical simulation of a midlatitude squall line in two dimensions. *J. Atmos. Sci.*, **46**, 3144-3176.

Houze, R. A., Jr., 1977: Structure and dynamics of a tropical squall-line system. *Mon. Wea. Rev.*, **105**, 1540-1567.

Randall, O. A., Harshvardhan, D. A. Dazlich, and T. G. Corsetti, 1989: Interactions among radiation, convection, and large-scale dynamics in a general circulation model. *J. Atmos. Sci.*, **46**, 1943-1970.

Weisman, M. L., and J. B. Klemp, 1984: The dependence of numerically simulated convective storms on vertical wind shear and buoyancy. *Mon. Wea. Rev.*, **110**, 504-520.




Fig. 5. Crosssections in the vertical-horizontal plane of induced vertical velocity and system-relative horizontal velocity in intervals of 1 and 4  $\text{m s}^{-1}$ , respectively, at 4 hour of simulation time. (a) and (b) 3-D simulation. (c), (d) 2-D simulation.

Technical Information Department, Lawrence Livermore National Laboratory  
University of California, Livermore, California 94551