

LOCKHEED MARTIN 

RECEIVED

AUG 01 1996

ES/ER/TM-162/R1

OSTI

**ENVIRONMENTAL  
RESTORATION  
PROGRAM**

**Preliminary Remediation Goals  
for Ecological Endpoints**

This document has been approved by the  
K-25 Site Technical Information Office  
for release to the public. Date: 7-26-96

ENERGY SYSTEMS



MANAGED BY  
LOCKHEED MARTIN ENERGY SYSTEMS, INC.  
FOR THE UNITED STATES  
DEPARTMENT OF ENERGY

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831; prices available from 423-576-8401 (fax 423-576-2865).

Available to the public from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

ES/ER/TM-162/R1

## **Preliminary Remediation Goals for Ecological Endpoints**

R. A. Efroymson  
G. W. Suter II  
B. E. Sample  
D. S. Jones

Date Issued—July 1996


Prepared by the  
Environmental Restoration Risk Assessment Program  
Lockheed Martin Energy Systems, Inc.  
Oak Ridge, Tennessee 37831

Prepared for the  
U.S. Department of Energy  
Office of Environmental Management  
under budget and reporting code EW 20

LOCKHEED MARTIN ENERGY SYSTEMS, INC.  
managing the  
Environmental Management Activities at the  
Oak Ridge K-25 Site Paducah Gaseous Diffusion Plant  
Oak Ridge Y-12 Plant Portsmouth Gaseous Diffusion Plant  
Oak Ridge National Laboratory  
under contract DE-AC05-84OR21400  
for the  
U.S. DEPARTMENT OF ENERGY


DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

**MASTER**




## PREFACE

This technical memorandum was prepared to present preliminary remediation goals (PRGs) for ecological endpoints for risk assessments and decision making at Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) sites. This work was performed under Work Breakdown Structure 1.4.12.2.3.04.05.02 (Activity Data Sheet 8304). Publication of this document meets an Environmental Restoration Risk Assessment Program milestone for FY 96. PRGs are upper concentration limits for specific chemicals in specific environmental media that are anticipated to protect human health or the environment. They can be used for multiple remedial investigations at multiple facilities.



## CONTENTS

PREFACE .....	iii
TABLES .....	vii
ACRONYMS .....	ix
EXECUTIVE SUMMARY .....	xi
1. INTRODUCTION .....	1
1.1 TOXICOLOGICAL BENCHMARKS AND ARARS .....	1
1.2 ENVIRONMENTAL MEDIA .....	2
1.3 LAND USE SCENARIOS .....	2
1.4 MODIFICATION OF PRGS .....	3
2. SURFACE WATER .....	3
3. SEDIMENT .....	10
4. SOIL .....	17
5. REFERENCES .....	23




## TABLES

1. Preliminary remediation goals for surface waters .....	4
2. Preliminary remediation goals for sediments .....	11
3. Preliminary remediation goals for pore water of sediments .....	15
4. Preliminary remediation goals for soils .....	18
5. Life history parameters used to estimate PRGs for wildlife .....	22
6. Summary of species-specific and final soil PRGs for wildlife .....	22
A.1. Soil PRG for red fox assumed to consume 81% small mammals, 10% plants and 9% worms .....	29
A.2. Soil PRG for white-tailed deer assumed to consume 100% plants .....	30
A.3. Soil PRG for white-footed mice assumed to consume 50% plants and 50% worms .....	31
A.4. Soil PRG for Short-tailed Shrews assumed to consume 100% worms .....	32
A.5. Soil PRG for American Woodcock assumed to consume 100% worms .....	33
A.6. Soil PRG for red-tailed hawk assumed to consume 100% small mammals .....	34



## ACRONYMS


ARARs	Applicable or Relevant and Appropriate Requirements
DQOs	data quality objectives
EPA	United States Environmental Protection Agency
FACR	final acute-chronic ratios
FAV	final acute values
LOAEL	lowest-observed-adverse-effects level
NAWQC	National Ambient Water Quality Criteria
NOAEL	no-observed- adverse-effects level
ORNL	Oak Ridge National Laboratory
PELs	Probable Effects Levels
PRGs	Preliminary Remediation Goals
RBRAO	risk-based remedial action objective
RGOs	Remedial Goal Options
RI/FS	remedial investigation/feasibility study
SQAGs	Sediment Quality Assessment Guidelines
TELs	Threshold Effects Levels



## EXECUTIVE SUMMARY

Preliminary remediation goals (PRGs) are useful for risk assessment and decision making at Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) sites. PRGs are upper concentration limits for specific chemicals in specific environmental media that are anticipated to protect human health or the environment. They can be used for multiple remedial investigations at multiple facilities. In addition to media and chemicals of potential concern, the development of PRGs generally requires some knowledge or anticipation of future land use.

In *Preliminary Remediation Goals for Use at the U. S. Department of Energy Oak Ridge Operations Office* (Energy Systems 1995), PRGs intended to protect human health were developed with guidance from *Risk Assessment Guidance for Superfund: Volume I—Human Health Evaluation Manual, Part B* (RAGS) (EPA 1991). However, no guidance was given for PRGs based on ecological risk. The numbers that appear in this volume have, for the most part, been extracted from toxicological benchmarks documents for Oak Ridge National Laboratory (ORNL) and have previously been developed by ORNL. The sources of the quantities, and many of the uncertainties associated with their derivation, are described in this technical memorandum.



# 1. INTRODUCTION

Preliminary remediation goals (PRGs) are useful for risk assessment and decision making at Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) sites. PRGs are upper concentration limits for specific chemicals in specific environmental media that are anticipated to protect human health or the environment. They can be used for multiple remedial investigations at multiple facilities. In addition to media and chemicals of potential concern, the development of PRGs generally requires some knowledge or anticipation of future land use. The development of PRGs at Oak Ridge National Laboratory (ORNL) is proceeding as two separate exercises among experts in environmental and human health sciences, but the goals are brought together during remedial investigations.

In *Preliminary Remediation Goals for Use at the U. S. Department of Energy Oak Ridge Operations Office*, PRGs intended to protect human health were developed with guidance from *Risk Assessment Guidance for Superfund: Volume I—Human Health Evaluation Manual, Part B* (RAGS). However, no guidance was given for PRGs based on ecological risk. The numbers that appear in this volume have, for the most part, been extracted from toxicological benchmarks documents for ORNL. The sources of the quantities, and many of the uncertainties associated with their derivation, are described in this technical memorandum.

PRGs are intended to correspond to minimal and acceptable levels of effects on the general ecological assessment endpoints as defined in the data quality objectives (DQO) process for ecological risk assessments on the Oak Ridge Reservation (Suter et al. 1994). In general, they correspond to small effects on individual organisms which would be expected to cause minimal effects on populations and communities. The PRGs may not be sufficiently protective of species of special concern which are based on effects on individual organisms (Suter et al. 1994). Remedial goals for such species should be developed ad hoc and should be based on no-observed-adverse-effects levels (NOAELs).

## 1.1 TOXICOLOGICAL BENCHMARKS AND ARARS

Toxicological benchmarks have previously been developed at ORNL for the initial screening of contaminants for potential consideration in risk assessments. Some of these are Applicable or Relevant and Appropriate Requirements (ARARs) for remedial action, and others are quantities derived from toxicity test endpoints. Although selected benchmarks are used as PRGs in various media, the two quantities should not be confused. The major differences are:

1. Benchmarks are specific to a receptor or endpoint that is to be protected. PRGs are medium-specific.
2. PRGs are single values for each combination of chemical and medium; benchmarks differ with the assessment endpoint.
3. Benchmarks are conservative, since they are designed to exclude or to screen out only those contaminants for which there is no potential ecological concern. PRGs are regulatory values or thresholds for significant effects.

The guidance document for human health PRGs (Energy Systems 1995) requires that remedial goals be based on ARARs or concentrations determined by risk assessment (EPA 1991). For ecological endpoints, the only federal or state ARARs are National Ambient Water Quality Criteria (NAWQC), available for more than a dozen contaminants in surface waters, and sediment quality criteria available for only five organic contaminants. The United States Environmental Protection Agency (EPA) guidance document provides no equations to protect ecological endpoints or suggested levels of protection analogous to the  $10^{-6}$  risk for human carcinogens (EPA 1991).

## 1.2 ENVIRONMENTAL MEDIA

Three environmental media are considered here: surface water, sediment (including pore water), and soil. Groundwater contamination has always been assumed to have greater consequences for human health than for nonhuman organisms. Data on microscopic and other small biota of groundwater are scarce. Therefore, ecologically-based groundwater PRGs are not presented in this technical memorandum. Although contaminants of potential concern at a site can be identified based on concentrations in food for wildlife or in the organism's tissues, ultimately one of the three media mentioned previously will be remediated. Therefore, the media examined do not include "foods" and are limited to surface water, sediments, and soil.

## 1.3 LAND USE SCENARIOS

A major difference between this document and the guidance provided in RAGS and used in the human health PRGs guidance report (Energy Systems 1995) is that this report lacks emphasis on land use scenarios. For human health, land use determines human activities which determine exposure. Exposure pathways for humans can change, for example, depending on whether the land is industrial or not. Bathing may occur in residential areas and not in industrial areas; ingestion of plants (by humans) may not occur in industrial areas; and inhalation of particulates should not be significant in residential areas. Therefore, because humans engage in different activities in different locations, exposure will depend on land use.

Plants and animals, however, tend to inhabit a particular location and engage in all activities on that particular site. If a site is current or future habitat, then the PRG applies. The streams that flow through agricultural, residential, or industrial lands have the potential to support invertebrates and fish, regardless of land use. Land use types will only indirectly influence aquatic life, for example, through nutrient inputs to a stream. Similarly, exposure pathways for wildlife are not expected to change, depending on land use, though the relative emphasis of one pathway over another may be somewhat altered. If a site contains no habitat, such as a parking lot, it should be screened out during the conceptual development phase for an operable unit (i.e., before a remedial investigation is undertaken).

For lower organisms that are immersed in a medium, the spatial scale is so small that issues of land use do not usually arise (an exception may be soil organisms, as discussed in the following text). The physical habitat for organisms in a stream need not be substantially changed when land uses change. In these cases, correlations between concentrations and effects are used more often than detailed exposure equations. It is notable that ARARs (NAWQC and sediment quality criteria) are not attached to any particular land use scenario. The emphasis for ecological PRG development is on summary statistics for a wide range of effects on a wide range of organisms in a wide range of laboratory and field environments.

Among organisms that are exposed to aquatic contaminants, land use is probably most important to piscivorous wildlife, such as osprey or mink. For some contaminants in water, PRGs are based on aquatic-feeding species. PRGs for water account for both bioaccumulation through the food chain and drinking water. Piscivores may not feed as frequently under industrial land use scenarios. However, this document recommends the same PRGs for water in all contexts because of the paucity of information on piscivore behavior.

A second exceptional case where land use may be important is during the development of PRGs for soils. Soil microbial, invertebrate, and plant communities will be dependent on the management and nutrient additions and extractions from soil. Therefore, PRGs presented for soil may be modified according to land use.

#### **1.4 MODIFICATION OF PRGS**

Non-ARARs-based PRGs may be modified during the remedial investigation/feasibility study (RI/FS) using site-specific data (EPA 1991). Such data may include:

1. land use assumptions;
2. exposure assumptions and habitat considerations (e.g., fraction of land that is suitable habitat);
3. environmental assumptions used for ORNL toxicological benchmarks (e.g., water hardness, soil pH, and organic content);
4. synergistic, antagonistic, or additive effects of pollutants;
5. impacts of contamination of one medium on another (EPA 1991);
6. impacts of remediation of one medium (such as sediments) on contamination of another medium (such as surface water);
7. effects of remediation on organisms and their habitat;
8. new contaminants of concern;
9. desirable level of protection.

In addition, Remedial Goal Options (RGOs), the clean-up goals recommended in the RI/FS, can contain objectives other than concentration limits in environmental media. Two examples are to (1) prevent a contaminated plume from intersecting a stream and (2) prevent toxicity in a standard toxicity test of the contaminated medium.

## **2. SURFACE WATER**

PRGs for surface waters were chosen by comparing the ORNL benchmarks for screening toxicity of contaminants to aquatic life (chronic NAWQC or secondary chronic values; Suter and

Tsao 1996) with those for toxicity to piscivorous wildlife (LOAEL; Sample et al. 1996). The lower of the two benchmarks is the PRG listed in Table 1. If the benchmarks and therefore the PRGs are not exceeded, the contaminant concentration in water probably presents no significant ecological hazard.

**Table 1. Preliminary remediation goals for surface waters**

Chemical	Water Concentration (mg/L)	Endpoint	Criterion
<i>Inorganic chemical</i>			
Aluminum	0.087	aquatic life	chronic NAWQC
Antimony	0.03	aquatic life	secondary chronic value
Arsenic III	0.19	piscivores	chronic NAWQC
Arsenic V	0.0031	aquatic life <sup>2</sup>	secondary chronic value
Barium	0.004	aquatic life <sup>2</sup>	secondary chronic value
Beryllium	0.00066	aquatic life <sup>2</sup>	secondary chronic value
Boron	0.0016	aquatic life	secondary chronic value
Cadmium	0.0011 <sup>1</sup>	aquatic life	chronic NAWQC
Chromium III	0.21 <sup>1</sup>	aquatic life	chronic NAWQC
Chromium VI	0.011	aquatic life	chronic NAWQC
Cobalt	0.023	aquatic life <sup>2</sup>	secondary chronic value
Copper	0.012 <sup>1</sup>	aquatic life	chronic NAWQC
Cyanide	0.0052	aquatic life <sup>2</sup>	chronic NAWQC
Iron	1.0	aquatic life <sup>2</sup>	chronic NAWQC
Lead	0.0032 <sup>1</sup>	aquatic life	chronic NAWQC
Lithium	0.014	aquatic life	secondary chronic value
Manganese	0.12	aquatic life	secondary chronic value
Mercury, inorg. or total	0.0013	aquatic life	secondary chronic value
Mercury, methyl	0.0000026	piscivores	from river otter LOAEL

Table 1. (continued)

Chemical	Water Concentration (mg/L)	Endpoint	Criterion
Molybdenum	0.37	aquatic life	secondary chronic value
Nickel	0.16 <sup>1</sup>	aquatic life	chronic NAWQC
Selenium	0.00039	piscivores	from river otter LOAEL
Silver	0.00036	aquatic life	secondary chronic value
Strontium	1.5	aquatic life <sup>2</sup>	secondary chronic value
Thallium	0.009	piscivores	from river otter LOAEL
Tin	0.073	aquatic life	secondary chronic value
Uranium	0.0026	aquatic life <sup>2</sup>	secondary chronic value
Vanadium	0.020	aquatic life	secondary chronic value
Zinc	0.11 <sup>1</sup>	aquatic life	chronic NAWQC
Zirconium	0.017	aquatic life <sup>2</sup>	secondary chronic value
<i>Organic Chemical</i>			
Acenaphthene	0.023	aquatic life <sup>2</sup>	chronic NAWQC
Acetone	1.5	aquatic life	secondary chronic value
Anthracene	0.00073	aquatic life <sup>2</sup>	secondary chronic value
Benzene	0.13	aquatic life	secondary chronic value
Benzidene	0.0039	aquatic life <sup>2</sup>	secondary chronic value
Benzo(a)anthracene	0.000027	aquatic life <sup>2</sup>	secondary chronic value
Benzo(a)pyrene	0.000014	aquatic life	secondary chronic value
Benzoic acid	0.042	aquatic life <sup>2</sup>	secondary chronic value
Benzyl alcohol	0.0086	aquatic life <sup>2</sup>	secondary chronic value
BHC, gamma (lindane)	0.00008	aquatic life <sup>2</sup>	chronic NAWQC
BHC (other)	0.0000040	piscivores	from river otter LOAEL

Table 1. (continued)

Chemical	Water Concentration (mg/L)	Endpoint	Criterion
Biphenyl	0.014	aquatic life <sup>2</sup>	secondary chronic value
Bis(2-ethylhexyl) phthalate	0.00012	aquatic life	from river otter LOAEL
2-Butanone	14	aquatic life <sup>2</sup>	secondary chronic value
Butylbenzyl phthalate	0.019	aquatic life <sup>2</sup>	secondary chronic value
Carbon disulfide	0.00092	aquatic life <sup>2</sup>	secondary chronic value
Carbon tetrachloride	0.0098	aquatic life <sup>2</sup>	secondary chronic value
Chlordane	0.000037	piscivores	from river otter LOAEL
Chlorobenzene	0.064	aquatic life <sup>2</sup>	secondary chronic value
Chloroform	0.028	aquatic life	secondary chronic value
DDD p,p'	$4.1 \times 10^{-8}$ <sup>3</sup>	piscivores	from belted kingfisher LOAEL
DDT	$4.1 \times 10^{-8}$ <sup>3</sup>	piscivores	from belted kingfisher LOAEL
Decane	0.049	aquatic life <sup>2</sup>	secondary chronic value
Diazinon	0.000043	aquatic life <sup>2</sup>	secondary chronic value
Dibenzofuran	0.0037	aquatic life <sup>2</sup>	secondary chronic value
1,2-Dichlorobenzene	0.014	aquatic life <sup>2</sup>	secondary chronic value
1,3-Dichlorobenzene	0.071	aquatic life <sup>2</sup>	secondary chronic value
1,4-Dichlorobenzene	0.015	aquatic life <sup>2</sup>	secondary chronic value
1,1-Dichloroethane	0.047	aquatic life <sup>2</sup>	secondary chronic value
1,2-Dichloroethane	0.91	aquatic life	secondary chronic value
1,1-Dichloroethene	0.025	aquatic life <sup>2</sup>	secondary chronic value
1,2-Dichloroethenes	0.59	aquatic life <sup>2</sup>	secondary chronic value
1,1-Dichloropropene	0.000055	aquatic life <sup>2</sup>	secondary chronic value
Di-n-butyl phthalate	0.001	piscivores	from belted kingfisher LOAEL

Table 1. (continued)

Chemical	Water Concentration (mg/L)	Endpoint	Criterion
Diethyl phthalate	0.21	aquatic life <sup>2</sup>	secondary chronic value
Endosulfan	0.000051	aquatic life <sup>2</sup>	secondary chronic value
Endrin	0.000061	aquatic life <sup>2</sup>	chronic NAWQC
Ethyl benzene	0.0073	aquatic life <sup>2</sup>	secondary chronic value
Fluoranthene	0.0062	aquatic life <sup>2</sup>	chronic NAWQC
Fluorene	0.0039	aquatic life <sup>2</sup>	secondary chronic value
Heptachlor	0.0000069	aquatic life	secondary chronic value
Hexachloroethane	0.012	aquatic life <sup>2</sup>	secondary chronic value
Hexane	0.00058	aquatic life <sup>2</sup>	secondary chronic value
2-Hexanone	0.099	aquatic life <sup>2</sup>	secondary chronic value
Methoxychlor	0.000019	aquatic life <sup>2</sup>	secondary chronic value
1-Methylnaphthalene	0.0021	aquatic life <sup>2</sup>	secondary chronic value
4-Methyl-2-pentanone	0.17	aquatic life <sup>2</sup>	secondary chronic value
2-Methylphenol	0.013	aquatic life <sup>2</sup>	secondary chronic value
Methylene chloride	2.2	aquatic life <sup>2</sup>	secondary chronic value
Naphthalene	0.012	aquatic life <sup>2</sup>	secondary chronic value
4-Nitrophenol	0.30	aquatic life <sup>2</sup>	secondary chronic value
N-Nitrosodiphenylamine	0.21	aquatic life <sup>2</sup>	secondary chronic value
2-Octanone	0.0083	aquatic life <sup>2</sup>	secondary chronic value
PCBs total	0.0000019 <sup>4</sup>	piscivores	from river otter LOAEL <sup>4</sup>
Aroclor 1016	0.00023 <sup>5</sup>	piscivores	from river otter LOAEL
Aroclor 1221	0.00028	aquatic life <sup>2</sup>	secondary chronic value
Aroclor 1232	0.00058	aquatic life <sup>2</sup>	secondary chronic value

Table 1. (continued)

Chemical	Water Concentration (mg/L)	Endpoint	Criterion
Aroclor 1242	0.000047	piscivores	from river otter LOAEL
Aroclor 1248	0.0000019	piscivores	from river otter LOAEL
Aroclor 1254	0.0000019	piscivores	from river otter LOAEL
Aroclor 1260	0.094	aquatic life <sup>2</sup>	secondary chronic value
Pentachlorobenzene	0.00047	aquatic life <sup>2</sup>	secondary chronic value
1-Pentanol	0.11	aquatic life <sup>2</sup>	secondary chronic value
Phenanthrene	0.0063	aquatic life <sup>2</sup>	secondary chronic value
Phenol	0.11	aquatic life <sup>2</sup>	secondary chronic value
2-Propanol	0.0075	aquatic life <sup>2</sup>	secondary chronic value
1,1,2,2-Tetrachloroethane	0.61	aquatic life <sup>2</sup>	secondary chronic value
Tetrachloroethene	0.098	aquatic life <sup>2</sup>	secondary chronic value
Toluene	0.0098	aquatic life	secondary chronic value
Tribromomethane	0.32	aquatic life <sup>2</sup>	secondary chronic value
1,2,4-Trichlorobenzene	0.11	aquatic life <sup>2</sup>	secondary chronic value
1,1,1-Trichloroethane	0.011	aquatic life <sup>2</sup>	secondary chronic value
1,1,2-Trichloroethane	1.2	aquatic life <sup>2</sup>	secondary chronic value
Trichloroethene	0.47	aquatic life	secondary chronic value
Vinyl acetate	0.016	aquatic life <sup>2</sup>	secondary chronic value
Vinyl chloride	0.782	piscivores <sup>5</sup>	from river otter LOAEL

Table 1. (continued)

Chemical	Water Concentration (mg/L)	Endpoint	Criterion
Xylene	0.013	aquatic life	secondary chronic value

## Notes:

- <sup>1</sup> Hardness dependent criterion for aquatic life benchmark normalized to 100 mg/L.
- <sup>2</sup> Toxic concentration benchmarks are not available for piscivorous wildlife. Therefore, the PRG cannot be assumed to protect wildlife.
- <sup>3</sup> Only a single value was available for DDT and metabolites, though different benchmarks were available for the protection of aquatic life.
- <sup>4</sup> The lowest available concentration for the protection of piscivores from any Aroclor (1248) was used.
- <sup>5</sup> Toxic concentration benchmarks are not available for aquatic life. Therefore, the PRG cannot be assumed to protect fish or aquatic invertebrates.

Since the NAWQC are ARARs for remedial action, they serve as the basis for screening contaminants in water. The chronic NAWQCs are EPA's calculation of final acute values (FAV) divided by final acute-chronic ratios (FACR), where the FAV is the fifth percentile of 48- to 96-hour median lethal concentration (LC50) values or equivalent median effective concentration (EC50) values for each criterion chemical. The FACR is the geometric mean of quotients of at least three LC50/CV ratios from tests of different families of aquatic organisms (Stephan et al. 1985). For several metals, NAWQC are functions of water hardness, and the default PRGs for those metals assume a water hardness of 100 mg/L. However, site-specific water hardness may be substantially different, thereby altering the magnitude or perhaps the direction of the difference between the aquatic life and piscivore toxicological benchmarks.

In this technical memorandum, as well as in the report by Suter and Tsao (1996), NAWQC are not included as potential PRGs for aquatic life if they are based on the protection of humans or other piscivores. This is because ecological PRGs should not be based on effects on humans, and the PRGs based on protection of aquatic life may be lower than the NAWQCs based on fish consumption. In addition, they are not used as potential PRGs for piscivorous wildlife because they are not as rigorously derived or as appropriate to wildlife as the values derived by Opresko et al. (1995).

Where NAWQC were not available, *secondary chronic values* were derived to be used as benchmarks for screening contaminants for toxicity to aquatic life (Suter and Tsao 1996). These values rely on fewer data than do the NAWQC. The method for calculating the secondary chronic value is described in EPA's *Proposed Water Quality Guidance for the Great Lakes System* (1993) and is explained by Suter and Tsao (1996).

For chemicals that are bioaccumulated by piscivores, benchmarks that protect these wildlife may be lower aqueous concentrations than those that protect the aquatic life within the stream. The benchmarks used for wildlife species that feed primarily on aquatic organisms were derived by Sample et al. (1996). The mammalian and avian species considered in the document are representative of wildlife found on the Oak Ridge Reservation. To obtain PRGs, lowest-observed-adverse-effects levels (LOAELs) rather than NOAELs are compared to surface water toxicological benchmarks because (1) NOAELs alone give no indication as to how much higher a concentration must be before adverse effects are observed (LOAELs are presumed to be the threshold levels at which effects become evident), (2) NOAELs often have more uncertainties associated with them than do LOAELs (see Opresko et al. 1995), and (3) LOAELs for effects on

individual wildlife are expected to correspond to no-effect or negligible-effect levels on wildlife populations. The equation used for calculating the LOAEL-based wildlife benchmarks is:

$$C_w = (LOAEL_w \times bw_w) / [W + (F \times BAF)] \quad (\text{Opresko et al. 1995}),$$

which is equivalent to those used by the EPA (1993) where:

$C_w$	=	the benchmark concentration in water.
$LOAEL_w$	=	the lowest observed adverse effects level (derived from LOAELs in individual studies),
$bw_w$	=	body weight of wildlife,
$W$	=	water consumption rate (kg/d),
$F$	=	food consumption rate (kg/d),
$BAF$	=	bioaccumulation factor (ratio of concentration of contaminant in fish tissue to concentration in water; L/kg).

For most of the analytes listed in Table 1, the chronic NAWQC or the secondary chronic value is the PRG. For several analytes, the PRG is based on the LOAEL for mink. However, one analyte, di-n-butyl phthalate, has a PRG that is derived from an avian LOAEL. For some analytes listed in Table 1, piscivore benchmarks were not available. Therefore, in these cases, the concentration cannot be assumed to protect piscivores, and the PRGs may change as the data gaps are filled.

### 3. SEDIMENT

Organisms that reside in sediments are exposed to different concentrations of contaminants from those in the water column. Chemicals in sediment may be present at higher concentrations and for longer time periods than chemicals dissolved in the surface water. Both the concentrations of chemicals in the solid phase of sediments and concentrations in the pore water are relevant to the exposure of benthic (sediment) organisms, and PRGs are presented for both media (Tables 2 and 3). If PRGs are available for both sediment and pore water, the PRG that is determined by the remedial investigation to be the best estimate of risk to sediment biota should take precedence. It is assumed that benthic organisms, including fish, are not significant constituents of the diets of mammalian and avian piscivores; therefore, piscivores are not determinants of PRGs for sediment, as they sometimes are for surface waters. If sediments are to be dredged and disposed of on land, PRGs for soil, as well as PRGs for sediments, should be considered. PRGs for sediments are taken from one of seven sources.

The lowest value of the following sediment toxicity benchmarks for each chemical is the PRG: (1) sediment quality criteria proposed by EPA (EPA 1993b-f); (2) sediment criteria based on the chronic NAWQC; (3) criteria calculated from the lowest chronic value for fish, daphnids, or other invertebrates in surface waters; 4) the NOAA Effects Range-Median (ER-M); or (5) the Florida Department of Environmental Protection Probable Effect Level (PEL). All of these are described at length by Jones et al. (1996), and the lowest chronic values are not used as the PRG if they were originally estimated from acute toxicity (Suter and Tsao 1996). If these criteria are not available, the PRG is the lower of (1) the sediment benchmark calculated from the secondary chronic value for aquatic toxicity or (2) the Ontario Ministry of the Environment Severe Effect Level. The secondary chronic value is often one or two orders of magnitude lower than the lowest chronic values; therefore, PRGs based on this value are likely to be more conservative than other PRGs.

The five sediment quality criteria proposed in 1993 by EPA (EPA 1993b-f) are potential ARARs for assessing sediment quality with respect to acenaphthene, dieldrin, endrin, fluoranthene, and phenanthrene at hazardous waste sites. These and the ER-Ms and PELs were the only potential PRGs for organic chemicals that were not calculated based on partitioning between water and sediment.

**Table 2. Preliminary remediation goals for sediments**

Chemical	Sediment Concentration (mg/kg)	Type of Benchmark <sup>1</sup>
<i>Organic chemical</i>		
Acenaphthene	0.089	PEL
Acenaphthylene	0.13	PEL
Acetone <sup>2</sup>	0.0091	LCV for daphnid
Aldrin	0.080	Ontario Ministry of the Environment-severe
Anthracene	0.25	PEL
Benzene	0.16	SCV
Benzidine <sup>2</sup>	0.0017	SCV
Benzo(a)anthracene	0.69	PEL
Benzo(a)pyrene	0.76	PEL
Benzyl alcohol <sup>2</sup>	0.0011	SCV
Biphenyl	1.1	SCV
Bis(2-ethylhexyl)phthalate	2.7	PEL
4-Bromophenyl phenyl ether	1.2	SCV
2-Butanone <sup>2</sup>	0.27	SCV
Carbon disulfide	0.00086	SCV
Carbon tetrachloride	2.0	LCV for fish
Chlordane	0.0048	PEL
Chlorobenzene	0.417	SCV
Chloroform	0.96	LCV for fish

Table 2. (continued)

Chemical	Sediment Concentration (mg/kg)	Type of Benchmark <sup>1</sup>
Chrysene	0.85	PEL
Decane	41	SCV
DDD p,p'	0.0078	PEL
DDE p,p'	0.027	ER-M
DDT	0.052	PEL
Diazinon	0.0019	SCV
Dibenzo(a,h)anthracene	0.14	PEL
Dibenzofuran	0.42	SCV
1,2-Dichlorobenzene	0.33	SCV
1,3-Dichlorobenzene	1.7	SCV
1,4-Dichlorobenzene	0.35	SCV
1,1-Dichloroethane	0.027	SCV
1,2-Dichloroethane	4.3	LCV for daphnid
1,1-Dichloroethylene	3.5	LCV for fish
1,2-Dichloroethylene	0.40	SCV
1,3-Dichloropropene	0.23	LCV for fish
Di-n-butyl phthalate	240	LCV for daphnid
Diethyl phthalate	0.61	SCV
Dieldrin	0.0043	PEL
Endosulfan	0.0055	SCV
Endrin	0.045	ER-M
Ethyl benzene	5.4	LCV for fish
Fluoranthene	1.5	PEL
Fluorene	0.14	PEL

Table 2. (continued)

Chemical	Sediment Concentration (mg/kg)	Type of Benchmark <sup>1</sup>
Heptachlor	13	LCV for fish
Hexachloroethane	1.0	SCV
Hexane	0.040	SCV
2-Hexanone <sup>2</sup>	0.023	SCV
Lindane (gamma BHC)	0.00099	PEL
Methoxychlor	0.019	SCV
Methylene chloride	18	LCV for fish
4-Methyl-2-pantanone <sup>2</sup>	15	LCV for fish
2-Methylphenol <sup>2</sup>	0.012	SCV
Naphthalene	0.39	PEL
2-Octanone <sup>2</sup>	0.018	SCV
PCBs total	0.18	PEL
Aroclor 1221	0.12	SCV
Aroclor 1232	0.60	SCV
Aroclor 1242	29	LCV for fish
Aroclor 1248	1.0	SCV
Aroclor 1254	72	LCV for fish
Aroclor 1260	63	LCV for fish
Pentachlorobenzene	0.70	SCV
1-Pentanol <sup>2</sup>	0.034	SCV
Phenanthrene	0.54	PEL
Phenol	0.032	chronic NAWQC
2-Propanol <sup>2</sup>	0.000084	SCV

Table 2. (continued)

Chemical	Sediment Concentration (mg/kg)	Type of Benchmark <sup>1</sup>
Pyrene	1.4	PEL
1,1,2,2-Tetrachloroethane	5.4	LCV for fish
Tetrachloroethylene	3.2	LCV for daphnid
Toluene	0.050	SCV
Tribromomethane	0.66	SCV
1,2,4-Trichlorobenzene	9.7	SCV
1,1,1-Trichloroethane	9.6	LCV for fish
1,1,2-Trichloroethane	9.8	LCV for fish
Trichloroethene	52	LCV for fish
Vinyl acetate	0.00084	SCV
Xylene	0.16	SCV
<i>Inorganic chemical</i>		
Arsenic	42	PEL
Cadmium	4.2	PEL
Chromium	160	PEL
Copper	110	PEL
Lead	110	PEL
Mercury	0.7	PEL
Nickel	43	PEL
Silver	1.8	PEL
Zinc	270	PEL

## Notes:

<sup>1</sup> PEL, Florida Department of Environmental Protection Probable Effects Level (Macdonald 1994); ER-M, NOAA Effects Range-Median (Long et al. 1995); SCV, secondary chronic value (Jones et al. 1996); LCV, lowest chronic value for daphnids, non-daphnid invertebrates, or fish; Ontario Ministry of the Environment - severe, severe effects level

<sup>2</sup> Denotes polar nonionic organic compounds, for which the equilibrium partitioning model is likely to provide a conservative model of exposure.

For nonionic organic chemicals for which octanol-water partition coefficients are available, sediment toxicity benchmarks were calculated based on equilibrium partitioning, assuming 1% organic carbon and using the benchmarks for surface waters (NAWQC, secondary chronic values, and lowest chronic values for fish, daphnids, and non-daphnid invertebrates). These benchmarks were considered as possible PRGs, with lower concentrations selected according to the priority discussed previously. An advantage of the equilibrium partitioning approach is that the PRG can be adapted to different sites by adjusting the organic carbon parameter. Both the sediment quality criteria and the equilibrium partitioning benchmarks have been used by ORNL to screen for contaminants of potential concern for ecological risk assessments (Jones et al. 1996). The equation originally used by EPA (1989) and then used by Jones et al. (1996) is:

$$SQB = f_{oc} \times K_{oc} \times WQB,$$

where:

$SQB$  = sediment quality benchmark,  
 $f_{oc}$  = mass fraction of organic carbon,  
 $K_{oc}$  = organic carbon-water partition coefficient,  
 $WQB$  = water quality benchmark.

The derivation of the equation is given by Jones et al. (1996). The biological assumptions of the equilibrium partitioning approach, according to Jones et al. (1996), are:

1. the sensitivities of benthic species and species tested to derive WQC, predominantly water column species, are similar;
2. the levels of protection afforded by WQC are appropriate for benthic organisms; and
3. exposures are similar regardless of feeding type or habitat (EPA 1993b).

Sediments and pore water are assumed to be in continual equilibrium (MacDonald 1994a).

**Table 3. Preliminary remediation goals for pore water of sediments (to be used with Table 1)<sup>2</sup>**

Chemical	Water Concentration (mg/L)	Criterion
<i>Inorganic chemical</i>		
Arsenic III	0.19	chronic NAWQC
Mercury, methyl	0.0000028	secondary chronic value
Selenium	0.005	chronic NAWQC
Thallium	0.012	secondary chronic value

Table 3. (continued)

Chemical	Water Concentration (mg/L)	Criterion
<i>Organic chemical</i>		
BHC (other than gamma)	0.0022	secondary chronic value
DDD p,p'	0.000011	secondary chronic value
DDT	0.000013	secondary chronic value
Di-n-butyl phthalate	0.035	secondary chronic value
PCBs total	0.00014	secondary chronic value
Aroclor 1242	0.000053	secondary chronic value
Aroclor 1248	0.000081	secondary chronic value
Aroclor 1254	0.000033	secondary chronic value
Xylene	0.013	secondary chronic value

## Notes:

<sup>1</sup> Hardness dependent criterion for aquatic life benchmark normalized to 100 mg/L.

<sup>2</sup> PRGs for pore water are presented in Table 1 except for surface water values that were based on risk in piscivores. PRGs for those chemicals are listed here and obtained from Suter and Tsao (1996).

PRGs for inorganic chemicals in sediments are taken from the Florida Sediment Quality Assessment Guidelines (SQAGs) (MacDonald 1994a). The SQAGs include Threshold Effects Levels (TELs), "the upper limit of the range of sediment contaminant concentrations dominated by no effects data entries . . . [and] not considered to represent significant hazards to aquatic organisms" and Probable Effects Levels (PELs), "the lower limit of the range of contaminant concentrations that are usually or always associated with adverse biological effects" (MacDonald 1994a). In this document, PELs are used as PRGs for several metals. The calculation used is:

$$PEL = \sqrt{EDS_m \times NEDS_H} ,$$

where  $EDS_m$  is the 50th percentile concentration in the effects data set, and  $NEDS_H$  is the 85th percentile concentration in the no effects data set. Few data exist on chronic effects of contaminants on organisms in sediments; therefore, many of the studies present acute responses.

The Florida SQAGs were designed for prioritizing risk management actions, interpreting and designing monitoring programs for sediment contamination, designing wetland restoration programs, supporting decisions by multiple parties relating to sediments, etc. They were not intended for use as sediment quality criteria (MacDonald 1994a). The SQAGs were designed for use in marine and estuarine systems only. In addition, factors that influence bioavailability of metals at a site, such as

acid volatile sulfide for divalent cations, are not taken into account by these guidelines or PRGs (MacDonald 1994a).

Jones et al. (1996) caution that the sediment benchmarks do not represent remedial goals, since the removal or other disturbance of sediment can affect habitat or cause toxic effects in surface water. Similarly, MacDonald (1994a) suggests that the Florida SQAGs should not be used directly as clean-up targets for hazardous sites without additional site-specific studies. The PRGs for sediments are not ideal and should be modified on a site-by-site basis. Nonetheless, they are the best and most current remedial goals available to protect nonhuman organisms and ecological systems in the absence of reliable sediment toxicity benchmarks.

Although sediments are usually identified for remediation on the basis of their bulk concentrations, in some cases pore water concentrations are the appropriate PRG because the toxicity of the sediment is more clearly associated with the pore water than bulk sediment contaminant levels. This circumstance will occur when the toxicity is primarily due to exposure to pore water, and variance in sediment properties causes the sediment/water distribution coefficient to be variable. Pore water PRGs would also be appropriate where ecological risks are associated with a contaminated groundwater plume that intersects or is predicted to intersect the bed of a stream or river. The PRGs for these cases are the potential PRGs for aquatic life in surface water (i.e., chronic NAWQCs and secondary chronic values). These values are presented in Table 1, except for those chemicals with aqueous PRGs based on wildlife risks. The values for these chemicals are presented in Table 3, since it is assumed that piscivores do not feed on sediment-associated organisms.

#### 4. SOIL

PRGs for soil were chosen by comparing the ORNL toxicological benchmarks for plants, microorganisms, and earthworms in soils to each other and to calculated PRGs for wildlife. ARARs for soils do not exist. Earthworms represent highly exposed invertebrates. Benchmarks for plants appear in *Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Terrestrial Plants* (Will and Suter 1995b); benchmarks for earthworms and microorganisms appear in *Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Soil and Litter Invertebrates and Heterotrophic Process* (Will and Suter 1995a). The procedure for calculating PRGs for wildlife endpoints is described in the following paragraphs. All benchmarks and all PRGs are based on one or more field, greenhouse, or growth chamber studies.

Benchmarks for the four types of organisms (wildlife, plants, microorganisms, earthworms) were compared, and the lowest value available is the PRG (Table 4). In media other than soil, if the benchmarks and therefore the PRGs are not exceeded, it is assumed that the chemical concentration in the medium presents no significant ecological hazard. In soils, the uncertainties associated with the PRGs are probably greater than in water or sediments. These uncertainties include:

1. For many chemicals in Table 4, toxicity to only one or two of the three types of organisms (plants, microorganisms, invertebrates) has been studied.
2. Will and Suter (1995a,b) have low confidence in most of the soil benchmarks because of a limited number of studies and/or biological endpoints for almost all contaminants.

3. Soil-earthworm (Sample et al. 1996a) and soil-plant (Efroymson et al. 1996) contaminant uptake factors do not account for soil and biota properties.

Although the confidence in the numbers in Table 4 is generally low, PRGs for soils are needed. As the toxicity of contaminants to additional organisms is investigated, these preliminary values will be modified. PRGs can only be based on toxicity to categories of organisms that have been studied; final remedial goals can incorporate safety factors to protect other populations.

**Table 4. Preliminary remediation goals for soils**

Chemical	Soil Concentration (mg/kg)	Endpoint <sup>1</sup>
<i>Inorganic chemical</i>		
Aluminum	50 <sup>6</sup>	plant <sup>3</sup>
Antimony	5 <sup>6</sup>	plant <sup>3,4,9</sup>
Arsenic	2.66	shrew
Barium	208	woodcock
Beryllium	10 <sup>6</sup>	plant <sup>3,4,9</sup>
Boron	0.5 <sup>6</sup>	plant <sup>3,9</sup>
Bromine	10 <sup>6</sup>	plant <sup>3,4,9</sup>
Cadmium	3 <sup>6</sup>	plant
Chromium	0.4 <sup>6</sup>	earthworm
Cobalt	20 <sup>6</sup>	plant <sup>3,9</sup>
Copper	50 <sup>7</sup>	earthworm
Fluorine	30 <sup>6</sup>	microorganism <sup>3,9</sup>
Iodine	4 <sup>6</sup>	plant <sup>3,4</sup>
Iron	200 <sup>6</sup>	microorganism <sup>2,3,9</sup>
Lead	50 <sup>7</sup>	plant
Lithium	2 <sup>6</sup>	plant <sup>3</sup>
Manganese	100 <sup>6</sup>	plant <sup>3</sup>

Table 4. (continued)

Chemical	Soil Concentration (mg/kg)	Endpoint <sup>1</sup>
Mercury	0.0185	woodcock
Molybdenum	2 <sup>6</sup>	plant <sup>3</sup>
Nickel	24	woodcock
Selenium	0.79	shrew
Silver	2 <sup>6</sup>	plant <sup>9</sup>
Technetium	0.2 <sup>6</sup>	plant <sup>3,4,9</sup>
Thallium	1 <sup>6</sup>	plant <sup>3,4</sup>
Tin	50 <sup>6</sup>	plant <sup>3,9</sup>
Titanium	1000 <sup>6</sup>	microorganism <sup>2,3,9</sup>
Uranium	5 <sup>6</sup>	plant <sup>3,4</sup>
Vanadium	2 <sup>6</sup>	plant <sup>3</sup>
Zinc	26.3	woodcock
<i>Organic chemical</i>		
Acenaphthene	20 <sup>6</sup>	plant <sup>3,4,9</sup>
Biphenyl	60 <sup>6</sup>	plant <sup>3,4,9</sup>
Chlorobenzene	40 <sup>6</sup>	earthworm <sup>2,4,9</sup>
3-Chlorophenol	10 <sup>6</sup>	earthworm <sup>2,4,9</sup>
Di-n-butyl phthalate	200 <sup>6</sup>	plant <sup>3,4,9</sup>
1,4-Dichlorobenzene	20 <sup>6</sup>	earthworm <sup>2,4,9</sup>
3,4-dichlorophenol	20 <sup>6</sup>	earthworm <sup>2,4,9</sup>
Diethyl phthalate	100 <sup>6</sup>	plant <sup>3,4,9</sup>
2,4-Dinitrophenol	20 <sup>7</sup>	plant <sup>3,4,9</sup>
Furan	600 <sup>6</sup>	plant <sup>3,4,9</sup>

Table 4. (continued)

Chemical	Soil Concentration (mg/kg)	Endpoint <sup>1</sup>
Hexachlorobenzene	1000 <sup>6</sup>	microorganism <sup>3,4,9</sup>
Hexachlorocyclopentadiene	10 <sup>6</sup>	plant <sup>3,4,9</sup>
4-nitrophenol	7 <sup>6</sup>	earthworm <sup>2,4,9</sup>
Pentachlorophenol	3 <sup>6</sup>	plant
Pentachlorobenzene	20 <sup>6</sup>	earthworm <sup>2,4,9</sup>
Phenol	30 <sup>6</sup>	earthworm <sup>9</sup>
PCBs	40 <sup>6</sup>	plant <sup>3,4,9</sup>
Styrene	300 <sup>6</sup>	plant <sup>3,4,9</sup>
1,2,3,4-Tetrachlorobenzene	10 <sup>6</sup>	earthworm <sup>2,4,9</sup>
2,3,4,5-Tetrachlorophenol	20 <sup>6</sup>	earthworm <sup>2,4,9</sup>
Toluene	200 <sup>6</sup>	plant <sup>3,4,9</sup>
1,2,3-Trichlorobenzene	20 <sup>6</sup>	earthworm <sup>2,4,9</sup>
1,2,4-Trichlorobenzene	20 <sup>6</sup>	earthworm <sup>2,4,9</sup>
2,4,5-Trichlorophenol	9 <sup>6</sup>	earthworm <sup>2,4,9</sup>
2,4,6-Trichlorophenol	10 <sup>6</sup>	earthworm <sup>2,4,9</sup>

## Notes:

- <sup>1</sup> The most sensitive type of organism among plants, earthworms, and microorganisms. The PRG is based on the toxicological benchmark concentration for this organism and does not consider effects on wildlife.
- <sup>2</sup> Toxic concentration benchmarks are not available for plants in soils. Therefore, the PRG cannot be assumed to protect plants.
- <sup>3</sup> Toxic concentration benchmarks are not available for earthworms. Therefore, the PRG cannot be assumed to protect earthworms.
- <sup>4</sup> Toxic concentration benchmarks are not available for microorganisms. Therefore, the PRG cannot be assumed to protect microorganisms.
- <sup>5</sup> The benchmark for methyl mercury in plants was compared to benchmarks for total mercury in earthworms and microorganisms.
- <sup>6</sup> Will and Suter (1995a,b) have low confidence in this value. The level of confidence refers to the benchmark chosen for the PRG and not to the relationship between it and the benchmarks not chosen.
- <sup>7</sup> Will and Suter (1995a,b) have moderate confidence in this value.
- <sup>8</sup> Will and Suter (1995a,b) have high confidence in this value.
- <sup>9</sup> Either soil-plant, soil-earthworm uptake factors or LOAELs were not available. Therefore, the PRG cannot be assumed to protect wildlife.

Remedial goals for soils should be modified based on the bioavailability of the contaminants of concern. The bioavailable fraction of a chemical in soil is probably lower than the total concentration. Toxicity tests in soil on which the PRGs are based sometimes begin with known concentrations of a chemical or may assume a relationship between what is extractable by an arbitrary solvent and what is bioavailable. The organic fraction and pH of soil are two major factors that influence the uptake of chemicals by plants. "Aged" organic contaminants may not be as available for uptake as freshly added chemicals. 2,4-Dinitrophenol is an example of a chemical that is more toxic to plants under acidic conditions (Will and Suter 1995b). The context of the studies from which the toxicological benchmarks for soil were derived is available in the Will and Suter reports (1995a,b), Sample et al. (1996b), and in greater detail in the original publications. As more is known about the bioavailability of contaminants in soils, the default PRGs should be modified.

The addition of PRGs to protect terrestrial wildlife is a new feature in this 1996 revision of the PRGs report. Wildlife PRGs for soil were derived by iteratively calculating exposure estimates using different soil concentrations and soil-to-biota contaminant uptake factors for the Oak Ridge Reservation. The soil concentrations were manipulated to produce an exposure estimate equivalent to the wildlife endpoint-specific and contaminant-specific LOAEL, which were obtained from Sample et al. (1996b). Uptake factors for plants were obtained from Efroymson et al. (1996) and for earthworms and small mammals from Sample et al. (1996a). Because different diets may dramatically influence exposures and sensitivity to contaminants varies among species, PRGs were developed for six species present on the Oak Ridge Reservation: short-tailed shrew, white-footed mouse, red fox, white-tailed deer, American woodcock, and red-tailed hawk. The PRGs for each of these species were compared, and the lowest value was selected as the final wildlife PRG. This PRG appears in Table 4 if the concentration in soil is lower than the toxicity benchmarks for earthworms, plants, or soil microbial processes. Estimates of oral exposure to contaminants were generated using the generalized exposure model (Sample and Suter 1994):

$$E_j = \sum_{i=1}^m p_{ik} \left( \frac{IR_i \times C_{ijk}}{BW} \right)$$

where:

- $E_j$  = total exposure to contaminant (j) (mg/kg/d),
- $m$  = total number of ingested media (e.g., food or soil),
- $IR_i$  = ingestion rate for medium (i) (kg/d or L/d),
- $p_{ik}$  = proportion of type (k) of medium (i) consumed (unitless),
- $C_{ijk}$  = concentration of contaminant (j) in type (k) of medium (i) (mg/kg or mg/L),
- $BW$  = body weight of endpoint species (kg).

PRGs were calculated for only those chemicals for which both uptake factors and LOAELs were available. The 90th percentile of the soil-to-biota uptake factor was used as a conservative estimate of the chemical concentrations in wildlife food types (earthworms, plants, or small mammals). Species-specific life history parameters needed to estimate exposure were obtained from Sample and Suter (1994) and are presented in Table 5. The model accounts for the ingestion of soil as well as food. Summaries of the derivation of PRGs for each species are presented in the appendix. Soil PRGs for each wildlife species and the recommended final PRG for protection of wildlife, generally, are presented in Table 6. For all chemicals the final PRG for protection of wildlife was based on the PRG for either short-tailed shrew or American woodcock (Table 6). This result is due to the large

quantity of soil ingested by these wildlife and the relatively high chemical uptake rates for their food (earthworms).

**Table 5. Life history parameters used to estimate PRGs for wildlife**

Species	Body Weight (kg)	Ingestion Rate (kg/d)		Percent of diet		
		Food	Soil	Earthworm	Plant	Small Mammal
Short-tailed Shrew	0.015	0.009	0.00117	100%	0%	0%
White-footed Mouse	0.022	0.0034	0.000068	50%	50%	0%
Red Fox	4.5	0.45	0.0126	9%	10%	81%
White-tailed Deer	56.5	1.74	0.0348	0%	100%	0%
American Woodcock	0.198	0.15	0.0156	100%	0%	0%
Red-tailed Hawk	1.126	0.109	0	0%	0%	100%

**Table 6. Summary of species-specific and final soil PRGs for wildlife**

Analyte	Preliminary Remedial Goal (mg/kg in soil)						
	Red Fox	White-tailed Deer	White-footed Mouse	Short-tailed Shrew	American Woodcock	Red-tailed Hawk	Final
Aluminum	1040	1920	1440	155	ND <sup>a</sup>	ND <sup>a</sup>	155
Arsenic	32.5	119	20	2.66	18.5	16500	2.66
Barium	900	700	1170	250	208	7000	208
Cadmium	62	77.5	33	5.4	4.05	1570	4.05
Chromium	72	1380	40.4	5.7	0.78	233	0.78
Copper	143	455	415	77	87.5	860	77
Lead	5010	10100	8050	1000	56	2630	56
Lithium	1280	3175	1670	199	ND <sup>a</sup>	ND <sup>a</sup>	199
Manganese	19000	6800	14100	4200	ND <sup>a</sup>	ND <sup>a</sup>	4200
Mercury	0.165	12.7	0.9	0.128	0.0185	0.89	0.0185
Molybdenum	32	122	16.4	2.33	21.3	36000	2.33
Nickel	560	4150	345	49.5	24	4750	24
Aroclor 1254	56 <sup>b</sup>	138	11.8	1.47	3.27	ND <sup>b</sup>	1.47
Aroclor 1260	0.88	138	0.63	0.089	0.19	3.55	0.089

Table 6. (continued)

Analyte	Preliminary Remedial Goal (mg/kg in soil)						
	Red Fox	White-tailed Deer	White-footed Mouse	Short-tailed Shrew	American Woodcock	Red-tailed Hawk	Final
Selenium	5.05	38	5.7	0.79	0.88	44.5	0.79
Thallium	13	15.9	30.8	2.1	ND <sup>a</sup>	ND <sup>a</sup>	2.1
Uranium	505	1380	800	62	ND <sup>a</sup>	ND <sup>a</sup>	62
Vanadium	231	170	237	32.6	ND <sup>a</sup>	ND <sup>a</sup>	32.6
Zinc	650	3950	1140	177	26.3	570	26.3

Notes:

<sup>a</sup> ND = LOAEL for birds not available for these chemicals.

<sup>b</sup> Soil-small mammal uptake factor not available. Red fox available from soil only. PRG for red-tailed hawk could not be calculated because soil ingestion=0;

PRGs for soil, more than for other media, are likely to be influenced by different land use scenarios. Uses of soil will affect the fraction of land that is suitable for habitat and the necessity of protecting various organisms. The PRGs in Table 4 and the calculations for wildlife assume that habitat is 100% available for the organisms in the assessed region. This assumption is reasonable for relatively immobile organisms such as plants, earthworms, and microorganisms. However, for wildlife, the role of habitat will be important for determining exposure. For example, if the availability of habitat at a site is minimal, use of the site by wildlife, and therefore contaminant exposure, is likely to be minimal.

## 5. REFERENCES

Efroymson, R. A., B. E. Sample, G. W. Suter II, and T. L. Ashwood. 1996. *Soil-plant Uptake Factors: Review and Recommendations for the Oak Ridge Reservation*. ES/ER/TM-198. Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Energy Systems (Lockheed Martin Energy Systems, Inc.). 1995. *Preliminary remediation goals for use at the U. S. Department of Energy Oak Ridge Operations Office*, ES/ER/TM-106, Risk Assessment Program, Health Sciences Research Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

EPA (United States Environmental Protection Agency). 1989. Briefing report to the EPA Science Advisory Board on the Equilibrium Partitioning Approach to Generating Sediment Quality.

EPA. 1991. *Risk Assessment Guidance for Superfund: Volume I - Human Health Evaluation Manual (Part B, Development of Risk-based Preliminary Remediation Goals)*, Office of Emergency and Remedial Response, U.S. Environmental Protection Agency, Washington, D.C. Pub. 9285.7-01B.

EPA. 1993a. Water quality guidance for the Great Lakes System and correction; Proposed rules, *Federal Register* 58(72):20802-21047.

EPA. 1993b. *Sediment Quality Criteria for the Protection of Benthic Organisms: Acenaphthene.*, EPA-822-R-93-013, United States Environmental Protection Agency, Washington, D. C.

EPA. 1993c. *Sediment Quality Criteria for the Protection of Benthic Organisms: Dieldrin*, EPA-822-R-93-015, United States Environmental Protection Agency, Washington, D. C.

EPA. 1993d. *Sediment Quality Criteria for the Protection of Benthic Organisms: Endrin*, EPA-822-R-93-016, United States Environmental Protection Agency, Washington, D. C.

EPA. 1993e. *Sediment Quality Criteria for the Protection of Benthic Organisms: Fluoranthene.*, EPA-822-R-93-012, United States Environmental Protection Agency, Washington, D. C.

EPA. 1993f. *Sediment Quality Criteria for the Protection of Benthic Organisms: Phenanthrene*, EPA-822-R-93-014, United States Environmental Protection Agency, Washington, D. C.

Etnier, E. L., E. P. McDonald, and L. M. Houlberg. 1993. *Applicable or relevant and appropriate requirements (ARARs) for remedial action at the Oak Ridge Reservation: A compendium of environmental laws*, ES/ER/TM-1/R2, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Jones, D. S., R. N. Hull, and G. W. Suter II. 1996. *Toxicological Benchmarks for Screening Contaminants of Potential Concern for Effects on Sediment-associated Biota: 1996 Revision*, ES/ER/TM-95/R2, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Long, E. R. and L. G. Morgan. 1990. *The potential for biological effects of sediment-sorbed contaminants tested in the National Status and Trends Program*, NOAA Technical Memorandum NOS OMA 58, National Oceanic and Atmospheric Administration, Seattle, Washington.

Long, E. R., D. D. MacDonald, S. L. Smith, and F. D. Calder. 1995. Incidence of Adverse Biological Effects within Ranges of Chemical Concentrations in Marine and Estuarine Sediments. *Environmental Management* 19(1), 81-97.

MacDonald, D. D., MacDonald Environmental Services Ltd. November 1994a. *Approach to the assessment of sediment quality in Florida coastal waters. Vol. 2 - Application of the sediment quality assessment guidelines*, Florida Department of Environmental Protection, Office of Water Policy, Tallahassee, Florida.

MacDonald, D. D., MacDonald Environmental Services Ltd. November 1994b. *Approach to the assessment of sediment quality in Florida coastal waters. Vol. 3 - Supporting documentation: Biological effects database for sediments*, Florida Department of Environmental Protection, Office of Water Policy, Tallahassee, Florida.

Sample, B. E., D. M. Opresko, and G. W. Suter II. 1996. *Toxicological Benchmarks for Wildlife: 1996 Revision*, ES/ER/TM-86/R3, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Sample, B. E., and G. W. Suter II. 1994a. *Estimating Exposure of Terrestrial Wildlife to Contaminants*. ES/ER/TM-125, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Sample, B. E., R. A. Efroymson, G. W. Suter II, and T. L. Ashwood. 1996b. *Soil-earthworm and Soil-small mammal Contaminant Uptake Factors: Review and Recommendations for the Oak Ridge Reservation*. ES/ER/TM-197. Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Stephan, C. E., D. I. Mount, D. J. Hansen, J. H. Gentile, G. A. Chapman, and W. A. Brungs. 1985. *Guidelines for deriving numerical national water quality criteria for the protection of aquatic organisms and their uses*, PB85-227049, National Technical Information Service, Springfield, Virginia.

Suter, G. W. II, B. E. Sample, D. S. Jones, T. L. Ashwood, and J. M. Loar. 1995. Approach and Strategy for Performing Ecological Risk Assessments for the U. S. Department of Energy's Oak Ridge Reservation: 1995 Revision, ES/ER/TM-33/R2, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Suter, G. W. II and C. L. Tsao. 1996. *Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Aquatic Biota: 1996 Revision*, ES/ER/TM-96/R2, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Will, M. E. and G. W. Suter II. 1995a. *Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Soil and Litter Invertebrates and Heterotrophic Process: 1995 Revision*, ES/ER/TM-126/RI, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Will, M. E. and G. W. Suter II. 1995b. *Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Terrestrial Plants: 1995 Revision*, ES/ER/TM-85/R2, Oak Ridge National Laboratory, Oak Ridge, Tennessee.



## **APPENDIX**



Table A.1. Soil PRG for red fox assumed to consume 81% small mammals, 10% plants and 9% worms

Analyte	Soil Conc (mg/kg)	90th Percentile Worm UF	90th Percentile Plant UF	90th Percentile Mammal UF	Estimated Worm Conc (mg/kg)	Estimated Plant Conc (mg/kg)	Estimated Mammal Conc (mg/kg)	Body Wt (kg)	Worm Ingestion (kg/d)	Plant Ingestion (kg/d)	Mammal Ingestion (kg/d)	Soil Ingestion (kg/d)	Worm Exposure (mg/kg/d)	Plant Exposure (mg/kg/d)	Mammal Exposure (mg/kg/d)	Soil Exposure (mg/kg/d)	Total Exposure (mg/kg/d)	LOAEL (mg/kg/d)	Form	HQ
Aluminum	1040	0.118	0.030	0.014	122.72	30.68	14.56	4.5	0.0405	0.045	0.3645	0.0126	1.10448	0.3068	1.17936	2.912	5.50254	5.515	AlCl3	1.00
Arsenic	32.5	0.811	0.032	0.008	26.351	1.04975	0.26	4.5	0.0405	0.045	0.3645	0.0126	0.237159	0.0104975	0.02106	0.091	0.3597165	0.360	Arsenite	1.00
Barium	900	0.160	0.237	0.061	144	213.12	55.26	4.5	0.0405	0.045	0.3645	0.0126	1.296	2.1312	4.47606	2.52	10.42326	10.5	barium hydroxide	1.00
Cadmium	62	6.410	1.114	0.132	397.4386	69.099	8.1592	4.5	0.0405	0.045	0.3645	0.0126	3.5769474	0.65099	0.6608952	0.1736	5.1024326	5.094	cadmium chloride	1.00
Chromium	72	8.331	0.666	0.221	599.8248	4.788	15.9264	4.5	0.0405	0.045	0.3645	0.0126	5.3984222	0.0474788	1.2900384	0.2016	6.3179416	6.94	Cr6	1.00
Copper	143	0.326	0.381	0.740	118.1752	54.4401	105.8486	4.5	0.0405	0.045	0.3645	0.0126	1.0635768	0.544401	8.5717366	0.4004	10.5821144	10.6	copper sulfate	1.00
Lead	5010	0.164	0.052	0.045	819.135	261.522	222.945	4.5	0.0405	0.045	0.3645	0.0126	7.372215	2.61522	18.058545	14.028	42.07358	42.25	lead acetate	1.00
Lithium	1280	0.217	0.034	0.033	277.248	43.52	41.856	4.5	0.0405	0.045	0.3645	0.0126	2.495232	0.4352	3.390336	3.584	9.904768	9.9	lithium carbonate	1.00
Manganese	19000	0.117	0.362	0.005	2226.8	6883.7	102.6	4.5	0.0405	0.045	0.3645	0.0126	20.0412	68.837	8.3106	53.2	150.3888	150	Mn3O4	1.00
Mercury	0.165	4.444	0.095	0.746	0.733326	0.0156575	0.1231725	4.5	0.0405	0.045	0.3645	0.0126	0.006599934	0.00015675	0.00976973	0.000462	0.017195657	0.017	Methyl Mercury	1.00
Molybdenum	32	2.091	0.085	0.010	66.9248	2.7136	0.3232	4.5	0.0405	0.045	0.3645	0.0126	0.023232	0.027136	0.0261792	0.0896	0.7452384	0.74	Chloride	1.00
Nickel	560	5.782	0.156	0.232	3228.2	87.304	129.808	4.5	0.0405	0.045	0.3645	0.0126	29.1438	0.87304	10.51448	1.568	42.09928	42.25	nickel hydroxide	1.00
Acroclor 1254	56	0.625	35.0168	0	0	4.5	0.0405	0.045	0.3645	0.0126	0.3151512	0	0	0	0.1568	0.4719512	0.474	n/a	1.00	
Acroclor 1260	0.88	12.381	5.220	10.89528	0	4.593864	4.5	0.0405	0.045	0.3645	0.0126	0.09805752	0	0.372102984	0.002464	0.472624504	0.474	assumed to be 1254	1.00	
Selenium	5.05	1.395	0.060	0.231	7.044245	0.300475	1.16655	4.5	0.0405	0.045	0.3645	0.0126	0.063398205	0.00300475	0.09449055	0.01414	0.175033505	0.174	Selenate (SeO4)	1.00
Thallium	13	0.000	0.023	0	0.2964	0	4.5	0.0405	0.045	0.3645	0.0126	0	0.002964	0	0.0364	0.039364	0.039	thallium sulfate	1.00	
Uranium	505	0.063	0.002	0.000	31.9665	0.7575	0.1515	4.5	0.0405	0.045	0.3645	0.0126	0.2876985	0.007575	0.0122715	1.414	1.721545	1.722	Uranyl acetate	1.00
Vanadium	231	0.088	0.084	2.377	4211.48	1545.115	4.5	4.5	0.0405	0.045	0.3645	0.0126	0.1833678	0.194271	0	0.6468	1.0244388	1.030	metavanadate (NaVO3)	1.00
Zinc	650	6.479	0.716	0	465.27	0	37.90312	4.6327	0	0.0405	0.045	0.3645	0.0126	125.154315	1.82	169.510335	169.0	zinc oxide	1.00	

**Table A.2. Soil PRG for white-tailed deer assumed to consume 100% plants**

Analyte	Soil Conc (mg/kg)	90th Percentile Plant UF	Estimated Plant Conc (mg/kg)	Body Wt (kg)	Food Ingestion (kg/d)	Soil Ingestion (kg/d)	Food Exposure (mg/kg/d)	Soil Exposure (mg/kg/d)	Total Exposure (mg/kg/d)	LOAEL (mg/kg/d)	Form	HQ
Aluminum	1920	0.030	56.64	56.5	1.74	0.0348	1.744311504	1.18258407	2.92689558	2.930	AlCl3	1.00
Arsenic	119	0.032	3.8437	56.5	1.74	0.0348	0.118372354	0.0732956	0.19166793	0.191	Arsenite	1.00
Barium	700	0.237	165.76	56.5	1.74	0.0348	5.104821239	0.43115044	5.53597168	5.6	barium hydroxide	1.00
Cadmium	77.5	1.115	86.37375	56.5	1.74	0.0348	2.660005752	0.0477345	2.70774027	2.706	cadmium chloride	1.00
Chromium	1380	0.067	91.77	56.5	1.74	0.0348	2.82619115	0.8499823	3.67617345	3.69	Cr+6	1.00
Copper	455	0.381	173.2185	56.5	1.74	0.0348	5.334516637	0.28024779	5.61476442	5.6	copper sulfate	1.00
Lead	10100	0.052	527.22	56.5	1.74	0.0348	16.23650973	6.22088496	22.4573947	22.44	lead acetate	1.00
Lithium	3175	0.034	107.95	56.5	1.74	0.0348	3.324477876	1.9555752	5.2800531	5.3	lithium carbonate	1.00
Manganese	6800	0.362	2463.64	56.5	1.74	0.0348	75.87139115	4.18331838	80.0597097	80	Mn3O4	1.00
Mercury	12.7	0.095	1.2065	56.5	1.74	0.0348	0.03715593	0.007322	0.0449782	0.045	Methyl Mercury Chloride	1.00
Molybdenum	122	0.085	10.3456	56.5	1.74	0.0348	0.318607858	0.0751434	0.39375122	0.39	MoO4	1.00
Nickel	4150	0.156	646.985	56.5	1.74	0.0348	19.92484779	2.55610619	22.480954	22.44	nickel sulfate hexahydrat	1.00
Aroclor 1254	138	0	56.5	1.74	0.0348	0	0.0849982	0.0849982	0.085	n/a	1.00	
Aroclor 1260	138	0	56.5	1.74	0.0348	0	0.0849982	0.0849982	0.085	assumed to be 1254	1.00	
Selenium	38	0.060	2.261	56.5	1.74	0.0348	0.0696308	0.0234053	0.0930361	0.093	Selenate (SeO4)	1.00
Thallium	15.9	0.023	0.36252	56.5	1.74	0.0348	0.01116433	0.009793	0.0209576	0.021	thallium sulfate	1.00
Uranium	1380	0.002	2.07	56.5	1.74	0.0348	0.06374867	0.8499823	0.91373097	0.915	Uranyl acetate	1.00
Vanadium	170	0.084	14.297	56.5	1.74	0.0348	0.440295991	0.10470796	0.54500496	0.547	sodium metavanadate	1.00
Zinc	3950	0.716	2827.41	56.5	1.74	0.0348	87.07421947	2.43292035	89.5071398	89.8	zinc oxide (NaVO3)	1.00

**Table A.3. Soil PRG for white-footed mice assumed to consume 50% plants and 50% worms**

Analytic	Soil Conc (mg/kg)	90th Percentile Worm UF	Estimated Plant UF	Estimated Worm Conc (mg/kg)	Estimated Plant Conc (mg/kg)	Body Wt (kg)	Worm Ingestion (kg/d)	Plant Ingestion (kg/d)	Soil Ingestion (kg/d)	Worm Exposure (mg/kg/d)	Plant Exposure (mg/kg/d)	Soil Exposure (mg/kg/d)	Total Exposure (mg/kg/d)	LOAEL (mg/kg/d)	Form	HQ
Aluminum	1440	0.118	0.030	169.92	42.48	0.022	0.0017	0.00068	13.13018182	3.282545455	4.450909091	20.86563636	20.8556	AlCl3	1.00	
Arsenic	20	0.811	0.032	16.216	0.646	0.022	0.0017	0.00068	1.253054545	0.049918182	0.061818182	1.364790909	1.362	Arsenite	1.00	
Barium	1170	0.160	0.237	187.2	277.056	0.022	0.0017	0.00068	14.46545455	21.40887273	3.616363636	39.49069091	39.5	hydioxide barium	1.00	
Cadmium	33	6.410	1.114	211.5399	36.7785	0.022	0.0017	0.00068	16.346265	2.841975	0.102	19.29024	19.264	cadmium chloride	1.00	
Chromium	40.4	8.331	0.056	336.568336	2.6866	0.022	0.0017	0.00068	26.0755509	0.207600909	0.124872727	26.34002873	26.24	Cr+6	1.00	
Copper	415	0.826	0.381	342.956	157.9905	0.022	0.0017	0.00068	26.50114545	12.20835682	1.282727273	39.99222955	40.0	copper sulfate	1.00	
Lead	8050	0.164	0.052	1316.175	420.21	0.022	0.0017	0.00068	101.7044318	32.47077273	24.88181818	159.0570227	159.77	lead acetate	1.00	
Lithium	1670	0.217	0.034	361.722	56.78	0.022	0.0017	0.00068	27.95124545	4.387545455	5.161818182	37.50060909	37.5	lithium carbonate	1.00	
Manganese	14100	0.117	0.362	1652.52	5108.43	0.022	0.0017	0.00068	127.6947273	394.7423182	43.58181818	566.0188636	567	Mn3O4	1.00	
Mercury	0.9	4.444	0.095	3.99996	0.8855	0.022	0.0017	0.00068	0.309087818	0.006606818	0.002781818	0.318476455	0.320	Mercury Chloride	1.00	
Molybdenum	16.4	2.091	0.085	34.29896	1.39072	0.022	0.0017	0.00068	2.650374182	0.107464727	0.050690909	2.808529818	2.81	MoO4	1.00	
Nickel	345	5.782	0.156	1994.9625	53.7855	0.022	0.0017	0.00068	154.1561932	4.156152273	0.066363636	159.3787091	159.77	nickel sulfate hexahydrate	1.00	
Aroclor 1254	11.8	0.625	7.37854	0	0.022	0.0017	0.00068	0.570159909	0	0.036472727	0.606632636	0.607	n/a	1.00		
Aroclor 1260	0.63	12.381		7.80003	0	0.022	0.0017	0.00068	0.602729591	0	0.001947273	0.604676864	0.607	assumed to be 1254	1.00	
Selenium	5.7	1.395	0.060	7.95093	0.33915	0.022	0.0017	0.00068	0.614390045	0.026207045	0.017618182	0.658215273	0.659	Selenite (SeO4)	1.00	
Thallium	30.8	0.000	0.023	0	0.70224	0.022	0.0017	0.00068	0	0.054264	0.092	0.149464	0.149	thallium sulfate	1.00	
Uranium	800	0.063	0.002	50.64	1.2	0.022	0.0017	0.00068	3.913090909	0.092727273	2.472727273	6.478545455	6.511	Uranyl acetate	1.00	
Vanadium	237	0.088	0.084	20.9034	19.9317	0.022	0.0017	0.00068	1.615262727	1.50176818	0.732545455	3.887985	3.894	Vanadate (NaVO3)	1.00	
Zinc	1140	6.479	0.716	7386.288	816.012	0.022	0.0017	0.00068	570.7386182	63.05547273	3.323636364	637.3377273	639.1	zinc oxide	1.00	

**Table A.4. Soil PRG for short-tailed shrews assumed to consume 100% worms**

Analyte	Soil Conc (mg/kg)	90th Percentile Worm UF	Estimated Worm Conc (mg/kg)	Body Wt (kg)	Food Ingestion (kg/d)	Soil Ingestion (kg/d)	Food Exposure (mg/kg/d)	Soil Exposure (mg/kg/d)	Total Exposure (mg/kg/d)	LOAEL (mg/kg/d)	Form	HQ
Aluminum	155	0.118	18.29	0.015	0.009	0.00117	10.974	12.09	23.064	22.952	AlCl3	1.00
Arsenic	2.66	0.811	2.156728	0.015	0.009	0.00117	1.2940368	0.20748	1.5015168	1.498	Arsenite	1.00
Barium	250	0.160	40	0.015	0.009	0.00117	24	19.5	43.5	43.5	barium hydroxide	1.00
Cadmium	5.4	6.410	34.61562	0.015	0.009	0.00117	20.769372	0.4212	21.190572	21.200	cadmium chloride	1.00
Chromium	5.7	8.331	47.48613	0.015	0.009	0.00117	28.491678	0.4446	28.936278	28.88	Cr+6	1.00
Copper	77	0.826	63.6328	0.015	0.009	0.00117	38.17968	6.006	44.18568	44.0	copper sulfate	1.00
Lead	1000	0.164	163.5	0.015	0.009	0.00117	98.1	78	176.1	175.83	lead acetate	1.00
Lithium	199	0.217	43.1034	0.015	0.009	0.00117	25.86204	15.522	41.38404	41.3	lithium carbonate	1.00
Manganese	4200	0.117	492.24	0.015	0.009	0.00117	295.344	327.6	622.944	624	Mn3O4	1.00
Mercury	0.128	4.444	0.5688832	0.015	0.009	0.00117	0.34132992	0.009984	0.35131392	0.352	Mercury Chloride	1.00
Molybdenum	2.33	2.091	4.872962	0.015	0.009	0.00117	2.9237772	0.18174	3.1055172	3.09	MoO4	1.00
Nickel	49.5	5.783	286.23375	0.015	0.009	0.00117	171.74025	3.861	175.60125	175.83	hexahydrate sulfate	1.00
Aroclor 1254	1.47	0.625	0.919191	0.015	0.009	0.00117	0.5515146	0.11466	0.6661746	0.6668	n/a	1.00
Aroclor 1260	0.089	12.381	1.101909	0.015	0.009	0.00117	0.6611454	0.006942	0.6680874	0.6688	be 1254 assumed to	1.00
Selenium	0.79	1.395	1.101971	0.015	0.009	0.00117	0.6611826	0.06162	0.7228026	0.725	Selenate (SeO4)	1.00
Thallium	2.1	0.000	0	0.015	0.009	0.00117	0	0.1638	0.1638	0.164	thallium sulfate	1.00
Uranium	62	0.063	3.9246	0.015	0.009	0.00117	2.35476	4.836	7.19076	7.165	Uranyl acetate	1.00
Vanadium	32.6	0.088	2.87532	0.015	0.009	0.00117	1.725192	2.5428	4.267992	4.285	metavanadat e (NaVO3)	1.00
Zinc	177	6.479	1146.8184	0.015	0.009	0.00117	688.09104	13.806	701.89704	703.3	zinc oxide	1.00

**Table A.5. Soil PRG for american woodcock assumed to consume 100% worms**

Analyte	Soil Conc (mg/kg)	90th Percentile Worm UF	Estimated Worm Conc (mg/kg)	Body Wt (kg)	Food Ingestion (kg/d)	Soil Ingestion (kg/d)	Food Exposure (mg/kg/d)	Soil Exposure (mg/kg/d)	Total Exposure (mg/kg/d)	LOAEL (mg/kg/d)	Form	HQ
Aluminum	0.118	0	0.198	0.15	0.0156	0	0	0	0	0	Al2(SO4)2	#DIV/0!
Arsenic	18.5	0.811	14.9998	0.198	0.15	0.0156	11.36348485	1.457575758	12.82106061	12.8	sodium arsenite	1.00
Barium	208	0.160	33.28	0.198	0.15	0.0156	25.21212121	16.38787879	41.6	41.7	barium hydroxide	1.00
Cadmium	4.05	6.410	25.961715	0.198	0.15	0.0156	19.66796591	0.319090909	19.98705682	20.00	cadmium chloride	1.00
Chromium	0.78	8.331	6.498102	0.198	0.15	0.0156	4.922804545	0.061454545	4.984259091	5.00	Cr+3 as CrK(SO4)2	1.00
Copper	87.5	0.826	72.31	0.198	0.15	0.0156	54.78030303	6.893939394	61.67424242	61.7	copper oxide	1.00
Lead	56	0.164	9.156	0.198	0.15	0.0156	6.936363636	4.412121212	11.34848485	11.30	lead acetate	1.00
Lithium	0.217	0	0.198	0.15	0.0156	0	0	0	0	0	#DIV/0!	#DIV/0!
Manganese	0.117	0	0.198	0.15	0.0156	0	0	0	0	0	Mn3C4	#DIV/0!
Mercury	0.0185	4.444	0.0822214	0.198	0.15	0.0156	0.062288939	0.001457576	0.063746515	0.064	Methyl Mercury	
Molybdenum	21.3	2.091	44.54682	0.198	0.15	0.0156	33.74759091	1.678181818	35.42577273	35.30	Dicyandiamide	1.00
Nickel	24	5.782	138.78	0.198	0.15	0.0156	105.1363636	1.890909091	107.02727277	107.00	sodium molybdate (MoO4)	1.00
PCB-1254	3.277	0.625	2.044731	0.198	0.15	0.0156	1.549038636	0.257636364	1.806675	1.800	n/a	1.00
PCB-1260	0.19	12.381	2.35239	0.198	0.15	0.0156	1.782113636	0.014969697	1.797083333	1.800	assumed to be 1254	1.00
Selenium	0.88	1.395	1.227512	0.198	0.15	0.0156	0.929933333	0.069333333	0.999266667	1.000	sodium selenite	1.00
Thallium	0.00	0	0.198	0.15	0.0156	0	0	0	0	0	#DIV/0!	
Uranium	0.063	0	0.198	0.15	0.0156	0	0	0	0	0	depleted metallic U	#DIV/0!
Vanadium	0.088	0	0.198	0.15	0.0156	0	0	0	0	0	vanadyl sulfate	#DIV/0!
Zinc	26.3	6.479	170.40296	0.198	0.15	0.0156	129.0931515	2.072121212	131.1652727	131.0	zinc sulfate	1.00

**Table A.6. Soil PRG for red-tailed hawk assumed to consume 100% small mammals**

Analyte	Soil Conc (mg/kg)	90th Percentile Mammal U/F	Estimated Mammal Conc (mg/kg)	Body Wt (kg)	Food Ingestion (kg/d)	Soil Ingestion (kg/d)	Food Exposure (mg/kg/d)	Soil Exposure (mg/kg/d)	Total Exposure (mg/kg/d)	LOAEL (mg/kg/d)	Form	HQ
Aluminum	0.014	0	1.126	0.109	0	0	0	0	0	0.0	Al2(SO4)2	#DIV/0!
Arsenic	16500	0.008	132	1.126	0.109	0	12.77797513	0	12.77797513	12.8	sodium arsenite	1.00
Barium	7000	0.061	429.8	1.126	0.109	0	41.60586146	0	41.60586146	41.7	hydroxide	1.00
Cadmium	1570	0.132	206.612	1.126	0.109	0	20.00062877	0	20.00062877	20.00	chloride	1.00
Chromium	233	0.221	51.5396	1.126	0.109	0	4.989179751	0	4.989179751	5.00	Cr+3 as CrK(SO4)2	1.00
Copper	860	0.740	636.572	1.126	0.109	0	61.62197869	0	61.62197869	61.7	copper oxide	1.00
Lead	2630	0.045	117.035	1.126	0.109	0	11.3293206	0	11.3293206	11.30	lead acetate	1.00
Lithium	0.033	0	0	1.126	0.109	0	0	0	0	0	Mn3O4	#DIV/0!
Manganese	0.005	0	0	1.126	0.109	0	0	0	0	0	Methyl Mercury	#DIV/0!
Mercury	0.89	0.746	0.664385	1.126	0.109	0	0.064314356	0	0.064314356	0.064	Dicyandiamide	1.00
Molybdenum	36000	0.010	363.6	1.126	0.109	0	35.19751332	0	35.19751332	35.30	sodium molybdate	1.00
Nickel	4750	0.232	1101.05	1.126	0.109	0	106.5847691	0	106.5847691	107.00	(MoO4)	1.00
PCB-1254		0	0	1.126	0.109	0	0	0	0	1.800	n/a	0.00
PCB-1260	3.55	5.220	18.532065	1.126	0.109	0	1.793956559	0	1.793956559	1.800	assumed to be 1254	1.00
Selenium	44.5	0.231	10.2795	1.126	0.109	0	0.993084813	0	0.993084813	1.000	sodium selenite	1.00
Thallium		0	0	1.126	0.109	0	0	0	0	0	depleted metallic U	#DIV/0!
Uranium	0.000	0	0	1.126	0.109	0	0	0	0	0.0	vanadyl sulfate	#DIV/0!
Vanadium		0	0	1.126	0.109	0	0	0	0	0.000	zinc sulfate	1.00
Zinc	570	2.377	1354.947	1.126	0.109	0	131.1627202	0	131.1627202	131.0		

## DISTRIBUTION

- 1. L. V. Asplund
- 2. R. R. Bonczek
- 3. M. Clauberg
- 4. J. Croswait
- 5. J. R. Duncan
- 6-8. R. E. Efroymson
- 9. K. M. Golden
- 10. R. K. Holmes
- 11. J. Jagoe
- 12-14. D. S. Jones
- 15-29. S. B. Kerr
- 30. J. M. Loar
- 31. D. M. Matteo
- 32. C. W. McGinn
- 33. D. B. Miller
- 34. J. H. Moyer
- 35. B. D. Nourse
- 36. D. M. Opresko
- 37. P. T. Owen
- 38. S. T. Purucker
- 39. A. S. Quist
- 40-42. B. E. Sample
- 43. D. M. Steinhauff
- 44-46. G. W. Suter
- 47. S. S. Talmage
- 48. M. F. Tardiff
- 49. C. J. E. Welsh
- 50. D. C. White
- 51. M. E. Will
- 52. ER Document Management Center—RC

- 53. A. Temeshy, Bechtel National Inc., 151 Lafayette Dr., Oak Ridge, TN 37831-0350
- 54. M. Leslie, CDM Federal Programs Corporation, 800 Oak Ridge Turnpike, Suite 500, Oak Ridge, TN 37830
- 55. D. Gonzales, Jacobs Engineering, 125 Broadway, Oak Ridge, TN 37830
- 56. G. Snyder, Jacobs, Engineering, 100 West 3rd St., Piketon, OH 45611
- 57. J. Tate, Jacobs, Engineering, 125 Broadway, Oak Ridge, TN 37830
- 58. D. Wilkes, Jacobs, Engineering, 175 Freedom Blvd., Kevil, KY 42053
- 59. P. Carter, Science Applications International Corporation, 11197 U.S. Route 23, Suite 200, Waverly, Ohio 45690
- 60. B. Kornaby, Science Applications International Corporation, P.O. Box 2501, 301, Laboratory Rd., Oak Ridge, TN 37831
- 61. S. Pack, Science Applications International Corporation, P.O. Box 2501, 301, Laboratory Rd., Oak Ridge, TN 37831
- 62. B. G. Blaylock, SENES, 102 Donner Drive, Oak Ridge, TN 37831
- 63. K. Hopper, Systematic Management Services, 117 Broadway Ave., Oak Ridge, TN 37830



**Please do not forward or discard this document.**

If this address is not correct for the designated  
addressee, please return this document to the

**ER Central Document Management Center  
Building K-1002, MS 7243  
Martin Marietta Energy Systems, Inc.  
P.O. Box 2003  
Oak Ridge, TN 37831**