

LA-UR- 99-520

*Approved for public release;
distribution is unlimited.*

CONF- 990303 --

Title: CLASSIFICATION OF LUNAR TERRANES USING
NEUTRON AND THORIUM GAMMA-RAY DATA

Author(s): William C. Feldman, NIS-1
David J. Lawrence, NIS-1
Sylvestre Maurice, Observatoire Midi-Pyrénées
Richard C. Elphic, NIS-1
Bruce I. Barraclough, NIS-1
A.B. Binder, Lunar Research Institute
P.G. Lucey, Hawai'i Institute of Geophysics and Planetology

Submitted to: Lunar and Planetary Science Conference
Houston, Texas
March 15-19, 1999

RECEIVED
APR 13 1999
OSTI

MASTER

Los Alamos
NATIONAL LABORATORY

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-36. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

CLASSIFICATION OF LUNAR TERRANES USING NEUTRON AND THORIUM GAMMA-RAY DATA.

W. C. Feldman¹, D. J. Lawrence¹, S. Maurice², R. C. Elphic¹, B. L. Barraclough¹, A. B. Binder³, and P. G. Lucey⁴, Los Alamos National Laboratory, Group NIS-1, MS D466, Los Alamos, NM 87545 (wfeldman@lanl.gov),
²Observatoire Midi-Pyrénées, Toulouse, FRANCE, ³Lunar Research Institute, 1180 Sunrise Dr., Gilroy, CA 95020 USA, ⁴Hawai'i Institute of Geophysics and Planetology, University of Hawai'i, Manoa, HI 96822 USA.

Introduction: A major scientific goal of the Lunar Prospector (LP) gamma-ray and neutron spectrometers is to classify all lunar terranes according to composition. A preliminary analysis of early data indicates this goal will be met for the major rock-forming elements on a spatial scale of about 200 km. The low-altitude phase of LP now in progress should allow reduction of this scale by about a factor of 10 for those elements that have sufficiently high measurable fluxes relative to their backgrounds. Most promising are the flux intensities of thermal, epithermal, and fast neutrons (which each average about 300 counts per 50 km of ground track) and 2.6 MeV gamma rays from thorium (which averages about 50 counts per 50 km of ground track). We therefore explore the information content of these measurables to classify the various lunar terrane types.

Theoretical Expectations: Computer simulations show that the ratio of lunar epithermal to thermal neutron flux is directly proportional to the macroscopic absorption cross section [1]. Estimates of these cross sections for the various Apollo and Luna samples [2] show that abundances of Fe, Ti, Gd, and Sm dominate [3,4]. Because the thermal neutron absorption cross section for Ti is about twice that for Fe, the ratio of epithermal to thermal neutron flux should primarily reflect the content of Fe+2Ti, with significant contributions from Gd+Sm in KREEP-rich terranes.

Simulations of lunar fast neutron fluxes (Gasnault et al., in preparation, 1999) show that their variation reflects only variations in Fe and Ti, in equal proportions. A combined analysis of all three neutron energy ranges should then be sufficient to separate abundance variations in Fe from those of Ti and Gd+Sm. Separation of these last two components can be implemented by adding an analysis of thorium gamma rays because of the measured strong correlation amongst all incompatible elements [2].

Results: Equal area $2^\circ \times 2^\circ$ maps of thorium abundances and of fast-neutron fluxes have been published previously [5,6]. The map for the ratio of epithermal to thermal neutrons is similar to these maps but is not shown here. All three maps show relatively high

intensities within the front-side Mare, smaller, but somewhat enhanced intensities within the South Pole Aitken basin, and very low intensities in the high-langs.




Figure 1: Ternary diagram for a classification of 2° by 2° equal area spatial pixels on the Moon according to measured epithermal/thermal and fast neutrons fluxes and to the LP GRS thorium abundances. Large red circles give measurements within Mare Humorum.




Figure 2: Ternary diagram for a classification of 2° by 2° equal area spatial pixels on the Moon according to measured epithermal/thermal and fast neutrons fluxes and to the LP GRS thorium abundances. Large red circles give measurements within Mare Marginis.

These three data sets were combined into a ternary diagram by first normalizing each measurable to the range spanning from 0 to 1 for each $2^\circ \times 2^\circ$ spatial

pixel on the Moon. Resultant triplets of normalized counts were then renormalized so that they add in quadrature to unity.

One ternary diagram for a region relatively high in Ti (Humorum) is shown in Fig. 1, and one for a region relatively low In Ti (Marginis) is shown in Fig. 2. Plotted in red are spatial locations within the selected region while those in black are for the rest of the Moon. A close inspection reveals the following: 1) Locations within LP ternary diagrams for a given region, cluster about well-defined centroids that vary from region to region. 2) Comparison of the horizontal positions of these centroids with Ti abundances as determined using Clementine Spectral Reflectance (CSR) data [7] shows that regions of higher Ti content locate closer to the Epithermal/Thermal neutron vertex, and those having lower Ti content locate closer to the Fast neutron vertex.

D.J. et al., (1998) Science, 281, 1484. [6] Feldman et al., (1998) Science, 281, 1489. [7] Lucey, P.G., (1998) JGR 103, 3679.




Figure 3: Correlation between titanium abundances determined using Clementine Spectral Reflectance data and the difference between normalized fast and epithermal/thermal neutron fluxes. The correlation line has significance given by $R=-0.4$

This trend is quantified in Fig. 3 by the correlation between Ti abundances as determined using the CSR data and the horizontal centroid positions in LP ternary diagrams. Inspection shows a weak, although significant correlation. Reasons for the substantial scatter that is very evident at low values of the <Fast - Epithermal/Thermal> neutron flux are not known but need to be investigated.

References: [1] Feldman, W.C. et al., (1991) GRL, 18, 2157. [2] Haskin, L., and Warren, P., (1991) in 'Lunar Sourcebook' Heiken, G., Vaniman, D., and French, B.M. eds., pp 357-474. [3] Elphic, R.C. et al., (1998) Science, 281,1493. [4] Lingenfelter, R. et al., (1972) Earth Planet. Sci. Lett. 16, 355. [5] Lawrence,