

JUN 14 1963

UCRL-10750

MASTER

University of California

Ernest O. Lawrence
Radiation Laboratory

ADIABATIC CHARGED-PARTICLE MOTION

Berkeley, California

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

ADIABATIC CHARGED-PARTICLE MOTION

Theodore G. Northrop

April 2, 1963

ADIABATIC CHARGED-PARTICLE MOTION

Theodore G. Northrop

Lawrence Radiation Laboratory
University of California
Berkeley, California

April 2, 1963

Abstract. The adiabatic theory of charged-particle motion is developed systematically in this review. We present the essentials of the theory without giving all the analysis in detail. The general expressions for guiding-center motion and particle energy change are given, with application to the Van Allen radiation and to Fermi acceleration. It is shown that Fermi acceleration and betatron acceleration should not be regarded as distinct processes. Modifications of the nonrelativistic theory necessary when the particle is relativistic are discussed. Proofs are given of the invariance to lowest order of the first and second adiabatic invariants for the case of static fields. Finally, applications are made to the theory of plasmas.

ADIABATIC CHARGED-PARTICLE MOTION¹

Theodore G. Northrop

Lawrence Radiation Laboratory
University of California
Berkeley, California

April 2, 1963

1. INTRODUCTION

The adiabatic approximation to charged-particle motion has been widely used in our attempts to understand the Van Allen radiation and to predict the results of high altitude nuclear explosions. It has also been used extensively in the theory of plasma confinement and stability in strong magnetic fields. A thorough understanding of the adiabatic predictions is therefore desirable, particularly since deviations from these predictions may be important in explaining what we observe. Our purpose in this review is to present what adiabatic theory says, without presenting all of the analysis in the greatest possible generality. Some of the analysis, especially for relativistic particles in time-dependent fields, becomes quite lengthy and will be omitted.

¹ This work was performed under the auspices of the U. S. Atomic Energy Commission.

† Many of the subjects presented here are amplified in a monograph by the author [Northrop, 1963].

2. THE GUIDING CENTER MOTION OF NONRELATIVISTIC PARTICLES

In a uniform magnetic field that is constant in time a charged particle moves in a helical path. The motion may be described exactly as motion about a circle whose center is moving along a line of force. If the field is not quite uniform and not quite time independent, one expects that the motion will not be quite helical; one also expects that something approximating helical motion will still be discernible, and therefore that a good approximation will contain gyration about a center that now may move at right angles to the line of force as well as along it. This expectation is indeed correct, and the equations governing this "guiding center" motion can be derived by following one's physical intuition. To do this let $\vec{r} = \vec{R} + \vec{p}$, where the vectors are defined in Figure 1. To correspond to the picture of rapid gyration about the guiding center, let $\vec{p} = p(\hat{e}_2 \sin \omega t + \hat{e}_3 \cos \omega t)$, where ω is the angular frequency of gyration $eB(\vec{R})/mc$, $B(\vec{R})$ is the magnetic field at \vec{R} , and $\hat{e}_2(\vec{R})$ and $\hat{e}_3(\vec{R})$ are unit vectors perpendicular to $\vec{B}(\vec{R})$ and to each other. If $\vec{R} + \vec{p}$ is now substituted into the equation of motion for the particle

$$\ddot{\vec{r}} = \frac{e \vec{r}}{c} \times \vec{B}(\vec{r}) + e \vec{E}(\vec{r}) \quad (1)$$

and an average is taken over a period of the gyration, the result after a little algebra with the unit vectors is [Hellwig, 1955; Northrop, 1961]

$$\ddot{\vec{R}} = \frac{e}{m} \left[\vec{E}(\vec{R}) + \frac{\vec{R}}{c} \times \vec{B}(\vec{R}) \right] - \frac{M}{m} \nabla B(\vec{R}) + \text{terms proportional to } \frac{m}{e} \quad (2)$$

-3-

Here M is the well-known magnetic moment $e\vec{p}^2\omega/2c = mv_{\perp}^2/2B$, where v_{\perp} is the particle velocity perpendicular to $\vec{B}(\vec{R})$. In (2) only terms through zero order in m/e have been kept; m/e can be used as the expansion parameter because if (1) is written in suitable dimensionless form, the dimensionless parameter that appears is the gyration radius divided by the dimensions of the system, and m/e is proportional to this ratio.

The component of \vec{R} perpendicular to $\vec{B}(\vec{R})$ in (2) is the guiding center velocity perpendicular to $\vec{B}(\vec{R})$. It is the so-called "drift velocity" and is obtained by taking the vector product of (2) with \vec{B} . We have

$$\dot{\vec{R}}_{\perp} = \frac{c\vec{E} \times \hat{\vec{e}}_1}{B} + \frac{Mc}{e} \frac{\hat{\vec{e}}_1 \times \nabla B}{B} + \frac{mc}{e} \frac{\hat{\vec{e}}_1 \times \ddot{\vec{R}}}{B} + O(\epsilon^2) \quad (3)$$

where ϵ is m/e , $\hat{\vec{e}}_1$ is \vec{B}/B , and all field quantities are evaluated at \vec{R} . There are three drift terms here. The first is the well-known " $\vec{E} \times \vec{B}$ " drift, and the second is the "gradient B " drift. The third term contains the "line curvature" drift, but it also contains quite a few other drifts, as will be developed below. All the drifts occur because the curvature of the particle trajectory is alternately larger and smaller as the particle goes around its "circle" of gyration; the gyration "circle" is not really quite a circle. This variation in the curvature produces a gradual drift to one side as illustrated in Figure 2. The cause of the alternately large and small curvature is different for each of the drifts. The " $\vec{E} \times \vec{B}$ " and " ∇B " drifts have been frequently described before [Alfvén, 1950; Spitzer, 1952]. The

six drifts that are contained in the last term of (3) also can be given geometric interpretations. That term may be expanded by writing \vec{R} as

$$\frac{d}{dt}(\vec{R}_\perp + \hat{e}_1 \vec{R} \cdot \hat{e}_1) = \hat{e}_1 \frac{dv_{||}}{dt} + v_{||} \frac{d\hat{e}_1}{dt} + \frac{d\vec{R}_\perp}{dt}, \text{ where } v_{||} \text{ is}$$

$\vec{R} \cdot \hat{e}_1(\vec{R})$, the component of guiding center velocity parallel to the line of force at \vec{R} . We only need $d\vec{R}_\perp/dt$ to zero order in ϵ , since the entire term is multiplied by ϵ in (3). By iteration of (3), we obtain $d\vec{R}_\perp/dt = \vec{u}_E(\vec{R})/dt + O(\epsilon)$, where \vec{u}_E is $c\vec{E} \times \hat{e}_1/B$.

Also, $d\hat{e}_1(\vec{R})/dt$ is needed. It is the rate of change of the unit vector as one follows the guiding center. This unit vector changes direction in a time-dependent magnetic field even in the absence of guiding center motion. In addition the guiding center sees a change in \hat{e}_1 as it moves in a field whose direction in space is not constant. Consequently, the total derivative $d\hat{e}_1/dt$ equals

$\frac{d\hat{e}_1}{dt} + v_{||} \frac{d\hat{e}_1}{ds} + \vec{u}_E \cdot \nabla \hat{e}_1 + O(\epsilon)$, where s is distance along the line of force. Similarly $d\vec{u}_E/dt$ equals $\vec{d}\vec{u}_E/dt + v_{||} \frac{d\vec{u}_E}{ds} + \vec{u}_E \cdot \nabla \vec{u}_E$.

With these substitutions, the total drift velocity becomes

$$\begin{aligned} \vec{R}_\perp &= \frac{\hat{e}_1}{B} \times \left(-c\vec{E} + \frac{Mc}{e} \nabla B + \frac{mc}{e} v_{||} \frac{d\hat{e}_1}{dt} + \frac{mc}{e} \frac{d\vec{u}_E}{dt} \right) \\ &= \frac{\hat{e}_1}{B} \times \left\{ -c\vec{E} + \frac{Mc}{e} \nabla B + \frac{mc}{e} \right. \\ &\quad \left. \left[v_{||} \frac{d\hat{e}_1}{dt} + v_{||}^2 \frac{d\hat{e}_1}{ds} + v_{||} \vec{u}_E \cdot \nabla \hat{e}_1 + \frac{\vec{d}\vec{u}_E}{dt} + v_{||} \frac{\vec{d}\vec{u}_E}{ds} + \vec{u}_E \cdot \nabla \vec{u}_E \right] \right\} + O(\epsilon^2), \end{aligned} \quad (4)$$

-5-

where all quantities are evaluated at \vec{R} . The term proportional to $\partial \hat{e}_1 / \partial s$ is the well-known "line curvature" drift. However, the other five terms in the square bracket, although possibly less familiar, should not be overlooked. In practical cases the electric fields are often so small that the four terms containing \vec{u}_E are negligible, and the field lines may change direction so slowly that the $\partial \hat{e}_1 / \partial t$ drift is small. But these five terms in the bracket are not necessarily small, and situations where each is of primary importance are known in plasma physics. For example, the term proportional to $\vec{u}_E \cdot \nabla \vec{u}_E$ is responsible for the shear, or Helmholtz, instability of a plasma [Northrop 1956, 1961]. Shears occur at the solar wind-geomagnetic field interface, where the solar plasma slides over the geomagnetic field.

The $\partial \hat{e}_1 / \partial t$ drift is an easy one to understand geometrically. If the direction of the magnetic field changes without a change in the particle velocity, then some of what was "parallel" velocity will become "perpendicular," and vice versa. In other words, if there is a change in the reference direction, with respect to which one defines parallel and perpendicular, then the respective components of velocity will change. It is easy to work out the details and see that there is a periodic variation (at the gyration frequency) in the curvature of the particle trajectory while the line of force changes direction. This leads to a drift, just as in the more familiar case of the $\vec{E} \times \vec{B}$ and ∇B drifts.

The component of (2) parallel to the magnetic field gives the parallel acceleration of the guiding center. The scalar product of (2) with $\hat{e}_1(\vec{R})$ is

$$\ddot{\vec{R}} \cdot \hat{\vec{e}}_1 = \frac{e}{m} \vec{E}_{||} - \frac{M}{m} \hat{\vec{e}}_1 \cdot \nabla B + O(\epsilon), \quad (5)$$

where $\vec{E}_{||}$ is $\vec{E}(\vec{R}) \cdot \hat{\vec{e}}_1(\vec{R})$. The parallel acceleration $dv_{||}/dt$ is $\frac{d}{dt}(\vec{R} \cdot \hat{\vec{e}}_1)$, which differs from $\ddot{\vec{R}} \cdot \hat{\vec{e}}_1$ by $\vec{R} \cdot \frac{d\hat{\vec{e}}_1}{dt}$; and since the latter equals $(\hat{\vec{e}}_1 \cdot \vec{v}_{||} + \vec{u}_E) \cdot \frac{d\hat{\vec{e}}_1}{dt} + O(\epsilon)$, then

$$\frac{dv_{||}}{dt} = \frac{e}{m} \vec{E}_{||} - \frac{M}{m} \frac{\partial B}{\partial s} + \vec{u}_E \cdot \frac{d\hat{\vec{e}}_1}{dt} + O(\epsilon). \quad (6)$$

The term $v_{||} \hat{\vec{e}}_1 \cdot \frac{d\hat{\vec{e}}_1}{dt}$ vanished because $\hat{\vec{e}}_1$ is a unit vector. The term $-(M/m)(\partial B/\partial s)$ is the usual mirror effect that produces reflection of particles and makes them oscillate north and south in the geomagnetic field, thus trapping them. The total time derivative $d\hat{\vec{e}}_1/dt$ may be expanded to $(\partial \hat{\vec{e}}_1/\partial t) + (v_{||} \partial \hat{\vec{e}}_1/\partial s) + \vec{u}_E \cdot \nabla \hat{\vec{e}}_1$, just as in the drift equation. This $\vec{u}_E \cdot \frac{d\hat{\vec{e}}_1}{dt}$ term is another example of an effect caused by a change in the reference direction. If the electric field is small, the term may be negligible.

3. ENERGY CHANCES

The kinetic energy W of a particle, averaged over a gyration is $(mv_{||}^2/2) + (mu_E^2/2) + MB$. This may be demonstrated, but it is really obvious: the first two terms are the energy of the guiding center motion and MB is the energy of rotation about the guiding center. The parallel energy $W_{||}$ is $mv_{||}^2/2$, and the average perpendicular energy W_{\perp} is $(mu_E^2/2) + MB$. The rate of change dW/dt of total kinetic energy, averaged over a gyration, can be deduced in a

formal fashion, but the result is so intuitively correct that the procedure will be omitted here. The result is

$$\frac{1}{e} \frac{dW}{dt} = \dot{\vec{R}} \cdot \vec{E}(\vec{R}, t) + \frac{M}{e} \frac{dB}{dt}(\vec{R}, t) + O(\epsilon^2), \quad (7)$$

where $\dot{\vec{R}}$ is $\hat{e}_1 v_{||} + \vec{R}_{\perp}$. The first term on the right side is energy increase resulting from the average particle motion in the electric field, while the second term is the induction effect or "betatron acceleration" caused by the curl of \vec{E} acting about the circle of gyration. Part of the energy increase given by (7) is fed into the parallel energy, and the rest into perpendicular energy. Simultaneously, energy is exchanged between parallel and perpendicular components by the mirror effect, the exchange occurring without a change in total kinetic energy. The process may be visualized as in Figure 3, where the partition of dW/dt between dW_{\perp}/dt and $dW_{||}/dt$ comes from the formal analysis. Note that $M dB/dt$ is only part of the perpendicular energy increase; $e\vec{R} \cdot \vec{E}$ contains the rest of the perpendicular energy increase plus the entire rate of increase of parallel energy.

4. FERMI ACCELERATION

Fermi acceleration [Fermi 1949, 1954; Teller 1954; Davis 1956; Parker 1958] is a special case of the adiabatic energy change of the preceding section. Fermi suggested that repeated collisions between a charged particle and moving clumps of magnetized plasma in space would

accelerate a few particles to extreme energies. In effect, the clumps act as massive particles with which the high energy particles attempt to establish kinetic equilibrium. The many particles in a clump, although of low energy, give it a very large mass. Thus, at thermal equilibrium, the high energy particles will have very high energies indeed. The statistics of these collisions will not be discussed here [see Teller 1954]; instead, details of a single Fermi-type collision will be interpreted in light of the preceding section.

Equation (7) applies to any adiabatic situation, but Fermi had in mind special ones--namely, those where there is a frame of reference (that of the clump) in which the magnetic field is static and there is no electric field. In the frame of the clump there is therefore no energy gain or loss by the particle. The collision is elastic and its net effect is to alter the velocity of the guiding center. In the earth's frame, with respect to which the clump is in motion, there may be an energy change, somewhat in analogy to a ball struck by a baseball bat. A particle will lose energy if the clump is overtaken by the particle, and it will gain if the clump overtakes the particle.

Suppose the earth is fixed at O in Figure 4 and that the clump is fixed in a frame O^* moving at velocity \vec{u} with respect to the earth. The rate of energy gain is, from (7) and (4)

$$\begin{aligned}\frac{dW}{dt} &= ev_{||} E_{||} + e\vec{R}_\perp \cdot \vec{E} + M \frac{\partial B}{\partial t} \\ &= ev_{||} E_{||} + M\vec{u}_E \cdot \nabla B + mv_{||} \vec{u}_E \cdot \frac{d\vec{E}}{dt} + m\vec{u}_E \cdot \frac{d\vec{u}_E}{dt} + M \frac{\partial B}{\partial t}.\end{aligned}\tag{8}$$

-9-

Quantities in (8) must now be expressed in terms of \vec{u} . For example, the electric field seen in O is $-\frac{\vec{u}}{c} \times \vec{B}^* \approx -\frac{\vec{u}}{c} \times \vec{B}$. The magnetic fields \vec{B} and \vec{B}^* are equal through order \vec{u}/c , i.e., nonrelativistically. The actual cosmic problem may have relativistic clump velocities, and relativistic energy for the colliding particle. Relativistic adiabatic motion will be reviewed in the next section, but the nonrelativistic case is adequate here for illustrative purposes.

The following relations also hold, as seen from the earth's frame of reference:

$$\frac{\partial \vec{B}}{\partial t} = -\vec{u} \cdot \nabla \vec{B} ,$$

$$\frac{d\hat{e}_1}{dt} = (v_{||} - u_{||}) \frac{\partial \hat{e}_1}{\partial s} ,$$

and

$$\vec{u}_E \cdot \frac{d\vec{u}_E}{dt} = -(v_{||} - u_{||}) u_{||} \vec{u}_\perp \cdot \frac{d\hat{e}_1}{ds} .$$

Substitution into (8) gives

$$\frac{1}{e} \frac{dW}{dt} = -\frac{M}{e} u_{||} \frac{\partial \vec{B}}{\partial s} + \frac{m}{e} (v_{||} - u_{||})^2 \vec{u}_\perp \cdot \frac{d\hat{e}_1}{ds} + O(\epsilon^2) . \quad (9)$$

If the magnetic field in the clump is such that the guiding center moves along a straight line of force, the last term in (9) is zero, and one then has what Fermi named "type a" acceleration. As seen from the clump frame, the particle moves into an increasing magnetic field (magnetic mirror) along a straight line of force, and reflects with no energy change. As viewed from the earth's frame there will be an energy change.

-10-

On the other hand if the field line along which the guiding center moves is curved, and if the magnitude of the field is constant along the line, the first term on the right side of (9) vanishes. The last term is then Fermi's "type b" acceleration. In either case, (9) may be integrated with respect to time to give the total energy change produced by the particle's collision with the clump. Types a and b really differ only in the mechanism whereby the guiding center velocity is reversed in the clump frame. In either case the energy change seen by the observer on the earth is $2\mu u_{||} (v_{||} - u_{||})$, where $v_{||}$ is the component of guiding center velocity parallel to the magnetic field after the collision (i.e., far from the clump) and $u_{||}$ is the component of \vec{u} parallel to that field. This energy change is naturally more easily obtained from the fact that the velocity in the static frame is merely reversed by the collision. But our purpose here has been to apply (7) in the frame of reference in which there is an energy change. Equation (9) can also be integrated over a collision without breaking it up into the special cases "a" and "b".

Fermi acceleration and betatron acceleration are sometimes invoked as distinct processes whereby a particle gains energy. However, they are not distinct. If one follows the fate of the $(M/e) \frac{\partial B}{\partial t}$ term in the transition from (8) to (9), he finds the term goes into forming $-\frac{M}{e} u_{||} (\frac{\partial B}{\partial s})$, which is the "type a" acceleration. Consequently, betatron acceleration should not be viewed as a process distinct from Fermi acceleration, since it is part of "type a". It is correct to distinguish between betatron acceleration and acceleration resulting

-11-

from guiding center motion in the electric field, since these appear as distinct terms in (9).

Pure betatron acceleration in space is improbable, since if there is a $\partial B / \partial t$, there will usually be an electric field at the guiding center, and the $\vec{R} \cdot \vec{E}$ term in (7) will be nonvanishing.

5. RELATIVISTIC ADIABATIC MOTION

If the particle has relativistic energy, (1) is replaced by

$$\frac{d\vec{p}}{dt} = \frac{d}{dt} \frac{m_0 \vec{r}}{(1 - \beta^2)^{1/2}} = \frac{e}{c} \vec{r} \times \vec{B}(\vec{r}) + e\vec{E}(\vec{r}), \quad (10)$$

where \vec{p} is the momentum, $\beta = v/c$, and m_0 is the rest mass. Three cases can be distinguished: when the electric field is zero, when its component \vec{E}_\perp perpendicular to \vec{B} is small, and when \vec{E}_\perp is large.

If there is no electric field, the force on the particle is always at right angles to the velocity, with the result that the energy is constant. Then $m_0/(1 - \beta^2)^{1/2}$ can be removed from under the d/dt in (10) and the equation is identical with the nonrelativistic one for a particle of mass $m_0/(1 - \beta^2)^{1/2}$. All the preceding nonrelativistic theory, with \vec{E} set equal to zero, now applies. In the following two equations the nonrelativistic guiding center equations are rewritten with $m_0/(1 - \beta^2)^{1/2}$ replacing m . The drift velocity is

$$\vec{R}_\perp = \frac{1}{(1 - \beta^2)^{1/2}} \frac{m_0 v_\perp^2}{2B} \frac{c}{e} \frac{\hat{e}_1 \times \nabla B}{B} + \frac{1}{(1 - \beta^2)^{1/2}} \frac{m_0 c}{e} v_\parallel^2 \frac{\hat{e}_1}{B} \times \frac{\partial \hat{e}_1}{\partial s} \quad (11)$$

and the parallel force is

$$\frac{m_0}{(1 - \beta^2)^{1/2}} \frac{dv_{||}}{dt} = - \frac{1}{(1 - \beta^2)^{1/2}} \frac{m_0 v_{\perp}^2}{2B} \frac{\partial B}{\partial s}. \quad (12)$$

Nonrelativistically, the magnetic moment is $M = mv_{\perp}^2/2B$.

Relativistically, the corresponding invariant is

$M_r = m_0 v_{\perp}^2 / 2B(1 - \beta^2) = p_{\perp}^2 / 2m_0 B$. It is not obvious that this is the correct generalization of M for relativistic energy. It is easy enough to verify for the simple case of a particle in a uniform, azimuthally symmetric, field that changes with time. The general case is not so easy to prove. The adiabatic invariants will be studied more in the next section.

The parallel force in (12) is now larger by $(1 - \beta^2)^{-1/2}$ than would be predicted by the nonrelativistic equation for the same rest mass. Similarly, the drifts in (11) are faster by the same factor. These effects are caused by the increased gyration radius resulting from the relativistic mass increase. For example, the increased gyration radius increases the amount of field inhomogeneity sampled by the particle, hence increases the ∇B drift. Similarly, the parallel force increases because the larger gyration radius subjects the particle to a greater convergence of the field lines, and it is this convergence that produces the mirror effect. As illustrated in Figure 5, it is the product of v_{\perp} and the radial component of \vec{B} that results in a parallel force.

-13-

If the electric field is sufficiently small (formally, of order ϵ), the four terms containing \vec{u}_E in (4) become of order ϵ^2 and may be dropped. The drift proportional to $\partial \vec{e}_1 / \partial t$ will also probably be negligible, since $\nabla \times \vec{E}$ and $\partial \vec{B} / \partial t$ are related by the Maxwell equation. Then only the three familiar drifts remain. One may surmise that the correct relativistic modification is obtained by adding $c \vec{E} \times \vec{B} / B^2$ to (11) and $e E_{||}$ to the parallel force in (12). This does in fact turn out to be the correct procedure, but it is not a deductive one, since (11) and (12) were derived by assuming no electric field. The relativistic case has been studied by Hellwig [1955] and by Vandervoort [1960] for \vec{E}_\perp large (i.e., of order 1), and the small \vec{E}_\perp results are a special case.

The relativistic rate of energy change for \vec{E}_\perp small is

$$\frac{dW}{dt} = e \vec{R} \cdot \vec{E} + M_r (1 - \beta^2)^{1/2} \frac{\partial B}{\partial t} . \quad (13)$$

Only the betatron term has been altered, a comparison with (7) shows.

The complete guiding center equations for large \vec{E}_\perp are rather long and will not be repeated here [see Vandervoort, 1960; Northrop, 1963]. Their principal features are corrections to existing terms of the small \vec{E}_\perp relativistic expressions above. Additionally, two new drift terms that are in the direction of \vec{E}_\perp appear. They are pure relativistic effects that have no analog in the small \vec{E}_\perp relativistic case. One of these two drifts can be explained by the change in direction of \vec{B} when a Lorentz transformation is made in the presence of an electric field. Basically, the drift is a result

of the change in the reference direction with respect to which parallel and perpendicular are defined. Some of what was parallel velocity is converted to perpendicular velocity.

6. THE ADIABATIC INVARIANTS

The magnetic moment. The emphasis so far has been on the guiding center motion and on energy changes. Not only are the guiding center equations useful, but also valuable are quantities that are constant over long periods of guiding center motion--i.e., any invariants of the adiabatic motion, or "adiabatic invariants." They are not exact invariants of the particle motion, any more than the guiding center equations are exact equations for the particle motion. Formal analysis [Kruskal, 1960; Northrop and Teller, 1960] shows that there are at the most three adiabatic invariants for the charged particle. Each one is really an asymptotic series in a smallness parameter ϵ --a series of the form:

constant = $a_0 + \epsilon a_1 + \epsilon^2 a_2 + \dots$ Systematic analysis [Gardner, 1959; Kruskal, 1960] is essential for obtaining higher order terms in the series. Historically, however, the forms of the lowest order invariants (i.e., the a_0 's) were deduced by physical insight and by consideration of special cases [Alfvén, 1950; Rosenbluth, 1955; Northrop and Teller, 1960]. The connection with more formal theory was made later. Such an evolutionary history is common in physical science. In this paper only invariance to lowest order (the a_0 's) will be proven.

The formal theories also show that the adiabatic invariant series are not the action integrals of the form $\oint p dq$, where p and q are canonical variables, but are instead Poincaré integral invariants

of the form $\sum_i \oint p_i dq_i$, where the number of terms in the sum is the number of degrees of freedom of the canonical system. However, the number of adiabatic invariants may vary from one to three, depending on the field geometry, as will become apparent shortly. In general, the number of invariants is less than or equal to the number of degrees of freedom of the system [Kruskal, 1960].

The first invariant is the magnetic moment, defined previously as $mv_{\perp}^2/2B$ for the nonrelativistic case; $mv_{\perp}^2/2B$ is really M_0 of the magnetic moment series: constant = $M_0 + \epsilon M_1 + \epsilon^2 M_2 + \dots$. The definition of v_{\perp} was glossed over slightly in the beginning of this review. If the component of \vec{E} perpendicular to \vec{B} is small, the $\vec{E} \times \vec{B}$ drift is much less than the particle velocity and the particle trajectory will be as in Figure 2. The motion is almost circular, and the v_{\perp} to be used in the magnetic moment is the velocity about the circle. When \vec{E}_{\perp} is this small, the last four drifts in (4) will probably be negligible. Suppose that \vec{E}_{\perp} is now increased. Eventually the trajectory will resemble a prolate cycloid as in Figure 6. There is no resemblance to circular motion in the laboratory frame, but in the frame moving at $\vec{E} \times \vec{B}$ the motion is approximately circular again as in Figure 2. It is the v_{\perp} in this drifting frame that should be used in $mv_{\perp}^2/2B$. Adiabatic theory therefore can still apply even when the perpendicular electric field is so large that the particle trajectory in the observer's frame shows no looping or resemblance to circular motion. One must only be careful to use the complete expressions in (4) and (6), and to define v_{\perp} properly.

The invariance of M is easy to demonstrate for simple cases, like a time dependent magnetic field with azimuthal symmetry and straight lines of force. A proof for the most general situation (general time-dependent magnetic field and large electric field) seems to be rather long [Kruskal, 1958; Gardner, 1959; Northrop, 1963). The most general case for which a simple proof seems to exist is the static one, where the energy is constant; a large curl-free electric field may be present. By conservation of energy,

$$\frac{d}{dt} \left(\frac{m v_{||}^2}{2} + \frac{m u_E^2}{2} + MB + e\phi \right) = 0, \quad (14)$$

where ϕ is the electrostatic potential. Recall that the invariance of M was not invoked in deriving the guiding center equations. Thus the value of $dv_{||}/dt$ from (6) can be used to convert (14) to

$$\frac{d(MB)}{dt} = -\frac{ed\phi}{dt} - \vec{m u}_E \cdot \frac{d\vec{E}}{dt} - mv_{||} \left(\frac{e}{m} E_{||} - \frac{M}{m} \frac{\partial B}{\partial s} + \vec{u}_E \cdot \frac{d\vec{e}_L}{dt} \right). \quad (15)$$

The total derivative $d\phi/dt$ equals $v_{||}(\partial\phi/\partial s) + \vec{R}_L \cdot \vec{\nabla}\phi$, where \vec{R}_L is given by (4). Putting it all together and doing a little vector algebra gives

$$\frac{d(MB)}{dt} = M \vec{u}_E \cdot \nabla B + M v_{||} \frac{\partial B}{\partial s} \equiv M \frac{dB}{dt}, \quad (16)$$

or

$$\frac{dM}{dt} = 0.$$

The next two higher terms in the magnetic moment series have also been derived [Kruskal, 1958; Gardner, 1962]. They are rather complicated.

The expression "nonadiabatic behavior" as applied to the magnetic moment has by custom come to mean any deviation of $mv_{\perp}^2/2B$ from constancy. However, it is actually the series $M_0 + \epsilon M_1 + \epsilon^2 M_2 + \dots$ that is the invariant of the particle motion, and not just M_0 . Therefore, M_0 can vary according to adiabatic theory. It seems preferable to define as nonadiabatic any behavior not predicted by the series. Since the series is asymptotic [Berkowitz and Gardner, 1959], and not convergent, it would not be surprising to see particle behavior that completely ignores the adiabatic predictions, even in low order, and this would be genuine nonadiabatic behavior. Examples of such motion are known [Garren et al., 1958; Northrop, 1963] for the magnetic moment.

The second or longitudinal invariant. Another invariant of the particle motion, or really of the guiding center motion, is

$$J = \oint p_{\parallel} ds, \quad (17)$$

where p_{\parallel} is mv_{\parallel} , the guiding center momentum parallel to the line of force. The invariant J exists if there is a mirror-type geometry, such that the guiding center oscillates back and forth along the lines of force while drifting slowly at right angles to them, as illustrated in Figure 7. For J to be constant, it is necessary that the drift be slow compared to v_{\parallel} --i.e., that \vec{E}_{\perp} be of order ϵ . The

integral is taken over a complete oscillation, the deviation of the guiding center from a line due to the drift during one oscillation being negligible if \vec{E}_\perp is small.

The earliest suggestion that J is an invariant appears to have come from Rosenbluth [Chew, Goldberger, and Low; 1955]. A proof of the invariance of J and some applications to laboratory magnetic-field configurations was given by Kadomtsev [1958] for a nonrelativistic particle in a static magnetic field. A proof that remains valid at relativistic energies, and that includes time-dependent fields has been given by Northrop and Teller [1960] along with applications to the Van Allen radiation. The proof of the invariance of J given below is for a nonrelativistic particle in a static field with no electric field; inclusion of nonstatic fields greatly increases the length of the proof. Therefore, only the results will be given for the time-dependent case. The time-dependent results will be needed to discuss the third invariant. Relativistic modifications do not seem to materially complicate the proofs.

To begin the proof of J , a curvilinear coordinate system will now be introduced. The three coordinates will be denoted by α , β , and s , where α and β are two parameters specifying the line of force, and s denotes position along the line. (Distinguish this β from v/c in a previous section.) A system of nonintersecting lines can be generated as the intersections of two families of surfaces $\alpha(\vec{r}) = \text{constant}$, and $\beta(\vec{r}) = \text{constant}$, where $\alpha(\vec{r})$ and $\beta(\vec{r})$ are two different functions of position. It is apparent that for a given system of lines the functions $\alpha(\vec{r})$ and $\beta(\vec{r})$ are not unique. Consider

the simple example of straight lines of force. They can be generated by the intersections of two families of planes, by a family of planes and one of cylinders, etc. Among the many possible pairs of functions $\alpha(\vec{r})$ and $\beta(\vec{r})$ for a given magnetic field, there is a subclass for which the vector potential \vec{A} is $\alpha \nabla B$ and \vec{B} therefore is $\nabla \alpha \times \nabla \beta$. That such a subclass exists is not quite obvious, but it is not difficult to prove. The utility of the subclass is that for it $|\nabla \alpha \times \nabla \beta|/B$ is constant everywhere, being unity, and this fact reduces the algebra involved in the proof.

In the absence of electric fields the energy W equals $mv_{||}^2/2 + MB$, so that

$$J(\alpha, \beta, M, W) = \oint \left\{ 2m[W - MB(\alpha, \beta, s)]^{1/2} ds \right\}. \quad (18)$$

The instantaneous rate of change of J due to the particle drift \vec{R}_\perp in Figure 7 is

$$\frac{dJ}{dt} = \frac{\partial J}{\partial \alpha} \frac{d\alpha}{dt} + \frac{\partial J}{\partial \beta} \frac{d\beta}{dt}. \quad (19)$$

Differentiation of the integral in (18) gives

$$\frac{\partial J}{\partial \alpha} = -m M \oint \frac{ds}{[2m(W - MB)]^{1/2}} \frac{\partial B(\alpha, \beta, s)}{\partial \alpha},$$

and

$$\frac{\partial J}{\partial \beta} = -m M \oint \frac{ds}{[2m(W - MB)]^{1/2}} \frac{\partial B(\alpha, \beta, s)}{\partial \beta}. \quad (20)$$

Because α and β are constant on a line of force, they are changed only by the drift velocity, and not by the parallel velocity. Therefore,

-20-

$d\alpha/dt = \vec{R} \cdot \nabla\alpha(\vec{R})$ and $d\beta/dt = \vec{R}_\perp \cdot \nabla\beta(\vec{R})$. Substituting \vec{R}_\perp from (14), with the electric field zero, gives

$$\frac{d\alpha}{dt} = \frac{\hat{e}_1}{B} \times \left(\frac{mc}{e} \nabla B + \frac{mc}{e} v_{||}^2 \frac{\partial \hat{e}_1}{\partial s} \right) \cdot \nabla\alpha \quad (21)$$

Consider now the quantity $(\partial\vec{R}/\partial\beta) \times \vec{B}$, where the guiding center position \vec{R} is a function of (α, β, s) :

$$\frac{\partial\vec{R}}{\partial\beta} \times \vec{B} = \frac{\partial\vec{R}}{\partial\beta} \times (\nabla\alpha \times \nabla\beta) = -(\nabla\alpha \cdot \frac{\partial\vec{R}}{\partial\beta}) \nabla\beta + (\nabla\beta \cdot \frac{\partial\vec{R}}{\partial\beta}) \nabla\alpha. \quad (22)$$

By implicit differentiation of $\alpha = \alpha[\vec{R}(\alpha, \beta, s)]$ one finds that

$$\nabla\alpha \cdot \frac{\partial\vec{R}}{\partial\beta} = 0, \quad \text{and that} \quad \nabla\beta \cdot \frac{\partial\vec{R}}{\partial\beta} = 1.$$

Thus, $(\partial\vec{R}/\partial\beta) \times \vec{B} = \nabla\alpha$, and (21) becomes

$$\frac{d\alpha}{dt} = \frac{\hat{e}_1}{B} \times \left(\frac{mc}{e} \nabla B + \frac{mc}{e} v_{||}^2 \frac{\partial \hat{e}_1}{\partial s} \right) \cdot \left(\frac{\partial\vec{R}}{\partial\beta} \times \vec{B} \right). \quad (23)$$

Interchanging the dot and cross, and expanding the triple vector product

$\hat{e}_1 \times (\frac{\partial\vec{R}}{\partial\beta} \times \vec{B})$ gives

$$\begin{aligned} \frac{d\alpha}{dt} &= - \left(\frac{mc}{e} \nabla B + \frac{mc}{e} v_{||}^2 \frac{\partial \hat{e}_1}{\partial s} \right) \cdot \left(\frac{\partial\vec{R}}{\partial\beta} - \hat{e}_1 \hat{e}_1 \cdot \frac{\partial\vec{R}}{\partial\beta} \right) \\ &= - \frac{mc}{e} \frac{\partial\vec{R}}{\partial\beta} \cdot \nabla B - \frac{mc}{e} v_{||}^2 \frac{\partial\vec{R}}{\partial\beta} \cdot \frac{\partial \hat{e}_1}{\partial s} + \frac{mc}{e} \hat{e}_1 \cdot \frac{\partial\vec{R}}{\partial\beta} \hat{e}_1 \cdot \nabla B \\ &= - \frac{mc}{e} \frac{\partial B(\alpha, \beta, s)}{\partial\beta} - \frac{mc}{e} v_{||}^2 \frac{\partial R(\alpha, \beta, s)}{\partial\beta} \frac{\partial \hat{e}_1(\alpha, \beta, s)}{\partial s} \\ &\quad + \frac{mc}{e} \hat{e}_1 \cdot \frac{\partial\vec{R}}{\partial\beta} \frac{\partial B(\alpha, \beta, s)}{\partial s}. \end{aligned} \quad (24)$$

-21-

In the second term on the right-hand side of (24), we have

$$\frac{\partial \vec{R}}{\partial \beta} \cdot \frac{\partial \hat{e}_1}{\partial s} = \frac{\partial}{\partial s} (\hat{e}_1 \cdot \frac{\partial \vec{R}}{\partial \beta}) , \text{ since } \hat{e}_1 \cdot \frac{\partial}{\partial s} \frac{\partial \vec{R}(\alpha, \beta, s)}{\partial \beta} = \hat{e}_1 \cdot \frac{\partial}{\partial \beta} \frac{\partial \vec{R}}{\partial s} , \text{ and}$$

$$\hat{e}_1 \cdot \frac{\partial}{\partial \beta} \frac{\partial \vec{R}}{\partial s} = \hat{e}_1 \cdot \frac{\partial \hat{e}_1}{\partial \beta} , \text{ which is zero. Therefore, the second term becomes}$$

$$- \frac{mc}{e} v_{||}^2 \frac{\partial}{\partial s} (\hat{e}_1 \cdot \frac{\partial \vec{R}}{\partial \beta}) = - \frac{mc}{e} v_{||} \frac{d}{dt} (\hat{e}_1 \cdot \frac{\partial \vec{R}}{\partial \beta}) . \text{ From (6),}$$

$$\frac{\partial B}{\partial s} = - \frac{m}{M} \frac{dv_{||}}{dt} , \text{ so that the last two terms in (24) combine to}$$

$$- \frac{mc}{e} \frac{d}{dt} (v_{||} \hat{e}_1 \cdot \frac{\partial \vec{R}}{\partial \beta}) . \text{ The instantaneous rate of change of } \alpha \text{ finally is}$$

$$\frac{d\alpha}{dt} = - \frac{Mc}{e} \frac{\partial B(\alpha, \beta, s)}{\partial \beta} - \frac{mc}{e} \frac{d}{dt} (v_{||} \hat{e}_1 \cdot \frac{\partial \vec{R}}{\partial \beta}) . \quad (25)$$

By a similar analysis, we have

$$\frac{d\beta}{dt} = \frac{Mc}{e} \frac{\partial B}{\partial \alpha} + \frac{mc}{e} \frac{d}{dt} (v_{||} \hat{e}_1 \cdot \frac{\partial \vec{R}}{\partial \alpha}) . \quad (26)$$

If $\partial B / \partial \beta$ from (25) is substituted into $\partial J / \partial \beta$ from (20), the result is

$$\begin{aligned} \frac{\partial J}{\partial \beta} &= \oint \frac{mds}{[2m(W - MB)]^{1/2}} \frac{e}{c} \left[\frac{d\alpha}{dt} + m \frac{d}{dt} (v_{||} \hat{e}_1 \cdot \frac{\partial \vec{R}}{\partial \beta}) \right] \\ &= \frac{e}{c} \oint \frac{ds}{v_{||}} \frac{d\alpha}{dt} . \end{aligned} \quad (27)$$

The integral of $m \frac{d}{dt} (v_{||} \hat{e}_1 \cdot \frac{\partial \vec{R}}{\partial \beta})$ has vanished because $ds/v_{||}$ is dt ,

and $v_{||}$ is zero at the reflection points. Equation (27) can be written as

$$\frac{\partial J}{\partial \beta} = \frac{e}{c} T \langle \dot{\alpha} \rangle, \quad (28)$$

where T is the time for a longitudinal oscillation, and the brackets denote the time average over an oscillation. Similarly,

$$\frac{\partial J}{\partial \alpha} = -\frac{e}{c} T \langle \dot{\beta} \rangle. \quad (29)$$

Equation (19) can then be written as

$$\frac{dJ}{dt} = \frac{e}{c} T \left[\langle \dot{\alpha} \rangle \dot{\beta} - \langle \dot{\beta} \rangle \dot{\alpha} \right]. \quad (30)$$

Now this quantity is not zero except under very special circumstances, so that J is not instantaneously being conserved by the guiding center motion. However, the rate of change of J averaged over a longitudinal oscillation is

$$\begin{aligned} \oint \frac{ds}{v_{||}} \frac{dJ}{dt} &= \frac{eT}{c} \left[\langle \dot{\alpha} \rangle \oint \frac{ds}{v_{||}} \dot{\beta} - \langle \dot{\beta} \rangle \oint \frac{ds}{v_{||}} \dot{\alpha} \right] \quad (31) \\ &= \frac{eT}{c} \left[\langle \dot{\alpha} \rangle \langle \dot{\beta} \rangle - \langle \dot{\beta} \rangle \langle \dot{\alpha} \rangle \right], \end{aligned}$$

which is identically zero, and this is the important fact for the long term motion.

Equations (28) and (29) are new equations of motion, with the guiding center oscillation averaged out; they are the analog of the guiding center equations of motion, which are the particle equations of motion with the particle gyration averaged out.

When Equations (28) and (29) are solved for $\langle \dot{\alpha} \rangle$ and $\langle \dot{\beta} \rangle$, they are at first sight suggestively canonical in form, with $J(\alpha, \beta, M, W)$ playing the role of Hamiltonian. But they are not quite canonical. In the first place the time of oscillation T is also a function of (α, β, M, W) . Furthermore there are the time averages of $\dot{\alpha}$ and $\dot{\beta}$, rather than the instantaneous values. The first difficulty can be overcome by differentiating $J = J(\alpha, \beta, M, W)$ implicitly with respect to α and β to yield $\partial J(\alpha, \beta, M, W)/\partial \beta = -(\partial J/\partial W) \partial W(\alpha, \beta, M, J)/\partial \beta$, etc. for $\partial J/\partial \alpha$. The factor $\partial J/\partial W$ is simply T , as can be verified from (18). Then

$$\langle \dot{\alpha} \rangle = -\frac{c}{e} \frac{\partial W(\alpha, \beta, M, J)}{\partial \beta},$$

and

$$\langle \dot{\beta} \rangle = \frac{c}{e} \frac{\partial W}{\partial \alpha}.$$
(32)

Except for the time averages, these are now canonical. It would seem that the matter of the time averages could be overlooked if one is interested only in the average guiding-center position, and therefore that the equations of motion can be regarded as canonical. If this is the case, any theorems in classical mechanics that come from the canonical equations should have an analog in the (α, β) space. Liouville's theorem comes to mind immediately, and it is possible to derive it [Northrop and Teller, 1960] for the density in (α, β) space by disregarding the time averages. To dispel any lingering doubts about the time averages, a more direct derivation can be made by using the expressions for the instantaneous values of $\dot{\alpha}$ and $\dot{\beta}$. The consequences of the Liouville theorem will be described shortly.

The third adiabatic invariant. As a guiding center oscillates between mirror points, it gradually changes lines of force. During its motion along a line, it instantaneously drifts towards a variety of adjacent lines, but there is one line towards which it moves on the average and this line is specified by (52). Thus a surface composed of lines on which J is constant is gradually traversed by the guiding center. Now it may happen that this surface is closed, so that the particle eventually returns to a line it traversed earlier. If so, there is a third periodicity and a third adiabatic invariant is to be expected. The surfaces seem to be closed for particles in the inner Van Allen belt. Such a surface (idealized) is sketched in Figure 8.

Note that if the particle is not trapped between mirrors, the longitudinal motion is not periodic and there is not even a second adiabatic invariant, nor is there a third. Only the magnetic moment exists. This illustrates the fact that the number of adiabatic invariants depends on the geometry and is less than or equal to the number of degrees of freedom.

To return to the Liouville theorem: it says that in the steady state in the absence of electric fields, contours of constant magnetic field are also constant guiding-center density contours on a longitudinal invariant surface (Figure 8).

The third adiabatic invariant is the flux Φ of \vec{B} enclosed by the surface of Figure 8. That this flux should be constant in a static situation is a trivial statement, much as the invariance of the magnetic moment in a uniform field is trivially true. But the flux is

also invariant if the field is time dependent, and this is the significant fact. The surface about which the particle precesses is not even well defined unless the particle traverses it in a time small compared to the time scale for fields to change. It is not surprising therefore that this rapid precession assumption is necessary to prove the invariance of ψ . From a practical standpoint, the time scale of field fluctuations must be slowest to conserve ψ ; they can be faster and still conserve J , and fastest of all without disturbing M , since the time scale then need only be long compared to the gyration period.

Proof of the invariance of ψ is reminiscent of the proof for J . It is necessary to extend equations (32) to include time-dependent fields. When the fields are time dependent, it is appropriate to generalize the quantity W used previously to a quantity K , defined by

$$K = \frac{mv_{||}^2}{2} + MB + e(\psi + \frac{\alpha}{c} \frac{\partial \beta}{\partial t}) \quad (33)$$

where ψ is the scalar potential for the electric field, so that \vec{E} is $-\nabla\psi - \frac{1}{c} \frac{\partial(\alpha\vec{v}\beta)}{\partial t}$. In a time-dependent field α and β are functions of both time and position. The second invariant is now defined by

$$J(\alpha, \beta, M, K, t) = \oint \left\{ 2m \left[K - e \left(\frac{\alpha}{c} \frac{\partial \beta}{\partial t} + \psi \right) - MB \right] \right\}^{1/2} ds, \quad (34)$$

where $\partial\beta/\partial t$ is to be expressed as a function of (α, β, s, t) . The generalization of equations (32) turn out to be

-26-

$$\langle \dot{\alpha} \rangle = -\frac{c}{e} \frac{\partial K}{\partial \beta} (\alpha, \beta, J, M, t),$$

$$\langle \dot{\beta} \rangle = \frac{c}{e} \frac{\partial K}{\partial \alpha},$$

$$\langle \dot{K} \rangle = \frac{\partial K}{\partial t},$$

and

(35)

$$I = T \frac{\partial K}{\partial J}.$$

The quantity $\langle K \rangle$ is related to the gain in energy averaged over a longitudinal oscillation.

The details of the proof that ϕ is invariant will not be given here [see Northrop and Teller, 1960]. One finds that $d\phi/dt$ is not zero as the particle drifts around the surface defined by the invariance of J (i.e., as it precesses around the earth); the average motion from line to line as given by equations (35) does not conserve ϕ . But if $d\phi/dt$ is averaged over a complete precession, the time average is zero. This is analogous to the situation with dJ/dt . The instantaneous rate of change of ϕ is

$$\frac{d\phi}{dt} = \frac{cT_p}{e} \left[\langle \dot{K} \rangle - \langle \langle \dot{K} \rangle \rangle \right] \neq 0, \quad (36)$$

where $\langle \langle \dot{K} \rangle \rangle$ means $\langle \dot{K} \rangle$ averaged over a precession, and T_p is the time for the particle to precess once around the surface. The right side of (36) obviously vanishes when averaged over the period T_p .

Before leaving the subject of the third invariant, several points should be discussed concerning motion of lines of force and the average (over a longitudinal oscillation) guiding-center drift. The "velocity"

of a line of force in a time-dependent field is not physically observable. We cannot see lines of force. One is therefore free to define line velocity, and it should be defined so as to enhance our visualization of how the magnetic-field pattern changes with time. One usually uses the picture in which a magnetic field has an intensity proportional to the line density one draws. As the field changes with time this picture remains valid if the lines are moved around at a "flux-preserving velocity." To define this velocity, suppose an arbitrary closed curve is drawn in space; now let each element of the curve move at a velocity $\vec{U}(\vec{r}, t)$. If the flux through the curve remains constant as the curve distorts, \vec{U} is said to be flux preserving. As shown by Newcomb [1958], \vec{U} must satisfy $\nabla \times (\vec{E} + \vec{U} \times \vec{B}/c) = 0$. This limits \vec{U} but does not determine it uniquely. One often chooses \vec{U} as $c\vec{E} \times \vec{B}/B^2$, which is acceptable if $\nabla \times \vec{E}_H$ is zero.

A more general definition of line velocity that is always acceptable (but not unique) is

$$\vec{U}(\vec{r}, t) = \left(\frac{\partial \beta}{\partial t} \nabla \alpha - \frac{\partial \alpha}{\partial t} \nabla \beta \right) \times \frac{\hat{e}_l}{cB} \quad (37)$$

It is not difficult to show that $\nabla \times (\vec{E} + \vec{U} \times \vec{B}/c)$ is zero for this choice of \vec{U} . Moreover, this choice has the advantage that $\frac{\partial \alpha}{\partial t} + \vec{U} \cdot \nabla \alpha$ is zero, and likewise for β . The significance of this is that as an observer moves at the line velocity, the (α, β) label on the line he is following remains unchanging with time.

A convenient space in which to visualize the invariant surfaces is a Cartesian (α, β, s) space, in which the field lines are straight

-28-

and parallel to the s axis, as in Figure 9. The choice of \vec{U} in (37) makes the lines of force fixed in this space; by contrast, a particle for which J (but not necessarily \vec{J}) is invariant moves in (α, β) space in accord with equations (35), and consequently does not remain attached to a line of force.

The picture developed so far of line motion is very appealing, but is not unique. To illustrate, suppose \vec{U} is defined by

$$\vec{U}(\vec{r}, t) = \frac{c}{eB} \hat{e}_1 \times \nabla K(\vec{r}, M, J, t) + \left(\frac{\partial \beta}{\partial t} \nabla \alpha - \frac{\partial \alpha}{\partial \beta} \nabla \beta \right) \times \frac{\hat{e}_1}{B}, \quad (38)$$

where K is to be regarded as a function of the specified variables via (34). This velocity can also be proved flux preserving. However, for it, we have

$$\begin{aligned} \frac{d\alpha}{dt} + \vec{U} \cdot \nabla \alpha &= \frac{c}{eB} (\hat{e}_1 \times \nabla K) \cdot \nabla \alpha \\ &= -\frac{c}{e} \frac{\partial K}{\partial \beta} (\alpha, \beta, M, J, t) = \langle \dot{\alpha} \rangle, \end{aligned} \quad (39)$$

and similarly for $\langle \dot{\beta} \rangle$. With this definition of line velocity, the line of force consequently moves at exactly the average particle drift velocity, and the particle remains attached to the line.

Either of the two pictures is acceptable, though definition (37) seems preferable since it does not depend on any particle parameters, while definition (38) depends on J and M . It is a little unappealing to use a definition of line velocity that depends on the particle under observation. One prefers to visualize the motion of field lines as

being intrinsic to the field and not dependent on particles. Furthermore, if two particles with different J and M are on the same line of force, there will be an ambivalence in the line velocity. Finally, if the electric fields are so large that all the drift terms in (4) must be retained, J is not conserved. Definition (38) is still flux preserving, but J is now a time-dependent parameter. And because the guiding center no longer shows a slow average drift, governed by equations (35), it is not possible to say that the particle follows the line of force on the average. The guiding center follows a trajectory in Figure 9 determined by (4) and (6). Under these circumstances, definition (37) for the line velocity certainly is superior to (38).

7. APPLICATION OF ADIABATIC THEORY TO PLASMAS

In the previous sections the motion of a single particle in a prescribed field has been studied. The adiabatic model may also apply to a plasma, where the density of positively and negatively charged particles is so large that their interactions are important in determining their motions. The field each particle moves in is the sum of (a) any "external" field and (b) those fields due to the motions and positions of all other particles. For the particle motion to be adiabatic, close collisions between charged particles must be infrequent (high plasma temperature and low density) so that a particle at no time feels a sudden force. Such self-consistent calculations are necessary to analyze the stability of plasma confinement in a given field configuration.

-30-

Newcomb [1963] has developed a method for using the first two adiabatic invariants in studying plasma stability. The change in energy of an equilibrium plasma under a prescribed displacement $\vec{\xi}(\vec{r})$ of the element of plasma at \vec{r} can be obtained from invariance of the magnetic moment and longitudinal invariant. If this energy change is positive for all possible $\vec{\xi}(\vec{r})$, the plasma is stable. If the change is negative for any $\vec{\xi}(\vec{r})$, it is unstable. It is plausible that the change in particle energies should be derivable from the first two invariants. The magnetic moment is associated with perpendicular energy, while the longitudinal invariant is associated with parallel velocity and energy. Changes in field energy under the perturbation must also be accounted for in obtaining the total change in energy.

The mechanism of these instabilities can be explained in terms of the adiabatic particle drifts. In the presence of the perturbation the drifts lead to charge accumulations whose electric fields drive the perturbation further in a typically regenerative fashion [see Rosenbluth and Longmire, 1957, and Northrop, 1961, for examples].

We can also apply adiabatic motion to the current density in a collisionless plasma. Each component (i.e., ions or electrons) of the plasma obeys the macroscopic momentum conservation equation

$$nm \frac{d\vec{V}}{dt} = - \nabla \cdot \vec{P} + ne \frac{\vec{V}}{c} \times \vec{B} + ne \vec{E}, \quad (40)$$

where \vec{V} is the average (over the velocity distribution) of the particle velocity \vec{v} , and \vec{P} is the pressure tensor defined as $\langle nm(\vec{v} - \vec{V})(\vec{v} - \vec{V}) \rangle$, where the brackets mean an average over the particle velocity distribution. The current density \vec{J} of that

-31-

component is $n e \vec{V}$, where n is the particle density. Solving (40) for \vec{V} , we obtain

$$\vec{V} = \vec{V}_{\parallel} + \frac{c \hat{e}_1 \times \nabla \cdot \vec{P}}{n e B} + \frac{c \vec{E} \times \hat{e}_1}{B} + \frac{mc}{eB} \hat{e}_1 \times \frac{d\vec{V}}{dt}. \quad (41)$$

Consider now a steady situation, where there is no electric field. Then \vec{V}_{\perp} is $c \hat{e}_1 \times \nabla \cdot \vec{P} / n e B$. This is just the east-west asymmetry effect of mirroring protons observed by Heckman and Nakano [1963]. They observed that at the inner edge of the inner Van Allen belt, more high energy protons are moving east than west; there is an average proton velocity \vec{V} towards the east. The pressure gradient is caused by the atmospheric density gradient, there being fewer particles at lower altitudes due to the greater loss to the atmosphere. At the outer edge of a radiation belt, where the density decreases with increasing radius (for whatever reason), the reverse asymmetry should appear, with more particles moving west than east.

The divergence of the pressure tensor can be expanded in the adiabatic case as [see Chew, Goldberger, and Low, 1956]

$$\nabla \cdot \vec{P} = \hat{e}_1 \left[\frac{\partial P_{\parallel}}{\partial s} - \frac{P_{\parallel} - P_{\perp}}{B} \frac{\partial B}{\partial s} \right] + \left[(P_{\parallel} - P_{\perp}) \frac{\partial \hat{e}_1}{\partial s} + \nabla P_{\perp} \right], \quad (42)$$

where P_{\parallel} is $nm \langle (v_{\parallel} - v_{\parallel})^2 \rangle$ and P_{\perp} is $\frac{1}{2} nm \langle v_{\perp}^2 \rangle$. In the east-west asymmetry experiment there would be a small contribution from the line curvature $\partial \hat{e}_1 / \partial s$ in addition to the one from the pressure gradient ∇P_{\perp} .

It is possible to prove from the Vlasov (collisionless Boltzmann) equation that

$$\vec{J}(\vec{r}, t) \equiv ne\vec{V} = Ne(\vec{R}_\perp + \vec{e}_L v_\parallel) + c \nabla \times \vec{M}, \quad (43)$$

where N is the number of guiding centers per unit volume at (\vec{r}, t) and \vec{M} is the total magnetic moment per unit volume of particles with guiding centers at \vec{r} . The brackets mean the average over particles with guiding centers at \vec{r} . The perpendicular component of (43) is easily derived from (41) and the guiding-center equations. However, the parallel component is rather difficult to prove formally [see Northrop, 1963], even though the entire expression (43) is intuitively correct. It says that the total current density in a plasma is the sum of the guiding center current and the current that results from the curl of the magnetic moment per unit volume.

8. NONADIABATIC EFFECTS

The application of adiabatic theory and the lowest order invariants to the Van Allen radiation has been outlined in previous sections. According to the theory, in the absence of collisions, particles would remain indefinitely in the geomagnetic field and repeatedly precess about their invariant surfaces. In practice all three invariants may not hold sufficiently well for this permanent trapping to occur. There is low temperature plasma permeating the magnetosphere about the earth, and the solar wind may produce disturbances that are propagated through this plasma. These

disturbances in turn may be sufficiently fast to affect one or more of the lowest order invariants. Even if the deviation of one of them from a constant is very small, this very small effect can operate over very long times in the geophysical case. The question becomes whether these effects are cumulative, or whether they are oscillatory and self-cancelling over a long period. If the motion is truly nonadiabatic, in the sense defined in Section 6, the effects may be cumulative and the particle may become lost from the geomagnetic field. For example, if the magnetic moment decreases continuously, the particle will eventually become lost in the atmosphere. However, if the motion is adiabatic, in the sense of being predicted by the first few terms of the invariant series, then the particle may still be permanently trapped, with the guiding center following a slightly different path from that predicted by the lowest order invariant. The distinction between these two possibilities, cumulative and oscillatory, may not always be sharp, though in one geometry it seemed to be quite sharp for the magnetic moment [see Garren, et al., 1958].

There would certainly be value in computing at least one higher term for the longitudinal and flux invariants. The consequences of the earth's rotation, coupled with the azimuthal asymmetry of its field, do not seem to be known except in the limit when \mathbf{T} is invariant. In this limit a particle precesses rapidly about its invariant surface, and the surface rotates slowly and rigidly with a 24-hour period. The next terms of the longitudinal and flux invariant series ought to describe the lowest order modification to this simple picture.

To conclude, it does not seem possible at present to make any general statements about nonadiabatic effects, other than that numerical computation is probably needed to study them. However, these effects may be important in the dynamics of the trapped radiation and therefore merit attention.

REFERENCES

Alfvén, H., Cosmical Electrodynamics, Clarendon Press, Oxford, 1950.

Berkowitz, J. and C. S. Gardner, On the Asymptotic Series Expansion of the Motion of a Charged Particle in Slowly Varying Fields, Commun. Pure Appl. Math. 12, 501, 1959.

Chew, G., M. Goldberger, and F. Low, The Boltzmann Equation and the One-Fluid Hydromagnetic Equations in the Absence of Particle Collisions, Proc. Roy Soc. (London), A236, 112, 1956.

Chew, G. F., M. Goldberger, and F. Low, The Individual Particle Equations of Motion in the Adiabatic Approximation, Los Alamos Scientific Laboratory Report IA-2055, Chapt. T-759, p. 9, 1955.

Davis, L., Jr., Modified Fermi Mechanism for the Acceleration of Cosmic Rays, Phys. Rev. 101, 351, 1956.

Fermi, E., On the Origin of the Cosmic Radiation, Phys. Rev. 75, 1169, 1949.

Fermi, E., Galactic Magnetic Fields and the Origin of Cosmic Rays, Astrophys. J. 119, 1, 1954.

Gardner, C. S. Adiabatic Invariants of Periodic Classical Systems, Phys. Rev. 115, 791, 1959.

Gardner, C. S., 1962, RCA Laboratories, Princeton, New Jersey, private communication. The work is quoted in Northrop [1963].

Garren, A., et al., Individual Particle Motion and the Effect of Scattering in an Axially Symmetric Magnetic Field, in Proc. Second United Nations Intern. Conf. Peaceful Uses Atomic Energy (United Nations, Geneva), Vol. 31, p. 65, 1958.

Heckman, H. and G. Nakano, East-West Asymmetry in the Flux of Mirroring Geomagnetically Trapped Protons, J. Geophys. Res. (to be published, April 15, 1963).

Hellwig, G., Über die Bewegung geladener Teilchen in Schwach veränderlichen Magnetfeldern, Z. Naturforsch. 10A, 508, 1955.

Kadomtsev, B., in Plasma Physics and the Problem of Controlled Thermonuclear Reactions, Akad. Nauk SSSR, Vol. III, 1958.

Kruskal, M. The Gyration of a Charged Particle, Princeton University, Project Matterhorn Report PM-S-33 (NYO-7903), March, 1958.

Kruskal, M., in The Theory of Neutral Ionized Gases, John Wiley & Sons, Inc., New York, 1960.

Newcomb, W., Motion of Magnetic Lines of Force, Ann. Phys. 3, 347, 1958.

Newcomb, W., Stability of a Collisionless Plasma, I and II, Ann. Phys. (to be published).

Northrop, T., Helmholtz Instability of a Plasma, Phys. Rev. 103, 1150, 1956.

Northrop, T., and E. Teller, Stability of the Adiabatic Motion of Charged Particles in the Earth's Field, Phys. Rev. 117, 215, 1960.

Northrop, T., The Guiding Center Approximation to Charged Particle Motion, Ann. Phys. 15, 79, 1961.

Northrop, T., The Adiabatic Motion of Charged Particles, John Wiley & Sons, Inc., New York (to be published, 1963).

Parker, E., Origin and Dynamics of Cosmic Rays, Phys. Rev. 109, 1328, 1958.

Rosenbluth, M., and C. Longmire, Stability of Plasma Confined by Magnetic Fields, Ann. Phys. 1, 120, 1957.

Spitzer, L., Jr., Equations of Motion for an Ideal Plasma, Astrophys. J. 116, 299, 1952.

Teller, E., Theory of Origin of Cosmic Rays, Rept. Progr. Phys. XVII, 154, 1954.

Vandervoort, P., The Relativistic Motion of a Charged Particle in an Inhomogeneous Electromagnetic Field, Ann. Phys. 10, 401, 1960.

FIGURE CAPTIONS

Fig. 1. The charged particle gyrates about its guiding center.

Fig. 2. A drift.

Fig. 3. Energy changes in a time-dependent field.

Fig. 4. Fermi acceleration.


Fig. 5. The mirror effect.

Fig. 6. Particle trajectory when \vec{E}_\perp is large.

Fig. 7. Mirror geometry needed for existence of second adiabatic invariant.


Fig. 8. An invariant surface for a particle trapped in the earth's field.

Fig. 9. A line of force in (α, β, s) space.



MU-29958

Fig. 1.



MU-29959

Fig. 2.






Fig. 3. Energy changes in a time-dependent field.




MU-29960

Fig. 4.




MU-29961

Fig. 5.




MU-29962

Fig. 6.




MU-29963

Fig. 7.



MU-29964

Fig. 8.



MU-29965

Fig. 9.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.