

82-1801

ORNL-5002

MASTER

Bioaccumulation Factors for Radionuclides in Freshwater Biota

Environmental Sciences Division Publication No. 783



OAK RIDGE NATIONAL LABORATORY
OPERATED BY UNION CARBIDE CORPORATION • FOR THE U.S. ATOMIC ENERGY COMMISSION

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

Printed in the United States of America. Available from
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road, Springfield, Virginia 22161
Price: Printed Copy \$7.60; Microfiche \$2.25

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the Energy Research and Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

ORNL-5002
UC-11 — Environmental
and Earth Sciences

Contract No. W-7405-eng-26

Environmental Sciences Division

BIOACCUMULATION FACTORS FOR RADIONUCLIDES
IN FRESHWATER BIOTA

Contributors:

Henry A. Vanderploeg
Dennis C. Parzyck
William H. Wilcox
James R. Kercher
Stephen V. Kaye

NOTICE
This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Energy Research and Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

NOVEMBER 1975

Research supported by the U. S. Nuclear Regulatory Commission under Interagency Agreement 40-494-75 with the U. S. Energy Research and Development Administration.

Environmental Sciences Division
Publication No. 783

OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37830
operated by
UNION CARBIDE CORPORATION
for the
U. S. ENERGY RESEARCH AND DEVELOPMENT ADMINISTRATION

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Rey

This report analyzes over 200 carefully selected papers to provide concise data sets and methodology for estimation of bioaccumulation factors for tritium and isotopes of strontium, cesium, iodine, manganese, and cobalt in major biotic components of freshwater environments. Bioaccumulation factors of different tissues are distinguished where significant differences occur. Since conditions in the laboratory are often unnatural in terms of chemical and ecological relationships, this review was restricted as far as possible to bioaccumulation factors determined for natural systems. Because bioaccumulation factors were not available for some shorter-lived radionuclides, a methodology for converting bioaccumulation factors of stable isotopes to those of shorter-lived radionuclides was derived and utilized.

The bioaccumulation factor for a radionuclide in a given organism or tissue may exhibit wide variations among bodies of water that are related to differences in ambient concentrations of stable-element and carrier-element analogues. To account for these variations, simple models are presented that relate bioaccumulation factors to stable-element and carrier-element concentrations in water. The effects of physicochemical form and other factors in causing deviations from these models are discussed. Bioaccumulation factor data are examined in the context of these models, and bioaccumulation factor relations for the selected radionuclides are presented.

SUMMARY

BIOACCUMULATION FACTOR CONCEPTS

The bioaccumulation factor for an organism or tissue i is the steady-state ratio of radionuclide concentration in the organism or tissue to that in water:

$$BF(R)_i = [R]_i / [R]_w$$

where

$BF(R)_i$ = bioaccumulation factor for radionuclide R in organism or tissue i ,

$[R]_i$ = radionuclide concentration ($\mu\text{Ci/g}$ fresh weight) in organism or tissue i , and

$[R]_w$ = radionuclide concentration in water ($\mu\text{Ci/g}$), a constant.

Bioaccumulation factors are used to predict radionuclide concentrations in whole organisms or their tissues from knowledge of the radionuclide concentration in water for chronic releases of radionuclides. Bioaccumulation factors for radionuclides were related to ambient concentrations of their stable or non-isotopic carrier element analogues according to three idealized patterns. The first pattern is that the bioaccumulation factor for radionuclide R in organism or tissue i , $BF(R)_i$, is constant regardless of stable-element or carrier-element concentration:

$$BF(R)_i = \text{const.} \quad (1)$$

The second pattern applies to an element that is homeostatically

maintained at a constant concentration in organism or tissue i:

$$BF(R)_i = \Sigma_i / [C]_w , \quad (2)$$

where

Σ_i = concentration of stable element in organism or tissue i, a constant ($\mu\text{g/g}$ fresh weight), and

$[C]_w$ = concentration of corresponding stable element in water ($\mu\text{g/g}$).

The third pattern applies to a radioisotope and its non-isotopic carrier element which is homeostatically maintained at a constant concentration in i:

$$BF(R)_i = \frac{q_i \Sigma_i^*}{[C^*]_w} , \quad (3)$$

where

q_i = discrimination coefficient,

Σ_i^* = concentration of non-isotopic carrier element in organism or tissue i, a constant ($\mu\text{g/g}$ fresh weight), and

$[C^*]_w$ = concentration of non-isotopic carrier element in water ($\mu\text{g/g}$).

The discrimination coefficient, q_i , is the ratio $([R]_i / [C^*]_i) / ([R]_w / [C^*]_w)$, where $[C^*]_i$ is the carrier element concentration in i. Equation (3) is often rewritten in the form:

$$\ln BF(R)_i = \ln q_i \Sigma_i^* - \ln [C^*]_w \quad (4)$$

Radionuclides exist in a wide variety of physicochemical forms in natural waters, and their different forms have different availabilities to the food chain. Sediments, too, may be source of radionuclides to biota, and sediment type can influence the availability. For those elements that are homeostatically maintained at constant concentrations in a given organism, the concentration of stable element in the organism is independent of concentration of stable element in water or its availability in prey, sediment, and different physicochemical forms in the water. In contrast, differences in availability of radionuclides in different sediment types and different physicochemical forms in water can lead to marked deviations from the idealized patterns of Equations (1) and (3).

When bioaccumulation factors were not available for radionuclides, they were estimated from bioaccumulation factors of the corresponding stable elements. Owing to physical decay, the bioaccumulation factor of a short-lived radionuclide is much less than that of the stable element or longer-lived nuclides. Bioaccumulation factors of shorter-lived radionuclides were estimated from bioaccumulation factors of the stable element according to the relation:

$$BF(R)_i = \frac{k}{k+\lambda} \quad BF(C)_i , \quad (5)$$

where

k = elimination coefficient of C in i (day^{-1}),

λ = radioactive decay constant (day^{-1}), and

$BF(C)_i$ = bioaccumulation factor of corresponding stable element
in i .

BIOACCUMULATION FACTORS

Cesium

Potassium is a non-isotopic carrier element for cesium because of their chemical similarities and the greater abundance of potassium in water. Further, since potassium concentration is homeostatically maintained at constant concentrations in animals, the bioaccumulation factor of cesium is given by Equation (3) or (4). Unlike this pattern for animals, potassium concentration of water has only a small effect on the cesium bioaccumulation factor in algae.

The primary mode of accumulation of cesium and potassium in aquatic animals is from the food chain. Absorption efficiency of potassium and cesium from food is high. In animals the excretion coefficient of potassium is about 3 times larger than the excretion coefficient of cesium. As a result, q_i increases by a factor of about 3 with each trophic level. If potassium concentrations in the predators and prey are about equal, the cesium bioaccumulation factor increases by a factor of 3 with each trophic level.

Because cesium is strongly adsorbed by suspended particulate materials, especially clays, the proportion of cesium in the soluble phase decreases with increasing suspended solids concentrations. Potassium, too, is sorbed but to a much lesser degree. Since algae accumulate cesium, potassium, and other elements from the soluble phase, the availabilities to the food chain of cesium and of cesium

relative to potassium decrease with increasing suspended solids concentrations. Thus, discrimination coefficients and bioaccumulation factors decrease with increasing suspended solids concentrations.

On the basis of data available, we recommend the bioaccumulation factors given in Table 1.

Strontium

Calcium is a non-isotopic carrier element for strontium because of their chemical similarities and the greater abundance of calcium in water. Further, since calcium concentration is homeostatically maintained at constant concentrations in animals, the bioaccumulation factor of strontium is given by Equation (3) or (4). Unlike this pattern for animals, calcium concentration of water has only a small effect on the strontium bioaccumulation factor in algae. The primary mode of uptake of strontium and calcium in animals (as well as plants) is from water. As a result, trophic level has little effect on the discrimination coefficient and the bioaccumulation factor of strontium. Further, the discrimination coefficient is relatively independent of calcium concentration in water. Calcium and strontium concentrations in bones and shells are higher than in other tissues of animals.

Strontium has physicochemical properties similar to calcium and, like calcium, appears mainly as free ions in water. As a result of this and the fact that the discrimination coefficient is independent of calcium concentration in water, the discrimination coefficient varies little among sites. This implies that the product $q_j \Sigma_j^*$

Table 1. Recommended bioaccumulation factor relations for long-lived isotopes of cesium^a

Taxon/ Functional Group	Tissue	Environment	Recommended Bioaccumulation Factor Relation
Piscivorous fishes	All tissues	Clear waters ^b	$1.5 \times 10^4 / [K]_w$ ^c
Piscivorous fishes	All tissues	Turbid waters ^d	$3 \times 10^3 / [K]_w$
Non-piscivorous fishes	All tissues	Clear waters	$5 \times 10^3 / [K]_w$
Non-piscivorous fishes	All tissues	Turbid waters	$1 \times 10^3 / [K]_w$
Algae	Whole	All waters	10^3
Aquatic vascular plants	Whole	All waters	10^3
Emergent vascular plants	Whole	All waters	10^3
Molluscs	Shell Soft tissues	All waters All waters	10^2 10^3
Invertebrates other than molluscs	Whole	All waters	10^3
Amphibians	All tissues	All waters	10^4
Waterfowl	All tissues	All waters	3×10^3

^aTo convert to bioaccumulation factors of ^{136}Cs use conversion factors in Table 3.1.1.

^bSuspended solids concentration less than 50 ppm.

^c $[K]_w$ = stable potassium concentration of water in ppm.

^dSuspended solids concentration greater than 50 ppm.

appearing in the numerator of Equation (3) may be treated as a constant and that a regression of $\ln BF(Sr)_i$ versus $\ln [Ca]_w$, where $[Ca]_w$ is calcium concentration in water, will have a slope near -1 and a high correlation coefficient.

On the basis of the data available, we recommend the strontium bioaccumulation factors given in Table 2 and shown in Figure 1.

Tritium

Tritium was included in this report because of a concern that the relative kinetics of tritium and protium resulting from tritium's heavier mass ("isotope effect") might lead to a preferential accumulation of tritium over protium. Experiments in aquatic systems indicate that this does not occur and that the bioaccumulation factor for tritium is less than or about equal to the bioaccumulation factor for protium, which has a bioaccumulation factor approximately equal to 1. We recommend that a bioaccumulation factor of 1 be used for all tissues of all aquatic biota (Table 3).

Iodine

Iodine is highly concentrated by the thyroid tissue of vertebrates. As a result, the bioaccumulation factor for iodine in the thyroid tissue of fishes is very high. Recommended bioaccumulation factors are given in Table 4.

Manganese

Manganese is homeostatically maintained at constant concentrations in vertebrates and some invertebrates. Thus, the bioaccumulation

Table 2. Recommended bioaccumulation factor relations for strontium

Taxon/ Functional Group	Tissue	Environment	Recommended Bioaccumulation Factor Relation
Fishes	Bone	All waters	Figure 1
	Flesh	All waters	Figure 1
Algae	Whole	All waters	2×10^3
Vascular plants (aquatic and emergent)	Whole	All waters	2×10^2
Molluscs	Shell	All waters	$6.8 \times 10^4 / [\text{Ca}]_w^a$
	Soft tissues	All waters	3×10^2

^a $[\text{Ca}]_w$ = stable calcium concentration of water in ppm.

ORNL-DWG 75-4447R




Figure 1 Bioaccumulation factors for strontium in freshwater fishes as a function of calcium concentration in water.

Table 3. Recommended bioaccumulation factor for tritium

Taxon/ Functional Group	Tissue	Environment	Recommended Bioaccumulation Factor
All organisms	All tissues	All waters	1

Table 4. Recommended bioaccumulation factors for stable iodine, iodine-129, and iodine-131

Taxon/ Functional Group	Tissue	Environment	Recommended Bioaccumulation Factor Relation
<u>Stable Iodine and Iodine-129</u>			
Fishes	Muscle	All waters	50
	Ovary (eggs)		800
	Thyroid tissue		290,000
Crustacea		All waters	340
Phytoplankton		All waters	800
<u>Iodine-131</u>			
Fishes	Muscle	All waters	40
	Ovary (eggs)		800 ^a
	Thyroid tissue		110,000
Aquatic insect larvae	Whole	All waters	400
Molluscs	Soft tissues	All waters	50
	Shell		400
Algae	Whole	All waters	260
Macrophytes	Whole	All waters	120

^aDefault value based on bioaccumulation factor of stable iodine.

factor of manganese in fishes is inversely proportional to manganese concentration in water [Equation (2)]. Recommended bioaccumulation factors are given in Table 5.

Cobalt

Cobalt in solution has a strong tendency to form complexes with dissolved organic matter. Since metals complexed with organic molecules have significantly lower availabilities to plants and animals than free ions, cobalt bioaccumulation factors would be expected to decrease with increasing eutrophy of water. Cobalt bioaccumulation factors, which in this report are based on cobalt concentrations in the soluble phase of water, conform to this expected pattern.

Absorption efficiency of cobalt from food is very low in fishes. This may explain the relatively low bioaccumulation factors of cobalt in fishes. Recommended bioaccumulation factors for cobalt are given in Table 6.

Table 5. Recommended bioaccumulation factor relations for manganese

Taxon/ Functional Group	Tissue	Environment	Recommended Bioaccumulation Factor Relation
Fishes	Whole	All waters	$6.7/[\text{Mn}]_w$ ^a
	Flesh	All waters	$0.32/[\text{Mn}]_w$
Algae	Whole	All waters	10^4
Submerged macrophytes	Whole	All waters	10^4
Floating-leaf macrophytes	Whole	All waters	10^3
Emergent vascular plants	Whole	All waters	10^3
Molluscs: Bivalves	Soft	All waters	10^5
	Shell		3×10^4
Snails	Soft		2×10^3
	Shell		10^4
	Whole		10^4
Crustaceans	Whole	All waters	10^4
Insect larvae	Whole	All waters	10^3

^a $[\text{Mn}]_w$ = stable manganese concentration in water in ppm.

Table 6. Recommended bioaccumulation factors for cobalt
(based on isotope concentrations in water after
filtration to remove particulates)

Taxon/ Functional Group	Tissue	Environment	Recommended Bioaccumulation Factors
Fishes	Flesh ^a	Mesotrophic and oligo-trophic waters	320
	Whole ^a		440
	Flesh ^a	Eutrophic waters	27
	Whole		44
Algae	Whole	All waters	10^4
Emergent vascular plants	Whole	All waters	10^3
Submerged- leaf macrophytes	Whole	Mesotrophic and oligo-trophic waters	10^4
		Eutrophic waters	2×10^3
Floating-leaf macrophytes	Whole	Mesotrophic and oligo-trophic waters	10^3
		Eutrophic waters	4×10^2
Molluscs: Bivalves	Soft	Mesotrophic and oligo-trophic waters	10^4
		Eutrophic waters	4×10^2
Snails	Soft	All waters	10^4
Insect larvae	Whole	All waters	10^4
Tubificid worms	Whole	Eutrophic waters	5×10^2
Crustaceans	Whole	All waters	10^3

^aBioaccumulation factors of other tissues of fishes may be estimated from relative concentrations of cobalt given in Table 3.6.2.

CONTENTS

	<u>Page</u>
1. Introduction	1
1.1 Scope	1
1.2 Objectives	1
1.3 Format	2
2. Bioaccumulation Factor Concepts	5
2.1 Idealized Bioaccumulation Factor Patterns	5
2.2 Physicochemical Forms of Radionuclides and Availability to Biota	9
2.3 Bioaccumulation Factors and Availability of Radionuclides	10
2.4 Bioaccumulation Factors for Short-Lived Radionuclides	11
2.5 Experimental Error in Determination of Bioaccumulation Factors	14
2.5.1 Analytical Error	14
2.5.2 Use of Literature Values for Water Concentrations	15
2.5.3 Filtration of Water Samples	16
2.5.4 Determination of Organism Concentration Based on Dry Weight	17
2.5.5 Averaging Isotope Concentrations of Different Tissues	17
2.5.6 Determinations Made Under Nonsteady-State Conditions	18
2.6 Use of Bioaccumulation Factors in Radiation Dose Estimation	19
2.6.1 Dose to Biota	19
2.6.2 Dose to Man	20
3. Bioaccumulation Factors for Cesium, Strontium, Tritium, Iodine, Manganese and Cobalt	25
3.1 Cesium	25
3.1.1 Cesium Metabolism	25
3.1.2 Environmental Cesium	28
3.1.3 Review of Cesium Bioaccumulation Factors	32
3.1.3.1 Cesium Bioaccumulation Factors for Fishes	32
3.1.3.2 Cesium Bioaccumulation Factors for Algae	44

CONTENTS (continued)

	<u>Page</u>
3.1.3.3 Cesium Bioaccumulation Factors for Vascular Aquatic Plants	48
3.1.3.4 Cesium Bioaccumulation Factors for Emergent Vascular Plants	50
3.1.3.5 Cesium Bioaccumulation Factors for Invertebrates	50
3.1.3.6 Cesium Bioaccumulation Factors for Amphibians	50
3.1.3.7 Cesium Bioaccumulation Factors for Waterfowl	50
3.2 Strontium	63
3.2.1 Strontium Metabolism	63
3.2.2 Environmental Strontium	64
3.2.3 Review of Strontium Bioaccumulation Factors	64
3.2.3.1 Strontium Bioaccumulation Factors for Fishes	66
3.2.3.2 Strontium Bioaccumulation Factors for Algae	67
3.2.3.3 Strontium Bioaccumulation Factors for Aquatic and Emergent Macrophytes	78
3.2.3.4 Strontium Bioaccumulation Factors for Molluscs	78
3.3 Tritium	87
3.3.1 Hydrogen Bioaccumulation	87
3.3.2 Pathways of Hydrogen Entry into Aquatic Organisms	88
3.3.3 Compartment Size and Turnover Rates	90
3.3.4 Tritium Concentration in Aquatic Food Chains	93
3.3.4.1 Tissue-Water Tritium	93
3.3.4.2 Tissue-Bound Tritium	93
3.3.4.2.1 Exchangeable Tissue-Bound Tritium	94
3.3.4.2.2 Nonexchangeable Tissue-Bound Tritium	96
3.4 Iodine	103
3.4.1 Environmental Iodine	103

CONTENTS (continued)

	<u>Page</u>
3.4.2 Iodine Metabolism	103
3.4.3 Review of Iodine Bioaccumulation Factors	105
3.4.3.1 Stable Iodine Bioaccumulation Factors for Fish Tissue	105
3.4.3.1.1 Stable Iodine Bioaccumulation Factors for Fish Muscle	109
3.4.3.1.2 Stable Iodine Bioaccumulation Factors for Fish Thyroids and Ovaries	110
3.4.3.2 Iodine-131 Bioaccumulation Factors for Fish Tissue	114
3.4.3.2.1 Iodine-131 Bioaccumulation Factors for Fish Muscle	114
3.4.3.2.2 Iodine-131 Bioaccumulation Factors for Fish Thyroids	117
3.4.3.3 Stable Iodine Bioaccumulation Factors for Plants	118
3.4.3.4 Iodine-131 Bioaccumulation Factors for Plants	118
3.4.3.5 Stable Iodine Bioaccumulation Factors for Invertebrates	118
3.4.3.6 Iodine-131 Bioaccumulation Factors for Invertebrates	121
3.5 Manganese	129
3.5.1 Environmental Manganese	129
3.5.2 Manganese Metabolism	130
3.5.3 Review of Manganese Bioaccumulation Factors	131
3.5.3.1 Manganese Bioaccumulation Factors for Fishes	131
3.5.3.2 Manganese Bioaccumulation Factors for Plants	136
3.5.3.3 Manganese Bioaccumulation Factors for Invertebrates	140
3.5.3.3.1 Manganese Bioaccumulation Factors for Molluscs	140
3.5.3.3.2 Manganese Bioaccumulation Factors for Invertebrates Other than Molluscs	145

CONTENTS (continued)

	<u>Page</u>
3.6 Cobalt	151
3.6.1 Cobalt Metabolism	151
3.6.2 Environmental Cobalt and Availability	153
3.6.3 Review of Cobalt Bioaccumulation Factors	154
3.6.3.1 Cobalt Bioaccumulation Factors for Fishes	155
3.6.3.2 Cobalt Bioaccumulation Factors for Plants	166
3.6.3.3 Cobalt Bioaccumulation Factors for Invertebrates	169
Bibliography	179
Appendix	211

TABLES

<u>Table</u>		<u>Page</u>
3.1.1	Biological half-lives and elimination coefficients for cesium and $[k/(k+\lambda)]$'s for ^{136}Cs	26
3.1.2	Sorption of ^{134}Cs to various concentrations of clay suspended in distilled water	29
3.1.3	Percentage of cesium in water in the particulate phase in some freshwater ecosystems	30
3.1.4	Mean ^{137}Cs bioaccumulation factors for whole fishes from Finnish lakes	34
3.1.5	Bioaccumulation factors for cesium as related to potassium and suspended solids concentrations	36
3.1.6	Bioaccumulation factors for cesium in fishes	37
3.1.7	Bioaccumulation factors for cesium in freshwater algae . . .	45
3.1.8	Bioaccumulation factors for cesium in aquatic vascular plants	49
3.1.9	Bioaccumulation factors for cesium in emergent vascular plants	51
3.1.10	Bioaccumulation factors for cesium in aquatic invertebrates	52
3.1.11	Bioaccumulation factors for ^{137}Cs in amphibians	54
3.1.12	Bioaccumulation factors for ^{137}Cs in waterfowl	55
3.2.1	Percentage of radiostrontium in water in the particulate phase in some freshwater ecosystems	65
3.2.2	Bioaccumulation factors and discrimination coefficients for strontium in fishes	68
3.2.3	Parameters from linear regressions between $\text{BF}(\text{Sr})_i$ and $[\text{Ca}]_w$ for bone and flesh of fishes	75
3.2.4	Bioaccumulation factors and discrimination coefficients for strontium in algae	76
3.2.5	Bioaccumulation factors and discrimination coefficients for strontium in aquatic and emergent macrophytes	79

TABLES (continued)

<u>Table</u>		<u>Page</u>
3.2.6	Bioaccumulation factors and discrimination coefficients for strontium in mollusc shells	80
3.3.1	Component size and source contribution for a hypothetical fish	92
3.3.2	Steady-state concentration in organic groups common in living tissue	95
3.4.1	Iodine concentrations in surface waters, the ocean, and geologic sources	104
3.4.2	Elimination rates for iodine-131 in aquatic vertebrates . . .	106
3.4.3	Stable iodine in freshwater fish muscle	107
3.4.4	Stable iodine in anadromous and marine fish muscle	108
3.4.5	Total iodine present in the thyroid and eggs of five Rainbow and California steelhead trout	112
3.4.6	Thyroid stable iodine bioaccumulation factors for marine fish	113
3.4.7	Stable iodine bioaccumulation factor for fish ovaries (eggs) derived from relative concentrations of stable iodine in muscle and ovaries	115
3.4.8	Iodine-131 bioaccumulation factors for muscle in freshwater animals	116
3.4.9	Iodine-131 bioaccumulation factors for freshwater plants	119
3.4.10	Stable iodine bioaccumulation factors for invertebrates . .	120
3.4.11	Iodine-131 bioaccumulation factors for invertebrates . . .	122
3.5.1	Bioaccumulation factors for manganese in freshwater fishes	132
3.5.2	Manganese concentrations and coefficients of variation in water and freshwater fishes	135
3.5.3	Bioaccumulation factors for manganese in aquatic plants . .	138

TABLES (continued)

<u>Table</u>		<u>Page</u>
3.5.4	Bioaccumulation factors for manganese in molluscs	141
3.5.5	Bioaccumulation factors for manganese in invertebrates other than molluscs	146
3.6.1	Biological elimination rates for ⁶⁰ Co in various aquatic organisms	152
3.6.2	Relative steady-state concentrations of ⁶⁰ Co in tissues of <i>Ictalurus melas</i> (black bullheads) from White Oak Lake	156
3.6.3	Bioaccumulation factors for cobalt in fishes from eutrophic environments (based on filtered water concentrations)	157
3.6.4	Bioaccumulation factors for cobalt in fishes from mesotrophic environments (based on filtered water concentrations)	161
3.6.5	Mean bioaccumulation factors for cobalt in fishes from mesotrophic and eutrophic environments	163
3.6.6	Bioaccumulation factors for cobalt in aquatic plants	167
3.6.7	Mean cobalt bioaccumulation factors for submerged and floating-leaf vascular plants from Perch Lake and Lake Maggiore (based on filtered water concentrations) . . .	170
3.6.8	Bioaccumulation factors for cobalt in invertebrates (based on filtered water concentrations)	171

FIGURES

<u>Figure</u>	<u>Page</u>
3.1.1 Bioaccumulation factors for ^{137}Cs and stable cesium in freshwater fishes as a function of potassium concentration in water	43
3.2.1 Bioaccumulation factors for strontium in freshwater fishes as a function of calcium concentration in water	74
3.3.1 Pathways of hydrogen entry into the hydrogen components of an aquatic animal	89
3.3.2 Three-link food chain diagram with organisms compartmentalized	97
3.5.1 Bioaccumulation factors for manganese in freshwater fishes as a function of manganese concentration in water	137
3.6.1 Concentration of stable cobalt in fish flesh in various environments	165

1. INTRODUCTION

1.1 Scope

This report analyzes over 200 carefully selected papers to provide concise data sets and methodology for estimation of bioaccumulation factors for selected radionuclides in major biotic components of freshwater environments. The papers were critically reviewed from the standpoint of experimental techniques for each of the selected radionuclides. Since conditions in the laboratory are often unnatural in terms of chemical and ecological relationships, this review is restricted as far as possible to bioaccumulation factors determined for natural systems. Cesium, strontium, iodine, manganese, and cobalt were chosen based on an ORNL survey of releases from 16 light-water-cooled nuclear power stations. This survey showed that the major radiological impact on the aquatic biota and man comes from these radionuclides (Kaye, 1973). Although tritium contributes only a minor fraction of the dose from liquid effluents, it is nonetheless treated too because of a concern that it may be concentrated in passage through food chains. Bioaccumulation factors for freshwater environments are of immediate concern since most nuclear plants (operable, under construction, or planned) are located adjacent to bodies of freshwater.

1.2 Objectives

The objective of this report is to present bioaccumulation factors for the selected radionuclides in the different tissues of major biotic components of freshwater systems. The bioaccumulation factor of a radionuclide in an organism or tissue may exhibit wide variations among sites

that are related to differences in stable-element and carrier-element concentrations and other limnological variables. For this reason, bioaccumulation factor relations rather than single bioaccumulation factors are presented to account for site-specific variations.

1.3 Format

General discussions of important aspects of element bioaccumulation are presented in Section 2. We present simple models to illustrate the relation between bioaccumulation factors for radionuclides and the concentrations of their stable-element and carrier-element analogues. The effects of physicochemical form and other factors in causing deviations from these models are also discussed along with methodology for converting stable-element bioaccumulation factors to bioaccumulation factors for shorter-lived radionuclides. Section 3 reviews the literature on bioaccumulation factors for the six selected radionuclides and gives recommended bioaccumulation factor relations based on the literature values.

Reference

Kaye, S. V., Assessing potential radiological impacts to aquatic biota in response to the National Environmental Policy Act (NEPA) of 1969. IN Symposium on Environmental Behavior of Radionuclides Released in the Nuclear Industry, IAEA Symposium, Aix-en-Provence, France, 14-18 May 1973.

2. BIOACCUMULATION FACTOR CONCEPTS

The bioaccumulation factor of an organism or tissue *i* is the steady-state ratio of radionuclide concentration in the organism or tissue to that in water:

$$BF(R)_i = [R]_i/[R]_w$$

where

$BF(R)_i$ = bioaccumulation factor for radionuclide *R* in organism or tissue *i*,

$[R]_i$ = radionuclide concentration ($\mu\text{Ci/g}$ fresh weight) in organism or tissue *i*, and

$[R]_w$ = radionuclide concentration in water ($\mu\text{Ci/g}$), a constant.

Bioaccumulation factors are used to predict radionuclide concentrations in whole organisms or their tissues from knowledge of the radionuclide concentration in water for chronic releases of radionuclides. Bioaccumulation factors also appear in expressions that are used to predict the transient dynamics of radionuclide concentration in organisms (Vanderploeg et al., 1974). Therefore, an understanding of variables that affect bioaccumulation factors is central to understanding steady-state and transient dynamics of radionuclide concentration in aquatic organisms.

2.1 Idealized Bioaccumulation Factor Patterns

Bioaccumulation factors for radionuclides are often portrayed as being related to ambient concentrations of their stable or non-isotopic

carrier element analogues according to three idealized patterns (e.g., Fleishman, 1973; Peterson, 1970). The first pattern is that the bioaccumulation factor for radionuclide R in organism or tissue i, $BF(R)_i$, is constant regardless of stable-element or carrier-element concentrations:

$$BF(R)_i = \text{const.} \quad (2.1.1)$$

The second pattern is that the concentration of an element in an organism or tissue is homeostatically maintained at a constant concentration despite different concentrations of the element in water. This implies that the bioaccumulation factor of a homeostatically controlled stable element is inversely proportioned to its concentration in the water:

$$BF(C)_i = \Sigma_i / [C]_w , \quad (2.1.2)$$

where Σ_i = concentration of stable element in organisms or tissue i, a constant ($\mu\text{g/g}$ fresh weight), and $[C]_w$ = concentration of stable element in water ($\mu\text{g/g}$).

Assuming that the specific activity, that is, the ratio of radionuclide concentration to the stable element concentration ($\mu\text{Ci/g}$ stable element) in i is equal to that of the water, the bioaccumulation factor for the radionuclide, $BF(R)_i$, is also inversely proportional to the concentration of the stable element:

$$BF(R)_i = \Sigma_i / [C]_w . \quad (2.1.3)$$

Taking the natural logarithm of each side of Equation (2.1.2) and (2.1.3) results in the linear relation:

$$\ln BF(R)_i = \ln BF(C)_i = \ln \Sigma_i - \ln [C]_w . \quad (2.1.4)$$

Equation (2.1.4) implies that the plot of $\ln BF(R)_i$ versus $\ln [C]_w$ has a slope of -1. The third pattern is that the bioaccumulation factor for the radionuclide, $BF(R)_i$, is inversely proportional to the concentration of a non-isotopic carrier element in water, a non-isotopic carrier element being nearly chemically similar to but occurring in higher concentrations than the stable-element analogue. The derivation of this relationship follows.

The bioaccumulation factor for radionuclide R is related to $BF(C^*)_i$, the bioaccumulation factor for the carrier element C^* , by

$$BF(R)_i = BF(C^*)_i \frac{BF(R)_i}{BF(C^*)_i} , \quad (2.1.5)$$

which may be written as

$$BF(R)_i = BF(C^*)_i \frac{(R/C^*)_i}{(R/C^*)_w} , \quad (2.1.6)$$

where $(R/C^*)_i$ and $(R/C^*)_w$ are ratios of radionuclide concentration to carrier element concentration found in the organism or tissue and in the water, respectively. Assume that the non-isotopic carrier element is homeostatically controlled in the organism, that is:

$$BF(C^*)_i = \Sigma_i^* / [C^*]_w , \quad (2.1.7)$$

where

Σ_i^* = concentration of non-isotopic carrier element in organism or tissue i , a constant ($\mu\text{g/g}$ fresh weight), and

$[C^*]_w$ = concentration of non-isotopic carrier element in water ($\mu\text{g/g}$).

Combining Equations (2.1.6) and (2.1.7) and letting $q_i = (R/C^*)_i/(R/C^*)_w$,

$$BF(R)_i = \frac{q_i \Sigma_i^*}{[C^*]_w} . \quad (2.1.8)$$

Taking the natural logarithm of each side of Equation (2.1.8),

$$\ln BF(R)_i = \ln q_i \Sigma_i^* - \ln [C^*]_w . \quad (2.1.9)$$

Since Σ_i^* is a constant, a plot of $\ln BF(R)_i$ versus $\ln [C^*]_w$ will have a slope of -1 if q_i is constant.

If water were the immediate source of the radionuclide and carrier element to organism or tissue i , q_i would be called the "observed ratio" (Comar et al., 1956). Since water may not necessarily be the direct source of an element to an organism, we will designate q_i the discrimination coefficient.

On the basis of empirical data on ^{137}Cs and its carrier element potassium, Fleishman (1973) suggested that q_i is not a constant but rather a function of $[C^*]_w$. It can be shown mathematically that if an animal accumulates the carrier and nuclide from water and food, q_i becomes a function of $[C^*]_w$ (see Appendix).

Because the number of sites where steady-state bioaccumulation factors can be determined from the release of radionuclides is limited, bioaccumulation factors for the radionuclides are often estimated from

distributions of their stable element analogues. Therefore, the requirement of equal specific activities in organisms and water that we have for Equation (2.1.3) applies as well to Equations (2.1.1) and (2.1.8) or any relation when stable element data are used to estimate bioaccumulation factors.

2.2 Physicochemical Forms of Radionuclides and Availability to Biota

Radionuclides exist in a wide variety of physicochemical forms in natural waters and their different physicochemical forms have different availabilities to aquatic biota. The major physicochemical forms of trace metals that Gibbs (1973) reports for rivers illustrate this diversity:

1. Dissolved ionic species and inorganic associations,
2. Complexes with organic molecules in solution,
3. Adsorbed to solids,
4. Precipitates and coprecipitates on solids (metallic coatings),
5. Incorporated in solid biological materials, and
6. Incorporated in crystalline structures.

Both unicellular and multicellular algae, which generally form the base of the aquatic food web, accumulate elements from the soluble phase. Likewise, some radionuclides are directly accumulated by aquatic animals from the soluble phase. Aquatic vascular plants, which may be important as food to consumers in some systems, accumulate elements from the soluble phase of water and also from the interstitial water of sediment. It has also been demonstrated that radionuclides and metals complexed

with organic molecules may have significantly lower availabilities to algae and animals than free ions (Timofeeva et al., 1960).

Aquatic animals, especially filter feeders, may accumulate radionuclides from the suspended phase. The availability of different forms of radionuclides in the suspended phase has not been studied. It can be said that adsorbed (exchangeable) radionuclides and radionuclides in solid biological materials are available to some aquatic consumers. Elements in the crystal structure of clay minerals and other lithogenous detritus are almost completely unavailable.

Radionuclides enter food webs not only from the water but also from bottom sediments, and availability of radionuclides to the food web varies with sediment type. Benthic invertebrates accumulate radionuclides from bottom sediments. Fishes may accumulate radionuclides indirectly from bottom sediment by ingestion of these invertebrates and also directly by incidental ingestion of sediment with prey (Gallegos et al., 1970). Absorption of radionuclides from ingested sediment varies with the nature of the sediment (Gallegos et al., 1970; Eyman and Kitchings, 1974). In addition, different sediments have different capacities for sorption of radionuclides (Friend, 1963; Lomenick and Gardiner, 1965).

2.3 Bioaccumulation Factors and Availability of Radionuclides

For those elements that are homeostatically maintained at constant concentrations in a given organism, the concentration of stable element in the organism is independent of concentrations of stable element in water or its availability from prey, sediment, and different physico-chemical forms of the water. If the radionuclide in the organism has

the same specific activity as that in water, the bioaccumulation factor for the radionuclide will be given by Equation (2.1.3). In contrast, differences in availability of radionuclides in different sediments and different physicochemical forms in water can lead to marked deviations from the idealized pattern of a constant bioaccumulation factor, Equation (2.1.1).

For the bioaccumulation factor pattern of Equation (2.1.8), we pointed out that q_i is a function of $[C^*]_w$. We note too that if sediment is an important source of radionuclide to the animal, the concentration of carrier in sediment relative to food and water will have an influence on q_i (see Appendix). Moreover, variations in the proportions of radionuclide and carrier element appearing in various physicochemical forms can lead to variations in q_i since the radionuclide or carrier could be preferentially accumulated by the food web.

2.4 Bioaccumulation Factors for Short-Lived Radionuclides

Because of radioactive decay, the bioaccumulation factor for a short-lived radionuclide is expected to be less than the corresponding bioaccumulation factor of the stable element. Since the bioaccumulation factors of some stable elements have been more commonly measured than the bioaccumulation factors of short-lived radionuclides, a methodology is presented to predict the bioaccumulation factor for the short-lived radionuclides from the bioaccumulation factor of the stable element, $BF(C)_i$. The methodology requires knowledge of the elimination coefficient, k , in organism or tissue i .

Treating organism or tissue i as a single compartment, the rate of change in the amount of radionuclide R in i , R_i , (μCi) is

$$\frac{dR_i}{dt} = I_T - R_i (k + \lambda), \quad (2.4.1)$$

and the steady-state amount of stable element C is given by

$$C_i = \frac{I_T'}{k} \quad (2.4.2)$$

where

I_T = rate of input of radionuclide from all sources ($\mu\text{Ci/day}$),

I_T' = rate of input of the corresponding stable element from all sources ($\mu\text{g/day}$),

k = elimination coefficient at steady-state concentration of C in i (day^{-1}), and

λ = radioactive decay constant (day^{-1}).

Assuming constant weight of the organism or tissue, the steady-state concentration of radionuclide R and stable element C are given by

$$[R]_i = \frac{I_T}{B(k + \lambda)} \quad (2.4.3)$$

$$[C]_i = \frac{I_T'}{Bk} \quad (2.4.4)$$

where B = biomass or fresh weight of the organism or tissue i (g).

Assuming all uptake is from water,

$$[R]_i = \frac{a [R]_w}{B(k + \lambda)} \quad (2.4.5)$$

$$[C]_i = \frac{a [C]_w}{Bk} , \quad (2.4.6)$$

where a = uptake coefficient for element C from water (day⁻¹),

$[R]_w$ = radionuclide concentration in water ($\mu\text{Ci/g}$), and

$[C]_w$ = stable element concentration in water ($\mu\text{g/g}$).

Dividing Equation (2.4.5) by $[R]_w$ and Equation (2.4.6) by $[C]_w$ gives

$$BF(R)_i = \frac{a}{B(k + \lambda)} \quad (2.4.7)$$

$$BF(C)_i = \frac{a}{Bk} \quad (2.4.8)$$

which implies that

$$BF(R)_i = \frac{k}{k + \lambda} \quad BF(C)_i \quad (2.4.9)$$

for uptake from water. The derivation of Equation (2.4.9) parallels that given by Peterson (1970). Thus if the radioactive decay constant, λ , is significant relative to the elimination coefficient, k , $BF(R)_i$ will be significantly less than $BF(C)_i$. If there are intervening steps in the food chain between the organism or tissue and water, radioactive decay will occur at each step and further diminish $BF(R)_i$. Since the elimination coefficients of prey (or forage) are generally much larger than the elimination coefficients of their respective predators, Equation (2.4.9) will generally not greatly overpredict $BF(R)_i$. Since Equation (2.4.9) will not greatly overpredict the $BF(R)_i$ and since information on food web structure is needed to derive a more accurate $BF(R)_i$, we suggest that Equation (2.4.9) be used to convert bioaccumulation factors for stable elements or long-lived nuclides to bioaccumulation factors for

short-lived radionuclides. Tables of k and $[k/(k+\lambda)]$ values are given when the bioaccumulation factors for shorter-lived isotopes must be estimated from the bioaccumulation factors for stable elements.

2.5 Experimental Error in Determination of Bioaccumulation Factors

In earlier sections we treated the effects of variations in certain environmental conditions on bioaccumulation factors. Variations also result from experimental error, that is, errors resulting from artifacts of analysis, evaluation and presentation of data. Below we discuss some important sources of experimental error and the precautions which can be taken to minimize them. These important sources of experimental error are:

- a. Analytical error,
- b. Use of literature values for water concentrations,
- c. Filtration of water samples,
- d. Determination of organism concentration based on dry weight,
- e. Averaging isotopic concentrations of different tissues, and
- f. Determinations made under nonsteady-state conditions.

2.5.1 Analytical Error

The concentrations of radionuclides or stable elements in most environmental samples are generally so low that serious problems can be encountered in their measurement, especially in water. Eight laboratories participated in an intercalibration study involving spiked freshwater samples (Maletskos, 1972). The results were given as a ratio of the reported values to the "correct" values \pm the

standard deviation. The results were: $^{54}\text{Mn} = 0.91 \pm 0.20$, $^{60}\text{Co} = 0.97 \pm 0.40$, $^{137}\text{Cs} = 1.05 \pm 0.27$, and $^{90}\text{Sr} = 1.09 \pm 0.46$. Iodine and tritium were not included in this study. Although it is not likely that errors in analysis of biological material would parallel those occurring in water analysis, it is obvious that analytical error can result in a significant variation in measured bioaccumulation factors. Significant error may also be associated with measuring stable element concentrations in water, especially for cesium, cobalt and manganese.

Unfortunately, it is not generally possible to evaluate the accuracy of measurements from particular studies. In deriving our bioaccumulation factor relations, we implicitly assumed that variations in bioaccumulation factors arising from analytical error were small in relation to variations caused by environmental variables and that deviations caused by analytical error were randomly distributed.

2.5.2 Use of Literature Values for Water Concentrations

Some of the variation in reported bioaccumulation factors can be attributed to the use of average literature values for stable isotope concentration in water. This is especially true in freshwater environments where element concentrations are more variable than in marine environments. Thompson et al. (1972) derived some average bioaccumulation factors by determining the average stable element concentration in organisms from a number of freshwater habitats and then dividing this average concentration by the average stable element concentration of a number of freshwater habitats. Often the water values were taken from studies different from those giving the concentrations in the

organisms. Thus, some error may have been introduced into the average bioaccumulation factor values.

In selecting or calculating bioaccumulation factors for this report only those studies that collected water and organism samples from the same locality are included.

2.5.3 Filtration of Water Samples

Some experimenters have reported bioaccumulation factors based on concentrations of isotopes from filtered as well as unfiltered water samples. Since a significant fraction of some elements in water may be in the suspended phase - in some cases greater than 90% - bioaccumulation factors based on concentrations in filtered samples may be much larger than bioaccumulation factors based on unfiltered concentrations.

To understand the significance of filtered or unfiltered bioaccumulation factors for prediction at proposed nuclear power plants, it is necessary to discuss the nature of the effluent and how radionuclide concentrations in the water at the proposed reactor sites are estimated. Radionuclides in the reactor effluent are in the soluble phase. Concentrations in the body of water of concern are usually estimated from predicted concentrations in the effluent and from water turnover rates. Actual concentrations in the water may be somewhat lower because the bottom sediment may serve as a sink for radionuclides. Since radionuclides may become distributed between soluble and suspended phases, the "predicted concentration" approximates the unfiltered concentration of the radionuclide in the water. Assuming that the specific activity

of the radionuclide in the soluble and the suspended phases are equal, an unfiltered bioaccumulation factor will most closely predict the radionuclide concentration in the organism. Generally, filtered bioaccumulation factors will overpredict the radionuclide concentration in organisms.

2.5.4 Determination of Organism Concentration Based on Dry Weight

Bioaccumulation factors expressed in terms of dry weight of the organism are higher than those calculated using wet weight concentrations. Thus, care is taken to ensure that bioaccumulation factors listed in this report are on a fresh weight basis. When dry weight bioaccumulation factors were given in a publication, they were converted to fresh weight bioaccumulation factors if conversion factors between fresh weight and dry weight were available in the same publication. In some cases where conversion factors were not available in the same publication, conversion factors from other sources were used.

2.5.5 Averaging Isotope Concentrations of Different Tissues

In reporting or evaluating bioaccumulation factors care must be taken to specify what tissue or tissues of the organism were analyzed. Certain elements have an affinity for different tissues. For example, strontium concentrates to a higher degree in bony tissues, and iodine concentrates to a higher degree in the thyroid than in other tissues of the organism. Thus, any bioaccumulation factor is incomplete without reference to the specific tissue analyzed. The whole-body bioaccumulation factors, which are often reported, may be different from those of any particular tissue.

Another potential source of uncertainty associated with whole-body bioaccumulation factors is that the organism's gut may contain sediment when the whole-body bioaccumulation factor is determined. The concentration of metals, such as cesium, manganese, cobalt, and strontium in sediment may be significantly higher than the concentrations found in the tissues of the organism. Sediment contamination is most likely a significant factor for benthic invertebrates.

2.5.6 Determinations Made Under Nonsteady-State Conditions

Measurements of organism concentrations in situations where a steady-state condition is not present can result in low or high bioaccumulation factors depending on the circumstance. If an organism is transferred from an uncontaminated environment to an environment with a fixed level of a radioisotope, the time necessary for the establishment of a nearly steady-state concentration is a function of the biological half-time of that element in the organism and the radioactive half-life. Obviously, organism concentrations analyzed prior to the establishment of a steady state will result in low bioaccumulation factors. If the environmental concentration is subjected to sudden variations, the subsequent changes in the organism concentrations may lag behind that of the environment. Low bioaccumulation factors would be expected if determinations are made shortly after a sudden rise in environmental concentration, or after an organism moves from an area of low isotope concentration to an area of high concentration. The opposite situations would result in high bioaccumulation factors.

2.6 Use of Bioaccumulation Factors in Radiation Dose Estimation

The two principal uses of bioaccumulation factors for aquatic biota are: (1) calculation of radiation dose to biota from radio-nuclides deposited in tissues, and (2) calculation of radiation dose to man from consumption of contaminated biota. In both situations the radiation exposure is within the body and not exterior to it.

2.6.1 Dose to Biota

The internal absorbed dose (rads/year) to organism i from an internally deposited radionuclide is estimated from the energy deposited per gram of tissue and can be expressed mathematically as:

$$D_i = [R]_w BF(R)_i E K \text{ (rads/year)} \quad (2.6.1)$$

where

D_i = internal absorbed dose (rads/year) to organism i from a radionuclide uniformly deposited in its tissues,

$[R]_w$ = radionuclide concentration in water ($\mu\text{Ci/g}$),

$BF(R)_i$ = bioaccumulation factor for radionuclide R in organism i ,

E = effective absorbed energy (MeV) of the radionuclide for the physical dimensions appropriate for the organism i , and

K = conversion factor to convert $\mu\text{Ci/g}$ to rads/year.

There is no radioactive decay correction in the above equation because the concentration of the radionuclide in water is assumed to remain constant and the concentration in the biota is in a steady-state with

the water. In actual practice, the average annual concentration of the radionuclide in the receiving water at the point of interest is used in dose calculations.

The variable K in the above equation is defined as a conversion factor, but it can represent any type of internal dosimetry model. The variables $[R]_w$, $BF(R)_i$, and E are always required, no matter how specific or general the conversion factor K.

2.6.2 Dose to Man

A bioaccumulation factor is used to estimate the concentration of a radionuclide in the aquatic food (organism or tissue i) consumed by man. The intake rate of a radionuclide ($\mu\text{Ci/day}$) by man is calculated from

$$I_i = [R]_w BF(R)_i G_i (\mu\text{Ci/day}) \quad (2.6.2)$$

where

I_i = man's daily dietary intake rate of a radionuclide ($\mu\text{Ci/day}$) contained in aquatic food organism or tissue i,

$[R]_w$ = radionuclide concentration in water ($\mu\text{Ci/g}$),

$BF(R)_i$ = bioaccumulation factor for radionuclide R in organism or tissue i, and

G_i = man's daily intake rate of organism or tissue i (g/day).

In most exposure situations each organism is contaminated by more than one radionuclide, and man's dietary intake usually includes more

than one type of aquatic biota. Therefore, a summation must be performed over all radionuclides and biota to estimate man's total intake of radioactivity.

The internal dose to a reference organ or to the total body of man is calculated using the estimated radionuclide intake rate ($\mu\text{Ci}/\text{day}$) for each radionuclide in addition to the following dosimetry factors: fraction of the intake deposited in the organ of reference, the effective elimination constant for each radionuclide in the organ of reference, the effective absorbed energy of each radionuclide in the organ of reference, and the mass of the reference organ.

References

Comar, C. L., R. H. Wasserman, and M. M. Nold, Strontium-calcium discrimination factors in the rat, Proc. Soc. Exptl. Biol. Med. 92, 859-863 (1956).

Eyman, L. D. and J. T. Kitchings, Availability of ^{137}Cs to fishes from ingested, IN Proceedings of the XIX Congress of the International Association of Limnology (SIL), Winnipeg, Canada, August 22-29 (1974). (in press)

Fleishman, D. J., Accumulation of artificial radionuclides in freshwater fish, IN Radioecology, V. M. Klechkovskii, G. G. Polikarpov, and R. M. Aleksakhin (eds.), pp. 347-369, John Wiley and Sons, New York, 1973.

Friend, A. G., The aqueous behavior of strontium-85, cesium-137, zinc-65, and cobalt-60 as determined by laboratory type studies, IN Transport of Radionuclides in Freshwater Systems, B. H. Kornegay, et al. (eds.), pp. 43-60, TID-7664 (1963).

Gallegos, A. F., F. W. Wicker, and T. E. Hakonson, Accumulation of radiocesium in rainbow trout via a non-foodchain pathway, IN Proceedings of Fifth Annual Health Physics Society Midyear Topical Symposium: Health Physics Aspects of Nuclear Siting, Vol. 2, pp. 477-497, AEC CONF-701106 (1970).

Gibbs, R. J., Mechanisms of trace metal transport in rivers, Science 180, 71-73 (1973).

Lomenick, T. F., and D. A. Gardiner, The occurrence and retention of radionuclides in the sediments of White Oak Lake, Health Phys. 11, 567-577 (1965).

Maletskos, C. J., Intercalibration and intercomparison study of radionuclide concentrations in Lake Michigan samples, New England Deaconess Hospital, Progress Report October-March, 1972, COO-3371-1, USAEC Contract AT(11-1)-3371.

Peterson, H. T., Jr., Stable element and radionuclide reconcentration in aquatic ecosystems. I. Stable element effects in the equilibrium situation, Fifth Annual Health Physics Society Midyear Topical Symposium, Idaho Falls, Idaho, November 3-6, 1970.

Thompson, S. E., C. A. Burton, D. J. Quinn, and Y. C. Ng, Concentration factors of chemical elements in edible aquatic organisms, Lawrence Livermore Laboratory, Univ. California, Livermore, 1972.

Timofeev - Resovskii, N. V., E. A. Timofeeva - Resovskaya, G. A. Milyutina, and A. B. Getsova, Coefficients of accumulation of radioisotopes of sixteen different elements by freshwater organisms and the effect of EDTA on some of them, Dokl. Nauk SSSR 132, 369-372 (1960).

Vanderploeg, H. A., R. S. Booth, and F. H. Clark, A specific activity and concentration model applied to cesium movement in an oligotrophic lake, IN Proceedings of Symposium on Mineral Cycling in Southeastern Ecosystems, Augusta, Georgia, May 1-3, 1974 (in press).

3. BIOACCUMULATION FACTORS FOR CESIUM, STRONTIUM, TRITIUM, IODINE, MANGANESE AND COBALT

3.1 CESIUM

3.1.1 Cesium Metabolism

Because of their chemical similarities, cesium and potassium are metabolized somewhat similarly in organisms. In freshwater ecosystems, stable cesium, ^{133}Cs , is an element having concentrations ranging between 0.011 and 1.2 ppb for lakes and rivers (Copeland and Ayers, 1970; Kolehmainen, 1972; Kolehmainen and Nelson, 1969; Merlini et al., 1967; Nelson, 1967). Stable potassium is more abundant in freshwater than cesium and concentrations usually range between 0.2 ppm and 10 ppm.

The bioaccumulation factor for cesium in plants and animals is independent of cesium concentrations found in natural bodies of water (Gertz, 1973; King, 1964; Kolehmainen et al., 1968; Kolehmainen, 1972; Williams and Swanson, 1958). On the other hand, the bioaccumulation factor for potassium in animals is inversely proportional to the potassium concentration in water because the potassium concentration in animals is homeostatically maintained (Kolehmainen, 1972; Kolehmainen and Nelson, 1969; Peterson, 1970). Because of the chemical similarities and relative abundances of cesium and potassium, the bioaccumulation factor for cesium in animals is related to the potassium concentration in water according to Equation (2.1.8). Unlike this pattern for animals, the potassium concentration in water has only a small effect on the cesium bioaccumulation factor in some algae (Williams, 1970) and no effect in other algae (Gertz, 1973).

Cesium elimination from a plant or an animal may be approximated by a single elimination-rate coefficient, k . Table 3.1.1 gives approximate

Table 3.1.1 Biological half-times and elimination coefficients for cesium and $[k/(k+\lambda)]$ values for ^{136}Cs

Taxon/functional category	Biological half-time (days)	k (days ⁻¹)	$\frac{k}{k+\lambda}$ for ^{136}Cs	Reference
Unicellular algae	1	0.69	0.93	1.50
Multicellular algae	2	0.35	0.88	1.47
Aquatic vascular plants:				
floating	20	0.035	0.40	1.37
rooted	60 ^a	0.012	0.18	1.37
Zooplankton	5	0.14	0.74	1.46
Benthic insect larvae	7	0.10	0.67	1.25
Clams	40	0.017	0.24	1.20
Fishes	100	0.0069	0.12	1.13, 1.23, 1.24, 1.46

^aSince rooted plants may accumulate part of their burden of cesium from interstitial water of bottom sediments, time to steady-state may be much greater than if uptake were entirely from water above sediments.

biological half-times and elimination rate coefficients values for cesium in different taxa or functional groups. These coefficients must be considered as approximate since they may be influenced by such factors as temperature, size, growth rates, feeding rates, and species differences (Vanderploeg et al., 1974). Most bioaccumulation factors, except those for algae, were derived entirely from stable cesium or from ^{137}Cs (30-year radioactive half-life) concentrations. Cesium-134, having a 2.2-year half-life, will not have an appreciably different bioaccumulation factor from that of stable cesium or ^{137}Cs . Cesium-136, which has a 13-day radioactive half-life, will have an appreciably lower bioaccumulation factor than those of the longer-lived isotopes in some biota; therefore, Table 3.1.1 includes a set of approximate $[\text{k}/(\text{k}+\lambda)]$ values to convert bioaccumulation factors for stable cesium, or ^{137}Cs , to bioaccumulation factors for ^{136}Cs .

The primary mode of accumulation of cesium and potassium in fishes is generally thought to be via absorption from food (Kolehmainen, 1972); absorption from food may also be a primary mode of uptake for many aquatic invertebrates. However absorption of cesium from ingested sediments may be significant in some systems (Gallegos et al., 1970). In fishes (Kolehmainen, 1972) and other animals (Pendleton et al., 1965), the absorption efficiency of potassium and cesium from food varies but is often high--nearly 100 percent for some foods. The excretion coefficient, k^* , of potassium is on the order of 3 to 4.5 times greater than the excretion coefficient, k , of cesium in fishes (Kolehmainen, 1972) and in terrestrial animals (Fujita et al., 1966, Pendleton et al., 1965). Pendleton pointed out that this will result in a cesium to potassium ratio in the

predator that is higher than that of its prey by a factor of about three, assuming that absorption efficiency of cesium is the same as that for potassium. Assuming that the potassium concentration in the predator is the same as that in its prey, the bioaccumulation factor for cesium should increase by about a factor of three with each trophic level. Note that the variable k^*/k appears in our general expression for q_i (see Appendix).

3.1.2 Environmental Cesium

In solution, cesium and potassium, like other alkali metals, exist primarily as free ions and do not form inorganic complexes. Because cesium is strongly adsorbed by suspended particulate materials, especially clays, these materials may remove it from the soluble phase to varying degrees. Potassium, too, is sorbed, but its distribution coefficient, K_d , on suspended sediment (that is, the concentration of element sorbed to sediment per concentration in water) is many times smaller than that of cesium (Reynolds and Gloyne, 1964).

Table 3.1.2 shows that the percentage of cesium in the particulate phase increases with clay particle concentration. For freshwater ecosystems the fraction of cesium in water in the suspended phase ranges between 19 and 92 percent, and this fraction appears to be roughly correlated with concentration of suspended solids (Table 3.1.3). Because potassium is less strongly sorbed than cesium, the percentages of potassium appearing in the suspended phase are very much smaller than those for cesium in these bodies of water. Since algae accumulate cesium, potassium, and other elements from the soluble phase, the availabilities to the food chain of cesium and of cesium relative to potassium become a function of suspended solids concentration and the distribution coefficients of cesium and potassium on

Table 3.1.2 Sorption of ^{134}Cs to various concentrations of clay suspended in distilled water
(Garder and Skullberg 1964)

Concentration of clay (ppm)	Percent of cesium sorbed	K_d ^a
16	20	15,600
32	31	13,600
64	43	11,800
128	58	10,800
256	65	7,300

^aConcentration of element (cesium) sorbed to clay per concentration in water.

Table 3.1.3 Percentage of cesium in water in the particulate phase in some freshwater ecosystems

Location	Mean percent in particulate phase	Suspended solids (ppm) or nature of water body	Separation method	Reference
White Oak Lake, Tennessee	69	Turbid reservoir	Ultracentrifugation: Particles $> 0.7 \mu$	1.44
Clinch River, Tennessee ^a	82-92	25-185	Ultracentrifugation: Particles $> 0.7 \mu$	1.44
Tennessee River, Tennessee ^a	19-30	9-22	Ultracentrifugation: Particles $> 0.7 \mu$	1.44
Lake Maggiore, Italy	28	Mesotrophic Lake	0.7- μ filter	1.35
Experimental pond ^b	58	21	0.45- μ filter	1.10

^aRange of means of three river stations.

^bBased on five water values sampled between 24 and 80 days after introduction of radionuclides.

the solids. Thus, the q_i values and bioaccumulation factors of cesium become functions of suspended-solids concentrations and the distribution coefficients. Since the proportions of cesium and of cesium relative to potassium in the soluble phase are expected to vary greatly among environments, the bioaccumulation factors for cesium in algae and the q_i values and bioaccumulation factors of animals may vary greatly among bodies of water having the same potassium concentrations. We shall see in the following sections that much of the scatter in bioaccumulation factors for cesium is apparently related to the proportion of cesium in the particulate phase.

The food web accumulates cesium not only from the soluble phase of water but also from suspended and bottom sediments. Filter feeders may accumulate cesium adsorbed to particulate matter. Benthic invertebrates obtain cesium adsorbed to ingested bottom sediments. Fishes may accumulate cesium from bottom sediment by ingestion of these invertebrates and also by ingestion of sediment with prey (Eyman and Kitchings, 1974; Gallegos et al., 1970). Gallegos et al. (1970) reported that concentration of ^{137}Cs in trout in a montane lake was higher than could be explained by the trouts' accumulation from their prey. Their experiments indicated that this difference could be explained by the trout ingesting small amounts of sediment. In fishes, absorption efficiency of cesium from ingested clay varies with clay type: 8% for illite, 65% for kaolinite, and 85% for montmorillonite (Eyman and Kitchings, 1974). Since cesium is available from ingested sediments, differences in sediment type among water bodies may be another contributor to the variance in bioaccumulation factors among bodies of water.

3.1.3 Review of Cesium Bioaccumulation Factors

This section presents bioaccumulation factors and bioaccumulation factor relations for the following categories of aquatic biota:

1. Fishes
2. Algae
3. Macrophytes (vascular aquatics)
4. Emergent vascular plants
5. Invertebrates--molluscs, insects, and crustaceans
6. Amphibians
7. Waterfowl

Except for algae, all bioaccumulation factors for cesium were derived from distributions of stable cesium or from steady-state distributions of ^{137}Cs in natural or semi-natural bodies of water. Since ecological relationships in aquaria do not completely represent the natural system, bioaccumulation factors derived from laboratory experiments have limited predictive value. Moreover, in Table 3.1.1, it is seen that the biological half-times of ^{137}Cs in some biota range between approximately 40 and 100 days. Laboratory experiments have not been run for a sufficient time period in many cases for these longer-term components to attain steady state.

3.1.3.1 Cesium Bioaccumulation Factors for Fishes

Bioaccumulation factors have usually been reported for flesh or whole bodies of fishes. Since concentrations of cesium and potassium do not vary greatly among organs (Nelson, 1967), flesh or whole-body bioaccumulation factors apply to other organs as well.

Potassium concentration in freshwater fishes is about 3 mg/g fresh weight and varies little among species (Kolehmainen, 1972; Peterson, 1970). The range of potassium concentrations in fishes from both freshwater and marine environments is from 2.1 to 4.5 mg/l, (Fleishman, 1973;

Kolehmainen, 1972; Kolehmainen and Nelson, 1969). The range of potassium concentrations in freshwater and marine fishes is small considering that potassium concentration in the water range between 0.2 and 380 ppm. Clearly, the potassium concentrations are homeostatically maintained. Thus, the bioaccumulation factor for potassium is given by Equation (2.1.7). Since the bioaccumulation factor for potassium is given by Equation (2.1.7), the bioaccumulation factor for cesium is given by Equation (2.1.8). Owing to the variable availabilities of cesium relative to potassium in different environments, the value q_i varies from one environment to another.

Bioaccumulation factors calculated from concentrations of fallout-derived ^{137}Cs in water and fishes collected in Finnish lakes (Kolehmainen et al., 1967; Kolehmainen et al., 1968) are given in Table 3.1.4. Within lakes, the ratio of highest bioaccumulation factor to lowest bioaccumulation factor ranges between 2.4 and 7.1. As expected, the highest bioaccumulation factors are seen for high trophic-level fishes, namely, perch and pike, which are piscivores. The highest bioaccumulation factors are seen for the oligotrophic lake having a potassium concentration in water of 1 ppm. Note that the bioaccumulation factors in the oligotrophic lake are very much higher than the bioaccumulation factors in the other two lakes having the same potassium concentration but higher concentrations of organic matter. The difference appears to be related to the lower concentration of organic matter in the oligotrophic lake. Most probably, however, this difference is related to the lower suspended solids concentration (both organic and inorganic solids) expected in the oligotrophic lake. The bioaccumulation factors for the fishes from the lake having a water concentration of 3.5 ppm potassium are more than 25 times lower than that of the oligotrophic lake having a

Table 3.1.4 Mean ^{137}Cs bioaccumulation factors for whole fishes from Finnish Lakes (Kolehmainen et al. 1967, 1968)^a

Potassium concentration (ppm)	Lake type ^b	KMnO ₄ consumption (ppm) ^c	Bioaccumulation Factors						
			Perch (<i>Perca fluviatilis</i>) ^d	Pike (<i>Esox lucius</i>)	Burbot (<i>Lota vulgaris</i>)	Roach (<i>Leuciscus rutilus</i>)	Bream (<i>Abramis brama</i>)	Whitefish (<i>Coregonus spp.</i>)	Highest BF Lowest BF
0.8	0	32	11,000	5,600	4,400	3,100	---	2,500	4.4
0.9	D-0	65	9,900	5,500	3,300	1,800	---	---	5.5
1.0	0	6	---	14,000	---	5,200	---	3,700	3.8
1.0	D-0	120	---	2,900	2,800	1,200	800	---	3.6
1.0	E	43	1,800	2,100	2,200	900	---	---	2.4
1.8	D-E	45	5,000	2,900	2,500	1,200	700	---	7.1
3.5	E	75	300	430	480	200	75	---	4.0

^aBased on whole fish concentrations derived from 1964 and 1965 data; ^{137}Cs concentration in water (unfiltered) from 1964 data.

^b0 = oligotrophic; D-0 = dystrophic-oligotrophic; E = eutrophic.

^cA measure of organic content of water.

^dLarge perch.

potassium concentration of 1 ppm. If the value of q_j were constant, the bioaccumulation factors in the 3.5-ppm lake would be 3.5 times lower than the bioaccumulation factors in the 1-ppm lake. Thus, the potassium concentration in the water alone is not very useful for predicting the bioaccumulation factors for cesium in fishes.

The conclusion that the potassium concentration in the water alone is not very useful for predicting cesium bioaccumulation factors is corroborated by the data in Table 3.1.5. The table gives bioaccumulation factors for cesium as related to potassium and suspended solids concentrations. The bioaccumulation factors for cesium in fishes were taken from Table 3.1.6. The highest bioaccumulation factors are seen for bodies of water of low turbidity. The bioaccumulation factors of bluegills and catfish found in the clear water body with a potassium concentration of 1.4 ppm are more than six and seven times the respective bioaccumulation factors of bluegills and catfish from the very turbid water body with a potassium concentration of 1.3 ppm. Also, the highest bioaccumulation factor for largemouth bass is in a lake of high potassium concentration and low turbidity.

Most investigators have reported bioaccumulation factors that are calculated from concentration of cesium isotopes in unfiltered water samples; however, some have reported bioaccumulation factors based on concentration in the water after particulate material has been filtered off. We shall denote a "filtered" bioaccumulation factor by BF^* . Assuming all particulate material is filtered out, a BF^* represents an upper-limit bioaccumulation factor for that and other environments of similar potassium concentration and sediment type since cesium on suspended sediments is to

Table 3.1.5 Bioaccumulation factors for cesium as related to potassium and suspended solids concentrations^a

Potassium concentration in water (ppm)	Suspended solids concentration (ppm)	Bioaccumulation factors of Cesium						Reference
		Bluegill (<i>Lepomis macrochirus</i>)		Largemouth bass (<i>Micropterus salmoides</i>)		Catfish (<i>Ictalurus</i>)		
		¹³⁷ Cs	Stable Cs	¹³⁷ Cs	Stable Cs	¹³⁷ Cs	Stable Cs	
1.3	Very high (25-185)		140			160		1.36 1.44
1.4	Low (1.3)	900		1200		1200		1.19 1.34
1.8	High	270	310	350	620			1.27
9.8	Low			1600	570			1.7

^aBased on concentrations of cesium in unfiltered water samples.

Table 3.1.6 Bioaccumulation factors for cesium in fishes

Species	Tissue	K concentration (ppm)	BF, fallout ¹³⁷ Cs	BF*, fallout ¹³⁷ Cs ^a	BF, chronic release of ¹³⁷ Cs	BF*, chronic release of ¹³⁷ Cs	BF, stable cesium	BF*, stable cesium	Water treatment method	Feeding habits	Location	Suspended solids concentration (ppm)	Reference
<u>Salmo trutta</u>	F ^b	0.4	3,900								Lake Trawsfynydd, England	Low ^c	1.42
<u>Perca fluviatilis</u>	F	0.4	5,800								Lake Trawsfynydd, England	Low	1.42
<u>Salmo trutta</u>	F	0.3	3,000								English river	Low	1.42
<u>Salmo trutta</u>	F	0.4	1,400								English river	Low	1.42
<u>Salmo trutta</u>	F	1.9	920								English river	Low	1.42
<u>Salmo trutta</u>	F	3.8	940								English river	Low	1.42
<u>Salmo trutta</u>	F	4.1	260								English river	Low	1.42
<u>Salmo trutta</u>	F	4.1	380								English river	Low	1.42
<u>Pomoxis annularis</u>	F	1.3			500	6,400	0.7 μ ^e		P		Clinch River, Tennessee	25-185 ^f	1.36
<u>Aplodinotus grunniens</u>	F	1.3			350	4,400	0.7 μ				Clinch River, Tennessee	25-185	1.36
<u>Roccus chrysops</u>	F	1.3			640	8,000	0.7 μ		P		Clinch River, Tennessee	25-185	1.36
<u>Ictalurus punctatus</u>	F	1.3			160	2,000	0.7 μ				Clinch River, Tennessee	25-185	1.36
<u>Lepomis macrochirus</u>	F	1.3			140	1,700	0.7 μ				Clinch River, Tennessee	25-185	1.36
<u>Dorosoma cepedianum</u>	W ^g	1.8		310	810	400		0.7 μ			White Oak Lake, Tennessee	High ^h	1.27
<u>Notemigonus crysoleucas</u>	W	1.8		420	1,100	510		0.7 μ			White Oak Lake, Tennessee	High	1.27
<u>Carassius auratus</u>	W	1.8		200	530	690		0.7 μ			White Oak Lake, Tennessee	High	1.27
<u>Lepomis microlophus</u>	W	1.8		180	460	320		0.7 μ			White Oak Lake, Tennessee	High	1.27
<u>Lepomis macrochirus</u>	W	1.8		270	710	310		0.7 μ			White Oak Lake, Tennessee	High	1.27
<u>Chaenobryttus gulosus</u>	W	1.8		240	630	430		0.7 μ			White Oak Lake, Tennessee	High	1.27

Table 3.1.6 (continued)

Species	Tissue	K concentration (ppm)	BF, fallout ¹³⁷ Cs	BF*, fallout ¹³⁷ Cs	BF, chronic release of ¹³⁷ Cs	BF*, chronic release of ¹³⁷ Cs	BF, stable cesium	BF*, stable cesium	Water treatment method	Feeding habits	Location	Suspended solids concentration (ppm)	Reference
<u><i>Micropterus salmoides</i></u>	W	1.8			350	910	620		0.7 μ	P	White Oak Lake, Tennessee	High	1.27
<u><i>Salmo gairdneri</i></u>	F	1.7		12,000					A-ER [†]		East Twin Lake, Colorado	Low	1.12
<u><i>Micropterus salmoides</i></u>	W	9.8		1,625		570			P		Wintergreen Lake, Michigan	Low	1.7
<u><i>Perca flavescens</i></u>	W	9.8		1,200		1,600			P		Wintergreen Lake, Michigan	Low	1.7
<u><i>Hybrid sunfish</i></u>	W	9.8		500		250					Wintergreen Lake, Michigan	Low	1.7
<u><i>Erimyzon suetta kennelerii</i></u>	W	9.8		790		300					Wintergreen Lake, Michigan	Low	1.7
<u><i>Perca fluviatilis</i></u>	W	2.1	2,900			1,100			P		Lake Maggiore, Italy		1.2,1.3
<u><i>Perca fluviatilis</i></u>	W	2.6	1,400			140			P		Lake Varese, Italy		1.2,1.3
<u><i>Perca fluviatilis</i></u>	W	2.1	2,400			660			P		Lake Comabbio, Italy		1.2,1.3
<u><i>Perca fluviatilis</i></u>	W	1.3	5,100			930			P		Lake Monate, Italy		1.2,1.3
<u><i>Scardinius erythrophthalmus</i></u>	W	2.1	1,300			560					Lake Maggiore, Italy		1.2,1.3
<u><i>Scardinius erythrophthalmus</i></u>	W	2.6	890			140					Lake Varese, Italy		1.2,1.3
<u><i>Scardinius erythrophthalmus</i></u>	W	2.1	1,000			370					Lake Comabbio, Italy		1.2,1.3
<u><i>Scardinius erythrophthalmus</i></u>	W	1.3	1,400			350					Lake Monate, Italy		1.2,1.3

Table 3.1.6 (continued)

Species	Tissue	K concent- ration (ppm)	BF, fallout ^{137}Cs	BF*, fallout ^{137}Cs	BF, chronic release of ^{137}Cs	BF*, chronic release of ^{137}Cs	BF, stable cesium	BF*, stable cesium	Water treatment method	Feeding habits	Location	Suspended solids concentration (ppm)	Reference
<u>Lepomis gibbosis</u>	W	2.1	790				440				Lake Maggiore, Italy		1,2,1,3
<u>Lepomis gibbosis</u>	W	2.6	420				110				Lake Varese, Italy		1,2,1,3
<u>Lepomis gibbosis</u>	W	2.1	690				420				Lake Comabbio, Italy		1,2,1,3
<u>Lepomis gibbosis</u>	W	1.3	2,800				790				Lake Monate, Italy		1,2,1,3
<u>Lepomis macrochirus</u>	F	1.4 ^j			900					P	Par Pond, South Carolina	1.3	1.19
<u>Micropterus</u> <u>salmoides</u>	F	1.4			1,200					P	Par Pond, South Carolina	1.3	1.19
<u>Ictalurus natalis</u>	F	1.4			1,200						Par Pond, South Carolina	1.3	1.19
<u>Oncorhynchus</u> <u>kisutch</u>	F	1.6					2,700	0.45 μ	P		Lake Michigan	Low	1.5
<u>Perca flavescens</u>	F	1.6					3,300	0.45 μ	P		Lake Michigan	Low	1.5
<u>Salvelinus</u> <u>namaycush</u> <u>namaycush</u>	F	1.6					2,400	0.45 μ	P		Lake Michigan	Low	1.5
<u>Salmo trutta</u>	F	1.6					3,400	0.45 μ	P		Lake Michigan	Low	1.5
<u>Coregonus</u> <u>clypeiformis</u>	F	1.6					1,500	0.45 μ			Lake Michigan	Low	1.5
<u>Pomolobus</u> <u>pseudoharengus</u>	W	1.6					1,400	0.45 μ			Lake Michigan	Low	1.5
<u>Osmerus mordax</u>	W	1.6					1,900	0.45 μ			Lake Michigan	Low	1.5
<u>Notropis hudsonius</u>	W	1.6					2,100	0.45 μ			Lake Michigan	Low	1.5

Table 3.1.6 (continued)

Species	Tissue	K concentration (ppm)	BF, fallout ¹³⁷ Cs	BF*, fallout ¹³⁷ Cs	BF, chronic release of ¹³⁷ Cs	BF*, chronic release of ¹³⁷ Cs	BF, stable cesium	BF*, stable cesium	Water treatment method	Feeding habits	Location	Suspended solids concentration (ppm)	Reference
<u>Cyprinus carpio</u>	W	2.6				500					Carp pond		1.53

^aA BF* is a bioaccumulation factor calculated from isotope concentration in water after filtration.

^bF = flesh.

^cLow \leq 30 ppm suspended solids.

^dp = piscivorous.

^ePore size of filter.

^fStruxness et al. (1967).

^gW = whole fish.

^hHigh $>$ 30 ppm suspended solids.

ⁱA-ER = anion-exchange resin.

^jPersonal communication.

some degree available to the food web. Whenever possible, we calculated both unfiltered and filtered bioaccumulation factors.

The unfiltered and filtered bioaccumulation factors in Table 3.1.6 are categorized according to source of isotope: fallout-derived ^{137}Cs , chronic release of ^{137}Cs , and stable cesium. In fishes from Italian lakes (Table 3.1.6), Bortoli et al. (1967) reported that bioaccumulation factors calculated from distributions of fallout-derived ^{137}Cs were on the average four times higher than bioaccumulation factors calculated from stable cesium distributions. Clearly, fallout-derived ^{137}Cs is somehow more available to the food web than stable cesium. A greater proportion of fallout ^{137}Cs , which enters watersheds from the atmosphere as a soluble radionuclide, may be in the soluble phase or on exchangeable sites (adsorbed) of clay particles. The recently introduced ^{137}Cs may not be in isotopic equilibrium with the stable cesium on the clay particles, especially stable cesium at non-exchangeable sites. According to Eyman's study of Wintergreen Lake, Michigan (1972), bioaccumulation factors for ^{137}Cs were generally higher than bioaccumulation factors for stable cesium. In contrast, however, Kolehmainen's study (1972) of White Oak Lake (Table 3.1.6), showed that bioaccumulation factors for ^{137}Cs originating from a chronic release were about the same as bioaccumulation factors derived from stable cesium data. This may result from the time history of ^{137}Cs input into White Oak Lake. During the years preceding Kolehmainen's study, ^{137}Cs input was higher than during the study. We also note that the stable cesium concentration in water was measured on water that had received gross filtration to remove larger particles. All these patterns would suggest that the bioaccumulation factor for ^{137}Cs near a power station is somewhat

higher--roughly by a factor of two--than the bioaccumulation factor for stable cesium. The magnitude of the difference may depend on suspended solids concentrations.

We have shown that the bioaccumulation factor for cesium is highly variable from one environment to another and that much of this variation derives from differing proportions of radiocesium relative to potassium in the soluble phase and possibly sediment type. Unfortunately, the data necessary for accurate quantification of bioaccumulation factor relations are not available. Approximate relations, however, can be stated.

To obtain these relations we plotted the bioaccumulation factor for cesium in fishes against the potassium concentration in water on logarithmic graph paper, as shown in Figure 3.1.1. Unfiltered bioaccumulation factors were used so as to better predict radionuclide concentration in the organism. For turbid environments a filtered bioaccumulation factor may be more than 10 times greater than an unfiltered bioaccumulation factor since more than 90 percent of the cesium in water may be in the particulate phase. Fishes were grouped into two categories: piscivorous fishes and non-piscivorous fishes. By inspection we chose the relation

$$BF(Cs)_i = 1.5 \times 10^4 / [K]_w \quad (3.1.1)$$

as an upper-bound relation for piscivorous fishes and

$$BF(Cs)_i = 5 \times 10^3 / [K]_w \quad (3.1.2)$$

as an upper-bound relation for non-piscivorous fishes, where $[K]_w$ has units of ppm. Equations (3.1.1) and (3.1.2) are applicable to environments of low turbidity since they will greatly overpredict bioaccumulation




Figure 3.1.1 Bioaccumulation factors for ^{137}Cs and stable cesium in freshwater fishes as a function of potassium concentration in water. (All data in Tables 3.1.4 and 3.1.6 were plotted except (1) stable cesium bioaccumulation factors from studies in which ^{137}Cs bioaccumulation factors were reported and (2) BF*s.)

factors for fishes from turbid environments. Dividing Equations (3.1.1) and (3.1.2) by five gives rough approximations for the respective relations for fishes in turbid waters; that is,

$$BF(Cs)_i = 3 \times 10^3 / [K]_w \quad (3.1.3)$$

for piscivorous fishes from turbid waters and

$$BF(Cs)_i = 1 \times 10^3 / [K]_w \quad (3.1.4)$$

for non-piscivorous fishes from turbid waters. We may consider waters having greater than 50 ppm suspended solids as being turbid based on the data in Table 3.1.3. Alternatively, bioaccumulation factors for particular species may be estimated from the raw data in Tables 3.1.4 and 3.1.6. The bioaccumulation factor selected should preferably come from an environment having the same suspended solids and potassium concentrations as the environment to which it is applied. Even if this is done, a degree of uncertainty remains because of the unspecified effect of sediment type on bioaccumulation factors as well as error associated with measurement and experimental design.

3.1.3.2 Cesium Bioaccumulation Factors for Algae

As can be seen from the bioaccumulation factors for Cladophora glomerata and Pithophora oedegonia (Table 3.1.7), the potassium concentration in water has little effect on the bioaccumulation factor for cesium in multicellular algae. Recently, Gertz (1973) has shown that the potassium concentration has no effect on the bioaccumulation factor for cesium in Chlamydomonas reinhardtii, a unicellular green alga. He found that the

Table 3.1.7 Bioaccumulation factors for cesium in freshwater algae^a

Taxon/functional category	K concentration (ppm)	Bioaccumulation factors				Location	Source of ¹³⁷ Cs	Reference			
		¹³⁷ Cs		Stable Cs							
		BF	BF* ^b	BF	BF*						
Blue-green algae:											
<u>Plecotonema boryanum</u>	3.6	100 ^c						1.22			
Unidentified	1.4	1,200				Par Pond, South Carolina	Chronic-release	1.19			
Unidentified	—	3,400				Connecticut River	Chronic-release	1.1			
Diatoms:											
<u>Navicula seminulum</u>	14	130 ^c				Lab study		1.22			
Mixed species ^d	1.6			2900 ^e		Lake Michigan		1.4			
Euglenophyta:											
<u>Euglena intermedia</u>	8	700				Lab study		1.51			
Green algae, multicellular:											
<u>Chara</u> sp.	—	20				Lab study		1.45			
<u>Chara</u> <i>aspera</i>	—	18				Lab study		1.45			
<u>Chara</u> <i>braunii</i>	1	60				Experimental channel receiving river water	Chronic-release	1.16			
<u>Chara</u> <i>fragilis</i>	—	1,000				Lake Bol shoe Missavo, Russia		1.33			
<u>Chara</u> <i>fragilis</i>	—	36				Lab study		1.45			
<u>Chara</u> <i>tomentosa</i>	—	80				Lake Bol shoe Missavo, Russia		1.33			
<u>Cladophora</u> <i>fracta</i>	—	120				Lab study		1.45			
<u>Cladophora</u> <i>glomerata</i>	0.1	170				Lab study		1.48			
<u>Cladophora</u> <i>glomerata</i>	30	59				Lab study		1.48			
<u>Cladophora</u> <i>glomerata</i>	—	160				Lab study		1.45			
<u>Chlorella</u> <i>pyrenoidosa</i>	8	150				Lab study		1.51			
<u>Gonium</u> <i>pectorale</i>	10	140				Lab study		1.51			
<u>Mougeotia</u> sp.	9.8	330	500			Wintergreen Lake, Michigan	Fallout	1.7			

Table 3.1.7 (continued)

Taxon/functional category	K concentration (ppm)	Bioaccumulation factors				Location	Source of ¹³⁷ Cs	Reference			
		¹³⁷ Cs		Stable Cs							
		BF	BF* ^b	BF	BF*						
Green algae, multicellular:											
<u><i>Nitella hialina</i></u>	—	170				Lake Bol shoe Missavo, Russia		1.33			
<u><i>Oedogonium</i> sp.</u>	—	1,200				1.1 hectare pond	Chronic-release	1.39			
<u><i>Oedogonium</i> sp.</u>	0.3	3,000				Valkealampi Lake, Finland	Spike input	1.32			
<u><i>Oedogonium vulgare</i></u>	1	790				Lab study		1.51			
<u><i>Pithophora oedogonia</i></u>	0.1	170				Lab study		1.48			
<u><i>Pithophora oedogonia</i></u>	30	56				Lab study		1.48			
<u><i>Rhizoclonium hieroglyphicum</i></u>	1	1,500				Lab study		1.51			
<u><i>Scenedesmus acuminatus</i></u>	—	39				Lab study		1.45			
<u><i>Scenedesmus quadricauda</i></u>	—	28				Lab study		1.45			
<u><i>Spirogyra communis</i></u>	13	220				Lab study		1.51			
<u><i>Spirogyra crassa</i></u>	—	28				Lab study		1.45			
<u><i>Spirogyra ellipsospora</i></u>	1	340				Lab study		1.51			
<u><i>Spirogyra</i> sp.</u>	—	190				Lab study		1.45			
<u><i>Spirogyra</i> sp.</u>	—	380				Lake Bol shoe Missavo, Russia		1.33			
<u><i>Spirogyra</i> sp.</u>	1	150				Experimental channel receiving river water	Chronic-release	1.15			
<u><i>Stigeoclonium lubricum</i></u>	14	89 ^c				Lab study		1.22			
<u><i>Tolipellopsis stelligera</i></u>	—	220				Lake Bol shoe Missavo, Russia		1.33			
<u><i>Ulothrix</i> sp.</u>	—	1,400				Lab study		1.52			
<u><i>Ulothrix</i> sp.</u>	—	460				Isere River	Fallout	1.52			
<u><i>Vaucheria walzii</i></u>	1	500				Experimental channel receiving river water	Chronic-release	1.15			

Table 3.1.7 (continued)

Taxon/functional category	K concentration (ppm)	Bioaccumulation factors				Location	Source of ¹³⁷ Cs	Reference	
		¹³⁷ Cs BF	¹³⁷ Cs BF* ^b	Stable Cs BF	Stable Cs BF*				
Green algae, multicellular:									
Mixed species	—	1,500-4,000				Concrete-lined pond	Spike input	1.39	
Mixed species	1.8	230	600 ^f			White Oak Lake, Tennessee	Chronic-release	1.27	
Green algae, unicellular:									
<u>Chlamydomonas</u> sp.	8	52				Lab study		1.51	
<u>Chlamydomonas</u> <u>moewussi</u>	—	130-370				Lab study		1.26	
<u>Oocystis</u> <u>elliptica</u>	10	670				Lab study		1.51	
Functional categories:									
Phytoplankton	1.6			1900 ^e		Lake Michigan		1.4	

^aDry-weight bioaccumulation factors were divided by 10 to convert them to wet-weight bioaccumulation factors.

^bA BF* is a bioaccumulation factor calculated from isotope concentration in water after filtration.

^cAverage over all temperatures.

^dIncludes samples where more than 80% of the cells were diatoms.

^eGeometric mean of bioaccumulation factors calculated from each pair of filtered (0.45 μ pore size) water concentrations and non-zero "corrected" concentrations in phytoplankton given in Appendix II of Copeland and Ayers (1970).

^fWater ultracentrifuged — equivalent to 0.7 μ pore-size filter.

cesium bioaccumulation factor was decreased by increasing sodium concentration in water. Both Harvey (1969, 1970) and Gertz (1973) reported that the cesium bioaccumulation factor is unaffected by non-lethal temperatures.

Both the nutrient concentration and general health of the algae influence the cesium bioaccumulation factor for algae, since the bioaccumulation factor is higher at higher phosphate concentrations (Gertz, 1973) and the bioaccumulation factor of dead cells is lower than that of live cells (Williams, 1970; Williams, 1960). Moreover, the cesium bioaccumulation factor for multicellular algae is higher in flowing water than in still water (Watts and Harvey, 1963).

Most bioaccumulation factors for algae in Table 3.1.7 are near or less than 10^3 . We therefore recommend that for algae in general a bioaccumulation factor of 10^3 be used.

3.1.3.3 Cesium Bioaccumulation Factors for Vascular Aquatic Plants

Some vascular aquatic plants may obtain a significant fraction of their body content of cesium and other elements from root uptake of elements from interstitial water of bottom sediments. Diffusion of elements into interstitial water from water overlying sediments is slow. This may explain why the bioaccumulation factor of fallout-derived ^{137}Cs in *Nuphar* sp. (Table 3.1.8) is much lower than the corresponding bioaccumulation factor of stable cesium. Since most bioaccumulation factors in Table 3.1.8 are less than or about 10^3 , we recommend that a bioaccumulation factor of 10^3 be used for aquatic vascular plants.

Table 3.1.8 Bioaccumulation factors for cesium in aquatic vascular plants^a

Species	K concentration in water (ppm)	BF, fallout ¹³⁷ Cs	BF, spike input of ¹³⁷ Cs	BF, chronic release of ¹³⁷ Cs	BF, stable Cs	BF*, stable Cs ^b	Location	Reference
<u><i>Azolla fuliculoides</i></u>	—		250				Concrete-lined pond	1.39
<u><i>Ceratophyllum demersum</i></u>	—		400-1,000				Concrete-lined pond	1.39
<u><i>Ceratophyllum</i> sp.</u>	9.8	370			490		Wintergreen Lake, Michigan	1.7
<u><i>Elodea canadensis</i></u>	—		1,000				Concrete-lined pond	1.39
<u><i>Elodea canadensis</i></u>		390					Isere River	1.52
<u><i>Elodea canadensis</i></u>	2.6				1,400		200-m ² carp pond	1.53
<u><i>Lemna minor</i></u>	—		500				Concrete-lined pond	1.39
<u><i>Lemna minor</i></u>	—	500					Isere River	1.52
<u><i>Nuphar luteum</i></u>	3.5	130					Finnish lakes	1.30
<u><i>Nuphar luteum</i></u>	1.8	410					Finnish lakes	1.30
<u><i>Nuphar luteum</i></u>	1.0	1,000					Finnish lakes	1.30
<u><i>Nuphar luteum</i></u>	0.8	1,500					Finnish lakes	1.30
<u><i>Nuphar luteum</i></u>	0.6	1,100					Finnish lakes	1.30
<u><i>Nuphar</i> sp.</u>	9.8	770			3,800		Wintergreen Lake, Michigan	1.7
<u><i>Nymphaea lutea</i></u>	2.1				280	390	Lake Maggiore, Italy	1.3,1.35
<u><i>Potamogeton pectinatus</i></u>	—		700				Concrete-lined pond	1.39

^aDry-weight bioaccumulation factors were multiplied by 0.15 to convert them to wet-weight bioaccumulation factors.

^bA BF* is a bioaccumulation factor calculated from isotope concentration in water after filtration.

3.1.3.4 Cesium Bioaccumulation Factors for Emergent Vascular Plants

Most bioaccumulation factors for emergent vascular plants (Table 3.1.9) are less than or near 10^3 ; we therefore recommend that for this group in general a bioaccumulation factor of 10^3 be applied.

3.1.3.5 Cesium Bioaccumulation Factors for Invertebrates

There appears to be much interspecific variation in bioaccumulation factors for crustaceans (Table 3.1.10). Nevertheless, most bioaccumulation factors in Table 3.1.10 are near 10^3 , therefore, a bioaccumulation factor of 10^3 is recommended for invertebrates in general. However, bioaccumulation factor for shells of invertebrates may be much lower.

3.1.3.6 Cesium Bioaccumulation Factors for Amphibians

The few bioaccumulation factors collected for amphibians (Table 3.1.11) suggest that a bioaccumulation factor of 10^4 be used.

3.1.3.7 Cesium Bioaccumulation Factors for Waterfowl

The few bioaccumulation factors collected for waterfowl (Table 3.1.12) suggest that a bioaccumulation factor of 3×10^3 be used.

Table 3.1.9 Bioaccumulation factors for cesium in emergent vascular plants

Species/part	K concentration in water (ppm)	BF, fallout ¹³⁷ Cs	BF, spike input of ¹³⁷ Cs	BF, chronic release of ¹³⁷ Cs	BF, stable Cs	Location	Reference
<u><i>Equisetum fluviatile</i></u> ^a	3.5	480				Finnish lakes	1.30
<u><i>Equisetum fluviatile</i></u> ^a	1.8	1,100				Finnish lakes	1.30
<u><i>Equisetum fluviatile</i></u> ^d	1.0	1,600				Finnish lakes	1.30
<u><i>Equisetum fluviatile</i></u> ^d	0.9	1,200				Finnish Lakes	1.30
<u><i>Equisetum fluviatile</i></u> ^d	0.8	7,400				Finnish Lakes	1.30
<u><i>Phragmites communis</i></u>	2.6				1,200	200-m ² carp pond	1.53
<u><i>Polygonum lapathifolium</i></u> : seeds	—			240		1.1 hectare pond	1.39
<u><i>Polygonum persicaria</i></u> : leaves	—		600			Concrete-lined pond	1.39
seeds			400				
<u><i>Scirpus acutus</i></u> : culms	—		90			Concrete-lined pond	1.38,1.39
seeds			400			Concrete-lined pond	
leaves			100			Concrete-lined pond	
roots			400			Concrete-lined pond	
seeds				70		1.1 hectare pond	
<u><i>Scirpus americanus</i></u> : culms	—		50			Concrete-lined pond	1.39
seeds			300				
<u><i>Typha latifolia</i></u> : leaves	—		250			Concrete-lined pond	1.38,1.39
seeds			100				
roots			250				

^aDry-weight bioaccumulation factors were multiplied by 0.20 to convert them to wet-weight bioaccumulation factors.

Table 3.1.10 Bioaccumulation factors for cesium in aquatic invertebrates

Taxon	K concentration in water (ppm)	BF, fallout ¹³⁷ Cs	BF*, fallout ¹³⁷ Cs ^a	BF, spike input ¹³⁷ Cs	BF, chronic release of ¹³⁷ Cs	BF*, chronic release of ¹³⁷ Cs	BF, stable Cs	BF*, stable Cs	Location	Reference
Crustaceans:										
<u>Gammarus lacustris</u>	1.7			1,000				340 ^b	East Twin Lake, Colorado	1.12,1.18
<u>Hyalella azteca</u>	—				11,000				Concrete-lined pond	1.39
<u>Mysis relicta</u>	1.6						60 ^c		Lake Michigan	1.6
Zooplankton (primarily copepods)	1.6						600 ^d		Lake Michigan	1.4
Zooplankton (<u>Daphnia</u> and <u>Cyclops</u>)	1.7						340 ^b		East Twin Lake, Colorado	1.18
Insect larvae:										
<u>Chaoborus</u> sp.	9.8	830					1,600		Wintergreen Lake, Michigan	1.7
<u>Chironomid</u> larvae	1.8				640	1,600			White Oak Lake, Tennessee	1.27
<u>Erythemis</u> <u>callipata</u>	—			800					Concrete-lined pond	1.39
<u>Ischnura</u> sp.	—			800					Concrete-lined pond	1.39
Snails:										
<u>Limnea</u> <u>stagnalis</u>	2.6						1,300		200-m ² carp pond	1.53
<u>Radix</u> <u>japonica</u>	—			600					Concrete-lined pond	1.39

Table 3.1.10 (continued)

Taxon	K concentration in water (ppm)	BF, fallout ¹³⁷ Cs	BF*, fallout ¹³⁷ Cs ^a	BF, spike input ¹³⁷ Cs	BF, chronic release of ¹³⁷ Cs	BF*, chronic release of ¹³⁷ Cs	BF, stable Cs	BF*, stable Cs	Location	Reference
Clams:										
<u>Lampsiliis radiata</u>	1.4								Small stream, South Carolina	1.20
soft tissues					>220 ^e					
shell					>25					

^aA BF* is a bioaccumulation factor calculated from isotope concentration in water after filtration.

^bSpike input of stable cesium.

^cWater concentration taken from Copeland and Ayers (1970).

^dBioaccumulation factor equals geometric mean of "uncorrected" concentrations of cesium in zooplankton from Appendix II of Copeland and Ayers (1970) divided by mean cesium concentration in water.

^eOn the basis of biological half-life given by the author, the clams had not fully attained steady-state with the ¹³⁷Cs concentration of water into which they were placed.

Table 3.1.11 Bioaccumulation factors
for ^{137}Cs in amphibians
(Pendleton and Hanson,
1958)

Species/Tissue	BF
Bullfrog tadpole (<u>Rana catesbeiana</u>):	
entire	2,600
gut	4,500
flesh	1,000
Spadefoot toad tadpole (<u>Scaphiopus hammondi</u> <u>intermontanus</u>):	
entire	6,000
Bullfrog adult:	
muscle	8,000

Table 3.1.12 Bioaccumulation factors for ^{137}Cs in waterfowl (Pendleton and Hanson, 1958)

Species/Tissue	BF
American coot (<u>Fulica a. americana</u>):	
muscle	1,800
liver	2,200
bone	800
Common mallard (<u>Anas platyrhynchos</u>):	
muscle	2,000
liver	2,500
bone	700
Ruddy duck (<u>Oxyura jamaicensis rubida</u>):	
muscle	2,200
liver	2,800
bone	900

References for Section 3.1

The numbers in the left-hand margin correspond to the reference numbers used in the tables of section 3.1.

- 1.1 Blanchard, R. L., and B. Kahn, The fate of radionuclides discharged from a PWR nuclear power station into a river, Symposium on Environmental Behavior of Radionuclides Released in the Nuclear Industry, France, May 14-18, 1973, IAEA/SM-172/26.
- 1.2 Bortoli, M. de, P. Gaglione, A. Malvicini, and C. Polvani, Concentration factors for strontium and caesium in fish of the lakes in the region of Varese (Northern Italy) Minerva, Fisiconucleare, 324-331 (1967).
- 1.3 Bortoli, M. de, P. Gaglione, A. Malvicini, and E. Van Der Stricht, Environmental Radioactivity Report EUR-3554e, Ispra (1966).
- 1.4 Copeland, R. A., and J. C. Ayers, Trace element distribution in water, sediment, phytoplankton, zooplankton, and benthos of Lake Michigan: A baseline study with calculations of concentration factors and buildup of radioisotopes in the food web, Environmental Research Group Special Report No. 1, Great Lakes Research Division, Univ. Michigan, Ann Arbor, 1970.
- 1.5 Copeland, R. A., R. H. Beethe, and W. W. Prater, Trace element distribution in Lake Michigan fish: A baseline study with calculations of concentration factors and equilibrium radioisotopes distribution, Environmental Research Group Special Report No. 2, Great Lakes Research Division, Univ. Michigan, Ann Arbor, March, 1973.
- 1.6 Edgington, D. N., E. Harvey, and M. S. Torrey, A preliminary study of the chemical composition of the opossum shrimp, Mysis relicta, IN Radiological and Environmental Research Division Ann. Report, R. E. Rowland, ANL 7960, Part III, 1972.
- 1.7 Eyman, L. D., Cesium-137 and stable cesium in a hypereutrophic lake, Ph.D. Thesis, Michigan State University, East Lansing, 1972, 98 pp.
- 1.8 Eyman, L. D., and J. T. Kitchings, Availability of ¹³⁷Cs to fishes from ingested clays, IN Proceedings of the XIX Congress of the International Association of Limnology (SIL), Winnipeg, Canada, August 22-29, 1974 (in press).
- 1.9 Fleishman, D. J., Accumulation of artificial radionuclide in freshwater fish, IN Radioecology, V. M. Klechkovskii, G. G. Polikarpov, and R. M. Aleksakhin (eds.), pp. 347-369, John Wiley and Sons, New York, 1973.

References for Section 3.1 (continued)

- 1.10 Friend, A. G., The aqueous behavior of strontium-85, cesium-137, zinc-65, and cobalt-60 as determined by laboratory type studies, IN Transport of radionuclides in freshwater systems, B. H. Kornegay, W. A. Vaughn, D. K. Jamison, and J. M. Morgan (eds.), pp. 43-60, TID-7664 (1963).
- 1.11 Fujita, M., J. Iwamoto, and M. Kondo, Comparative metabolism of cesium and potassium in mammals--Interspecies correlation between body weight and equilibrium level, *Health Phys.* 12, 1237-1247 (1966).
- 1.12 Gallegos, A. F., Radiocesium kinetics in the components of a montane lake ecosystem, Ph.D. Thesis, Colorado State Univ., Fort Collins, December 1969.
- 1.13 Gallegos, A. F., and F. W. Whicker, Radiocesium retention by rainbow trout as affected by temperature and weight, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 361-371 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.
- 1.14 Gallegos, A. F., F. W. Whicker, and T. W. Hakonson, Accumulation of radiocesium in rainbow trout via a non-foodchain pathway, Fifth Annual Health Physics Society Mid-Year Topical Symposium, Idaho Falls, Idaho, 1970.
- 1.15 Garder, K., and O. Skulberg, An experimental investigation on the accumulation of radioisotopes by freshwater biota, *Arch. Hydrobiol.* 62, 50-69 (1966).
- 1.16 Garder, K., and O. Skulberg, Sorption phenomena of radionuclide to clay particles in river water, *Int. J. Air Wat. Poll.* 8, 229-241 (1964).
- 1.17 Gertz, S. M., The fate of a radionuclide in the aquatic environment--A mathematical equation representing radionuclide uptake by algae, Ph.D. Thesis, Drexel University, Philadelphia, 112 pp., 1973.
- 1.18 Hakonson, T. E., A. F. Gallegos, and F. W. Whicker, Use of cesium-133 and activation analysis for measurement of cesium kinetics in a montane lake, IN Radionuclides in Ecosystems, D. J. Nelson (ed.), pp. 344-348 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.
- 1.19 Harvey, R. S., Uptake of radionuclides by freshwater algae and fish, *Health Phys.* 10, 243-247 (1964).

References for Section 3.1 (continued)

- 1.20 Harvey, R. S., Uptake and loss of radionuclides by the freshwater clam Lampsilis radiata, *Health Phys.* 17, 149-154 (1969).
- 1.21 Harvey, R. S., Effects of temperature on the sorption of radio-nuclides by a blue-green algae, IN *Symposium on Radioecology*, D. J. Nelson and F. C. Evans (eds.), pp. 266-269 (1969), *Proceedings of the Second National Symposium on Radioecology*, Ann Arbor, Michigan, May 15-17, 1967.
- 1.22 Harvey, R. S., Temperature effects on the sorption of radio-nuclides by freshwater algae, *Health Phys.* 19, 293-297 (1970).
- 1.23 Hasanen, E., S. Kolehmainen, and J. K. Miettinen, Biological half-time of cesium-137 in three species of freshwater fish: Perch, roach and rainbow trout, IN *Radioecological concentration process*, B. Aberg and F. P. Hungate (eds.), pp. 921-924 (1967), *Proceedings of the International Symposium*, Stockholm, April 25-29, 1966.
- 1.24 Kevern, N. L., Feeding rate of carp estimated by a radioisotopic method, *Trans. Amer. Fisheries Soc.* 95(4), 363-371 (1966).
- 1.25 Kevern, N. R., N. A. Griffith, and T. Grizzard, Biological half-life of cesium-134 in carp and two aquatic insects, *Health Physics Division Annual Progress Report for the Period Ending July 31, 1964, Radiation Ecology Section*, pp. 101-102, ORNL-3697 (1964).
- 1.26 King, S. F., Uptake and transfer of cesium-137 by Chlamydomonas, Daphnia, and bluegill fingerlings, *Ecology* 45, 852-859 (1964).
- 1.27 Kolehmainen, S. E., Balances of cesium-137, stable cesium and potassium, in bluegill, Lepomis macrochirus Raf., and other fish in White Oak Lake, *Health Phys.* 23, 301-315 (1972).
- 1.28 Kolehmainen, S., E. Hasanen, and J. K. Miettinen, Cesium-137 levels in fish of different limnological types of lakes in Finland during 1963, *Health Phys.* 12, 917-922 (1966).
- 1.29 Kolehmainen, S., E. Hasanen, and J. K. Miettinen, Cesium-137 in fish, plankton, and plants in Finnish lakes, IN *Radioecological concentration process*, B. Aberg and F. P. Hungate (eds.), pp. 913-919 (1967), *Proceedings of the International Symposium*, Stockholm, April 25-29, 1966.
- 1.30 Kolehmainen, S., E. Hasanen, and J. K. Miettinen, Cesium-137 in the plants, plankton and fish of the Finnish lakes and factors affecting its accumulation, IN *Proceedings of the First International Congress of Radiation Protection*, W. S. Snyder (ed.), pp. 407-415 (1968), Permagon Press, Oxford.

References for Section 3.1 (continued)

- 1.31 Kolehmainen, S. E., and D. J. Nelson, Balances of cesium-137, stable cesium, and the feeding rates of bluegill (Lepomis macrochirus Raf.) in White Oak Lake, ORNL-4445 (1969).
- 1.32 Kolehmainen, S., Romantschuk, S. Takatalo, and J. K. Miettinen, Pollution experiments with Cs-137 in lakes of two limnologically different types. IN Congrès international sur la radio-protection du milieu devant le développement des utilisations pacifiques de l'énergie nucléaire, pp. 181-200, Toulouse, 14-16 Mars 1967, Société Française de Radio-protection, Paris (1968).
- 1.33 Kulikov, N. V., Radioecology of freshwater plants and animals, IN Radioecology, V. M. Klechkovskii, G. G. Polikarpov, and R. M. Aleksakhin (eds.), pp. 323-337 (1973), Wiley, New York.
- 1.34 Marshall, J. S., and J. H. LeRoy, Iron, manganese, cobalt, and zinc cycles in a South Carolina reservoir, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 465-473 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.
- 1.35 Merlini, M., F. Giradi, and G. Pozzi, Activation analysis in studies of an aquatic ecosystem, IN Nuclear activation techniques in the life sciences, pp. 615-629 (1967), Proceedings of the Symposium held by the IAEA in Amsterdam, May 8-12, 1967, also STI/PUB/155, SM-91/7.
- 1.36 Nelson, D. J., Cesium, cesium-137, and potassium concentrations in white crappie and other Clinch River fish, IN Symposium on radioecology, D. J. Nelson and F. C. Evans (eds.), pp. 240-248 (1969), Proceedings of the Second National Symposium on Radioecology, Ann Arbor, Michigan, May 15-17, 1967.
- 1.37 Nelson, D. J., N. A. Griffith, and S. A. Rucker, Radionuclide excretion studies, IN Health Physics Division Annual Progress Report for Period Ending July 31, 1969, pp. 119-120, ORNL-4446 (1970).
- 1.38 Pendleton, R. C., Effects of some environmental factors on bio-accumulation of cesium-137 in an aquatic community, IN Hanford Biology Research Annual Report for 1958, J. J. Davis (ed.), pp. 42-46 (1959).
- 1.39 Pendleton, R. C., and W. C. Hanson, Absorption of cesium-137 by components of an aquatic community, IN Proceedings of the Second International Conference on Peaceful Uses of Atomic Energy, Vol. 18, 419-422 (1958), U.N. Geneva, September 1-3, 1958.

References for Section 3.1 (continued)

1.40 Pendleton, R. C., C. W. Mays, R. D. Lloyd, and B. W. Church, A trophic level effect on cesium-137 concentration, *Health Phys.* 11, 1503-1510 (1965).

1.41 Peterson, H. I., Jr., Stable element and radionuclide re-concentration in aquatic systems. I. Stable element effects on equilibrium situation, IN Proceedings of the Fifth Annual Health Physics Society Midyear Topical Symposium, *Health Physics Aspects of Nuclear Siting*, Vol. 2, pp. 289-328, AEC CONF 701106 (1970).

1.42 Preston, A., D. F. Jefferies, and J. W. R. Dutton, The concentration of cesium-137 and strontium-90 in the flesh of brown trout taken from rivers and lakes in the British Isles between 1961-1966, The variables determining the concentrations and their use in radiological assessment, *Water Res.* 1, 475-496 (1967).

1.43 Reynolds, T. D., and E. F. Gloyna, Uptake and release of radio-nuclides by stream sediments, IN Proceedings of the Second International Conference on Advances in Water Pollution Research, Vol. 1, O. Jaag (ed.), pp. 151-172 (1964), Pergamon Press, New York.

1.44 Struxness, E. G., P. H. Carrigan, Jr., M. A. Churchill, K. E. Cowser, R. J. Morton, D. J. Nelson, and F. L. Parker, Comprehensive report of the Clinch River study, U.S. AEC Report ORNL-4035, Oak Ridge National Laboratory, Oak Ridge, Tennessee, 1967.

1.45 Timofeyeva-Resovskaya, Ye. A., Distribution of radioisotopes in the main components of freshwater bodies (Monograph), *Tr. Inst. Biol. Akad. Nauk SSSR, Ural'skiy filia* 30, 1-78 (JPRS 21.816) 1963.

1.46 Vanderploeg, H. A., R. S. Booth, and F. H. Clark, A specific activity and concentration model applied to cesium movement in an oligotrophic lake, IN *Mineral Cycling in Southeastern Ecosystems*, F. G. Howell (ed.), (in press), Symposium held 1-3 May 1974, Augusta, Georgia.

1.47 Watts, J. R., and R. S. Harvey, Uptake and retention of ¹³⁷Cs by a blue-green alga in continuous flow and batch culture systems, *Limnol. Oceanogr.* 8, 45-49 (1963).

1.48 Williams, L. G., Concentration of strontium-85 and cesium-137 from water solutions by Cladophora and Pithophora, *J. Phycol.* 6, 314-316 (1970).

1.49 Williams, L. G., Uptake of cesium-137 by cells and detritus of Euglena and Chlorella, *Limnol. Oceanogr.* 5, 301-311 (1960).

References for Section 3.1 (continued)

- 1.50 Williams, L. G., and Q. Pickering, Direct and food-chain uptake of cesium-137 and strontium-85 in bluegill fingerlings, *Ecology* 42, 205-206 (1961).
- 1.51 Williams, L. G., and H. D. Swanson, Concentration of cesium-137 by algae, *Science* 127, 187-188 (1958).
- 1.52 Wlodek, S., The behavior of cesium-137 in freshwater reservoirs with particular reference to the reactions of plant communities to contamination, IN Radioecological concentration process, B. Aberg and F. P. Hungate (eds.), pp. 897-912 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1971.
- 1.53 Wlodek, S., M. Bysiek, and D. Grzyboska, Autochthonous caesium and behavior of allochthonous caesium in a fish pond, *Ecologia Polska, Seria A*, Vol. 18(30), 1-11 (1970).

3.2 STRONTIUM

3.2.1 Strontium Metabolism

As a result of their chemical similarities and the greater abundance of calcium in nature, calcium is a non-isotopic carrier for strontium. Since calcium is homeostatically maintained at constant concentrations in fishes (Agnedal, 1967; Ophel and Judd, 1973; Peterson, 1970; Reed and Nelson, 1973; Suzuki et al., 1972; Templeton and Brown, 1964) and presumably other animals, the bioaccumulation factor for calcium in animals is given by Equation (2.1.7). The bioaccumulation factor for strontium is then given by Equation (2.1.8). These relations do not hold for algae since the calcium concentration in water does not appear to greatly influence the bioaccumulation factor for strontium in algae (Kevern, 1964a; Williams, 1970).

The primary uptake of calcium and strontium in fishes, and probably in many aquatic invertebrates, occurs directly from the water. The gill membranes of fishes are the primary sites of calcium and strontium uptake (Nelson, 1966; Suzuki et al., 1972; Templeton and Brown, 1964). Only about one-tenth of the calcium and strontium taken up by fishes is through the food chain (Agnedal, 1966). Since uptake from water is the primary pathway, the food chain dynamics of calcium and strontium are of little importance in the determination of bioaccumulation factors for strontium in aquatic organisms. Thus, the discrimination coefficient q_i in Equation (2.1.8) is not a function of trophic level in fishes because the primary source of strontium to fishes is water. The discrimination coefficient has also been shown to

be independent of the calcium concentration in water (Agnedal, 1967; Ophel and Judd, 1973; Peterson, 1970; Reed and Nelson, 1973; Suzuki et al., 1972; Templeton and Brown, 1964).

3.2.2 Environmental Strontium

Strontium is commonly found in nature with calcium, its closest chemical analogue (Nelson, 1961). Strontium concentrations in freshwater rivers and lakes may range between 0.004 and 0.23 ppm (Nelson, 1967; Ophel et al., 1972; Suzuki et al., 1972; Templeton and Brown, 1964). The more abundant element calcium is found in freshwater rivers and lakes in concentrations ranging from 1.9 to 114 ppm (Austin, 1963; Beninson et al., 1966; Nelson, 1967; Ophel et al., 1972; Suzuki et al., 1972; Templeton and Brown, 1964). Strontium has physicochemical properties similar to calcium and, like calcium, appears mainly in ionic form in water (Templeton and Brown, 1964). Strontium and calcium are not strongly sorbed by suspended particulate materials in the water. The fraction of strontium present in the suspended phase is given for a number of freshwater ecosystems in Table 3.2.1. The fraction of strontium in the suspended phase ranges between 0.9 and 10%. Since q_i is independent of $[Ca]_w$ and because both calcium and strontium have nearly the same physicochemical forms in water, the discrimination coefficient is not expected to vary much among sites.

3.2.3 Review of Strontium Bioaccumulation Factors

Bioaccumulation factors are discussed for the following categories of aquatic biota:

Table 3.2.1 Percentage of radiostrontium in water in the particulate phase in some freshwater ecosystems

Water body	Mean % in particulate phase	Suspended solids (ppm)	Separation method	Reference
White Oak Lake, Tennessee ^a	2	31 (mean)	Ultracentrifugation: Particles >0.7 μ	2.24
Clinch River, Tennessee ^a	2-9	25-185	Ultracentrifugation: Particles >0.7 μ	2.24
Tennessee River, Tennessee ^a	9-10	9-22	Ultracentrifugation: Particles >0.7 μ	2.24
Experimental Pond ^b	0.9	21	0.45- μ filter	2.7

^aRange of mean of three river stations.

^bBased on five water values sampled between 24 and 80 days after introduction of radionuclides.

1. Fishes
2. Algae
3. Aquatic and Emergent Macrophytes
4. Molluscs.

The bioaccumulation factors were examined for taxa in each of the above categories. For all taxa except algae, the strontium bioaccumulation factors were derived only from concentrations in natural or semi-natural bodies of water.

3.2.3.1 Strontium Bioaccumulation Factors for Fishes

A regression analysis of $\ln BF(Ca)_i$ versus $\ln [Ca]_W$ for brown trout (Templeton and Brown, 1964) and for three species of fish from Swedish lakes (Agnedal, 1966) yielded a linear plot with a slope of approximately -1 and a high degree of negative correlation (Peterson, 1970). The bioaccumulation factor for calcium can thus be given by Equation (2.1.7). Since calcium is a non-isotopic carrier for strontium, the bioaccumulation factor for strontium is given by Equation (2.1.8). The analyses by Peterson (1970) of data from Templeton and Brown (1964) and Agnedal (1967) clearly showed that the q_i values for freshwater fishes are independent of the concentrations of calcium in the water. This implies that the regression of $\ln BF(Sr)_i$ on $\ln [Ca]_W$ has a slope of about -1.

The strontium bioaccumulation factors, calcium concentrations in fishes and water, and q_i values for strontium in fish bone and flesh are given in Table 3.2.2 for studies that reported calcium concentration in water. Only data based on several measurements of strontium in fishes and water are listed. Some studies have

listed values for the strontium discrimination coefficients in bone and no bioaccumulation factors. In these studies, the discrimination coefficients were calculated from strontium and calcium concentrations in ashed bone. The ratio between ash weight and fresh weight in bone is not constant for all species or for all ashing procedures. Since a fresh weight cannot be assumed for a given ashed weight, bioaccumulation factors for strontium could not be calculated from these studies.

Because q_j is not a function of feeding habits, we decided to derive a relation between the strontium bioaccumulation factor and $[Ca]_W$ for all fish data combined. To do this, regressions of $\ln BF(Sr)_j$ versus $\ln [Ca]_W$ for all fishes combined were made. Results of these regressions for fish flesh and bone are given in Table 3.2.3 and in Figure 3.2.1. The slope of the regression for fish flesh is not significantly different ($P > 0.05$) from the expected slope of -1. The slope of the regression for bone, although significantly different from -1 ($P < 0.05$), is not appreciably different. To predict the bioaccumulation factor of strontium in fishes we recommend use of Figure 3.2.1 or the relation given in Table 3.2.3.

3.2.3.2 Strontium Bioaccumulation Factors for Algae

The bioaccumulation factors for strontium in algae are given in Table 3.2.4 for both field and laboratory studies. In a laboratory study, Kevern (1964) showed that the bioaccumulation factor for strontium in the unicellular green alga Oocytis eremosphaeria was independent of the calcium concentration in the water. In Table 3.2.4, the multicellular algae Pithophora and Cladophora glomerata (Williams, 1970) show only a small decrease in the bioaccumulation factor for strontium accompanying a very large increase in calcium

Table 3.2.2 Bioaccumulation factors and discrimination coefficients for strontium in fishes

Species	Tissue	Ca Concentration		BF(Sr) _i	q _i	Location	Reference
		Tissue (ppm)	Water (ppm)				
Perch	Bone	23100	2.0	4,030	0.35	Lake Langsjon, Sweden	2.1
	Muscle	480		92	0.38		
Perch	Bone	31000	40.0	170	0.22	Lake Storacksen, Sweden	2.1
Perch	Bone	31600	46.0	160	0.23	Lake Erken, Sweden	2.1
	Muscle	390		3	0.35		
Perch	Bone	43600	54.0	130	0.16	Lake Glisstjarn, Sweden	2.1
	Muscle	430		1.3	0.16		
Perch	Bone	39400	4.4	1,840	0.21	Lake Ulkesjon, Sweden	2.1
Perch	Bone	40000	12	410	0.12	Lake Viggen, Sweden	2.1
Perch	Bone	34200	18	330	0.17	Lake Orlangen, Sweden	2.1
Perch	Bone	39800	26	380	0.25	Lake Magelungen, Sweden	2.1
Perch	Bone	31700	63	100	0.20	Lake Sovdesjon, Sweden	2.1
Perch	Bone	30200	63	110	0.23	Lake Nittsjon, Sweden	2.1
Pike	Bone	42500	2.0	4,930	0.23	Lake Langsjon, Sweden	2.1
	Muscle	910		125	0.27		

Table 3.2.2 (continued)

Species	Tissue	Ca Concentration		BF(Sr) _i	q _i	Location	Reference
		Tissue (ppm)	Water (ppm)				
Pike	Bone	65400	40.0	290	0.18	Lake Storacksen, Sweden	2.1
	Muscle	470		3	0.25		
Pike	Bone	51000	54.0	180	0.19	Lake Glisstjarn, Sweden	2.1
	Muscle	440		1.7	0.21		
Pike	Bone	47000	26	310	0.17	Lake Magelungen, Sweden	2.1
	Muscle	590		6	0.26		
Pike	Bone	35100	2.0	8,810	0.50	Lake Rogen, Sweden	2.1
Pike	Bone	44500	4.4	2,190	0.22	Lake Ulkesjon, Sweden	2.1
Pike	Bone	42000	4.5	1,770	0.19	Lake M. Mollesjon Sweden	2.1
Pike	Bone	44500	12	480	0.13	Lake Viggen, Sweden	2.1
Pike	Bone	51200	18	450	0.16	Lake Orlangen, Sweden	2.1
Pike	Bone	20100	63	50	0.16	Lake Sovdesjon, Sweden	2.1
Pike	Bone	43100	63	120	0.18	Lake Nittsjon, Sweden	2.1
Roach	Bone	38600	2.0	9,170	0.48	Lake Langsjon, Sweden	2.1
	Muscle	880		198	0.45		
Roach	Bone	45300	40.0	390	0.35	Lake Storacksen, Sweden	2.1
	Muscle	930		9	0.39		
Roach	Bone	47000	46.0	310	0.30	Lake Erken, Sweden	2.1
	Muscle	800		7	0.41		
Roach	Bone	37300	54.0	240	0.35	Lake Glisstjarn, Sweden	2.1
	Muscle	550		3	0.30		

Table 3.2.2 (continued)

Species	Tissue	Ca Concentration		BF(Sr) _i	q _i	Location	Reference
		Tissue (ppm)	Water (ppm)				
Roach	Bone	30600	26	420	0.36	Lake Magelungen, Sweden	2.1
	Muscle	580		10	0.45		
Roach	Bone	38700	12	730	0.23	Lake Viggen, Sweden	2.1
Roach	Bone	19500	18	350	0.32	Lake Orlangen, Sweden	2.1
Roach	Bone	25800	63	120	0.29	Lake Sovdesjon, Sweden	2.1
Roach	Bone	50200	63	270	0.34	Lake Nittsjon, Sweden	2.1
<u>Salmo trutta</u>	Bone	47800	0.84	34,559	0.61	Loch Glutt, United Kingdom	2.26
	Muscle	203		198	0.82		
<u>Salmo trutta</u>	Bone	80000	4.90	6,260	0.38	Windermere, United Kingdom	2.26
	Muscle	140		9.8	0.34		
<u>Salmo trutta</u>	Bone	80400	4.50	3,194	0.18	River Prysor, United Kingdom	2.26
	Muscle	289		32	0.50		
<u>Pomoxis annularis</u>	Bone		27		0.29	Clinch River, Tennessee	2.17
	Flesh	135		1.00	0.20		
<u>Aplodinatus grunniens</u>	Bone		27		0.21	Clinch River, Tennessee	2.17
	Flesh	122		0.86	0.19		
<u>Roccus chrysops</u>	Bone		27		0.21	Clinch River, Tennessee	2.17
	Flesh	106		0.74	0.19		
<u>Ictalurus punctatus</u>	Bone		27		0.20	Clinch River, Tennessee	2.17
	Flesh	89		0.73	0.22		
<u>Lepomis macrochirus</u>	Bone		27		0.20	Clinch River, Tennessee	2.17
	Flesh	157		0.79	0.13		

Table 3.2.2 (continued)

Species	Tissue	Ca Concentration			q_i	Location	Reference
		Tissue (ppm)	Water (ppm)	$BF(Sr)_i$			
<u>Perca</u> <u>flavescens</u>	Whole Bone	5.7		472 3,225		Perch Lake, Ontario	2.20
<u>Ictalurus</u> <u>nebulosus</u>	Whole Bone	5.7		981 8,000		Perch Lake, Ontario	2.20
<u>Semotilus</u> <u>margarita</u>	Whole	5.7		456		Perch Lake, Ontario	2.20
<u>Lipomis</u> <u>gibbosus</u>	Whole	5.7		1,250		Perch Lake, Ontario	2.20
<u>Perca</u> <u>fluviatalis</u>	Whole	18.9		130		Lake Maggiore, Italy	2.4
<u>Perca</u> <u>fluviatalis</u>	Whole	9.5		110		Lake Varese, Italy	2.4
<u>Perca</u> <u>fluviatalis</u>	Whole	24.2		120		Lake Comabbio, Italy	2.4
<u>Perca</u> <u>fluviatalis</u>	Whole	31.4		300		Lake Monate, Italy	2.4
<u>Scardinius</u> <u>erythrophthalmus</u>	Whole	18.9		180		Lake Maggiore, Italy	2.4
<u>Scardinius</u> <u>erythrophthalmus</u>	Whole	9.5		210		Lake Varese, Italy	2.4
<u>Scardinius</u> <u>erythrophthalmus</u>	Whole	24.2		180		Lake Comabbio, Italy	2.4
<u>Scardinius</u> <u>erythrophthalmus</u>	Whole	31.4		380		Lake Monate, Italy	2.4

Table 3.2.2 (continued)

Species	Tissue	Ca Concentration			BF(Sr) _i	q _i	Location	Reference
		Tissue (ppm)	Water (ppm)					
<u>Lepomis gibbosus</u>	Whole	18.9		110			Lake Maggiore, Italy	2.4
<u>Lepomis gibbosus</u>	Whole	9.5		90			Lake Varese, Italy	2.4
<u>Lepomis gibbosus</u>	Whole	24.2		90			Lake Comabbio, Italy	2.4
<u>Lepomis gibbosus</u>	Whole	31.4		280			Lake Monate, Italy	2.4
<u>Lepomis gibbosus</u>	Bone	3.6 ^a		2,400			Par Pond, South	2.9
	Flesh			<48			Carolina	
<u>Micropterus salmoides</u>	Bone	3.6 ^a		1,700			Par Pond, South	2.9
	Flesh			<48			Carolina	
<u>Ictalurus natalis</u>	Bone	3.6 ^a		2,100			Par Pond, South	2.9
	Flesh			<48			Carolina	
<u>Perca flavescens</u>	Bone	6.3			0.21		Perch Lake, Ontario	2.21
<u>Ictalurus nebulosus</u>	Bone	6.3			0.54		Perch Lake, Ontario	2.21
<u>Perca flavescens</u>	Bone	26.7			0.18		Lake Huron, Ontario	2.21
<u>Catostomos catostomos</u>	Bone	26.7			0.21		Lake Huron, Ontario	2.21
<u>Dorosoma cepedianum</u>	Bone	26.7			0.25		Lake Huron, Ontario	2.21

Table 3.2.2 (continued)

Species	Tissue	Ca Concentration		$BF(Sr)_i$	q_i	Location	Reference
		Tissue (ppm)	Water (ppm)				
<u><i>Cyprinus carpio</i></u>	Bone	26.7		0.53		Lake Huron, Ontario	2.21
<u><i>Carassius auratus</i></u>	Bone	10.8		0.86		Lake Toyanogata, Japan	2.25
<u><i>Carassius auratus</i></u>	Bone	10.2		0.78		Lake Mikata, Japan	2.25
<u><i>Carassius auratus</i></u>	Bone	11.8		0.92		Lake Barato, Japan	2.25

^aPersonal communication.




Figure 3.2.1 Bioaccumulation factors for strontium in freshwater fishes as a function of calcium concentration in water.

Table 3.2.3. Parameters from linear regressions between $\ln BF(Sr)_i$ and $\ln [Ca]_W$ for bone and flesh of fishes. [To predict $BF(Sr)_i$ use the relation $BF(Sr)_i = \exp(\text{intercept} + \text{slope} \times \ln [Ca]_W)$.]

Tissue	Intercept \pm SE	Slope \pm SE	R^2
Bone	9.59 \pm 0.18	-1.15 \pm 0.06	0.901
Flesh	5.18 \pm 1.11	-1.21 \pm 0.37	0.736

Table 3.2.4 Bioaccumulation factors and discrimination coefficients for strontium in algae^a

Species	Ca concentration in water (ppm)	BF(Sr) _i	q _i	Location	Reference
Mixed blue-green algae	3.6 ^b	600		Par Pond, South Carolina	2.9
<u>Cladophora</u> <u>glomerata</u>	0.1	2,200		Laboratory	2.28
<u>Cladophora</u> <u>glomerata</u>	1.0	2,100		Laboratory	2.28
<u>Cladophora</u> <u>glomerata</u>	30	1,000		Laboratory	2.28
<u>Cladophora</u> sp.	23		1.8	Lough Neagh, U.K.	2.26
<u>Navicula</u> <u>seminulum</u>	17	70 ^c		Laboratory	2.12
<u>Nitella</u> sp.	23		0.38	Lough Neagh, U.K.	2.26
<u>Pithophora</u> <u>oedogonia</u>	0.1	2,100		Laboratory	2.28
<u>Pithophora</u> <u>oedogonia</u>	1.0	2,300		Laboratory	2.28
<u>Pithophora</u> <u>oedogonia</u>	30	1,280		Laboratory	2.28
<u>Plectonema</u> <u>boryanum</u>	39	240 ^c		Laboratory	2.12

Table 3.2.4 (continued)

Species	Ca concentration in water (ppm)	$BF(Sr)_i$	q_i	Location	Reference
<u><i>Spirogyra</i></u> sp.	5.7	120		Perch Lake, Ontario	2.20
<u><i>Spirogyra</i></u> sp.	5.4	900		Experimental Channel	2.8
<u><i>Stigeoclonium</i></u> <u><i>tubricum</i></u>	10.5	120 ^c		Laboratory	2.12
<u><i>Vaucheria</i></u> <u><i>walzii</i></u>	5.4	1,400		Experimental Channel	2.8

^aDry-weight bioaccumulation factors were divided by 10 to convert them to wet-weight bioaccumulation factors.

^bPersonal communication.

^cAverage over all temperatures.

concentration in the water. Thus it may be concluded that the bioaccumulation factor for strontium in algae is relatively independent of the calcium concentration in the water. Until additional data are available, a strontium bioaccumulation factor of 2×10^3 is recommended for algae.

3.2.3.3 Strontium Bioaccumulation Factors for Aquatic and Emergent Macrophytes

The bioaccumulation factors for aquatic and emergent macrophytes are found in Table 3.2.5. Discrimination coefficients are given when available. The data do not permit us to infer whether the strontium bioaccumulation factor is dependent on the calcium concentration in the water. On the basis of these few data, we recommend a bioaccumulation factor of 2×10^2 for aquatic and emergent macrophytes.

3.2.3.4 Strontium Bioaccumulation Factors for Molluscs

Table 3.2.6 gives the experimental data obtained on bioaccumulation of strontium in mollusc shells. The data in Table 3.2.6 and Equation (2.1.8) were used to calculate a bioaccumulation factor relation for strontium in mollusc shells. Mean values of q_i , the discrimination coefficient, and Σ_i^* , the concentration of calcium in mollusc shells, are taken from Table 3.2.6 to derive the relation. For mollusc shells, the strontium bioaccumulation factor relation derived was

$$BF(Sr)_i = \frac{6.8 \times 10^4}{[Ca]_w} ,$$

where $[Ca]_w$ has units of ppm.

Table 3.2.5 Bioaccumulation factors and discrimination coefficients for strontium in aquatic and emergent macrophytes

Species	Ca concentration in water (ppm)	BF(Sr) _i	q _i	Location	Reference
<u>Braseria schreberi</u>	6.3		0.55	Perch Lake, Ontario	2.21
<u>Ceratophyllum demersum</u>	5.7	220		Perch Lake, Ontario	2.20
<u>Elodea canadensis</u>	23		1.2	Lough Neagh, U.K.	2.26
<u>Fontinalis</u> sp.	6.0	240	1.2	Perch Lake, Ontario	2.20, 2.21
<u>Myriophyllum spicatum</u>	2.4		1.2	Devoke Water, U.K.	2.26
<u>Myriophyllum spicatum</u>	23		1.5	Lough Neagh, U.K.	2.26
<u>Nuphar variegatum</u>	6.3		1.8	Perch Lake, Ontario	2.21
<u>Nymphaea odorata</u>	6.3		0.90	Perch Lake, Ontario	2.21
<u>Pontederia cordata</u>	6.3		1.4	Perch Lake, Ontario	2.21
<u>Potamogeton natans</u>	2.4		1.5	Devoke Water, U.K.	2.26
<u>Potamogeton pectinatus</u>	23		1.7	Lough Neagh, U.K.	2.26
<u>Potamogeton perfoliatus</u>	23		0.84	Lough Neagh, U.K.	2.26
<u>Potamogeton pusillus</u>	6.0	190	2.0	Perch Lake, Ontario	2.20, 2.21
<u>Scirpus fluitans</u>	2.4		0.57	Dover Water, U.K.	2.26
<u>Scirpus subterminalis</u>	5.7	30		Perch Lake, Ontario	2.20
<u>Sparangium fluctuans</u>	5.7	150		Perch Lake, Ontario	2.20
<u>Sparangium</u> sp.	2.4		0.89	Devoke Water, U.K.	2.26
<u>Utricularia vulgaris</u>	5.7	160		Perch Lake, Ontario	2.20

Table 3.2.6 Bioaccumulation factors and discrimination coefficients for strontium in mollusc shells

Species	Ca Concentration		BF(Sr) _i	q _i	Location	Reference
	Shell (g/g)	Water (ppm)				
<u>Unioninae</u>						
<u>Elliptio dilatatus</u>	0.40	27	2,988	0.20	Clinch River, Tennessee	2.16
<u>Elliptio crassidens</u>	0.40	27	3,781	0.25	Clinch River, Tennessee	2.16
<u>Pleurobema cordatum</u>	0.40	27	2,913	0.20	Clinch River, Tennessee	2.16
<u>Fusconaia subrotunda</u>	0.40	27	2,671	0.18	Clinch River, Tennessee	2.16
<u>Lampsilinae</u>						
<u>Actinonaias carinata gibba</u>	0.40	27	2,691	0.18	Clinch River, Tennessee	2.16
<u>Ligumia recta latissima</u>	0.40	27	2,640	0.18	Clinch River, Tennessee	2.16
<u>Lampsilis ovata</u>	0.40	27	3,246	0.22	Clinch River, Tennessee	2.16
<u>Unioninae</u>						
<u>Quadrula mantenevra</u>	0.40	19	2,567	0.12	Tennessee River, Tennessee	2.16
<u>Quadrula pustulosa</u>	0.40	19	3,217	0.15	Tennessee River, Tennessee	2.16
<u>Elliptio crassidens</u>	0.40	19	3,117	0.15	Tennessee River, Tennessee	2.16
<u>Pleurobema cordatum</u>	0.40	19	3,767	0.18	Tennessee River, Tennessee	2.16

Table 3.2.6 (continued)

Species	Ca Concentration		BF(Sr) _i	q _i	Location	Reference
	Shell (g/g)	Water (ppm)				
<u>Unioninae</u> (cont'd)						
<u>Amblema costata</u>	0.40	19	3,202	0.15	Tennessee River, Tennessee	2.16
<u>Megalonaia gigantea</u>	0.40	19	2,995	0.14	Tennessee River, Tennessee	2.16
<u>Cycloraias tuberculata</u>	0.40	19	3,235	0.15	Tennessee River, Tennessee	2.16
<u>Lamsilinae</u>						
<u>Plagiola lineolata</u>	0.40	19	3,170	0.15	Tennessee River, Tennessee	2.16
Mean Σ_i	0.40		Mean q_i	0.17		

There is not a great deal of experimental data available on bioaccumulation of strontium in the soft tissue of molluscs. A recommended bioaccumulation factor for strontium in the soft tissue of molluscs was made on the basis of the limited experimental data that are available (Brungs, 1965; Harvey, 1969; Merlini et al., 1967; Nelson, unpublished data; Ophel, 1963). The recommended strontium bioaccumulation factor for the soft tissue of molluscs is 3×10^2 . It must be noted however that this value is given for bodies of water with calcium concentrations of approximately 20 to 30 ppm. The bioaccumulation factor is of course a function of the calcium and strontium concentration in the particular body of water and bioaccumulation factors as high as 720 were found for waters of very low calcium content (Ophel, 1963).

References for Section 3.2

The numbers in the left-hand margin correspond to the reference numbers used in the tables of section 3.2.

- 2.1 Agnedal, P. O., Calcium and strontium in Swedish waters and fish and accumulation of strontium-90, IN Radioecological concentration process, B. Aberg and F. P. Hungate (eds.), pp. 879-896 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1966.
- 2.2 Austin, J. H., Strontium uptake by algae, IN Transport of radionuclides in freshwater systems, B. H. Kornegay, W. A. Vaughan, D. K. Jamison, and J. M. Morgan (eds.), pp. 231-245, TID-7664 (1963).
- 2.3 Beninson, D. E., E. van der Elst, and D. Cancio, Biological aspects in the disposal of fission products into surface waters, IN Disposal of radioactive wastes into seas, oceans, and surface waters, pp. 337-354, Proceedings of International Atomic Energy Agency Symposium, Vienna, 1966, STI/PUB/126.
- 2.4 Bortoli, M. de, P. Gaglione, A. Malvicini, and C. Polvani, Concentration factors for strontium and caesium in fish of the lakes in the region of Varese (Northern Italy), Minerva Fisiconucleare 324-331 (1967).
- 2.5 Bortoli, M. de, P. Gaglione, A. Malvicini, and E. van der Stricht, Environmental Radioactivity Report, EUR-3554e, Ispra (1966).
- 2.6 Brungs, W. A., Experimental uptake of strontium-85 by freshwater organisms, *Health Phys.* 11, 41-46 (1965).
- 2.7 Friend, A. G., The aqueous behavior of strontium-85, cesium-137, zinc-65, and cobalt-60 as determined by laboratory type studies, IN Transport of radionuclides in freshwater systems, B. H. Kornegay, W. A. Vaughan, D. K. Jamison, and J. M. Morgan (eds.), pp. 43-60, TID-7664 (1963).
- 2.8 Gardner, K., and O. Skulberg, An experimental investigation on the accumulation of radioisotopes by freshwater biota, *Arch. Hydrobiol.* 62, 50-69 (1966).
- 2.9 Harvey, R. S., Uptake of radionuclides by freshwater algae and fish, *Health Phys.* 10, 243-247 (1964).

References for Section 3.2 (continued)

- 2.10 Harvey, R. S., Effects of temperature on the sorption of radionuclides by a blue-green algae, IN Symposium on Radioecology, D. J. Nelson and F. C. Evans (eds.), pp. 266-269 (1969), Proceedings of the Second National Symposium on Radioecology, Ann Arbor, Michigan, May 15-17, 1967.
- 2.11 Harvey, R. S., Uptake and loss of radionuclides by the freshwater clam Lampsilis radiata (Gmel.), Health Phys. 17, 149-154 (1969).
- 2.12 Harvey, R. S., Temperature effects on the sorption of radionuclides by freshwater algae, Health Phys. 19, 293-297 (1970).
- 2.13 Kevern, N. R., Strontium and calcium uptake by the green alga Oocystis eremosphaeria, Science 145, 1445-1446 (1964).
- 2.14 Kevern, N. R., N. A. Griffith, G. M. Rosenthal, Jr., and D. J. Nelson, Uptake and accumulation of strontium and calcium by algae and snails, Health Physics Division Annual Progress Report for the Period Ending July 31, 1964, Radiation Ecology Section, pp. 96-101, ORNL-3697 (1964a).
- 2.15 Merlini, M., F. Girardi, and G. Pozzi, Activation analysis in studies of an aquatic ecosystem, IN Nuclear activation techniques in the life sciences, pp. 615-629 (1967), Proceedings of the Symposium held by the IAEA in Amsterdam, May 8-12, 1967, also STI/PUB/155, SM-91/7.
- 2.16 Nelson, D. J., The strontium and calcium relationships in Clinch and Tennessee River molluscs, IN Radioecology, V. Schultz and A. W. Klement, Jr. (eds.), pp. 203-211 (1963), Proceedings of the First National Symposium on Radioecology, Fort Collins, Colorado, September 10-15, 1961.
- 2.17 Nelson, D. J., The prediction of strontium-90 uptake in fish using data on specific activities and biological half-lives, IN Radioecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 843-858 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1966.
- 2.18 Nelson, D. J., Ecological behavior of radionuclides in the Clinch River and Tennessee River, IN Proceedings of Reservoir Fishery Resources Symposium, pp. 169-187, Athens, Georgia, April 5-7, 1967.

References for Section 3.2 (continued)

- 2.19 Ophel, I. L., The fate of radiostrontium in a freshwater community, IN Radioecology, V. Schultz and A. W. Klements, Jr. (eds.), pp. 213-216, Reinhold, New York (1963).
- 2.20 Ophel, I. L., C. D. Fraser, and J. M. Judd, Strontium concentration factors in biota and bottom sediments of a freshwater lake, EUR-4800 (May 1972).
- 2.21 Ophel, I. L., and J. M. Judd, Strontium-calcium relationships in aquatic food chains, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 221-225 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.
- 2.22 Peterson, H. T., Jr., Stable element and radionuclide re-concentration in aquatic ecosystems. I. Stable element effects in the equilibrium situation, Fifth Annual Health Physics Society Midyear Topical Symposium, Idaho Falls, Idaho, November 3-6, 1970.
- 2.23 Reed, J. R., and D. J. Nelson, Radiostrontium uptake in blood and flesh in bluegills (Lepomis macrochirus), IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 226-233 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.
- 2.24 Struxness, E. G., P. H. Carrigan, M. A. Churchill, K. E. Cowser, R. J. Morton, D. J. Nelson, and F. L. Parker, Comprehensive report of the Clinch River Study, ORNL-4035 (April 1967).
- 2.25 Suzuki, Y., R. Nakamura, and T. Ueda, Accumulation of strontium and calcium in freshwater fishes of Japan, Japan Radiat. Res. Soc. 13, 199-207 (1972).
- 2.26 Templeton, W. L., and V. M. Brown, The relationship between concentrations of calcium, strontium and strontium-90 in wild brown trout, Salmo trutta L., and the concentrations of the stable elements in some waters of the United Kingdom, and the implications in radiological health studies, Int. J. Air Water Poll. 8, 49-75 (1964).
- 2.27 Timofeeva, N. A., and N. V. Kulikov, The role of freshwater plants in the accumulation of ⁹⁰Sr and its distribution over the components of a reservoir, IN Radioecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 835-841 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1966.

References for Section 3.2 (continued)

2.28 Williams, L. G., Concentration of strontium-85 and cesium-137 from water solutions by Cladophora and Pithophora, *J. Phycol.* 6, 314-316 (1970).

3.3 TRITIUM

Tritium was included in this report because of a concern that the relative kinetics of tritium and protium resulting from tritium's heavier mass ("isotope effect") might lead to a preferential accumulation of tritium over protium. Experiments in aquatic systems indicate that this does not occur and that the bioaccumulation factor for tritium is less than or about equal to the bioaccumulation factor for protium, which has a bioaccumulation factor approximately equal to 1. This section discusses the movement and distribution of protium in aquatic organisms and reviews experiments that compare the behavior of tritium and protium in aquatic systems. In the discussion the term hydrogen is used to refer to both isotopes of the element collectively.

3.3.1 Hydrogen Bioaccumulation

Hydrogen atoms in living organisms can be separated into two major categories or pools. The first pool, tissue-water hydrogen (TWH), is defined as all hydrogen atoms present in water molecules within the organism. The second pool, tissue-bound hydrogen (TBH), is defined as all hydrogen present in organic molecules, such as proteins, fats, and carbohydrates. As an illustration, a fish consists of approximately 75% water and 25% dry tissue. The percentages of water and dry tissue, however, do not accurately reflect the relative sizes of the two hydrogen pools because on a weight basis water contains 11% hydrogen, while dry fish muscle contains 8%. As a result the tissue-water hydrogen constitutes 80% of the total fish hydrogen and the tissue-bound hydrogen constitutes 20%. Because of this lower hydrogen concentration in dry

tissue, the bioaccumulation factor for protium in fish is approximately 0.93. Subsequent discussions focus on tritium and the degree to which its heavier mass may cause it to have a different bioaccumulation factor from that of protium.

3.3.2 Pathways of Hydrogen Entry into Aquatic Organisms

Tritium enters aquatic systems in the form of tritiated water, that is, HTO , where T represents tritium; H, protium; and O, oxygen. Tritiated water behaves like HOH and mixes rapidly with the tissue water of aquatic organisms. From the tissue water pool tritium can enter hydrogen sites in organic tissue. These hydrogens comprise the tissue-bound hydrogen pool which may be further subdivided into two components termed exchangeable (ET) and nonexchangeable (NET), which have different rates of turnover and different modes of uptake and loss (Figure 3.3.1). The exchangeable component consists of those hydrogens bonded to OH , COOH , NH , SH , and ortho and para positions of phenol groups. These hydrogens undergo a relatively rapid chemical exchange reaction with hydrogens of water molecules in the tissue water. This exchange is not dependent on enzymatic reactions and occurs in metabolically inactive tissues, such as wood, as well as metabolically active tissue. The nonexchangeable component consists predominantly of hydrogens bonded to carbon atoms of organic tissue. Movement of hydrogen into and out of this component is dependent on various enzymatic reductions and oxidations of tissue organics. Various metabolic reactions reduce organic molecules by incorporating hydrogen from tissue water into stable carbon-hydrogen bonds in

ORNL-DWG 74-4996R2




Figure 3.3.1 Pathways of hydrogen entry into the hydrogen components of an aquatic animal.

intermediary products used to synthesize lipids and proteins. Hydrogen is liberated from this nonexchangeable pool by oxidation reactions. These oxidations enzymatically remove nonexchangeable hydrogen from tissue organics and ultimately combine the hydrogens with oxygen to form tissue water. The turnover of this pool is much slower than the exchangeable pool. For a discussion of reactions that incorporate hydrogen from tissue water into the nonexchangeable pool, see Smith and Taylor (1969).

A second pathway for hydrogen entry into the nonexchangeable component, which applies to animals, consists of the ingestion and incorporation of intact food molecules containing nonexchangeable hydrogen. The absence of this pathway in plant uptake and its importance in animals and food chain transfers is discussed in Section 3.3.4.

3.3.3 Compartment Size and Turnover Rates

Tissue-water components constitute from 80 to 93% of aquatic plant weight and averages 75% in aquatic vertebrates. Tritium has been shown to move rapidly from ambient water into tissue water, with bioaccumulation factors approximating unity and biological half-times measured in minutes for small unicellular algae to hours for large aquatic macrophytes and aquatic vertebrates (Elwood, 1973; Harrison and Koranda, 1973; Stewart et al., 1973). Tritium in emergent portions of rooted macrophytes, such as cattails, may fail to reach steady-state with ambient water possibly because of a mixing of water in leaves with less contaminated moisture in the air (Raney and Vaadia, 1965).

Relative sizes of exchangeable and nonexchangeable components have not been determined for aquatic organisms. Data from small mammal experiments, indicating that exchangeable hydrogen constitutes 30% and nonexchangeable hydrogen 70% of the tissue bound component, were used as tissue-bound compartment sizes in constructing Table 3.3.1 and Figure 3.3.1 (Pinson and Langham, 1957; Siri and Evers, 1961).

Data on turnover rates of tritium in the bound components of aquatic organisms are sparse. Elwood (1973) and Patzer (1973), working on goldfish and mosquito fish, respectively, demonstrated 8-day half-times for tritium elimination from the combined tissue-bound component. For aquatic snails, a 62-hr half-time for 70% of the tissue-bound component has been demonstrated (Stewart et al., 1973). A typical study on rats demonstrated half-times of 22 days for 50% of the tissue-bound tritium and 130 days for the other 50% (Thompson and Ballou, 1956). Half-times in humans characteristically show a short tissue-bound component of 30 days and a longer component of 300 days (Pinson and Langham, 1957). In one case Pinson and Langham observed a 2020 day half-time in a chronically exposed human. Although aquatic organisms have demonstrated longer half-times for tissue-bound tritium than for tissue-water tritium, no measurements of elimination rates for exchangeable and nonexchangeable tritium have been reported for aquatic organisms exposed to tritium in their food and water under controlled conditions. Results of experiments cited above indicate some of the tissue-bound components may require a significant fraction of the life of the organism to equilibrate

Table 3.3.1 Component size and source contribution for a hypothetical fish

Components	% Source contribution		% Body hydrogen in each component
	Water	Organic food	
Tissue water			80
Tissue bound			
Exchangeable	100	0	6 ^b
Nonexchangeable	60 ^a	40 ^a	14 ^b

^aData of Patzer (1973).

^bData of Siri and Evers (1961).

with environmental tritium levels. As a result, this nonexchangeable tritium component does not reflect current exposure levels unless long-term steady-state condition has been maintained under constant environmental levels.

3.3.4 Tritium Concentration in Aquatic Food Chains

3.3.4.1 Tissue-Water Tritium

Bioaccumulation factors reported for tissue-water tritium in aquatic organisms range from 0.58 to 1.1 with a mean value less than 1 (Bruner, 1973; Elwood, 1973; Harrison and Koranda, 1973; Stewart et al., 1973; Weinberger, 1953; Weinberger and Porter, 1953). Thus, on the basis of these empirical studies, unity represents a conservative approximation of the tissue-water tritium bioaccumulation factor for aquatic organisms.

3.3.4.2 Tissue-Bound Tritium

Both laboratory and field measurements indicate that the bioaccumulation factor for the total tissue-bound component is less than 1. The tissue-bound tritium concentrations in plants and invertebrates have been summarized from sixteen references by Bruner (1973). In only two of these cases were tissue-bound tritium concentrations higher than ambient water (Cohen and Kneip, 1973; Koranda, 1965). These two measurements were made in areas of pulsed tritium inputs in the vicinity of nuclear power facilities or test sites and probably do not represent steady-state

situations. Other field and laboratory measurements on tissue-bound tritium in plants and vertebrates showed tritium bioaccumulation factors below unity (Elwood, 1973; Harrison and Koranda, 1973; Kanazawa et al., 1972; Porter, 1954; Rosenthal and Stewart, 1973; Stewart et al., 1973; Strand et al., 1973; Weinberger, 1953; Weinberger and Porter, 1953). Most studies did not include measurements of exchangeable tritium and nonexchangeable tritium as separate components. Thus, on the basis of empirical data alone, it is not possible to determine a separate bioaccumulation factor for exchangeable tritium and nonexchangeable tritium, or to determine the degree to which the specific activities of certain organic molecules might vary around the mean value for the total tissue-bound component. For this reason subsequent discussions focus on physical and biochemical aspects of the two tissue-bound components and their effects on the bioaccumulation factor for exchangeable tritium and nonexchangeable tritium, as well as individual molecules.

3.3.4.2.1 Exchangeable Tissue-Bound Tritium

Movement of tritium into and out of the exchangeable component involves chemical exchange reactions as discussed in Section 3.3.2. A variety of organic groups common in living tissue have been studied to determine the steady-state concentration obtained via exchange with tritiated water (Weston, 1973). The results, which are summarized in Table 3.3.2, indicate that tritium is not likely to pool in exchangeable

Table 3.3.2 Steady-state concentration in organic groups common in living tissue

Organic group	α^{-1} ^a	Reference
OH Groups		
CH ₃ COOH	1.0	3.4
CH ₃ OH	0.98	3.9
Cellulose	1.43	3.14
NH Groups		
NH ₃	0.94	3.8
HCONH ₂	0.97	3.1
Ribonuclease	1.0	3.6
DL-polyalanine	1.2	3.7
SH Groups		
H ₂ S	0.30	3.16
CH ₃ SH	0.30	3.11
CH Groups		
HC-CCH ₃	0.78	3.15
CH ₃ SH	0.64	3.30

^a α^{-1} = (Specific Activity organic group)/(Specific Activity water). Since α^{-1} is based on specific activities it is independent of variations in hydrogen content, and must be multiplied by the percent hydrogen in the tissue and divided by the hydrogen content of water (11%) to produce a "dry weight" bio-accumulation factor for the organic group.

components because in most cases the specific activity* of the molecules is less than that of the water.


A food chain is diagrammed in Figure 3.3.2 with organisms compartmentalized as in Figure 3.3.1 and each consumer utilizing the organism below him as the food source. In both Figure 3.3.1 and 3.3.2 there is no pathway to accommodate the food-chain transfer of exchangeable tritium even though this obviously occurs. The effect of food-chain transfer of exchangeable tritium is unimportant because 1) the exchangeable tritium of each organism has ready access to the same ambient tritium concentrations via exchange with tissue water, and 2) most aquatic organisms eat only a small fraction of their body weight per day thus the turnover of exchangeable tritium appears rapid in comparison.

3.3.4.2.2 Nonexchangeable Tissue-Bound Tritium

At the base of the food chain in Figure 3.3.2, photosynthesis, along with other reduction reactions, incorporates tritium from tissue water into the nonexchangeable component of plants. Data on the total tissue-bound component in plants indicate that there is a 2 to 55% discrimination against incorporation of tritium relative to protium (Bruner, 1973; Harrison and Koranda, 1973; Rosenthal and Stewart, 1973; Weinberger, 1953; Weinberger and Porter, 1953). These determinations, however, did not differentiate between the two tissue-bound components.

*Specific activity is used here to mean the ratio of radioactive to stable atoms of an element in a sample.

ORNL-DWG 74-4997R2



TWT = TISSUE WATER TRITIUM

ET = EXCHANGEABLE TRITIUM

NET = NONEXCHANGEABLE TRITIUM

Figure 3.3.2 Three-link food chain diagram with organisms compartmentalized.

Rambeck and Bassham (1973) and Kanazawa et al. (1972) investigated the effects of oxidation and reduction reactions on specific activities of individual organic molecules produced during the Krebs Cycle in algae. After five generations in a closed system the average specific activity of the individual molecules investigated was 30% less than the water to which they were exposed. Only two out of the 14 molecules investigated showed specific activity ratios greater than 1, citrate 1.8 and fumarate 1.2. These results, although limited to 2 species of algae and the Krebs Cycle, indicate that most organic molecules discriminate against tritium incorporation into nonexchangeable sites and that significant increases in the specific activity of certain molecules due to the kinetics of nonexchangeable tritium are not likely. Further studies are needed to verify these results for various series of metabolic reactions in other organisms.

Consumer organisms obtain nonexchangeable tritium by assimilation of intact tritiated food molecules as well as by reduction of tissue organics with tissue-water hydrogens. It can be calculated from Patzer's (1973) data on herbivorous fish that the nonexchangeable component receives approximately 60% of its hydrogen from tissue-water hydrogen and 40% from food. Data are not available on the exact degree to which the specific activity of the nonexchangeable component is affected by food-chain transfers of nonexchangeable tritium in food molecules, or by various oxidations and reductions.

Although empirical data on bioaccumulation factors for nonexchangeable tritium in organisms are wanting, it is possible to establish a maximum bioaccumulation factor for the nonexchangeable component by using the tissue-bound compartment sizes from Table 3.3.1, an average specific activity ratio for the exchangeable component from Table 3.3.2, and bioaccumulation factors determined empirically for the total tissue-bound component in numerous freshwater organisms. Based on Table 3.3.2, 0.7 represents the best estimate of an average bioaccumulation factor for the exchangeable component. Thus, the empirically-derived bioaccumulation factor of less than 1 for the total tissue-bound component, consisting of 30% exchangeable tritium and 70% nonexchangeable tritium, requires that the bioaccumulation factor of the nonexchangeable component be no greater than 1.1.

References for Section 3.3

The numbers in the left-hand margin correspond to the reference numbers used in the tables of section 3.3.

- 3.1 Brokskii, A. E., "The exchange of hydrogen with deuterium in solution," *Trans. Faraday Soc.* 33, 1180 (1937).
- 3.2 Bruner, H. D., *Distribution of tritium between the hydrosphere and invertebrates*, IN *Tritium*, A. A. Moghissi and M. W. Carter (eds.), pp. 303-313 (1973), Messinger Graphics, Phoenix, TID CONF-710809.
- 3.3 Cohen, L. K., and T. J. Kneip, *Environmental tritium studies at a PWR power plant*, IN *Tritium*, A. A. Moghissi and M. W. Carter (eds.), pp. 639-646 (1973), Messinger Graphics, Phoenix, TID CONF-710809.
- 3.4 Eidinoff, M. L., et al., "Equilibrium between protium, deuterium, and tritium in the system: Hydrogen-acetic acid; isotopic fractionation factors in a catalytic hydrogenation," *J. Amer. Chem. Soc.* 74, 5280 (1952).
- 3.5 Elwood, J. W., *Tritium behavior in fish from a chronically contaminated lake*, IN *Radionuclides in ecosystems*, D. J. Nelson (ed.), pp. 435-439 (1973), *Proceedings of the Third National Symposium on Radioecology*, Oak Ridge, Tennessee, May 10-12, 1971.
- 3.6 Englander, S. W., "A hydrogen exchange method using tritium and sephadex: Its application to ribonuclease," *Biochem.* 2, 798 (1963).
- 3.7 Englander, S. W., and A. Paulsen, *Hydrogen-tritium exchange of the random chain polypeptide*, *Biopolymers* 7, 379-393 (1969).
- 3.8 Gutman, J. R., and M. Wolfsberg, "Equilibrium in the exchange of tritium between ammonia and hydrogen and the zero-point energy difference between NH_3 and NH_2T ," *J. Chem. Phys.* 33, 1592 (1960).
- 3.9 Halford, J. O., and E. Pecherer, "Exchange of deuterium between methanol and water. Vibrations of the hydroxyl group in methanol and methanol-d. The entropy of methanol," *J. Chem. Phys.* 6, 571 (1938).
- 3.10 Harrison, F. L., and J. J. Koranda, *Tritiation of aquatic animals in an experimental freshwater pool*, IN *Radionuclides in ecosystems*, D. J. Nelson (ed.), pp. 425-434 (1973), *Proceedings of the Third National Symposium on Radioecology*, Oak Ridge, Tennessee, May 10-12, 1971.

References for Section 3.3 (continued)

- 3.11 Hobden, F. W., et al., "Determination and calculation of the equilibrium constants for isotopic exchange in the systems *n*-amyl alcohol-water and ethylthiol-water. Vapor pressures and raman spectra of *n*-amyl deuterio-alcohol and ethyl deuteriothiol," *J. Chem. Soc.* 61 (1939).
- 3.12 Kanazawa, T., K. Kanazawa, and J. A. Bassham, Tritium incorporation in the metabolism of *Chlorella pyrenoidosa*, *Environ. Sci. Technol.* 6, 638-642 (1972).
- 3.13 Koranda, J. J., Preliminary studies of the persistence of tritium and carbon-14 in the Pacific Proving Grounds, *Health Phys.* 11, 1445-1457 (1965).
- 3.14 Lang, A. R. G., and S. G. Mason, Tritium exchange between cellulose and water: Accessibility measurements and effects of cyclic drying, *Can. J. Chem.* 38, 373-387 (1960).
- 3.15 Mahadevan, E. G., and E. V. Monse, "Equilibrium constant for the deuterium exchange between methylacetylene and water," *J. Chem. Phys.* 43, 2556 (1965).
- 3.16 Marx, D. (Summarized by), "Etude des Constantes des Equilibres Isotopiques du Deuterium entre l'Eau et les Hydrures des Metalloides de la Deuxieme Famille," *C.E.A. Report 1382*, Saclay, France, 1960.
- 3.17 Patzer, R. G., A. A. Moghissi, and D. N. McNelis, Accumulation of tritium in various species of fish reared in tritiated water, IN *Proceedings of IAEA/NEA/WHO Symposium on Environmental Behavior of Radionuclides Released in the Nuclear Industry*, Aix-en-Provence, France, May 14-18, 1973, IAEA/SM-172/29, Vienna, 1973.
- 3.18 Pinson, E. A., and W. H. Langham, Physiology and toxicology of tritium in man, *J. Appl. Physiol.* 10, 108-126 (1957).
- 3.19 Porter, J. W., Incorporation of hydrogen isotopes into various fractions of rapidly growing algae, IN *Biology Research Annual Report 1953*, pp. 77-78, Hanford Atomic Products Operation, Richland, Washington, HW-30437 (January 1954).
- 3.20 Rambeck, W. A., and J. A. Bassham, Tritium incorporation and retention in photosynthesizing algae, *Biochem. Biophys. Acta* 304, 725-735 (1973).

References for Section 3.3 (continued)

- 3.21 Raney, F., and Y. Vaadia, Movement and distribution of THO in tissue water and vapor transported by shoots of Helianthus and Nicotiana, *Plant Phys.* 40, 383-388 (1965).
- 3.22 Rosenthal, G. M., Jr., and M. L. Stewart, Tritium incorporation in algae and transfer in simple aquatic food chains, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 440-444 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.
- 3.23 Siri, W., and J. Evers, Tritium exchange in biological systems, IN IAEA Proceedings of the Symposium on the Detection and Use of Tritium in the Physical and Biological Sciences, Vol. 2, pp. 71-84 (1961).
- 3.24 Smith, T. E., and R. T. Taylor, Incorporation of tritium from tritiated water into carbohydrates, lipids, and nucleic acids, USAEC Report UCRL-50781 (November 26, 1969).
- 3.25 Stewart, M. L., G. M. Rosenthal, and J. R. Kline, Tritium discrimination and concentration in freshwater microcosms, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 452-459 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 11-12, 1971.
- 3.26 Strand, J. A., W. L. Templeton, and E. G. Tangen, Accumulation and retention of tritium (tritiated water) in embryonic and larval fish, and radiation effect, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 445-451 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.
- 3.27 Thompson, R. C., and J. E. Ballou, Studies of metabolic turnover with tritium as a tracer. V. The predominantly non-dynamic state of body constituents in the rat, *J. Biol. Chem.* 223, 795-809 (1956).
- 3.28 Weinberger, D., Metabolism of hydrogen isotopes by algae, IN Biology Research, Annual Report 1952, pp. 72-74, Hanford Atomic Products Operation, Richland, Washington, HW-28636 (1953).
- 3.29 Weinberger, D., and J. W. Porter, Incorporation of tritium oxide into growing Chlorella pyrenoidosa cells, *Science* 117, 636-638 (1953).
- 3.30 Weston, R. E., Jr., Kinetic and equilibrium isotope effects of tritium substitution, IN Tritium, A. A. Moghissi and M. W. Carter (eds.), pp. 289-300 (1973), Messinger Graphics, Phoenix, TID-CONF-710809.

3.4 IODINE

3.4.1 Environmental Iodine

The stable iodine content of freshwater ranges from 0.4 to 10.0 ppb depending upon the water source and the environment through which the water travels. Streams originating from glaciers or runoff are low in iodine, less than 1 ppb, while those originating from springs and ground water sources are higher, up to 5 ppb. Livingston (1963) calculated a median river water concentration of 2 ppb. Table 3.4.1 lists some representative stable iodine concentrations from various freshwater, marine, and geologic sources. Iodine in oceans is approximately 50 ppb and apparently originates from the erosion of land masses (Bowen, 1966). The dominant physicochemical form of iodine and its relative biological availability in freshwater is uncertain (Winchester, 1970).

3.4.2 Iodine Metabolism

The thyroid removes inorganic iodine from the blood and uses it in the production of thyroxine, a hormone necessary for maintaining a variety of metabolic and growth functions. The thyroid releases thyroxine to the blood stream where it combines with large proteins to form protein-bound iodine. Thyroxine is split from the large protein and enters the body tissues where it is degraded. Inorganic iodine produced by the degradation of thyroxine re-enters the blood and is eliminated by urinary excretion. Some evidence indicates that an unspecified portion of this may be reabsorbed by the thyroid (Chavin and Bouwman, 1965).

Table 3.4.1 Iodine concentrations in surface waters, the ocean, and geologic sources

Water Sample	Average Concentration (ppb)	Range (ppb)	Reference
German lake	3.5	2.5 - 4.9	4.18
Canadian lake	1.54	0.49 - 2.9	4.21
Lake Michigan	1.0	----	4.10
Lake Michigan	5.0	----	4.39
Michigan streams	1 or less	----	4.39
Missouri spring	4.7 or less	----	4.12
New Jersey spring	5.2 or less	----	4.12
Swiss rivers	0.56	0.4 - 1.3	4.15
Panama rivers	1 or less	1.0 - 0.6	4.14
River Average	2.0	< 1.0 - 10	4.28
Oceans	50	----	4.47
Soils	----	600 - 6000	4.47
Rock			
Igneous	500	----	4.5
Limestone	1200	----	4.5
Shale	2200	----	4.5
Coal	6000	----	4.5

Although there are numerous studies on the physiological role of iodine, few of these investigations provide data on elimination rates and turnover times for iodine in aquatic organisms. The few studies that were found are summarized in Table 3.4.2. The rapid component (C-1) is taken to represent plasma clearance through urinary excretion, while the slower component (C-2) is attributed to the thyroid.

3.4.3 Review of Iodine Bioaccumulation Factors

3.4.3.1 Stable Iodine Bioaccumulation Factors for Fish Tissue

Only the thyroid tissue and ovaries (eggs) show great iodine bioaccumulation. No studies were found that provided data allowing direct calculation of bioaccumulation factors for the thyroid tissue because thyroid stable iodine measurements for freshwater fish were not expressed as concentrations. Data were available on thyroid iodine concentrations in some marine species and these marine data are cited to corroborate our indirectly estimated freshwater thyroid bioaccumulation factors. Muscle bioaccumulation factors of freshwater and marine fish are very similar despite the much higher iodine content of oceans (Tables 3.4.3 and 3.4.4). Organic and inorganic iodine show the same percent distribution in the serum of estuarine flounder and freshwater whitefish (Hickman, 1962). The similar iodine bioaccumulation factors for muscle and the similar iodine metabolism in fish from freshwater and marine habitats provide some justification for comparing bioaccumulation factors of marine and freshwater species.

References on stable iodine in fish ovaries did not report comparable iodine water concentrations, thus, ovary bioaccumulation factors

Table 3.4.2 Elimination rates for iodine-131 in aquatic vertebrates

Species	Biological Half-time for Iodine (days)	Administered Activity (%)	Elimination Coefficient (day ⁻¹)	Reference
<u>Salmo gairdneri</u>				
C-1	2.0	----	.347	4.13
C-2	8.6	----	.086	4.13
<u>Salmo gairdneri</u>				
C-1	1.7	----	.408	4.22,4.50
C-2	---	----	----	----
<u>Cyprinus carpio</u>				
C-1	---	----	----	----
C-2	14.0	----	.050	4.36
<u>Carassius auratus</u>				
C-1	0.8	----	.866	4.36
C-2	14.0	----	.050	4.36
<u>Perch (sp.)</u>				
C-1	1.5	----	.460	4.42
C-2	---	----	----	----
<u>Micropogon undulatus</u>				
C-1	0.25	59	2.770	4.2
C-2	2.25	40	.310	4.2
C-3	24.0	1	.028	4.2
<u>Rana pipiens</u>				
C-1	---	----	----	----
C-2	56.0	----	.012	4.50
<u>Hyla versicolor</u>				
C-1	5.0	----	.140	4.25
C-2	---	----	----	----
<u>Taricha granulosa</u>				
C-1	2.0	74	.350	4.50
C-2	210.0	26	.003	4.50

Table 3.4.3 Stable iodine in freshwater fish muscle

Species	Muscle Concentration ($\mu\text{g}/\text{kg}$)	Water Concentration ($\mu\text{g}/\text{liter}$)	Stable Iodine BF ^a	Location	Reference
<i>Salmo fario</i>	36	----	36	Switzerland	4.16 ^b
<i>Salmo fario</i>	48	----	48	Germany	4.4 ^b
<i>Salmo fario</i>	24	----	24	New Zealand	4.20 ^b
<i>Salmo fario</i>	50	----	50	New Zealand	4.20 ^b
<i>Salvelinus namaycush</i>	10	----	10	Lake Erie	4.45 ^b
<i>Luciopimelodus pati</i>	30	----	30		4.32 ^b
<i>Pimelodus albidus</i>	40	----	40		4.32 ^b
<i>Lepomis incisor</i>	40	----	40	Mississippi River	4.45 ^b
<i>Micropterus salmoides</i>	50	----	50	Potomac River	4.45 ^b
<i>Pomoxis annularis</i>	10	----	10	Mississippi River	4.45 ^b
<i>Perca flavescens</i>	20	----	20	Potomac River	4.45 ^b
<i>Perca fluviatilis</i>	20	----	20	Switzerland	4.16 ^b
Freshwater species in general	40	----	40		4.33 ^b
Carp	17	----	17		4.6
<i>Micropterus salmoides</i>	30	----	30		4.6
<i>Salvelinus namaycush</i>	31	----	31		4.6
River perch	40	----	40		4.6
<i>Salmo gairdneri</i>	15	1	15	Black River, Michigan	4.39
<i>Salmo gairdneri</i>	40	----	40	Pacific Coast Stream	4.26
<i>Oncorhynchus kisutch</i>	120	1	120	Lake Michigan	4.11 ^c
<i>Perca flavescens</i>	120	1	120	Lake Michigan	4.11 ^c
<i>Salvelinus namaycush</i>	170	1	170	Lake Michigan	4.11 ^c
<i>Salmo trutta</i>	110	1	110	Lake Michigan	4.11 ^c
<i>Coregonus clupeaformis</i>	140	1	140	Lake Michigan	4.11 ^c
Mean	50 \pm 45				

^aTissue concentration reported without comparable water values was divided by 1 $\mu\text{g}/\text{liter}$ to approximate a conservative bioaccumulation factor.

^bCited in Vinogradov (1953).

^cTissue concentrations represent averages calculated from Copeland's raw data that clearly specified that the tissue analyzed was muscle.

Table 3.4.4 Stable iodine in anadromous and marine fish muscle

Species	Muscle Concentration (μ g/100 g)	Water Concentration (μ g/liter)	Stable Iodine BF	Remarks	Reference
Sea trout	320	-----		Anadromous	4.6
<u>Salmo</u> <u>salar</u>	341	-----		Anadromous	4.6
<u>Salmo</u> <u>gairdneri</u> (sea run)	380	-----		Anadromous	4.26
<u>Oncorhynchus</u>	296	-----	53	Anadromous	4.26
Mackerel	371	-----	74	Marine	4.6
Herring	520	-----	104	Marine	4.6
<u>Roccus</u> <u>americanus</u>	742	-----	148	Marine	4.6
Cod	1463	-----	293	Marine	4.6
<u>Hippoglossus</u>	520	-----	104	Marine	4.6
<u>Melanogrammus</u> <u>aeglefinus</u>	318	-----	64	Marine	4.6
<u>Clupea</u> <u>harengus</u>	600	-----	120	Marine	4.31a
<u>Melanogrammus</u> <u>aeglefinus</u>	513	-----	103	Marine	4.34a
Mean BF				118 \pm 72	

^aReference cited in Vinogradov (1953).

could not be directly calculated from the studies. However, we were able to indirectly estimate an average ovary bioaccumulation factor for stable iodine from studies which reported both ovary and muscle iodine contents in the fish collected. From these data an ovary:muscle ratio was calculated and multiplied times the average muscle bioaccumulation factor to estimate an average ovary bioaccumulation factor for stable iodine.

Data on stable iodine in fish muscle were adequate to determine a reliable bioaccumulation factor. Due to the limited data on iodine in fish thyroid and ovaries, and the manner in which these data were reported, our estimates for thyroid and ovary bioaccumulation factors represent approximations.

3.4.3.1.1 Stable Iodine Bioaccumulation Factors for Fish Muscle

Stable iodine bioaccumulation factors for fish muscle from various species of freshwater fish are listed in Table 3.4.3. Bioaccumulation factors from a variety of investigators varied from 10 to 50 while determinations by Copeland et al. (1973) ranged from 120 to 170. By using a 5 ppb Lake Michigan iodine concentration, as reported by Robertson and Chaney (1953), with Copeland's tissue iodine concentrations, a range of bioaccumulation factors can be calculated for Copeland's samples that agrees well with ranges reported by other investigators. However, Winchester (1970), reported 1 ppb or less as the iodine concentration in Lake Michigan, thus, indicating that Copeland's water concentrations are correct and that higher tissue iodine concentration and higher bioaccumulation factors for Lake Michigan fish cannot be explained by an underestimate of iodine concentration in water. At this time it is the opinion of the authors that although Copeland's

data are higher than the others, we must accept their bioaccumulation factors of 120 to 170 for stable iodine in freshwater fish muscle until other data clarify the issue. From data in Table 3.4.3 we calculate an average bioaccumulation factor of 52 for stable iodine in fish muscle with a range from 10 to 170.

Based on clear decreases in iodine tissue concentrations from marine to anadromous to freshwater species, it does not appear that homeostatic control is a factor in controlling muscle iodine levels in fish (Table 3.4.4). Anadromous fish sampled during their spawning run show stable iodine tissue concentrations higher than strictly freshwater species (Table 3.4.4). This is evidently due to their recent exposure to higher iodine concentrations in oceans. Fontain et al. (1948), found 1280 $\mu\text{g/g}$ of iodine in the serum of Atlantic salmon at the beginning of their migration spawn. At the end of migration, serum iodine had dropped to 280 $\mu\text{g/g}$. It was not determined if iodine in other tissues showed a similar decline. The drop in serum iodine could be due to lower freshwater iodine concentrations and a lack of homeostatic control, or to a combination of physiological factors.

3.4.3.1.2 Stable Iodine Bioaccumulation Factors for Fish Thyroids and Ovaries

In rainbow trout and most teleosts the thyroid is not assembled into a single gland but, instead, is scattered throughout the lower jaw. In some cases thyroid tissue may occur in the eyes and kidneys as well (Baker et al., 1955). The removal of diffuse thyroidal tissue without portions of nonthyroidal tissue is difficult. As a result,

iodine concentrations are either not expressed on a unit weight basis or qualified to the extent that the tissue may contain nonthyroid material (Short et al., 1969).

Robertson and Chaney (1953) analyzed sections of carefully dissected Lake Michigan rainbow trout tissue excised from the floor of the mouth from the first to the fourth gill arch. The iodine content was expressed as total micrograms of iodine in the total "thyroid mass" (Table 3.4.5.). The weights of these samples were not reported. No determinations were made of the stable thyroidal iodine content, expressed per unit of exclusively thyroid tissue of freshwater fish. Using 0.003 g/100 g of fish as an approximation of the mass of the rainbow trout thyroid (Spector, 1956), we calculated an approximate thyroid iodine concentration of 269 $\mu\text{g/g}$ by using the average weights of the trout used in Robertson and Chaney's (1953) study. Dividing this thyroid concentration by Copeland's Lake Michigan iodine concentration of 1 ppb gives an estimate of 270,000 for the trout thyroid bioaccumulation factor. To provide some verification of this thyroid bioaccumulation factor, an average thyroid bioaccumulation factor for marine fish of 770,000 was calculated using data compiled from Vinogradov (1953) in Table 3.4.6. Considering the approximate nature of our estimates, these two values are fairly close and tend to support the freshwater thyroid bioaccumulation factor.

Hickman (1962) measured inorganic iodine in the serum and thyroids of estuarine flounder and found that the serum levels of iodine increased on a seasonal cycle as the salinity of the water increased. However, in the case of thyroidal iodine, only inorganic iodine showed a clear

Table 3.4.5 Total iodine present in the thyroid and eggs of five rainbow and California steelhead trout (in micrograms)^a

Rainbow Trout		Steelhead Trout ^b	
Thyroid	Eggs	Thyroid	Eggs
3.5	38.6	367	1170
0.5	18.6	755	2323
12.3	41.3	377	1814
11.9	70.0	---	1749
16.0	55.0	---	1065
10.0 ^c	44.0 ^c	500 ^c	1624 ^c

^aRobertson and Chaney (1953).

^bSteelhead trout are anadromous rainbow trout (Salmo gardneri). They were secured from the mouths of California rivers up to 150 miles upstream.

^cMean values.

Table 3.4.6 Thyroid stable iodine bioaccumulation factors for marine fish

Species	Thyroid Iodine (ppm)	Thyroid BFA	Reference ^b
<u>Gadus aeglefinus</u>	2,800	56,000	4.9,4.30
<u>Gadus morrhua</u>	11,600	232,000	4.9,4.31
<u>Salmo eriox</u>	173,600	3,500,000	4.30,4.31
<u>Salmo salar</u>	9,600	192,000	4.9,4.31
<u>Scomber scombrus</u>	32,200	644,000	4.29,4.30
<u>Sebastes marinus</u>	46,200	924,000	4.29,4.31
<u>Pleuronectes platessa</u>	7,800	156,000	4.9,4.31
<u>Molva vulgaris</u>	23,800	476,000	4.9
Mean		772,500	

^aCalculated using 50 ppb as marine water iodine concentration (Table 3.4.1).

^bCited in Vinogradov (1953).

response to environmental iodine fluctuations. The protein-bound iodine, which constitutes 95% of thyroidal iodine, varied much less and with no apparent pattern. Protein-bound iodine was, in fact, highest in September when environmental iodine was at a low point. Thyroid iodine concentration appears to fluctuate little in response to environmental iodine. Whether this is due to homeostatic control, slower turnover rates, or other factors is uncertain.

From Table 3.4.5 it is apparent that the ovaries (eggs) contain most of the iodine present in female rainbow trout. Due to the manner in which the data were reported and to uncertainties concerning analytical procedures, we calculated an ovary:muscle concentration ratio for stable iodine (Table 3.4.7) and then multiplied this ratio by the average muscle bioaccumulation factor from Table 3.4.3 to estimate a stable iodine bioaccumulation factor of 800 for ovaries.

3.4.3.2 Iodine-131 Bioaccumulation Factors for Fish Tissue

3.4.3.2.1 Iodine-131 Bioaccumulation Factors for Fish Muscle

The few experiments reporting iodine-131 levels in fish muscle are listed in Table 3.4.8. Experiments by Short et al. (1969) indicate that the food web is the principal mode of uptake. One study by Hunn and Reineke (1964) indicates that iodine can be accumulated directly from the water. In Short's study, an aquarium and a lake were both tagged with iodine-131. After 29 days fish in the lake showed a muscle bioaccumulation factor of 100, while those in the aquaria, without their natural food sources, showed a bioaccumulation factor of 1.7. It is possible, although experimental procedures were

Table 3.4.7 Stable iodine bioaccumulation factor for fish ovaries (eggs) derived from relative concentrations of stable iodine in muscle and ovaries

Species	Ovary/muscle Ratio	Ovary ^a BF	Reference
<u>Salmo gardneri</u>	15	750	4.39
<u>Salmo gardneri</u>	14	700	4.39
<u>Salmo fario</u>	29	1450	4.16
<u>Perca fluviatalis</u>	6	300	4.16
Mean Ovary BF		800	

^a Ovary bioaccumulation factor = Ovary/muscle ratio times average muscle bioaccumulation factor from Table 3.4.3.

Table 3.4.8 Iodine-131 bioaccumulation factors for muscle in freshwater animals

Species	Iodine-131 BF	Location	Reference
Carp <u><i>Cyprinus</i></u> <u><i>carassius</i></u> (whole fish)	25	Lake	4.27
<u><i>Prochilodus</i></u>	0.5	Aquaria	4.3
<u><i>Pimelodus</i></u>	0.8	Aquaria	4.3
<u><i>Cichlasoma</i></u>	0.5	Aquaria	4.3
Miscellaneous aquatic 17 species	20 (mean)	Aquaria	4.43
leech gastropods crustaceans insect larvae carp tadpoles			
Carp <u><i>Cyprinus</i></u> <u><i>carpio</i></u>	12	Aquaria	4.44
<u><i>Salmo gairdneri</i></u>	100	Lake	4.40
Mean Fish Muscle BF	39 ^a		

^aSee Sec. 3.4.3.2.1.

not described, that the low bioaccumulation factors reported by Beninson (1966) for fish in aquaria studies were due to the absence of a contaminated food source. The low bioaccumulation factors do not appear reliable in view of Short's study and the much higher stable iodine bioaccumulation factors reported in Table 3.4.8. An average iodine-131 bioaccumulation factor of 39 for fish muscle is calculated for fish muscle from Table 3.4.8 (Beninson's data omitted).

Because of the 8-day radioactive half-life of iodine-131, the turnover rate of iodine in a specific tissue and the length of the primary contamination pathway control the degree to which bioaccumulation factors for iodine-131 are less than bioaccumulation factors for stable iodine. Iodine-131 absorbed directly from the water will have less time to decay than iodine-131 obtained from food. Section 2.4 and Equation (2.4.9) describe a method for calculating bioaccumulation factors for radioactive isotopes based on their stable element bioaccumulation factors. Using Equation (2.4.9), the average stable element bioaccumulation factor for fish muscle (Table 3.4.3), and an average fast component elimination coefficient of 0.43 (Table 3.4.2), we calculate an iodine-131 bioaccumulation factor of 43. Based on the similarity between this calculated bioaccumulation factor and the average bioaccumulation factor determined from empirical studies, we recommend an average bioaccumulation factor of 40 for iodine-131 in fish muscle.

3.4.3.2.2 Iodine-131 Bioaccumulation Factors for Fish Thyroids

Short et al. (1969) removed the lower terminus of the gill arch, including the dorsal aorta and surrounding muscle, from lake trout

removed from an iodine-131 tagged lake after 29 days of exposure. An iodine-131 bioaccumulation factor of 12,000 was calculated for these "thyroid" samples which contained unspecified amounts of other tissue. This value appears low when compared to the thyroid bioaccumulation factor of 270,000 for stable iodine (Sec. 3.4.3.1.2). Since no freshwater iodine-131 bioaccumulation factors have been determined for thyroids free from attached nonthyroidal tissue, we recommend an iodine-131 thyroid bioaccumulation factor of 110,000. An 11-day elimination coefficient for the long component in trout (Table 3.4.2) and an average stable iodine bioaccumulation factor for thyroid tissue of 270,000 were used in calculating the recommended thyroid bioaccumulation factor. Further research is needed to clarify this issue.

3.4.3.3 Stable Iodine Bioaccumulation Factors for Plants

The only stable iodine determination found for freshwater plants was an average bioaccumulation factor of 800, calculated by Copeland and Ayers (1970) for samples of phytoplankton (green and bluegreen algae) taken from Lake Michigan.

3.4.3.4 Iodine-131 Bioaccumulation Factors for Plants

Bioaccumulation factors for rooted and floating macrophytes are combined in Table 3.4.9 since they are within the same range. Iodine-131 bioaccumulation factors for freshwater algae from one study averaged 255 (Short et al., 1969).

3.4.3.5 Stable Iodine Bioaccumulation Factors for Invertebrates

Table 3.4.10 lists stable iodine bioaccumulation factors for a small number of aquatic invertebrates. In insects, and probably

Table 3.4.9 Iodine-131 bioaccumulation factors for freshwater plants

Species	Iodine-131 BF	Remarks	Reference
<u>Macrophytes</u>			
<u>Bacopa</u>	130	Tank	4.3
<u>Cabomba</u>	178	Tank	4.3
<u>Elodea</u>	145	Tank	4.3
<u>Ceratophyllum</u>	178	Tank	4.3
<u>Echinodorus</u>	162	Tank	4.3
<u>Ceratophyllum</u>	95	Tank	4.44
<u>Myriophyllum</u>	100	Tank	4.44
<u>Utricularia</u>	209	Tank	4.44
<u>Lemna minor</u>	71	Tank	4.44
<u>L. trisulca</u>	154	Tank	4.44
<u>Potamogeton</u>	26	Tank	4.44
<u>Elodea canadensis</u>	134	Tank	4.44
<u>Stratiotes</u>	60	Tank	4.44
<u>Hydrocharis</u>	165	Tank	4.44
<u>Carex</u>	101	Tank	4.44
<u>Sphagnum</u>	90	Lake	4.27
<u>Nuphar</u>	60	Lake	4.27
Mean of 32 Plant Species	93	Tank	4.44
Mean Macrophyte BF	119		
<u>Algae</u>			
<u>Nitella</u>	380	Lake	4.40
<u>Spirogyra and</u> <u>Oedogonium</u>	130	Lake	4.40
Mean Algae BF	255		

Table 3.4.10 Stable iodine bioaccumulation factors for invertebrates

Organism	Concentration in Tissue (ppb)	Concentration in Water (ppb)	Stable Iodine BF	Tissue	Reference
<u>Crustacea</u>					
Shrimp (freshwater)	60 ^a	0.6 ^b	10	---	4.14
Snail	150	0.6 ^b	250	---	4.14
Shrimp (<i>Mysis</i>)	565 ^c	1	565	---	4.10
Amphipod (<i>Pontoporeia</i>)	390 ^c	1	390	---	4.10
Benthos (mostly crustacean)	500	1	500	---	4.10
<u>Mollusca</u>					
Clam (<i>Dreissensia</i>)	----	----	1000	Shell	4.49
Mean Crustacean BF				343	

^aA factor of 10 was used to convert dry weight values to wet weight values reported here.

^bWater concentrations represent drinking water measurements of a local Panama city.

^cCalculated from Copeland's (1970) raw data using samples of benthos said to consist of 100% *Mysis* or *Pontoporeia*.

crustacea as well, most of the iodine appears to be concentrated in the integument. Iodine-131 studies indicate that little iodine is lost from insects except when the integument is discarded (Odum and Golley, 1963; Van Hook and Crossley, 1969). The average bioaccumulation factor for crustacea, the only group of invertebrates for which stable iodine values of soft tissues or whole animals were found, is 343.

3.4.3.6 Iodine-131 Bioaccumulation Factors for Invertebrates

Table 3.4.11 lists iodine-131 bioaccumulation factors for various invertebrates. An average bioaccumulation factor value is recorded for aquatic insects and molluscs. The difference in the lake bioaccumulation factor of 140 and the aquaria bioaccumulation factor of 600 reported for the amphipod is attributed to incomplete mixing of lake waters (Short et al., 1969).

Table 3.4.11 Iodine-131 bioaccumulation factors for invertebrates

Organism	Iodine-131 BF	Tissue	Remarks	Reference
Crustacea:				
<u>Crayfish</u> (<i>Pacifastacus</i>)	10	Muscle	Lake	4.40
	62	Soft parts	Lake	4.40
	240	Carapace	Lake	4.40
<u>Amphipod</u>	600	Whole	Tank	4.40
	140	Whole	Lake	4.40
<u>Aegla</u>			Tank	4.3
<u>Littoral plankton</u>	530		Lake	4.40
<u>Limnetic plankton</u>	500		Lake	4.40
Aquatic Insects:				
<u>Dragon fly larvae</u> (<i>Leucorrhinia</i>)	151	?	Tank	4.48
<u>May fly larvae</u> (<i>Cloeon</i>)	690	?	Tank	4.48
<u>Caddis fly larvae</u> (<i>Glyphotaelius</i>)	163	?	Tank	4.48
<u>Drone fly larvae</u> (<i>Eristalis</i>)	600	?	Tank	4.48
Mean BF for Aquatic Insects	400			
Worm:				
<u>Herpobdella</u>	10	?		4.38, 4.44
Sponge:				
<u>Spongilla</u>	200	?	Lake	4.27
Molluscs:				
<u>Diplodon</u>	11	?	Tank	4.3
<u>Ampullaria</u>	23	?	Tank	4.3
<u>Planorbis</u>	134	?	Tank	4.3
<u>Margaritifera</u>	10	Muscle	Tank	4.40
	53	Soft parts	Lake	4.40
<u>Limnaea</u>	23	?	Tank	4.38, 4.44
<u>Radix</u>	14	?	Tank	4.38, 4.44
<u>Anisus</u>	53	?	Tank	4.38, 4.44
<u>Planorbis</u>	70	?	Tank	4.38, 4.44
<u>Dreissensia</u>	140	Gills	Tank	4.19
	70	Mantle	Tank	4.19
	40	Viscera	Tank	4.19
	20	Byssus	Tank	4.19
	400	Shell	Tank	4.19
<u>Pond snail</u>	20	?	Tank	4.43
Mean BF for Molluscs	50			

References for Section 3.4

The numbers in the left-hand margin correspond to the reference numbers used in the tables of section 3.4.

- 4.1 Baker, K. F., O. Berg, A. Gorbman, R. F. Nigrelli, and M. Gordon, Functional thyroid tumors in the kidneys of platyfish, *Cancer Res.* 15, 118-123 (1955).
- 4.2 Baptist, J. P., D. E. Hoss, and C. W. Lewis, Retention of Cr-51, Fe-59, Co-60, Zn-65, Sr-85, Nb-95, In-114, and I-131 by the Atlantic croaker (Micropogon undulatus), *Health Phys.* 18, 141-148 (1970).
- 4.3 Beninson, D. E., E. Vander Elst, and D. Cancio, Biological aspects in the disposal of fission products into surface waters, IN Disposal of radioactive wastes into seas, oceans, and surface waters, pp. 337-354, Proceedings of International Atomic Energy Agency Symposium, Vienna, 1966, STI/PUB/126.
- 4.4 Bleyer, B., Zur Kenntnis der Iodes als Biogenes Element, *Biochem. Z.* 170, 256 (1926).
- 4.5 Bowen, H. J. M., Trace elements in biochemistry, Academic Press Inc., New York, 1966.
- 4.6 Causeret, J., Fish as a source of mineral nutrition, IN Fish as food, G. Borgstrom (ed.), pp. 205-234, Academic Press, New York, 1962.
- 4.7 Chavin, W., Thyroid distribution and function in the goldfish, Carassius auratus L., *J. Exp. Zool.* 133, 259-280 (1956).
- 4.8 Chavin, W., and B. Bouwman, Metabolism of iodine and thyroid hormone synthesis in the goldfish, Carassius auratus L., *Gen. comp. Endocrin.* 5, 493-503 (1965).
- 4.9 Closs, K., Über das Vorkommen des Jods im Meere und in Meeresorganismen, *Arch. Math. Naturv.* 40, 1-150 (1931), IN Vinogradov 1953.
- 4.10 Copeland, R. A., and J. C. Ayers, Trace element distribution in water, sediment, phytoplankton, zooplankton, and benthos of Lake Michigan: A baseline study with calculations of concentration factors and buildup of radioisotopes in the food web, Environmental Research Group Special Report No. 1, Great Lakes Research Division, Univ. Michigan, Ann Arbor, 1970.

References for Section 3.4 (continued)

4.11 Copeland, R. A., R. H. Beethe, and W. W. Prater, Trace element distribution in Lake Michigan fish: A baseline study with calculations of concentration factors and equilibrium radioisotopes distribution, Environmental Research Group Special Report No. 2, Great Lakes Research Division, Univ. Michigan, Ann Arbor, March, 1973.

4.12 Dundee, H., and A. Gorbman, Utilization of radioiodine by the thyroid of neotenic salamander, Eurycea tynerensis Moore and Hughes, Physiol. Zool. 33, 58-63 (1960).

4.13 Eales, J. G., The influence of temperature on thyroid histology and radioiodine metabolism of yearling steelhead trout, Salmo gairdneri, Can. J. Zool. 42, 829-841 (1964).

4.14 Ewing, R. A., J. E. Howes, and R. B. Price, Bioenvironmental and radiological-safety feasibility studies Atlantic-Pacific Interoceanic Canal: Radionuclide and stable-element analysis of environmental samples from Routes 17 and 25, Battelle Memorial Institute, BMI-171-29 (1969), also IOCS-MEMO-BMI-33 (1969).

4.15 Fellenberg, Th. von, Das Vorkommen der Kreslauf und der Stoffwechsel des Jods, Ergebn. Physiol. 25, 176-363 (1926).

4.16 Fellenberg, Th. von, Untersuchungen über das Vorkommen von Jod in der Nature, Mitt. Lebensm. Hyg., Bern 14, 161 (1923).

4.17 Fontain, M., F. Lachiver, S. Leloup, and M. Olivereau, La Fonction thyroidienne du Saumon (Salmo salar L.) au cours de sa migration reproductrice, J. Physiol. Paris 40, 182-84 (1948).

4.18 Glasser, R., Zur Frage der Anreicherung von Radiojod (J-131) in der Muschel Dreissensia polymorpha Pall., Naturwissenschaften 47, 284 (1960).

4.19 Glasser, R., Über die Aufnahm von Radiojod aus wassriger Lösung durch die Muschel Dreissensia polymorpha Pall., Zool. Jb. Physiol. 69, 339-378 (1961). In Polikarpov 1966.

4.20 Hercus, C. E., and K. C. Roberts, The iodine content of foods, manures, and animal products in relation to the prophylaxis of endemic goitre in New Zealand, J. Hyg., Camb. 26, 49 (1927).

4.21 Hickman, C. P., Influence of environment on the metabolism of iodine in fish, Ann. N.Y. Acad. Sci., Gen. comp. Endocr. Sup. 1, 48-62 (1962).

References for Section 3.4 (continued)

4.22 Hunn, J. B., and P. O. Fromm, Uptake, turnover and excretion of iodine-131 by rainbow trout (Salmo gairdneri), Biol. Bull. 126, 282-290 (1964).

4.23 Hunn, J. B., and P. O. Fromm, In vivo uptake of radioiodine by rainbow trout, J. Water Poll. Contr. Assoc. 38, 1981-1985 (1966).

4.24 Hunn, J. B., and E. P. Reineke, Influence of iodine intake on iodine distribution in trout, Proc. Soc. exp. Biol. Med. 115, 91-93 (1964).

4.25 Hunt, E. L., and J. N. Dent, Iodine uptake and turnover in the frog tadpole, Physiol. Zool. 30, 87-91 (1957).

4.26 Jarvis, N. D., R. W. Clough, and E. D. Clark, Iodine content of Pacific Coast salmon, Univ. Washington Pub. Fish. 1, 109-140 (1925-28). In Robertson and Chaey 1953.

4.27 Kolehmainen, S., S. Takatalo, and J. K. Miettinen, A tracer experiment with iodine-131 in an oligotrophic lake, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 278-284 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

4.28 Livingston, D. A., Chemical composition of rivers and lakes, Geological Survey Professional Paper 440-G (1963).

4.29 Lunde, G., Über die Geochemie und Biochemie des Jods mit besonderer Berücksichtigung der norwegischen Kropfprophy laxe, Wien. klin. Wschr. 40, 1559 (1928). In Vinogradov 1953.

4.30 Lunde, G., J. Boe, and K. Closs, Iodine content of American marine animals, J. Cons. Perm. int. Explor. Mor. 5, 216 (1930). In Vinogradov 1953.

4.31 Lunde, G., K. Closs, H. Haaland, and S. O. Madsen, Jodinnholdet i norsk fisk og fiskerprodukter, Aarsberetn. Norg. Fisk., No. 4, 1-41 (1928). In Vinogradov 1953.

4.32 Mazzocco, P., Iodine in food in salta, Sem. Med., B. Aires 37, 364 (1930).

4.33 Monier-Williams, G. W., Trace elements in food, John Wiley and Sons, New York, 1950.

4.34 Nilson, H. W., and E. J. Coulson, The mineral content of the edible portions of some American fisheries products, Invest. Rep. U.S. Bur. Fish. 41, 1-7 (1939). In Vinogradov 1953.

References for Section 3.4 (continued)

4.35 Odum, E. P., and F. B. Golley, Radioactive tracers as an aid to the measurement of energy flow at the population level in nature, IN Radioecology, V. Schultz and A. W. Clement (eds.), p. 403 (1963), Proceedings of the National Symposium on Radioecology, Fort Collins, Colorado, September 10-15, 1961.

4.36 Ohmomo, Y., H. Suzuki, M. Sumiya, and M. Saiki, Studies on accumulation excretion and distribution of iodine-131 and cadmium-115m in freshwater fish and marine fish, J. Rad. Res. 13, 3 (1972).

4.37 Orr, J. B., and I. Leitch, Iodine in nutrition, A review of existing information up to 1927, United Kingdom Medical Research Council, Special Report Series No. 123, His Majesty's Stationery Office, London (1929).

4.38 Polikarpov, G. G., Radioecology of Aquatic Organisms (Reinhold Book Div., N.Y., 1966).

4.39 Robertson, O. H., and A. L. Chaney, Thyroid hyperplasia and tissue iodine content in spawning rainbow trout: A comparative study of Lake Michigan and California sea-run trout, Physiol. Zool. 26, 328-340 (1953).

4.40 Short, Z. F., R. F. Palumbo, P. R. Olson, and J. R. Donaldson, The uptake of iodine-131 by the biota of Fern Lake, Washington, in a laboratory and a field experiment, Ecology 50, 979-989 (1969).

4.41 Spector, W. S., Handbook of Biological Data, W. B. Saunders Co., Philadelphia (1956).

4.42 Takatalo, S., S. Kolehmainen, and J. K. Miettinen, Excretion of iodine from perch (Perca fluviatilis), IN Biological enrichment chains of radionuclides in Finnish lakes, pp. 104-111 (1968). Annual Report for the Period Sept. 1, 1967 to August 31, 1968, Department of Radiochemistry, Univ. Helsinki, Finland.

4.43 Timofeev-Resovskii, N. V., E. A. Timofeeva-Resovskaya, G. A. Milyutina, and A. B. Getsova, Coefficients of accumulation of radioisotopes of sixteen different elements by freshwater organisms and the effect of EDTA on some of them, Dokl. Akad. Nauk SSSR 132, 369-372 (1960).

4.44 Timofeyeva-Resovskaya, Ye. A., Distribution of radioisotopes in the main components of freshwater bodies (Monograph), Tr. Inst. Biol. Akad. Nauk SSSR, Ural'skiy filia 30, 1-78 (JPRS 21.816) 1963.

References for Section 3.4 (continued)

- 4.45 Tressler, D. K., and A. W. Wells, Iodine content of sea foods, Doc. U.S. Bur. Fish., No. 967 (1924).
- 4.46 Van Hook, R. I., Jr., and D. A. Crossley, Jr., Assimilation and biological turnover of cesium-134, iodine-131, and chromium-51 in brown crickets Acheta domesticus (L.), Health Phys. 16, 463-467 (1969).
- 4.47 Vinogradov, A. P., The elementary chemical composition of marine organisms, Sears Foundation for Marine Research, Yale Univ., New Haven, 1953.
- 4.48 Volkova, G. A., The build-up factor of the radioisotopes of some chemical elements in aquatic insects, Entomologicheskoye Obozreniye Vol. 17, 516-519 (1963), Trans. 63-41234 (November 25, 1963).
- 4.49 Wilke-Dorfurt, E., Über den Judgehalf der Schalen von Muscheltieren. Sum Chemischen Teil des Kropfproblems. II, Biochem. Zeit. 192, 74-82 (1928).
- 4.50 Willis, D. L., and B. B. Valett, Metabolism of iodine-131 in two amphibian species (Taricha granulosa and Rana pipiens), IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 394-400 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.
- 4.51 Winchester, J. W., Chemical equilibria of iodine in Great Lakes and waters, IN Proceedings of the Thirteenth Conference on Great Lakes Research, pp. 137-140, 1970, International Association for Great Lakes Research.

3.5 MANGANESE

3.5.1 Environmental Manganese

Manganese is found in six major forms in natural waters (Gibbs, 1973; Wangersky, 1963). In the suspended phase, manganese may be (1) a part of the crystalline structure of suspended material, (2) precipitated onto the suspended material probably as MnO_2 , (3) adsorbed onto solids by ion exchange, and (4) incorporated into solid biological materials. In the dissolved phase manganese may occur as one or more (5) dissolved ionic species, e.g., Mn^{++} or inorganic associations, and as (6) complexes with organic molecules in solution.

Gibbs (1973) found that 10% of the manganese in the Yukon River and 17% of the manganese in the Amazon River were in the dissolved phase; less than 1% of the manganese in either river was on ion exchange sites. In the Yukon River 46% of the manganese was precipitated as metallic coatings, 37% was in the crystalline structure and 7% was in biological material. In the Amazon River 50% of the manganese was on the precipitated metallic coatings, 27% was in the crystalline structure and 5% was in biological material.

In surface waters of five Alaskan lakes and Par Pond (South Carolina), 51 to 94% of the manganese was in the particulate phase (Marshall and LeRoy, 1973). In lakes, most of the manganese in the particulate phase is probably MnO_2 precipitates (Marshall and LeRoy, 1973). Dissolved manganese may be largely bound to organics since Mn^{++} will most likely oxidize to MnO_2 on surfaces under aerobic conditions (Gibbs, 1973; Wangersky, 1963). Physicochemical form of manganese affects biological availability (Gibbs, 1973): (1) manganese in the crystalline structure

of sediments is not available for uptake; (2) manganese in organic materials and manganese precipitates may be somewhat available; and (3) manganese in solution or at exchange sites may be very available. Since the proportions of these different physicochemical forms may vary considerably from site to site, the availability of manganese to organisms would be expected to vary from site to site. This picture is further complicated by strong seasonal cycles of manganese concentration in these forms (Marshall and LeRoy, 1973).

3.5.2 Manganese Metabolism

It was noted in Section 2.3 that the availability of a homeostatically-controlled element in different physicochemical forms had no effect on the concentration of that element in the organism. The bioaccumulation factor of a homeostatically-controlled stable element and the bioaccumulation factor of the radionuclide are given by Equation (2.1.4). Fortunately, the considerable site to site variability in manganese availability is not an issue for many organisms since manganese is homeostatically controlled in vertebrates and in some invertebrates. In a literature review Schroeder, Balassa, and Tipton (1966) concluded that: "An efficient homeostatic mechanism for manganese appears to operate in all vertebrate and most invertebrate animals." Bryan and Ward (1965) suggested that the marine lobster Homerus has a regulatory mechanism for manganese and further suggested that a similar mechanism should exist in molluscs and fishes. Cavallero and Merlini (1967) found only small site to site variations in manganese concentrations among aquatic vertebrates. Bortoli et al. (1969) found no differences in manganese concentrations among fishes from four

different Italian lakes and Lentsch et al. (1973) detected little or no variation in manganese concentrations in fishes from the Hudson River, despite great changes in manganese concentration in the water.

3.5.3 Review of Manganese Bioaccumulation Factors

3.5.3.1 Manganese Bioaccumulation Factors for Fishes

As indicated in Section 3.5.2 the evidence for homeostatic control of manganese concentration in fishes is strong. Manganese concentrations in fishes and water and the bioaccumulation factors for manganese in fishes are tabulated in Table 3.5.1. Table 3.5.2 gives the means and coefficients of variation for the data on manganese concentrations in water and fishes given in Table 3.5.1. The small coefficients of variation for fishes in Table 3.5.2 and the large range of bioaccumulation factors in Table 3.5.1 support the hypothesis of homeostatic control of manganese. It is also interesting to note that manganese concentrations in estuarine and marine fishes are about the same as those in freshwater fishes. The average muscle concentration of manganese in Pomatomus saltatrix (bluefish), a marine epipelagic carnivore, is 0.22 ppm wet weight and the average concentration of manganese in Antimora rostrata, a bathy-demersal fish, is 0.21 ppm (Cross et al., 1973). The concentration of manganese in the ocean is about 0.002 ppm. Manganese concentrations in five species of estuarine fishes was 4.7 ppm on a whole fish basis (Cross and Brooks, 1973). These values for marine and estuarine fishes fall within one standard deviation of the mean for freshwater fishes (Table 3.5.2).

Table 3.5.1 Bioaccumulation factors for manganese in freshwater fishes

Species	Tissue	Feeding habits	Stable element concentration in tissue (ppm)	Stable element concentration in water (ppb)	BF	Location	Reference
<u>Alburnus alborella</u> (bleak)	whole	plankton feeder	20.6 \pm 0.7(6)	9.0-100 ^a	2300-210	Five irrigated sites ^b	5.4
<u>Lepomis gibbosus</u>	whole		18.1 \pm 0.9(5)	9.0-100 ^a	2000-180	Five irrigated sites ^b	5.4
<u>Alburnus alborella</u>	whole	plankton feeder	12.7 \pm 0.8(11)	9.0-100 ^a	1400-130	Five irrigated sites ^c	5.4
<u>Lepomis gibbosus</u>	whole		8.1 \pm 0.3(21)	9.0-100 ^a	900-80	Five irrigated sites ^c	5.4
<u>Cobitis taenia</u>	whole	bottom feeder	5.2 \pm 0.4(4)	9.0-100 ^a	580-50	Five irrigated sites ^c	5.4
<u>Cyprinus carpio</u> (carp)	whole	bottom feeder	4.8 \pm 0.7(4)	9.0-100 ^a	530-50	Five irrigated sites ^c	5.4
<u>Tinca tinca</u> (tench)	whole	scavenger	3.9 \pm 0.3(6)	9.0-100 ^a	430-40	Five irrigated sites ^c	5.4
Average of <u>Perca fluviatilis</u> (perch), <u>Scardinius erythrophthalmus</u> (rudd), and <u>Lepomis gibbosus</u> ^c	whole		7.6	42 ^a	180	Lake Varese, Italy	5.2
	whole		8.6	23 ^a	374	Lake Camabbio	5.2
	whole		3.6	17 ^a	211	Lake Maggiore	5.2
	whole		5.2	8 ^a	650	Lake Monate	5.2
<u>Alosa pseudoharengus</u> (alewife)	whole	zooplankton feeder	4.6 \pm 0.3(19)	0.92 \pm 0.08(53) ^d	5000	Lake Michigan	5.6
	flesh		0.88 \pm 0.11(6)	0.92 \pm 0.08(53) ^d	960	Lake Michigan	5.6
<u>Coregonus macrourus</u> (whitefish, <u>Bonellia</u>)	flesh	plankton and benthic feeder	0.25 \pm 0.03	17 \pm 2 (4) ^a	15	Lake Maggiore	5.17
<u>Alburnus alborella</u> (bleak)	bone		25	17 \pm 2 (4) ^a	1500	Lake Maggiore	5.17
	flesh	plankton feeder	0.36 \pm 0.04	17 \pm 2 (4) ^a	21	Lake Maggiore	5.17
	bone		41	17 \pm 2 (4) ^a	2400	Lake Maggiore	5.17

Table 3.5.1 (continued)

Species	Tissue	Feeding habits	Stable element concentration in tissue (ppm)	Stable element concentration in water (ppb)	BF	Location	Reference
<u><i>Scardinius erythrophthalmus</i></u> (rudd)	flesh	omnivorous	0.35 \pm 0.4	17 \pm 2 (4) ^a	21	Lake Maggiore	5.17
	bone		39	17 \pm 2 (4) ^a	2300	Lake Maggiore	5.17
<u><i>Notropis hudsonius</i></u> (spottail shiner)	whole flesh	unknown benthic	3.3 \pm 0.3(15) 0.51 \pm 0.06(3)	0.92 \pm 0.08(53) ^d 0.92 \pm 0.08(53) ^d	3600 550	Lake Michigan	5.6
	flesh	benthic	0.34 \pm 0.04(5)	0.92 \pm 0.08(53) ^d	370	Lake Michigan	5.6
<u><i>Coregonus</i></u> sp. (chub)	flesh	zooplankton and benthic feeder	0.29 \pm 0.05(6)	0.92 \pm 0.08(53) ^d	315	Lake Michigan	5.6
<u><i>Coregonus artedii</i></u> (cisco)	flesh		0.5 \pm 0.1(3)	0.92 \pm 0.08(53) ^d	500	Lake Michigan	5.6
<u><i>Lepomis</i></u> sp. (sunfish)	flesh		0.25 \pm 0.03(13)	0.92 \pm 0.08(53) ^d	270	Lake Michigan	5.6
<u><i>Prosobium cylindraceum</i></u> (rand whitefish)	flesh	benthic	0.37 \pm 0.03(24)	0.92 \pm 0.08(53) ^d	400	Lake Michigan	5.6
<u><i>Osmerus mordax</i></u> (smelt)	whole flesh	plankton and benthic feeder	3.9 \pm 0.4(16) 0.27 \pm 0.05(9)	0.92 \pm 0.08(53) ^d 0.92 \pm 0.08(53) ^d	4200 290	Lake Michigan	5.6
<u><i>Perca flavescens</i></u> (yellow perch)	flesh	partially piscivorous	0.23 \pm 0.03(20)	0.92 \pm 0.08(53) ^d	250	Lake Michigan	5.6
<u><i>Oncorhynchus kisutch</i></u> (coho salmon)	flesh	piscivorous	0.21 \pm 0.02(24)	0.92 \pm 0.08(53) ^d	230	Lake Michigan	5.6
<u><i>Salvelinus namaycush</i></u> (Lake trout)	flesh	piscivorous	0.23 \pm 0.02(11)	0.92 \pm 0.08(53) ^d	250	Lake Michigan	5.6

Table 3.5.1 (continued)

Species	Tissue	Feeding habits	Stable element concentration in tissue (ppm)	Stable element concentration in water (ppb)	BF	Location	Reference
<u><i>Salmo gairdneri</i></u> (steelhead)	flesh	piscivorous	0.23 \pm 0.02(6)	0.92 \pm 0.08(53)	250	Lake Michigan	5.6
<u><i>Micropterus</i></u> <u><i>notomieu</i></u> (smallmouth bass)	flesh	piscivorous	0.25 \pm 0.02(3)	0.92 \pm 0.08(53)	270	Lake Michigan	5.6

^aUnfiltered value.^bMeasurements on fish made in July - August.^cMeasurements on fish made in September - November.^dFiltered value.

Table 3.5.2 Concentration and coefficients of variation of manganese concentration (ppm) in water and freshwater fishes (fresh weight).

	Water	Flesh	Whole Fishes
Arithmetic Mean	0.028 + 0.034	0.34 + .17	7.9 + 5.5
Coefficient of Variation	119%	50%	70%
N	7	16	14
Geometric Mean	0.014 \times 4.43	0.32 \times 1.45	6.7 \times 1.84
Coefficient of Variation	343%	45%	84%
N	7	16	14

Using the geometric means of stable manganese concentration in freshwater fishes in Table 3.5.2 and Equation (2.1.3),

$$BF(Mn)_i = 6.7/[Mn]_w \quad (3.5.1)$$

for whole fish and

$$BF(Mn)_i = 0.32/[Mn]_w \quad (3.5.2)$$

for fish flesh, where $[Mn]_w$ has units of ppm. These relations and data from Table 3.5.1 are plotted in Figure 3.5.1. Note that it is inconsequential whether filtered or unfiltered values are plotted on the graphs. However, for predictive purposes, an unfiltered value for the manganese concentration in water should be used (Section 2.5.3) unless the user is willing to accept the gross overestimates obtained by use of filtered concentrations.

We hypothesize that the differences in manganese concentration among species of fishes are owing to physiological demands rather than differences in concentration of manganese in the fishes' food. However, bottom feeders have lowest concentrations, plankton feeders have the highest, and piscivores have intermediate concentrations. These differences merit further research.

3.5.3.2 Manganese Bioaccumulation Factors for Plants

Table 3.5.3 summarizes the few data available on manganese bioaccumulation factors for freshwater plants. Harvey (1969, 1970), using a continuous flow, laboratory system to control temperature, showed that nonlethal temperatures have no effect on the manganese bioaccumulation factors for algae. On the basis of data in Table 3.5.2, we recommend




Figure 3.5.1 Bioaccumulation factors for manganese in freshwater fishes as a function of manganese concentration in water.

Table 3.5.3 Bioaccumulation factors for manganese in aquatic plants

Species	Description	Stable element concentration in plant (ppm)	Stable element concentration in water (ppm) Filtered Unfiltered	^{54}Mn concentration in plant (pCi/g)	^{54}Mn concentration in water (pCi/l)	BF ^a	Location	Reference
<u><i>Chlorophyta</i> (green algae and <i>Bacillariophyceae</i> (diatoms))</u>	phytoplankton	11+4 (14) ^b	0.00092+ 0.00008 (53)			12,000	Lake Michigan	5.5
<u>Mix of 1. <i>Lyngbya</i>, <i>Oscillatoria</i>, <i>Phormidium</i>; 2. <i>Cosmarium</i>; and 3. <i>Nitzchia</i>, <i>Melosira</i></u>	1. filamentous blue-green algae; 2. green algae; and 3. diatoms			11.0	0.48 ^c	23,000	Reactor discharge canal on Connecticut River	5.1
<u><i>Stigeoclonium lubricum</i></u>	filamentous green algae			2.3×10^5	4.5×10^4	5,000	Laboratory, continuous flow exp. (23°-32°C)	5.12
<u><i>Navicula seminulum</i></u>	unicellular diatom			0.70×10^5	4.5×10^4	1,600	Laboratory, continuous flow exp. (23°-32°C)	5.12
<u><i>Plectonema boryanum</i></u>	filamentous blue-green algae			1.4×10^5	4.5×10^4	3,200	Laboratory, continuous flow exp. (23°-32°C)	5.12
<u><i>Potamogeton perfoliatus</i></u>	submerged macro-phyte	250-2500	0.01-0.1			25,000 ^d	Hudson River	5.14
<u><i>Ceratophyllum</i> sp.</u>	rootless, submerged thin stemmed macro-phyte	81	0.003	0.01		8,100	Lake Maggiore, Italy, mesotrophic	5.19
<u><i>Lagarosiphon</i> sp.</u>		6	0.003	0.01		600	Lake Maggiore, Italy, mesotrophic	5.19
<u><i>Elodea</i> sp.</u>	submerged aquatic plant	105		0.026		4,000	Lake Maggiore, Italy, mesotrophic	5.17
<u><i>Potamogeton</i> sp. (pondweed)</u>	submerged floating leaves	46		0.026		18,000	Lake Maggiore, Italy, mesotrophic	5.17

Table 3.5.3 (continued)

Species	Description	Stable element concentration in plant (ppm)	Stable element concentration in water (ppm)		^{54}Mn concentration in plant (pCi/g)	^{54}Mn concentration in water (pCi/l)	BF ^a	Location	Reference
			Filtered	Unfiltered					
<i>Myriophyllum</i> sp. (millfoil)	mostly submerged, top of plant out of water	32		0.026			1,200	Lake Maggiore, Italy, mesotrophic	5.17
<i>Phragmites</i> sp. (reed grass)	emergent	28		0.026			1,100	Lake Maggiore, Italy, mesotrophic	5.17
<i>Nymphaea lutea</i> (water lily)	large floating leaves	9	0.003	0.01			900	Lake Maggiore, Italy, mesotrophic	5.19
<i>Najas</i> sp. (pondweed)	submerged	14		0.026			540	Lake Maggiore, Italy, mesotrophic	5.17
<i>Nuphar</i> sp. (water lily)	large floating leaves	5		0.026			190	Lake Maggiore, Italy, mesotrophic	5.17

^aBioaccumulation factors are all based on unfiltered concentrations in water unless only filtered values were reported.

^bMean \pm S.E. (no. of samples) of all non-zero "corrected" (Copeland and Ayers 1970) concentrations.

^cEstimate from release rates.

^dEstimate from nonlinear regression analysis.

that a value of 10^4 be used as the manganese bioaccumulation factor in algae. For submerged macrophytes we recommend a bioaccumulation factor of 10^4 . On the basis of the few data from Lake Maggiore, we recommend that a value of 10^3 be used for macrophytes with floating leaves and emergent vegetation.

3.5.3.3 Manganese Bioaccumulation Factors for Invertebrates

3.5.3.3.1 Manganese Bioaccumulation Factors for Molluscs

In bivalves, manganese concentrations in both the shell and soft tissues increase with size (Merlini, 1966; Merlini et al., 1965; Merlini et al., 1967). This fact explains part of the standard error associated with the mean concentrations of manganese given for bivalve tissues in Table 3.5.4. The data in Table 3.5.4 were averaged over all size classes. Manganese concentrations in the mantle and visceral sac may vary considerably with nature of the substrate for the same lake. This factor is in large part responsible for the large standard errors for Unio in Lake Maggiore since Unio was taken from sites having different substrates.

In snails, manganese concentrations do not vary greatly with size (Merlini, 1966; Merlini et al., 1965; Merlini et al., 1967). Moreover, only small differences in manganese concentration were found in the snail Viviparus with change in habitat within the same lake (Merlini et al., 1965). The data on bioaccumulation factors for manganese in molluscs (Table 3.5.4) suggest that $BF(Mn)_i$ is constant or that manganese concentrations are only partially homeostatically controlled. The bioaccumulation factor for Unio tissues in Lake Maggiore is about

Table 3.5.4 Bioaccumulation factors for manganese in molluscs

Taxon/Tissue	Stable element concentration in tissue (ppm)	Stable element concentration in water (ppm) ^d	Bioaccumulation Factor	Location	Reference
Bivalves:					
<u><i>Anodonta cygnea</i></u>					
Shell	419+50(2) ^b	0.016	26,000	Lake Maggiore	5.18
Gills	2450+750(2) ^b		150,000		
Mantle	720+190(2) ^b		45,000		
Visceral sac	155+35(2) ^b		9,700		
Muscle	30.5+12(2) ^b		1,900		
<u><i>Anodonta cygnea</i></u>					
Whole	341+36(4)	~0.1	3,400	Irrigation canal, Crescentino, Italy	5.4
Whole	206+7(4)	~0.1	2,100	Irrigation canal, Vercelli, Italy	5.4
<u><i>Lamellidens marginalis</i></u>		0.011 ^c		Lake Masunda, India	5.21
Labial palps	9872		900,000		
Gills	3210		300,000		
Mantle	1228		110,000		
Visceral sac	1220		110,000		
Muscle	834		80,000		
Foot	354		30,000		
Total soft	1247		110,000		
Shell	942		90,000		
<u><i>Quadrula pustulosa</i></u>		0.004 ^c			5.20
Shell	486		120,000		
<u><i>Unio mancus elongatus</i></u>		0.016		Lake Maggiore	5.16, 5.18, 5.19
Shell	382+43(12) ^d		24,000		
Gills	2100+260(7) ^d		130,000		
Mantle	1400+330(7) ^d		88,000		
Muscle	260+53 (7) ^d		16,000		
Visceral sac	370+96 (7) ^d		23,000		

Table 3.5.4 (continued)

Taxon/Tissue	Stable element concentration in tissue (ppm)	Stable element concentration in water (ppm) ^a	Bioaccumulation Factor	Location	Reference
<u><i>Unio mancus elongatus</i></u>					
Shell	870±57(3) ^b	0.033	26,000	Varese	5.16
Mantle	2400±300(3) ^b		71,000		
Visceral sac	600±58(3) ^b		18,000		
Snails:					
<u><i>Lymnea ovata</i></u>		0.016			
Shell	35±1(4)		2,200	Lake Maggiore	5.18, 5.19
Soft tissues	28±1(4)		1,800		
<u><i>Lymnea peregra</i></u>					
Whole	126±41(3)	~0.1	1,300	Irrigation canal, Casalino, Italy	5.4
Whole	85 (1)	~0.1	850	Irrigation canal, Vercelli, Italy	5.4
<u><i>Lymnea stagnalis</i></u>					
Whole	107±25(6)	~0.1	1,100	Irrigation canal, Casalino, Italy	5.4
Whole	70±6(5)	~0.1	700	Irrigation canal, Crescentino, Italy	5.4
Whole	110±14(15)	~0.1	1,100	Irrigation canal, Vercelli, Italy	5.4
<u><i>Physa acuta</i></u>					
Whole	25±2(16)	0.009	2,800	Irrigation canal, Cameri, Italy	5.4
Whole	85±13(2)	~0.1	850	Irrigation canal, Casalino, Italy	5.4
<u><i>Planorbis</i> sp.</u>					
Whole	599±221(3)	~0.1	6,000	Irrigation canal, Casalino, Italy	5.4
Whole	750±56(2)	~0.1	7,500	Irrigation canal, Vercelli, Italy	5.4

Table 3.5.4 (continued)

Taxon/Tissue	Stable element concentration in tissue (ppm)	Stable element concentration in water (ppm) ^a	Bioaccumulation Factor	Location	Reference
<u><i>Viviparus ater</i></u>					
Shell	88 \pm 3.7(19) ^d	0.016	5,500	Lake Maggiore	5.4
Soft tissues	12 \pm 0.46(19) ^d		750		
<u><i>Viviparus contectus</i></u>					
Whole	202 \pm 53(4)	\sim 0.1	2,000	Irrigation canal, Crescentino, Italy	5.4
Whole	92 \pm 8(6)	\sim 0.1	920	Irrigation canal, Vercelli, Italy	5.4

^aUnfiltered concentrations unless indicated otherwise.

^bAverage of means for size classes.

^cFiltered concentration.

^dAverage over all size classes and study sites.

the same as that in Lake Varese despite a two-fold difference in manganese concentration in the water. The whole-body bioaccumulation factor for Physa is about 3.4 times higher for the environment having a manganese concentration in water an order of magnitude lower than that of the other. However, we must not put too much emphasis on these results because the manganese concentrations in water, which are based on only a few measurements, can only be considered rough indicators of the concentration that the organisms would be exposed to over long periods of time since the manganese concentration in water has strong seasonal variability (Section 3.5.1). Further, as pointed out in Section 2.5.5, sediment contamination may affect whole-body concentrations of manganese in invertebrates.

For studies in which bioaccumulation factors are based on unfiltered values for the manganese concentration in water, the bioaccumulation factors for all soft tissues in bivalves except gills are less than 10^5 . Thus, we recommend that a bioaccumulation factor of 10^5 be used for soft tissues of bivalves (Table 3.5.4). Note that the value of this recommended bioaccumulation factor compares favorably with the ^{54}Mn bioaccumulation factor of 4.5×10^4 determined from the fallout data of Gaglione and Ravera (1964) for all soft tissues of Unio from Lake Maggiore. On the basis of data in Table 3.5.4, we recommend that a bioaccumulation factor of 3×10^4 be used for bivalve shells.

Based on the few data for snails we recommend that a bioaccumulation factor of 10^4 be used for snail shells and whole snails. For soft tissues of snails, we recommend a bioaccumulation factor of 2×10^3 .

3.5.3.3.2 Manganese Bioaccumulation Factors for Invertebrates other than Molluscs

On the basis of data in Table 3.5.5, we recommend that a bioaccumulation factor of 10^4 be used for crustaceans. It is apparent from Table 3.5.5 that there is considerable variation in manganese bioaccumulation factors among insect species. Owing to the lack of comprehensive data, we must recommend that a bioaccumulation factor of 10^3 be applied to insects in general.

Table 3.5.5 Bioaccumulation factors for manganese in invertebrates other than molluscs

Species	Description	Size class or life stage	Tissue	Stable element concentration in tissue (ppm)	Stable element concentration in water (ppb) ^a	BF	Location	Reference
Calanoid and cyclopoid copepods	zooplankton		whole	3.6+0.8(33) ^b	0.92+0.08(53) ^c	3,900	Lake Michigan	5.5
			whole	4.2+0.7(40) ^d		4,600	Lake Michigan	5.5
Amphipoda (mostly <i>Pontoporeia</i> sp.)	small shrimp		whole	7.4+2.1(17) ^b	0.92+0.08(53) ^c	8,000	Lake Michigan	5.5
<i>Ranatra linearis</i>	carnivorous aquatic hemipteran, feeding on tadpoles, insects and larvae	adult	whole	51.8 (1)	9	6,000	Cameri, Italy ^e	5.4
		adult	whole	333+86(2)	~100	3,000	Casalino, Italy	5.4
		adult	whole	354+50(2)	~100	4,000	Crescentino, Italy	5.4
<i>Notonecta glauca</i>	aquatic insect	adult	whole	6.9+0.8(2)	~100	70	Casalino, Italy	5.4
			whole	5.8 (1)	~100	60	Oldenico, Italy	5.4
<i>Hydrophilus piceus</i>	carnivorous aquatic insect	adult	whole	9.4+0.7(2)	9	1,000	Cameri, Italy	5.4
			whole	34.4+7.6(4)	~100	300	Casalino, Italy	5.4
			whole	76.9+1.9(3)	~100	800	Crescentino, Italy	5.4
			whole	44.7+6.6(4)	~100	400	Oldenico, Italy	5.4
			whole	17.3+0.2(4)	~100	200	Vercelli, Italy	5.4
<i>Dysticus marginalis</i>	aquatic insect	adult	whole	6.6 (1)	9	700	Cameri, Italy	5.4
			whole	14.7+7.9(2)	~100	100	Casalino, Italy	5.4
			whole	40.1 (1)	~100	400	Crescentino, Italy	5.4
<i>Haemopis</i> sp. (bloodworm)	parasitic leech found on fish and insects		whole	9.5	~100	100	Oldenico, Italy	5.4

^aUnfiltered values unless indicated otherwise.^bMean + S.E. (no. of samples) of all non-zero "corrected" (Copeland and Ayers 1970) concentrations.^cFiltered value.^dUncorrected value.^eIrrigation canal at this and following locations.

References for Section 3.5

The numbers in the left-hand margin correspond to the reference numbers used in the tables of section 3.5.

- 5.1 Blanchard, R. L., and B. Kahn, The fate of radionuclides released from a PWR Nuclear Power Station into a river, IN Symposium on environmental behavior of radionuclides released in the nuclear industry, Aix-en-Provence, France, 14-18 May, 1973, IAEA/SM-172/26.
- 5.2 Bortoli de, M. C., P. Gaglione, O. Ravera, and G. de Fisica, Fall-out manganese-54 and zinc-95 in water and fishes of four lakes in Northern Italy, Minerva Fisionucleare 13, 72-77 (1969).
- 5.3 Bryan, G. W., and E. Ward, The absorption and loss of radioactive and non-radioactive manganese by the lobster, Homarus vulgaris, J. Mar. Biol. Ass. U.K. 45, 65-95 (1965).
- 5.4 Cavalloro, R., and M. Merlini, Stable manganese and fallout radio-manganese in animals from irrigated ecosystems of the Po Valley, Ecology 48, 924-929 (1967).
- 5.5 Copeland, R. A., and J. C. Ayers, Trace element distribution in water, sediment, phytoplankton, zooplankton, and benthos of Lake Michigan: A baseline study with calculations of concentration factors and buildup of radioisotopes in the food web, Environmental Research Group Special Report Number 1, Great Lakes Research Division, Univ. Michigan, Ann Arbor, 1970.
- 5.6 Copeland, R. A., R. H. Beethe, and W. W. Prater, Trace element distribution in Lake Michigan fish: A baseline study with calculations of concentration factors and equilibrium radioisotope distributions, Environmental Research Group Special Report No. 2, Great Lakes Research Division, Univ. Michigan, Ann Arbor, 1973.
- 5.7 Cross, F. A., and J. H. Brooks, Concentrations of manganese, iron, and zinc in juveniles of five estuarine-dependent fishes, IN Radionuclides in Ecosystems, D. J. Nelson (ed.), pp. 769-775, Proc. of Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971 (1973).
- 5.8 Cross, F. A., L. H. Hardy, N. Y. Jones, and R. T. Barber, Relation between total body weight and concentrations of manganese, iron, copper, zinc, and mercury in white muscle of bluefish (Pomatomus saltatrix) and a bathy-demersal fish Antimora rostrata, J. Fish. Res. Bd. Can. 30, 1287-1290 (1973).

References for Section 3.5 (continued)

5.9 Gaglione, P., and O. Ravera, Manganese-54 concentrations in fall-out, water and *Unio* mussels of Lake Maggiore, 1960-63, *Nature* 204, 1215-1216 (1964).

5.10 Gibbs, R. J., Mechanisms of trace metal transport in rivers, *Science* 180, 71-73 (1973).

5.11 Harvey, R. S., Effects of temperature on the sorption of radio-nuclides by a blue-green algae, IN *Symposium on Radioecology*, D. J. Nelson and F. C. Evans (eds.), pp. 266-269, *Proceedings of the Second National Symposium on Radioecology*, Ann Arbor, Michigan, May 15-17, 1969 (1969).

5.12 Harvey, R. S., Temperature effects on the sorption of radio-nuclides by freshwater algae, *Health Phys.* 19, 293-297 (1970).

5.13 Hutchinson, G. E., *A treatise on limnology*, Vol. I, John Wiley and Sons, New York, 1957.

5.14 Lentsch, J. W., T. J. Kneip, M. E. Wrenn, G. P. Howells, and M. Eisenbud, Stable manganese and manganese-54 distributions in the physical and biological components of the Hudson River Estuary, IN *Radionuclides in Ecosystems*, D. J. Nelson (ed.), pp. 752-768, *Proceedings of the Third National Symposium on Radioecology*, Oak Ridge, Tennessee, May 10-12, 1971 (1973).

5.15 Marshall, J. S., and J. H. LeRoy, Iron, manganese, cobalt, and zinc cycles in a South Carolina Reservoir, IN *Radionuclides in Ecosystems*, D. J. Nelson (ed.), pp. 465-473, *Proceedings of the Third National Symposium on Radioecology*, Oak Ridge, Tennessee, May 10-12, 1971 (1973).

5.16 Merlini, M., The freshwater clam as a biological indicator of radiomanganese, IN *Radioecological Concentration Process*, B. Aberg, and F. P. Hungate (eds.), pp. 977-982, *Proceedings of the International Symposium*, Stockholm, April 25-29, 1966.

5.17 Merlini, M., C. Bigliocca, A. Berg, and G. Pozzi, Trends in the concentration of heavy metals in organisms of a mesotrophic lake as determined by activation analysis, IN *Nuclear techniques in environmental pollution*, pp. 447-458, *IAEA Symposium on the Use of Nuclear Techniques in the Measurement and Control of Environmental Pollution*, Salzburg, Austria, Oct. 25-30, 1970 (1971).

5.18 Merlini, M., F. Girardi, R. Pietra, and A. Brozgelli, The stable manganese content of molluscs from Lake Maggiore determined by activation analysis, *Limnol. Oceanogr.* 10, 371-378 (1965).

References for Section 3.5 (continued)

- 5.19 Merlini, M., F. Giradi, and G. Pozzi, Activation analysis in studies of an aquatic ecosystem, IN Nuclear Activation Techniques in the Life Sciences, pp. 615-629, Proceedings of the Symposium held by the IAEA in Amsterdam, May 8-12, 1967, also STI/PUB/155, SM-91/7 (1967).
- 5.20 Nelson, D. J., T. C. Rains, and J. A. Norris, High purity calcium carbonate in freshwater clam shell, Science 152, 1368-1370 (1966).
- 5.21 Patel, B., and G. R. Doshi, Lamellidens marginalis L. as indicator of manganese-54, Archivio di Oceanografia e Limnologic 17, 27-42 (1972).
- 5.22 Schroeder, H. A., J. J. Balassa, and I. H. Tipton, Essential trace metals in man: Manganese, J. Chron. Dis. 19, 545-571 (1966).
- 5.23 Wangersky, P. J., Manganese in ecology, IN Radioecology, V. Schultz and A. W. Klement, Jr. (eds.), pp. 499-508, First National Symposium on Radioecology, Sept. 10-15, 1961 (1963).

3.6 COBALT

3.6.1 Cobalt Metabolism

Cobalt is essential for some bacteria, some fungi, several species of blue green algae and several species of mammals (Bowen, 1966). Cobalt II activates enzymes. A principle use of cobalt III is in cyanocobalamin (vitamin B₁₂). Cobalamin is not synthesized by animals but is synthesized by bacteria and actinomycetes (Sherman, 1957). Cobalamin is required by some green algae and diatoms, many dinoflagellates and yellow-green algae, higher plants, insects, fish, birds, and mammals.

Absorption efficiency of cobalt from food is about 50% in snails (Nelson and Malone, 1969) and about 5% in fishes. These low absorption efficiencies are in part responsible for the decrease in the cobalt bioaccumulation factor with increasing trophic level that is sometimes observed (Kevern and Griffith, 1966; Lowman, 1963; Morton, 1965; OpheL and Fraser, 1973). This is because the ratio of intake rate to elimination rate governs isotopic concentration of cobalt in the organism [Equations (2.4.3) and (2.4.4)].

Elimination rates for cobalt are given in Table 3.6.1. Good estimates of turnover rates come from studies in which the organisms have come into steady-state with the source of radionuclide. In such studies, long time components contain most of the radionuclide. Because of its relatively short radioactive half-life (72 days), ⁵⁸Co will have a lower bioaccumulation factor than the bioaccumulation factors for ⁶⁰Co (5.27 year half-life) and stable cobalt. However, as can be seen from the biological half-times given in Table 3.6.1,

Table 3.6.1 Biological elimination rates for ^{60}Co in various aquatic organisms

Species	Description	Biological Half-Time (days)	Elimination Rate (10^{-2} days $^{-1}$)	Comment	Percent of Total Radioactivity	Reference
<u><i>Lemna minor</i></u> (duckweed)	macrophyte	19	3.6	from elimination after five month uptake	100	6.27
<u><i>Vallisneria americana</i></u> (wild celery)	macrophyte	51	1.4	from elimination after five month uptake	100	6.27
<u><i>Helisoma</i></u> sp.	snail	51	1.4	from elimination after five month uptake	90	6.30
<u><i>Goniobasis clavaeformis</i></u>	snail	4.5 320	15. 0.22	fast compartment slow compartment snails tagged by short term feeding	25 25	6.28 6.25
<u><i>Lampsilis radiata</i></u>	clam	4.8 277	14.4 0.25	fast compartment slow compartment	51 49	6.14 6.14
<u><i>Cambarus longulus</i></u>	crayfish	70	0.99	5 g animals, high level dose	90	6.35
		37	1.9	0.9 g animals, high level dose	90	6.35
<u><i>Diemictylus viridescens</i></u>	newt	158	0.44	five month uptake	90	6.30
<u><i>Ictalurus melas</i></u> (black bullhead)	fish	1.5 40.5 long, undetermined 3.5 3.5 long, undetermined	46. 1.7 undetermined 20. 2.0 undetermined	single feeding single feeding single feeding 4 days of water uptake 4 days of water uptake 4 days of water uptake	97.9 1.8 0.3 50 35 15	6.32 6.32 6.32 6.32 6.32 6.32
<u><i>Hydropsyche</i></u> sp.	Insect Trichoptera larva	105.3 38.4	0.66 1.8	uptake from water tag uptake from food ingestion	89.3 90	6.13 6.13

the ^{58}Co bioaccumulation factors are not expected to be greatly lower than the bioaccumulation factors for ^{60}Co or stable cobalt in many organisms. Considering the uncertainty in both biological half-times and in the bioaccumulation factors for ^{60}Co and stable cobalt, we will not distinguish between the bioaccumulation factors of ^{58}Co and the longer-lived isotopes.

3.6.2 Environmental Cobalt and Availability

The proportion of cobalt in the particulate phase varies greatly among bodies of water. In Par Pond, an oligotrophic reservoir of low turbidity, 22% of the stable cobalt in the water is in the particulate phase (Marshall and LeRoy, 1974). Greater than 98% of the stable cobalt in the Amazon and Yukon Rivers is particulate. The cobalt appears as adsorbed on organic solids and in the crystal structure of sediments. Fukai and Murray (1973) contrast the fraction of radio-cobalt appearing in the particulate phase in the Columbia River to that in the Clinch River, Tennessee; the Columbia had from 80 to 95% (at Vancouver, Washington) and the eutrophic Clinch, from 2 to 30%, despite the much higher suspended solids concentration in the latter river. In Perch Lake, a small dystrophic-eutrophic lake, essentially all the radiocobalt in the water is in the dissolved phase (Ophel and Fraser, 1973) in contrast to mesotrophic Lake Maggiore in which roughly 80% of stable cobalt is in the particulate phase (Merlini et al., 1967). These observations would suggest that the proportion of cobalt in the particulate phase increases with increase in suspended solids concentration and decreases with

increase in eutrophy. This latter correlation may be related to the tendency of cobalt to form associations with dissolved organic matter. Thus, increased dissolved organic matter concentrations in eutrophic waters may serve to keep radiocobalt in solution.

As indicated in Section 2.2, the soluble form of a radionuclide is generally more available to algae and subsequent trophic levels than the particulate form of the radionuclide. Thus, it would seem that the greater the proportion of soluble cobalt, the greater the availability. However, as indicated in Section 2.2, chelated radionuclides are less available to food webs than radionuclides appearing as free ions. For this reason, soluble cobalt in eutrophic environments is probably less available than soluble cobalt in oligotrophic or mesotrophic environments due to the tendency of cobalt to chelate or form other associations with dissolved organic matter. This expected pattern is seen in the review of the literature on cobalt bioaccumulation factors which follows.

3.6.3 Review of Cobalt Bioaccumulation Factors

Because most investigators have based their bioaccumulation factors on filtered concentrations of cobalt isotopes in water, we generally report filtered bioaccumulation factors. As indicated in Section 2.5.3 use of filtered bioaccumulation factors should generally lead to overprediction of radionuclide concentration in organisms. However, we noted that in nonturbid waters and in eutrophic waters, most of the radiocobalt in the water would be

in the soluble phase. In these waters, use of filtered bioaccumulation factors would not overpredict radiocobalt concentrations in organisms.

3.6.3.1 Cobalt Bioaccumulation Factors for Fishes

Researchers have reported cobalt bioaccumulation factors for flesh and whole bodies of fishes. The relative concentrations of ^{60}Co in black bullheads from White Oak Lake may be used to convert these bioaccumulation factors to bioaccumulation factors for other tissues. The relative steady-state concentrations are given in Table 3.6.2. The organ with the highest cobalt concentration is the kidney, which has a concentration 23 times that of the whole body. Note that the flesh to whole body ratio is 0.3, which compares well with the ratio of 0.37 determined for brown bullheads from Perch Lake, Ontario (Ophel and Fraser, 1973).

All bioaccumulation factors for cobalt in fishes were derived from steady-state concentrations of ^{60}Co or stable cobalt in natural bodies of water. Because cobalt bioaccumulation factors are expected to decrease with increasing eutrophy, the trophic states of the bodies of water were recorded. The bodies of water fell into two categories, eutrophic and mesotrophic; bioaccumulation factors for the former are given in Table 3.6.3, and bioaccumulation factors for the latter, in Table 3.6.4. Mean bioaccumulation factors for whole fishes and fish flesh are given in Table 3.6.5. As expected, lower cobalt bioaccumulation factors are seen for eutrophic environments. We recommend use of the bioaccumulation factors given

Table 3.6.2 Relative steady-state concentrations of ^{60}Co in tissues of Ictalurus melas (black bullheads) from White Oak Lake. Calculated from data of Reed (1971)

Tissue	Tissue Concentration
	Whole Body Concentration
Blood	4.3
Skin	0.92
Flesh	0.30
Liver	3.1
Stomach	0.29
Gut	4.0
Kidney	23
Bone	0.52
Gills	2.3

Table 3.6.3 Bioaccumulation factors for cobalt in fishes from eutrophic environments
(based on filtered water concentrations)

Species	Size class (in.)	Description	Tissue	Stable element concentration in tissue (ppm)	Stable element concentration in water (ppb)	BF	Location	Reference
<u>Perca flavescens</u> (yellow perch)	Adult (8.7-13)	piscivorous	whole			18	Perch Lake, Canada	6.29
	Adult young (2.4-3.9)	zooplankton and insect larvae	flesh whole			9 130		6.29 6.29
<u>Ictalurus nebulosus</u> (brown bullhead)	Adult (6.7-11.8)	omnivorous, plant material	whole			52		6.29
	Adult young (2.0-2.8)		flesh whole			19 63		6.29 6.29
<u>Leiomis gibbosus</u> (pumpkinseed sunfish)	Adult (4.7-7.9)	omnivorous, plant	whole			80		6.29
	young (2.0-3.2)	eating insect larvae	whole			94		6.29
<u>Semotilus margarita</u> (pearl dace)	Adult (2.8-4.7)	unknown	whole			18		6.29
<u>Notropis heterolepis</u> (blacknose shiner)	Adult (2.0-3.9)	unknown	whole			20		6.29
<u>Hybopsis plumbea</u> (lake chub)	Adult (2.0-3.9)	unknown	whole			38		6.29
<u>Dorosoma cepedianum</u> (gizzard shad)		piscivorous	whole fish less viscera			32	White Oak Lake, Tennessee	6.26
<u>Lepomis macrochirus</u> (bluegill)		chironomid larvae, ter- restrial insects, fish, carnivorous	whole fish less viscera			26		6.26

Table 3.6.3 (continued)

Species	Size class (in.)	Description	Tissue	Stable element concentration in tissue (ppb)	Stable element concentration in water (ppb)	BF	Location	Reference
<u><i>Carassius auratus</i></u> (goldfish)		sediment, algae, bryozoans, chironomids, plant remains, omnivorous	whole fish less viscera			49		6.26
<u><i>Cyprinus carpio</i></u> (carp)	large adults	low trophic level (algae, benthic detritovore)	whole			6.4	White Oak Lake	6.17
<u><i>Ictalurus melas</i></u> (black bullhead)	large adults	benthic omnivore	whole			70		6.17
<u><i>Dorosoma cepedianum</i></u> (gizzard shad)	large adults	partially piscivorous	whole			13		6.17
<u><i>Lepomis macrochirus</i></u> (bluegill)	large adults	carnivore	whole			5.0		6.17
<u><i>Micropterus salmoides</i></u> (largemouth bass)	large adults	piscivorous	whole			5.5		6.17
<u><i>Cyprinus carpio</i></u> (carp)		bottom feeder	flesh			14		6.5, 6.25
			total ^a			10	Clinch River, Tennessee	6.5, 6.25
<u><i>Carpiodes carpio</i></u> (carpsucker)		bottom feeder	flesh			25		6.5, 6.25
<u><i>Ictiobus bubalus</i></u> (smallmouth buffalo)		bottom feeder	flesh			7		6.5, 6.25
			total			17		6.5, 6.25
						7		6.5, 6.25

Table 3.6.3 (continued)

Species	Size class (in.)	Description	Tissue	Stable element concentration in tissue (ppm)	Stable element concentration in water (ppb)	BF	Location	Reference
Assorted fish ^b		sight feeders	flesh			5		6.5, 6.25
<u><i>Esox lucius</i></u> (northern pike)		piscivorous	flesh	0.07(0.03-0.11) ^c	3 (1-6) ^c	23	Illinois River	6.22
<u><i>Micropterus</i></u> <u><i>salmoides</i></u> (largemouth bass)		piscivorous	flesh	0.09(0.06-0.18)		30		6.22
<u><i>Morone chrysops</i></u> (white bass)		piscivorous	flesh	0.07(0.05-0.10)		23		6.22
<u><i>Lepisosteus</i></u> <u><i>platostomus</i></u> (shortnose gar)		piscivorous	flesh	0.15(0.11-0.18)		50		6.22
<u><i>Micropterus</i></u> <u><i>dolomieu</i></u> (smallmouth bass)		piscivorous	flesh	0.15(0.14-0.16)		50		6.22
<u><i>Ictiobus</i></u> <u><i>cyprinellus</i></u> (bigmouth buffalo)		omnivorous insect larvae, mollusks, algae, aquatic plants	flesh	0.081(0.06-0.12)		27		6.22
<u><i>Dorosoma</i></u> <u><i>cepedianum</i></u> (gizzard shad)		omnivorous insect larvae, mollusks, algae, aquatic plants	flesh	0.16(0.1-0.25)		53		6.22
<u><i>Moxostoma macro-</i></u> <u><i>lippidotum</i></u> (northern redhorse)		omnivorous insect larvae, mollusks, algae, aquatic plants	flesh	0.083(0.05-0.11)		28		6.22

Table 3.6.3 (continued)

Species	Size class (in.)	Description	Tissue	Stable element concentration in tissue (ppm)	Stable element concentration in water (ppb)	BF	Location	Reference
<u><i>Carpioles cyprinus</i></u> (quillback)		omnivorous insect larvae, mollusks, algae, aquatic plants	flesh	0.087(0.04-0.12)		29		6.22
<u><i>Cyprinus carpio</i></u> (carp)		omnivorous insect larvae, mollusks, algae, aquatic plants	flesh	0.068(0.03-0.1)		23		6.22

^aTotal fish consists of flesh and bone.

^bAssorted fish are all sight feeders; *Pomoxis annularis* (white crappie), *Lepomis macrochirus* (bluegill), *Roccus chrysops* (white bass), *Micropterus salmoides* (largemouth bass), *Stizostedion v. vitreum* (sauger), *Aplodinotus grunniens* (drum), *Ictalurus punctatus* (catfish).

^cRange of concentrations given in parentheses.

Table 3.6.4 Bioaccumulation factors for cobalt in fishes from mesotrophic environments
(based on filtered water concentrations)

Species	Size class (in.)	Description	Tissue	Stable element concentration in tissue (ppm)	Stable element concentration in water (ppb)(filtered)	BF	Location	Reference
<u><i>Alosa pseudoharengus</i></u> (alewife)	0.6-0.8	zooplankton feeder	whole	0.029 ± 0.014	0.19 ± 0.02	190	Lake Michigan	6.6, 6.20
<u><i>Alosa pseudoharengus</i></u> (alewife)	4.5-10	zooplankton feeder benthic feeder	whole	0.065 ± 0.005	0.19 ± 0.02	420	Lake Michigan	6.6, 6.7
<u><i>Osmerus mordax</i></u> (smelt)	5	zooplankton feeder	whole	0.19 ± 0.05	0.19 ± 0.02	1000	Lake Michigan	6.6, 6.7
<u><i>Alburnus alborella</i></u> (bleak)	4.3-5.1	plankton feeder	flesh	0.012 ± 0.002	0.02	600	Lake Maggiore	6.23,6.24
<u><i>Coregonus macrophthalmus</i></u> (whitefish, Bondella)	9.8	plankton and benthic feeder	flesh	0.0077 ± 0.0016	0.02	385	Lake Maggiore	6.23,6.24
<u><i>Osmerus mordax</i></u> (smelt)	>5	plankton and benthic feeder	flesh	0.043 ± 0.004	0.19 ± 0.02	230	Lake Michigan	6.6, 6.7
			whole	0.091 ± 0.011	0.19 ± 0.02	480	Lake Michigan	6.6, 6.7
<u><i>Percopsis omiscomaycus</i></u> (trout-perch)	2.4-3.7	benthic	whole	0.024 ± 0.002	0.19 ± 0.02	130	Lake Michigan	6.6, 6.20
<u><i>Notropis hudsonius</i></u> (spottail shiner)	2.2-2.6	benthic and zoo-plankton	whole	0.042 ± 0.015	0.19 ± 0.0	220	Lake Michigan	6.6, 6.20
<u><i>Notropis hudsonius</i></u> (spottail shiner)	4.5-5.5	benthic	whole	0.12 ± 0.03	0.19 ± 0.02	630	Lake Michigan	6.6, 6.7
<u><i>Notropis hudsonius</i></u> (spottail shiner)	7.5-8	unknown	flesh	0.041 ± 0.003	0.19 ± 0.02	220	Lake Michigan	6.6, 6.7
<u><i>Scardinus erythrophthalmus</i></u> (rudd)	1	omnivorous	flesh	0.0058 ± 0.0011	0.02	290	Lake Maggiore	6.23,6.24
<u><i>Perca flavescens</i></u> (yellow perch)	8-13	partially piscivorous	flesh	0.053 ± 0.0089	0.19 ± 0.02	280	Lake Michigan	6.6, 6.7

Table 3.6.4 (continued)

Species	Size class (in.)	Description	Tissue	Stable element concentration in tissue (ppm)	Stable element concentration in water (ppb) (filtered)	BF	Location	Reference
<u><i>Salvelinus namaycush</i></u> (lake trout)	>15	piscivorous	flesh	0.045 ± 0.0061	0.19 ± 0.02	236	Lake Michigan	6.6, 6.7
<u><i>Salmo trutta</i></u> (brown trout)	>12	piscivorous	flesh	0.044 ± 0.0064	0.19 ± 0.02	230	Lake Michigan	6.6, 6.7
<u><i>Oncorhynchus kisutch</i></u> (coho salmon)	6-27	piscivorous	flesh	0.044 ± 0.0036	0.19 ± 0.02	230	Lake Michigan	6.6, 6.7
<u><i>Prosobium cylindraceum</i></u> (round whitefish)	>13	benthic (>10 inches)	flesh	0.040 ± 0.0038	0.19 ± 0.02	210	Lake Michigan	6.6, 6.7
<u><i>Ictalurus melas</i></u> (black bullhead)		benthic omnivore	flesh	0.030	0.046 ^a	650	East Tennessee spring	6.32

^a D. J. Nelson, private communication.

Table 3.6.5 Mean bioaccumulation factors for cobalt in fishes from mesotrophic and eutrophic waters

Nature of body of water	Number of bodies of water	Bioaccumulation factors	
		Flesh	Whole
Mesotrophic	3	323 ± 47(11)	439 ± 115(7)
Eutrophic	4	26.6 ± 3.5(16)	43.8 ± 10.4(14)

in Table 3.6.5 for mesotrophic and eutrophic environments. In the absence of data for oligotrophic waters, we suggest using the mean value from the mesotrophic waters.

Lucas and Edgington (1970) have suggested that cobalt concentration may be under partial homeostatic control. The inverse correlation between the cobalt bioaccumulation factor and eutrophy could be due to partial homeostatic control if cobalt concentration in the water increased with eutrophy. In Figure 3.6.1 the cobalt concentrations in fish flesh are plotted against cobalt concentrations in water for three mesotrophic waters and one eutrophic water, the Illinois River. If the slope (exponent) of the logarithmic regression were 1, the cobalt bioaccumulation factor would be independent of the cobalt concentration in water. Since the slope is less than 1, either partial homeostatic control or eutrophy coincident with increasing cobalt concentration in water may be operating to diminish the slope. Lack of more extensive data on bioaccumulation factors from environments exhibiting a wide range of cobalt concentrations within each of a few distinct eutrophy categories prevent us from firmly deciding between the hypotheses of partial homeostatic control and eutrophy. In view of the known tendency of cobalt to form organic associations, the latter explanation seems more plausible.

Some workers have observed that the cobalt bioaccumulation factor in fishes decreases with increase in trophic level (Morton, 1965; Nelson et al., 1971; Ophel and Fraser, 1973). Other workers have not found a clear-cut correlation (Kevern and Griffith, 1966;

ORNL-DWG 74-7032




Figure 3.6.1 Concentration of stable cobalt in fish flesh in various environments.

Mathis and Cummings, 1973). For this reason we have specified the feeding habits of the fishes in Tables 3.6.3 and 3.6.4. Based on the limited and conflicting data at hand, however, we cannot specify a relation between the cobalt bioaccumulation factor in fishes and their feeding habits.

3.6.3.2 Cobalt Bioaccumulation Factors for Plants

In Table 3.6.6 are listed cobalt bioaccumulation factors for algae and vascular plants. Compared to fishes the cobalt bioaccumulation factors of plants are high. On the basis of the few data on algae in Table 3.6.6, we recommend a cobalt bioaccumulation factor of 10^4 for algae. The two species of emergent vascular plants exhibit widely differing cobalt bioaccumulation factors. The low bioaccumulation factor was calculated from ^{60}Co concentrations, and the high bioaccumulation factor was calculated from stable element concentrations. Possibly, because diffusion of elements in interstitial water of sediment is slow, the ^{60}Co concentration in interstitial water from which Pontederia obtained its ^{60}Co had not reached steady state with the ^{60}Co in the water column above. Based on the stable cobalt bioaccumulation factors for Phragmites, we recommend a bioaccumulation factor of 10^3 for emergent vascular plants.

Physical form of aquatic vascular plants may affect the cobalt bioaccumulation factor. Merlini et al. (1971) reported that floating leaf species have lower bioaccumulation factors than submerged species. Ophel and Fraser (1973) noted that the finely divided submerged species had the highest bioaccumulation factors, that other submerged

Table 3.6.6 Bioaccumulation factors for cobalt in aquatic plants

Taxon	BF (filtered)	BF (unfiltered)	Location	Reference
Algae:				
blue-green algae	30,000		Connecticut River	6.2
diatoms	1,500		Lake Michigan	6.6
<u>Navicula seminulum</u> (diatom)	1,100-2,000		Laboratory	6.16
<u>Nitella</u> sp. (green alga)	500		Perch Lake	6.29
<u>Plectonema boryanum</u> (filamentous blue-green)	250-620		Laboratory	6.16
<u>Stigeoclonium lubricum</u> (filamentous green)	2,800		Laboratory	6.16
Emergent vascular plants:				
<u>Phragmites</u> sp.	2,000	400	Lake Maggiore	6.23
<u>Pontederia cordata</u>	20		Perch Lake	6.29
Vascular plants, floating leaves:				
<u>Nuphar</u> sp.	800	160	Lake Maggiore	6.23
<u>Nuphar variegatum</u>	200		Perch Lake	6.29
<u>Nymphaea lutea</u>	900	200	Lake Maggiore	6.24
<u>Nymphaea odorata</u>	200		Perch Lake	6.29
<u>Potamogeton amplifolius</u>	600		Perch Lake	6.29
<u>Potamogeton natans</u>	500		Perch Lake	6.29
<u>Sparganium fluctuans</u>	300		Perch Lake	6.29

Table 3.6.6 (continued)

Taxon	BF (filtered)	BF (unfiltered)	Location	Reference
Vascular plants, submerged leaves:				
<u>Ceratophyllum demersum</u>	1,000		Perch Lake	6.29
<u>Ceratophyllum</u> sp.	5,000	1,000	Lake Maggiore	6.24
<u>Elodea</u> sp.	15,000	3,000	Lake Maggiore	6.23
<u>Myriophyllum</u> sp.	12,000	2,400	Lake Maggiore	6.23
<u>Nojas</u> sp.	2,400	500	Lake Maggiore	6.23
<u>Potamogeton amplipolius</u>	2,000		Perch Lake	6.29
<u>Potamogeton pusillus</u>	2,800		Perch Lake	6.29
<u>Potamogeton</u> sp.	10,000	2,000	Lake Maggiore	6.23
<u>Utricula vulgaris</u>	600		Perch Lake	6.29
Mixed species (<u>Potamogeton</u> , <u>Myriophyllum</u> , <u>Vallisneria</u>)	5,000 ^a		Hudson River	6.18

^aGeometric mean of five years of data.

species had intermediate bioaccumulation factors, and that floating plants had the lowest. In Table 3.6.7 we give mean cobalt bioaccumulation factors for submerged and floating vascular plants from Lake Maggiore and Perch Lake. In Lake Maggiore the cobalt bioaccumulation factors for submerged plants are significantly greater than those for floating plants.

In Table 3.6.7 the mean cobalt bioaccumulation factor for submerged vascular plants from mesotrophic Lake Maggiore is significantly greater than the mean bioaccumulation factor for dystrophic-eutrophic Perch Lake. This conforms to the expected result of decreased cobalt bioaccumulation factors with increased eutrophy. We will assume the mean values given in Table 3.6.7 for Lake Maggiore and Perch Lake to be representative of mesotrophic and eutrophic waters, respectively. Thus, the cobalt bioaccumulation factors for submerged and floating vascular plants in eutrophic waters are 2×10^3 and 4×10^2 , respectively. For mesotrophic and oligotrophic waters we recommend cobalt bioaccumulation factors of 10^4 for submerged vascular plants and 10^3 for floating vascular plants.

3.6.3.3 Cobalt Bioaccumulation Factors for Invertebrates

The few cobalt bioaccumulation factors available for invertebrates are listed in Table 3.6.8. The cobalt bioaccumulation factor for the bivalve Unio from mesotrophic Lake Maggiore is about 25 times larger than the bioaccumulation factors for other bivalves from the eutrophic Illinois River and the dystrophic-eutrophic Perch Lake. We attribute this difference to differences in degree of eutrophy. For bivalves in eutrophic environments we recommend

Table 3.6.7 Mean cobalt bioaccumulation factors for submerged and floating-leaf vascular plants from Perch Lake and Lake Maggiore (based on filtered water concentrations)

Location	Bioaccumulation factors		Significance of difference between plant types
	Submerged	Floating	
Lake Maggiore	8,900 \pm 2,300(5)	850 \pm 50(2)	P < 0.05
Perch Lake	1,600 \pm 500(4)	360 \pm 81(5)	P \approx 0.10
Significance of differences between lakes	P < 0.05	P \approx 0.06	

Table 3.6.8 Bioaccumulation factors for cobalt in invertebrates (based on filtered water concentrations)

Taxon	Tissue	Feeding habits	Stable element concentration in tissue (ppm)	Stable element concentration in water (ppb)	BF	Location	Reference
Bivalves:							
<u>Amblema plicata</u>	soft	filter feeder	0.7	3	200	Illinois River	6.2
<u>Elliptio</u> sp.	soft	filter feeder			330	Perch Lake	6.9
<u>Fusconaia flava</u>	soft	filter feeder	1.2	3	400	Illinois River	6.22
<u>Quadrula quadrula</u>	soft	filter feeder	0.8	3	300	Illinois River	6.22
<u>Unio mancus</u>	soft	filter feeder	0.18-0.20	0.02	9,000-10,000	Lake Maggiore	6.24
Crustaceans:							
<u>Cambarus</u> sp. (crayfish)	whole				1,600	Perch Lake	6.29
Copepods	whole	filter feeders	0.134	0.19	700	Lake Michigan	6.6
<u>Mysis relicta</u>	whole	detritus	0.034	0.19	200	Lake Michigan	6,6, 6.10
Insect larvae:							
<u>Argyractis</u> sp.	whole				23,000 ^a	Columbia River	6.8
<u>Glossoma</u> sp.	whole				11,000 ^a	Columbia River	6.8
Hydrobaeninae	whole	periphyton			5,000 ^a	Columbia River	6.8
<u>Hydropsyche cockerelli</u>	whole	phytoplankton			15,000 ^a	Columbia River	6.8
Tendipedinae	whole	detritus			3,000 ^a	Columbia River	6.8
Snails:							
<u>Amnicola</u> sp.	whole				4,400	Perch Lake	6.29
<u>Stagnicola nuttalliana</u>	soft				9,000 ^a	Columbia River	6.9

Table 3.6.8 (continued)

Taxon	Tissue	Feeding habits	Stable element concentration in tissue (ppm)	Stable element concentration in water (ppb)	BF	Location	Reference
Snails (continued):							
<u><i>Viviparus ater</i></u>	soft	detritus	0.24-0.51	0.02	12,000-26,000	Lake Maggiore	6.24
Tubificids:							
<u><i>Limnodrilus hoffmeisteri</i></u> and <u><i>Tubifex tubifex</i></u>	whole	detritus	1.6	3	500	Illinois River	6.22

^aBased on approximate ⁶⁰Co concentration in water of 2 pCi/l (Watson, personal communication).

a cobalt bioaccumulation factor of 4×10^2 . For bivalves in mesotrophic or oligotrophic waters we recommend a cobalt bioaccumulation factor of 10^4 . The few data available for insect larvae and snails suggest a cobalt bioaccumulation factor of 10^4 for both of these groups. The cobalt bioaccumulation factor for tubificids in the eutrophic Illinois River is 500, a value that we will assume is characteristic of eutrophic environments. The few data for crustaceans suggest a cobalt bioaccumulation factor of 10^3 .

References for Section 3.6

The numbers in the left-hand margin correspond to the reference numbers used in the tables of section 3.6.

- 6.1 Anderson, E. D. and L. L. Smith, Jr., A synoptic study of 30 fish species from western Lake Superior, Univ. Michigan, Agricultural Experiment Station, Technical Bulletin 279 (1971).
- 6.2 Blanchard, R. L. and B. Kahn, The fate of radionuclides discharged from a PWR nuclear power station into a river, Symposium on Environmental Behavior of Radionuclides Released in the Nuclear Industry, IAEA/SM-172/26 (1973).
- 6.3 Bowen, H. J. M., Trace elements in biochemistry, Academic Press, New York, 1966.
- 6.4 Carlander, K. D., Handbook of freshwater fishery biology, Vol. 1, Iowa State Univ. Press, Ames, Iowa (1969).
- 6.5 Churchill, M. A., J. S. Cragwall, Jr., R. W. Andrew, and S. L. Jones, Concentrations, total stream loads, and mass transport of radionuclides in the Clinch and Tennessee Rivers, ORNL-3721, Suppl. 1 (1965).
- 6.6 Copeland, R. A., and J. C. Ayers, Trace element distributions in water sediment, phytoplankton, zooplankton and benthos of Lake Michigan, Environmental Research Group Special Report No. 1, Great Lakes Research Division, Univ. Michigan, Ann Arbor (1972).
- 6.7 Copeland, R. A., R. B. Beethe, and W. W. Prater, Trace element distributions in Lake Michigan fish, Environmental Research Group Special Report No. 2, Great Lakes Research Division, Univ. Michigan, Ann Arbor (1973).
- 6.8 Davis, J. J., Accumulation of radionuclides by aquatic insects, IN Biological problems in water pollution, p. 211, U.S. Dept. Health, Education and Welfare, Madison, Wisconsin (1965).
- 6.9 Davis, J. J., R. W. Perkins, R. F. Palmer, W. C. Hanson, and J. F. Cline, Radioactive materials in aquatic and terrestrial organisms exposed to reactor effluent water, Proc. 2nd Intern. Conf. on The Peaceful Uses of Atomic Energy, United Nations, Geneva, Sept. 1-12, 1958, 18, 423-428 (1958).

References for Section 3.6 (continued)

- 6.10 Edgington, D. N., E. Harvey, and M. S. Torrey, A preliminary study of the chemical composition of the opossum shrimp, Mysis relicta, IN Radiological and Environmental Research Division Ann. Report, R. E. Rowland, ANL 7960, Part III (1972).
- 6.11 Friend, A. G., The aqueous behavior of strontium-85, cesium-137, zinc-65, and cobalt-60 as determined by laboratory type studies in transport of radionuclides in freshwater systems, Conf. in Austin, Texas, Jan. 30-Feb. 1, 1973, TID-7664.
- 6.12 Fukai, R., and C. M. Murray, Environmental behavior of radio-cobalt on radiosilver released from nuclear power stations into aquatic systems, IN Symposium on Environmental Behavior of Radionuclide Released in the Nuclear Industry, IAEA/SM-172/42 (1973).
- 6.13 Gammon, J. R., Determination of feeding rates in Trichoptera using ^{60}Co , IN Radiation Ecology Section, Health Physics Division, Annual Progress Report, ORNL-4446 (1970).
- 6.14 Harvey, R. S., Uptake and loss of radionuclides by the freshwater clam Lampsilis radiata (gmel.), Health Phys. 17, 149-154 (1969).
- 6.15 Harvey, R. S., Effects of temperature on the sorption of radionuclides by a blue-green alga, Proc. Second National Symposium on Radioecology, D. J. Nelson and F. C. Evans (eds.), CONF-676503 (1969).
- 6.16 Harvey, R. S., Temperature effects on the sorption of radio-nuclides by freshwater algae, Health Phys. 19, 293 (1970).
- 6.17 Kevern, N. R., and N. A. Griffith, Effect of trophic level on radionuclide accumulation by fish, IN Radiation Ecology Section, Health Physics Division, Annual Progress Report, ORNL-4007 (1966).
- 6.18 Lentsch, J. W., N. E. Wrenn, T. S. Kneip, and M. Eisenbud, Manmade radionuclides in the Hudson River Estuary, Fifth Ann. Health Phys. Soc. Midyear Topical Symposium, Idaho Falls, Idaho (1970).
- 6.19 Lowman, F. G., Iron and cobalt in ecology, IN Proc. First National Symposium in Radioecology, V. Schultz and A. W. Klement, Jr. (eds.), Reinhold Publ. Corp., New York (1963).

References for Section 3.6 (continued)

- 6.20 Lucas, H. F., Jr. and D. N. Edgington, Concentrations of trace elements in Great Lakes fishes, *J. Fish. Res. Bd. Can.* 27, 677 (1970).
- 6.21 Marshall, J. S. and J. H. LeRoy, Iron, manganese, cobalt, and zinc cycles in a South Carolina reservoir, IN *Proc. Third National Symposium on Radioecology*, D. J. Nelson (ed.), CONF-710501 (1973).
- 6.22 Mathis, B. J. and T. F. Cummings, Selected metals in sediments, water, and biota in the Illinois River, *J. Water Poll. Contr. Fed.* 45, 1573 (1973).
- 6.23 Merlini, Margaret, Carla Bigliocca, A. Berg, and G. Pozzi, Trends in the concentration of heavy metals in organisms of a mesotrophic lake as determined by activation analysis, IN *Proc. Symp. on Use of Nuclear Techniques in the Measurement and Control of Environmental Pollution*, IAEA/SM-142a/27 (1971).
- 6.24 Merlini, Margaret, F. Girardi, and G. Pozzi, Activation analysis in studies of an aquatic ecosystem, IN *Proc. Symp. on Nuclear Activation Techniques in the Life Sciences*, IAEA/SM-91/7 (1967).
- 6.25 Morton, R. J., Status report no. 5 on Clinch River study, ORNL-3721, Oak Ridge, Tennessee (1965).
- 6.26 Nelson, D. J., N. A. Griffith, J. W. Gooch, and S. A. Rucker, White Oak Lake Studies, IN *Ecological Sciences Division Annual Report*, July 31, 1970, ORNL-4634 (1971).
- 6.27 Nelson, D. J., N. A. Griffith, and S. A. Rucker, Radionuclide excretion studies, IN *Radiation Ecology Section, Health Physics Division, Annual Progress Report*, July 31, 1964. ORNL-4446 (1970).
- 6.28 Nelson, D. J. and C. R. Malone, Feeding rates of snails determined with ^{60}Co , IN *Radiation Ecology Section, Health Physics Division Annual Progress Report*, July 31, 1968, ORNL-4316 (1969).
- 6.29 Ophel, I. L., and C. F. Fraser, The fate of cobalt-60 in a natural freshwater ecosystem, IN *Proc. Third National Symposium on Radioecology*, D. J. Nelson (ed.), CONF-710501 (1973).

References for Section 3.6 (continued)

- 6.30 Patten, B. C., N. A. Griffith, S. A. Rucker, and D. J. Nelson, Aquarium experiments, IN Radiation Ecology Section, Health Physics Division Annual Progress Report, July 31, 1968, ORNL-4316 (1969).
- 6.31 Price, J. W., A study of the food habits of some Lake Erie fish, Bull. Ohio Biological Survey, New Series, Vol. II, No. 1, Ohio State Univ., Columbus (1963).
- 6.32 Reed, J. R., Uptake and excretion by ^{60}Co by black bullheads Ictalurus melas (Rafinesque), Health Phys. 21, 835 (1970).
- 6.33 Sherman, H. C., Essentials of nutrition, MacMillian Co., New York (1957).
- 6.34 Timofeev-Resovskii, N. V., E. A. Timofeeva-Resovskaya, G. A. Milyutinaya and A. B. Getsova, Coefficients of accumulation of radioactive isotopes of 16 different elements by freshwater organisms and the effect of EDTA on some of them, Doklady (Biol. Sci. Sect.) 132, 369 (1960).
- 6.35 Wiser, C. W. and D. J. Nelson, Uptake and elimination of ^{60}Co by crayfish, Amer. Midl. Natur. 72, 181 (1964).

BIBLIOGRAPHY

Abee, H. H., Biological phases of the Clinch River study, IN Clinch River study status report no. 5, R. J. Morton (ed.), pp. 89-106, ORNL-3721, TID-4500 (1965).

Agnedal, P. O., Calcium and strontium in Swedish waters and fish and accumulation of strontium-90, IN Radioecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 879-896 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1966.

Anderson, E. D. and L. L. Smith, Jr., A synoptic study of 30 fish species from western Lake Superior, Univ. Michigan, Agricultural Experiment Station, Technical Bulletin 279 (1971).

Andrushaitis, G. and Z. Kalnina, The cycling of radioactive sediments (strontium-90) in the lakes of Latvian SSR, Verh. Internat. Verein. Limnol. 18, 995-1000 (1972).

Austin, J. H., Strontium uptake by algae, IN Transport of radio-nuclides in freshwater systems, B. H. Kornegay, W. A. Vaughn, D. K. Jamison, and J. M. Morgan (eds.), pp. 231-245, TID-7664 (1963).

Ayers, J. C., Lake Michigan environmental survey final report, Special Report No. 49, Great Lakes Research Division, Univ. Michigan, Ann Arbor, 1970.

Baker, K. F., O. Berg, A. Gorbman, R. F. Nigrelli, and M. Gordon, Functional thyroid tumors in the kidneys of platyfish, *Cancer Res.* 15, 118-123 (1955).

Baptist, J. P., D. E. Hoss, and C. W. Lewis, Retention of Cr-51, Fe-59, Co-60, Zn-65, Sr-85, Nb-95, In-113m, and I-131 by the Atlantic croaker (Micropogon undulatus), *Health Phys.* 18, 141-148 (1970).

Bauer, G. C. H., A. Carlsson, and B. Lindquist, A comparative study on the metabolism of strontium-90 and calcium-145, *Acta. Physiol. Scand.* 35, 55-66 (1955).

Beninson, D. E., E. Van der Elst, and D. Cancio, Biological aspects in the disposal of fission products into surface waters, IN Disposal of radioactive wastes into seas, oceans, and surface waters, pp. 337-354, Proceedings of International Atomic Energy Agency Symposium, Vienna, 1966, STI/PUB/126.

Bergstrom, S. O. W., Strontium levels in Swedish fish, *Health Phys.* 21, 611-612 (1971).

Blanchard, R. L., and B. Kahn, The fate of radionuclides discharged from a PWR nuclear power station into a river, Symposium on Environmental Behavior of Radionuclides Released in the Nuclear Industry, France, May 14-18, 1973, IAEA/SM-172/26.

Blaylock, B. G., Chromosome aberrations in Chironomus riparius developing in different concentrations of tritiated water, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 1169-1173 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 11-12, 1971.

Bleyer, B., Zur Kenntnis der Iodes als Biogenes Element, Biochem. Z. 170, 256 (1926).

Bockrath, R., S. Person, and F. Funk, Calculated energy deposits from the decay of tritium and other radioisotopes incorporated into bacteria, Biophys. J. 8, 1027-1036 (1968).

Bogen, D. C., et al., Tritium intake in New York City, IN Tritium, A. A. Moghissi and M. W. Carter (eds.), pp. 639-646 (1973), Messinger Graphics, Phoenix, TID CONF-710809.

Bolin, B., Note on the exchange of iodine between the atmosphere, land and sea, Int. J. Air Poll. 2, 127-131 (1959).

Bond, V. P. and L. E. Feinendegen, Intranuclear tritiated thymidine: Dosimetric radiological and radiation protection aspects, Health Phys. 12, 1007-1020 (1966).

Bortoli, M. de, P. Gaglione, A. Malvicini, and C. Polvani, Concentration factors for strontium and cesium in fish of the lakes in the region of Varese (Northern Italy) Minerva, Fisiconucleare, 324-331 (1967).

Bortoli, M. de, P. Gaglione, A. Malvicini, and E. Van der Stricht, Environmental Radioactivity Report EUR-3554e, Ispra (1966).

Bortoli, M. de, P. Gaglione, O. Ravera, G. de Fisica, Fallout manganese-54 and zinc-65 in water and fishes of four lakes in northern Italy, Sanilaria e Protizone Controle Radiazione 13, 72-77 (1969).

Bovard, P. and A. Grauby, The fixation of radionuclides from atmospheric fallout in peat-bog Sphagnum sp., Polytrichum and Myriophyllum, IN Radioecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 533-537 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1966.

Bowen, H. J. M., Trace elements in biochemistry, Academic Press Inc., New York, 1966.

Bowers, C. Y., A. Segaloff, and B. Brown, Factors affecting the thyroid gland uptake of iodine-131 of the Rana catesbeiana tadpole, Endocrinology 65, 882-920 (1959).

Bradley, W. O., Radioactive iodine as an indicator of the uptake of iodine by the liver, gastrocnemius and thyroid of Rana pipiens, *Science* 117, 483-484 (1953).

Broda, E., Radioactive Isotopes in Biochemistry, American Elsevier Publishing Co., New York, 1960.

Brokskii, A. E., "The exchange of hydrogen with deuterium in solution," *Trans. Faraday Soc.* 33, 1180 (1937).

Bruner, H. D., Distribution of tritium between the hydrosphere and invertebrates, IN Tritium, A. A. Moghissi and M. W. Carter (eds.), pp. 303-313 (1973), Messinger Graphics, Phoenix, TID CONF-710809.

Brungs, W. A., Experimental uptake of strontium-85 by freshwater organisms, *Health Phys.* 11, 41-46 (1965).

Bryan, G. W., The accumulation of cesium-137 by brackish water invertebrates and its relation to the regulation of potassium and sodium, *J. Mar. Biol. Assoc. U. K.* 43, 541-565 (1963).

Bryan, G. W. and E. Ward, The absorption and loss of radioactive and nonradioactive manganese by the lobster, Homarus vulgaris, *J. Mar. Biol. Assoc. U. K.* 45, 65-95 (1965).

Bryan, G. W., A. Preston, and W. L. Templeton, Accumulation of radio-nuclides by aquatic organisms of economic importance in the United Kingdom, IN Disposal of radioactive wastes into seas, oceans, and surface waters, pp. 623-636, Proceedings of International Atomic Energy Agency Symposium, Vienna, 1966, STI/PUB/126.

Butler, H. L. and J. H. LeRoy, Observations on the biological half-life of tritium, *Health Phys.* 11, 283-285 (1965).

Cameron, A. T., Contributions to the biochemistry of iodine. I. The distribution of iodine in plant and animal tissues, *J. Biol. Chem.* 18, 335-381 (1914).

Cameron, A. T., Contributions to the biochemistry of iodine. II. The distribution of iodine in plant and animal tissues, *J. Biol. Chem.* 23, 1-39 (1915).

Carlander, K. D., Handbook of freshwater fishery biology, Vol. 1, Iowa State Univ. Press, Ames, Iowa.

Causeret, J., Fish as a source of mineral nutrition, IN Fish as food, G. Borgstrom (ed.), pp. 205-234, Academic Press, New York, 1962.

Cavallor, R. and M. Merlini, Stable manganese and fallout radio-manganese in animals from irrigated ecosystems of the Po Valley, *Ecology* 48, 924-928 (1967).

Chavin, W., Thyroid distribution and function in the goldfish, Carassius auratus L., J. Exp. Zool. 133, 259-280 (1956).

Chavin, W. and B. Bouwman, Metabolism of iodine and thyroid hormone synthesis in the goldfish, Carassius auratus L., Gen. Comp. Endocr. 5, 493-503 (1965).

Churchill, M. A., J. S. Cragwall, Jr., R. W. Andrew, and S. L. Jones, Concentrations, total stream loads, and mass transport of radionuclides in the Clinch and Tennessee rivers, ORNL-3721, Suppl. 1, 1965.

Closs, K., Über das Vorkommen des Jods im Meere und in Meeresorganismen, Arch. Math. Naturv. 40, 1-150 (1931), IN Vinogradov 1953.

Cohen, L. K., and T. J. Kneip, Environmental tritium studies at a PWR power plant, IN Tritium, A. A. Moghissi and M. W. Carter (eds.), pp. 639-646 (1973), Messinger Graphics, Phoenix, TID CONF-710809.

Comar, C. L., R. H. Wasserman, and M. M. Mold, Strontium-calcium discrimination factors in the rat, Proc. Soc. Exptl. Biol. Med. 92, 859-863 (1956).

Copeland, R. A., and J. C. Ayers, Trace element distribution in water, sediment, phytoplankton, zooplankton, and benthos of Lake Michigan: A baseline study with calculations of concentration factors and build-up of radioisotopes in the food web, Environmental Research Group Special Report No. 1, Great Lakes Research Division, Univ. Michigan, Ann Arbor, 1970.

Copeland, R. A., R. H. Beethe, and W. W. Prater, Trace element distribution in Lake Michigan fish: A baseline study with calculations of concentration factors and equilibrium radioisotopes distribution, Environmental Research Group Special Report No. 2, Great Lakes Research Division, Univ. Michigan, Ann Arbor, March, 1973.

Cross, F. A. and J. H. Brooks, Concentrations of manganese, iron, and zinc in juveniles of five estuarine-dependent fishes, IN Radio-nuclides in ecosystems, D. J. Nelson (ed.), pp. 769-775 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Cross, F. A., L. H. Hardy, N. Y. Jones, and R. T. Barber, Relation between total body weight and concentrations of manganese, iron, copper, zinc, and mercury in white muscle of bluefish (Pomatomus saltatrix) and a bathy-demersal fish Antimora rostrata, J. Fish. Res. Bd. Can. 30, 1287-1290 (1973).

Crowson, D. L., Man-made tritium, IN Tritium, A. A. Moghissi and M. W. Carter (eds.), pp. 23-27 (1973), Messinger Graphics, Phoenix, TID CONF-710809.

Cushing, C. E., Radiation ecology in freshwater communities, IN Man and aquatic communities, pp. 45-56, Seminar conducted by Water Resources Research Institute, Oregon State Univ., Corvallis, 1970.

Davis, J. J., Accumulation of radionuclides by aquatic insects, IN Biological problems in water pollution, pp. 211-215 (1965), Third Seminar, Research Branch, Robert A. Taft Sanitary Engineering Center, U.S. Department of Health, Education, and Welfare, Madison, Wisconsin, 1962.

Davis, J. J., Cesium and its relationships to potassium in ecology, IN Radioecology, V. Schultz and A. W. Klement, Jr. (eds.), pp. 539-556 (1963), First National Symposium on Radioecology, September 10-15, 1961.

Davis, J. J., and R. F. Foster, Bioaccumulation of radioisotopes through aquatic food chains, *Ecology* 39, 530-535 (1958).

Davis, J. J., R. W. Perkins, R. F. Palmer, W. C. Hanson, and J. F. Cline, Radioactive materials in aquatic and terrestrial organisms exposed to reactor effluent water, IN Waste treatment and environmental aspects of atomic energy, pp. 425-428, Vol. 18, Proceedings of the Second United Nations International Conference on the Peaceful Uses of Atomic Energy, September 1-13, 1958.

Dewey, W. C., R. M. Humphrey, and B. A. Jones, Comparisons of tritiated thymidine, tritiated water, and cobalt-60 gamma rays in inducing chromosome aberrations, *Rad. Res.* 24, 214-238 (1965).

Drury, D. E. and J. G. Eales, The influence of temperature on histological and radiochemical measurements of thyroid activity in the eastern brook trout, Salvelinus fontinalis Mitchell, *Can. J. Zool.* 46, 1-9 (1968).

Dundee, H. and A. Gorbman, Utilization of radioiodine by the thyroid of neotenic salamander, Eurycea tynerensis Moore and Hughes, *Physiol. Zool.* 33, 58-63 (1960).

Eales, J. G., The influence of temperature on thyroid histology and radioiodine metabolism of yearling steelhead trout, Salmo gairdneri, *Can. J. Zool.* 42, 829-841 (1964).

Edgington, D. N., E. Harvey, and M. S. Torrey, A preliminary study of the chemical composition of the opossum shrimp, Mysis relicta, IN Radiological and Environmental Research Division Ann. Report, R. E. Rowland, ANL 7960, Part III, 1972.

Eidinoff, M. L., et al., "Equilibrium between protium, deuterium, and tritium in the system: Hydrogen-acetic acid; isotopic fractionation factors in a catalytic hydrogenation," *J. Amer. Chem. Soc.* 74, 5280 (1952).

Elwood, J. W., Ecological aspects of tritium behavior in the environment, Nuclear Safety 12, 326-327 (1971).

Elwood, J. W., Tritium behavior in fish from a chronically contaminated lake, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 435-439 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Englander, S. W., "A hydrogen exchange method using tritium and sephadex: Its application to ribonuclease," Biochem. 2, 798 (1963).

Englander, S. W. and A. Poulsen, Hydrogen-tritium exchange of the random chain polypeptide, Biopolymers 7, 379-393 (1969).

Ettenhuber, E., Akkumulation von Radionukliden in Fischen, Isotopenpraxis 7, 75-84 (1971).

Evans, A. G., New dose estimates from chronic tritium exposures, Health Phys. 16, 57-63 (1969).

Ewing, R. A., J. E. Howes, and R. B. Price, Bioenvironmental and radio-logical-safety feasibility studies Atlantic-Pacific Interoceanic Canal: Radionuclide and stable-element analysis of environmental samples from Routes 17 and 25, Battelle Memorial Institute, BMI-171-29 (1969), also IOCS-MEMO-BMI-33 (1969).

Eyman, L. D., Cesium-137 and stable cesium in a hypereutrophic lake, Ph.D. Thesis, Michigan State University, East Lansing, 98 pp. (1972).

Eyman, L. D. and J. T. Kitchings, Availability of ^{137}Cs to fishes from ingested clays, IN Proceedings of the XIX Congress of the International Association of Limnology (SIL), Winnipeg, Canada, August 22-29, 1974 (in press).

Feldt, W., Research on the maximum radioactive burden of some German rivers, IN Environmental aspects of nuclear power stations, pp. 495-506, Proceedings of a Symposium organized by IAEA with USAEC, August 10-14, 1970.

Feldt, W. and J. Lange, Der Strontium-90 und Cesium-137 Gehalt von Brassen aus Norddeutschen Binnengewässern, Z. Lebensm. Unters. und Forsh. 177, 103-113 (1962).

Feinendegen, L. E., Tritium-labeled molecules in biology and medicine, Academic Press, New York, 1967.

Fellenberg, Th. Von., Untersuchungen über das Vorkommen von Jod in der Natur, Mitt. Lebensm. Hyg., Bern 14, 161 (1923).

Fellenberg, Th. Von., Das Vorkommen der Kreslauf und der Stoffwechsel des Jods, Ergebn. Physiol. 25, 176-363 (1926).

Findenegg, G. R., H. Paschinger, and E. Broda, Investigation of the light dependence of the uptake of rubidium, zinc, cobalt, lead and cerium by Chlorella by a flow method, *Planta (Berl.)* 99, 163-173 (1971).

Fitch, F. W., K. Dzpga, H. Benensohn, and P. V. Harper, The use of iodine-125 as a label in immunologic studies, IN Second Annual Oak Ridge Radioisotope Conference, Oak Ridge, Tennessee, April 19-22, TID-7689 (1964).

Fleishman, D. J., Accumulation of artificial radionuclide in freshwater fish, IN Radioecology, V. M. Klechkovskii, G. G. Polikarpov, and R. M. Aleksakhin (eds.), pp. 347-369, John Wiley and Sons, New York, 1973.

Fontain, M., F. Lachiver, S. Leloup, and M. Olivereau, La Fonction thyroidienne du Saumon (Salmo salar L.) au cours de sa migration reproductrice, *J. Physiol. Paris* 40, 182-184 (1948).

Foster, R. F., Tritium oxide absorption and retention in the body water of some aquatic organisms, IN Biology research annual report for 1954, H. A. Kornberg (ed.), pp. 98-103, Hanford Atomic Products Operation, Richland, Washington, HW-35917 (1955).

Foster, R. F., Relationships between the concentration of radionuclides in Columbia River water and fish, IN Transport of radionuclides in freshwater systems, B. H. Kornegay, W. A. Vaughn, D. K. Jamison, and J. M. Morgan (eds.), pp. 269-293, TID-7664 (1963).

Friend, A. G., The aqueous behavior of strontium-85, cesium-137, zinc-65, and cobalt-60 as determined by laboratory type studies, IN Transport of radionuclides in freshwater systems, B. H. Kornegay, W. A. Vaughn, D. K. Jamison, and J. M. Morgan (eds.), pp. 43-60, TID-7664 (1963).

Friend, A. G., A. H. Story, C. R. Henderson, and K. A. Bush, Behavior of certain radionuclides released into freshwater environments, U.S. Department of Health, Education, and Welfare, Annual Report 1959-1960, Public Health Service Publ. No. 999-RH-13.

Fujita, M., J. Iwamoto, and M. Kondo, Comparative metabolism of cesium and potassium in mammals--Interspecies correlation between body weight and equilibrium level, *Health Phys.* 12, 1237-1247 (1966).

Fukai, R. and C. N. Murray, Environmental behavior of radiocobalt and radiosilver released from nuclear stations into aquatic systems, IN Symposium on Environmental Behavior of Radionuclides Released in the Nuclear Industry, France, May 14-18, 1973, IAEA/SM-172/42.

Funk, F. and S. Person, Cytosine to thymine transitions from decay of cytosine-5-tritium in bacteriophage S13, *Science* 166, 1629-1631 (1969).

Gaglione, P. and O. Ravera, Manganese-54 concentrations in fallout, water and Unio mussels of Lake Maggiore, 1960-1963, *Nature* 204, 1215-1216 (1964).

Gallegos, A. F., Radiocesium kinetics in the components of a montane lake ecosystem, Ph.D. Thesis, Colorado State Univ., Fort Collins, December 1969.

Gallegos, A. F., and F. W. Whicker, Radiocesium retention by rainbow trout as affected by temperature and weight, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 361-371 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Gallegos, A. F., F. W. Whicker, and T. E. Hakonson, Accumulation of radiocesium in rainbow trout via a non-foodchain pathway, IN Proceedings of Fifth Annual Health Physics Society Midyear Topical Symposium: Health Physics Aspects of Nuclear Siting, Vol. 2, pp. 477-497, AEC CONF-701106 (1970).

Gammon, J. R., Determination of feeding rates in Trichoptera using ⁶⁰Co, IN Radiation Ecology Section, Health Physics Division, Annual Progress Report, ORNL-4446 (1970).

Garder, K., and O. Skulberg, Sorption phenomena of radionuclide to clay particles in river water, *Int. J. Air Wat. Poll.* 8, 229-241 (1964).

Garder, K., and O. Skulberg, An experimental investigation on the accumulation of radioisotopes by freshwater biota, *Arch. Hydrobiol.* 62, 50-69 (1966).

Gertz, S. M., The fate of a radionuclide in the aquatic environment--A mathematical equation representing radionuclide uptake by algae, Ph.D. Thesis, Drexel University, Philadelphia, 112 pp., 1973.

Gibbs, R. J., Mechanisms of trace metal transport in rivers, *Science* 180, 71-73 (1973).

Gibbs, W., E. Wilson, H. Hodges, and C. C. Lushbaugh, Comparative study of thyroid and whole body retention in box turtles, IN Second Annual Oak Ridge Radioisotope Conference, pp. 20-25, Oak Ridge, Tennessee, April 19-22, 1964, TID-7689.

Glasser, R., Sur Frage der Anreicherung von Radiojod (J-131) in der Muschel Dreissensia polymorpha Pall., *Naturwissenschaften* 47, 284 (1960).

Glasser, R., Über die Aufnahm von Radiojod aus wasseriger Lösung durch die Muschel Dreissensia polymorpha Pall., *Zool. Jb. Physiol.* 69, 339-378 (1961). In Polikarpov 1966.

Gloyna, E. P., Y. A. Yousef, and T. J. Padden, Radioactivity transport in water-continuous release of radionuclides in a small scale ecosystem, Department of Civil Engineering, Univ. Texas, Austin, ORO-490-21, Report No. EHE-71-1, CRER-75 (1971).

Gorbman, A., M. Clements, and R. O'Brien, Utilization of radioiodine by invertebrates, with special study of several Annelida and Mollusca, *J. Exp. Zool.* 127, 75-92 (1954).

Grimm, R. M., Iodine content of some marine algae, *Pacific Sci.* 6, 318-319 (1952).

Gustafson, P. F., Comments on radionuclides in aquatic ecosystems, IN Radioecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 853-858 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1966.

Gustafson, P. F., Cesium-137 in freshwater fish during 1954-1965, IN Symposium on radioecology, D. J. Nelson and F. C. Francis (eds.), pp. 249-257 (1969), Proceedings of the Second National Symposium on Radioecology, Ann Arbor, Michigan, May 15-17, 1967.

Gustafson, P. F., S. S. Brar, and S. E. Muniak, Cesium-137 in edible freshwater fish, *Nature* 211, 843-844 (1966).

Gustafson, P. F., A. Jarvis, S. S. Brar, D. N. Nelson, and S. M. Muniak, Investigations of cesium-137 in freshwater ecosystems, pp. 315-327, AEC Report ANL-7136 (1965).

Gutman, J. R. and M. Wolfsberg, "Equilibrium in the exchange of tritium between ammonia and hydrogen and the zero-point energy difference between NH_3 and NH_2T ," *J. Chem. Phys.* 33, 1592 (1960).

Hakonson, T. E., A. F. Gallegos, and F. W. Whicker, Use of cesium-133 and activation analysis for measurement of cesium kinetics in a montane lake, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 344-348 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Halford, J. O. and E. Pecherer, "Exchange of deuterium between methanol and water. Vibrations of the hydroxyl group in methanol and methanol-d. The entropy of methanol," *J. Chem. Phys.* 6, 571 (1938).

Hampson, M. A., Uptake of radioactivity by aquatic plants and location in the cells. I. The effect of pH on the strontium-90 and yttrium-90, *J. Exp. Bot.* 18, 17-33 (1967).

Hannerz, L., Fallout cesium-137 in fish and plankton from Lake Malar and the Baltic, Excerpt *Acta Radiol. Sup.* 254, 22-28 (1966).

Hannerz, L., The role of feeding habits in the accumulation of fallout cesium-137 in fish, Institute of Freshwater Research, Drottningholm 48, 112-119 (1968).

Hanson, W. C., Iodine in the environment, IN Radioecology, V. Schultz and A. W. Klement, Jr. (eds.), pp. 581-601 (1963), Proceedings of the First National Symposium on Radioecology, Fort Collins, Colorado, September 10-15, 1961.

Harrison, F. L. and J. J. Koranda, Tritiation of aquatic animals in an experimental freshwater pool, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 425-434 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Harvey, R. S., Uptake of radionuclides by freshwater algae and fish, Health Phys. 10, 243-247 (1964).

Harvey, R. S., Uptake and loss of radionuclides by the freshwater clam Lampsilis radiata, Health Phys. 17, 149-154 (1969).

Harvey, R. S., Effects of temperature on the sorption of radionuclides by a blue-green algae, IN Symposium on radioecology, D. J. Nelson and F. C. Evans (eds.), pp. 266-269 (1969), Proceedings of the Second National Symposium on Radioecology, Ann Arbor, Michigan, May 15-17, 1967.

Harvey, R. S., Temperature effects on the sorption of radionuclides by freshwater algae, Health Phys. 19, 293-297 (1970).

Harvey, R. S., Temperature effects on the maturation of midges (Tendipedidae) and their sorption of radionuclides, Health Phys. 20, 613-616 (1971).

Hasanen, E. and J. K. Miettinen, Cesium-137 content of freshwater algae and fish in Finland, Nature 200, 1018-1019 (1963).

Hasanen, E., S. Kolehmainen, and J. K. Miettinen, Biological half-time of cesium-137 in three species of freshwater fish: Perch, roach and rainbow trout, IN Radioecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 921-924 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1966.

Hatch, F. T. and J. A. Mazrimas, Tritiation of animals from tritiated water, Radiat. Res. 50, 339-357 (1972).

Helvey, T. C., The national concentration of deuterium in honey, Science 117, 276-277 (1953).

Henderson, C. and R. F. Foster, Studies of smallmouth bass (Micropterus dolomieu) in the Columbia River near Richland, Washington, Trans. Am. Fish. Soc. 86, 112-127 (1956).

Hercus, C. E. and K. C. Roberts, The iodine content of foods, manures, and animal products in relation to the prophylaxis of endemic goitre in New Zealand, *J. Hyg., Camb.* 26, 49 (1927).

Hickman, C. P., Influence of environment on the metabolism of iodine in fish, *Ann. N.Y. Acad. Sci., Gen. Comp. Endocr. Sup.* 1, 48-62 (1962).

Hiyama, Y., M. Saiki, H. Kamada, T. Koyanagi, and Y. Kurosawa, Hazard evaluation of radioactive contamination of the sea and decontamination of radioactivity in freshwater, *Proceedings of the International Congress on the Peaceful Uses of Atomic energy*, Vol. 14, pp. 97-106 (1965), P/849, 1964.

Hoar, W. S. and D. J. Randall (eds.), *Fish Physiology*, Academic Press, New York, 1969.

Hobden, F. W., et al., "Determination and calculation of the equilibrium constants for isotopic exchange in the systems *n*-amyl alcohol-water and ethylthiol-water. Vapor pressures and raman spectra of *n*-amyl deuterio-alcohol and ethyl deuteriothiol," *J. Chem. Soc.* 61 (1939).

Holcomb, C. M., C. G. Jackson, Jr., M. M. Jackson, and S. Kleinbergs, Occurrence of radionuclides in the exoskeleton of turtles, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 385-389 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Horton, J. H., J. C. Corey, and R. M. Wallace, Tritium loss from water exposed to the atmosphere, *Environ. Sci. Technol.* 5, 338-343 (1971).

Hunn, J. B. and P. O. Fromm, Uptake, turnover and excretion of iodine-131 by rainbow trout (*Salmo gairdneri*), *Biol. Bull.* 126, 282-290 (1964).

Hunn, J. B. and P. O. Fromm, In vivo uptake of radioiodine by rainbow trout, *J. Water Poll. Contr. Assoc.* 38, 1981-1985 (1966).

Hunn, J. B. and E. P. Reineke, Influence of iodine intake on iodine distribution in trout, *Proc. Soc. Exp. Biol. Med.* 115, 91-93 (1964).

Hunt, E. L. and J. N. Dent, Iodine uptake and turnover in the frog tadpole, *Physiol. Zool.* 30, 87-91 (1957).

Hutchinson, G. E., *A treatise on limnology*, Vol. I, John Wiley and Sons, New York, 1957.

Ichikawa, R., Uptake of potassium and cesium from water by the sockeye salmon, *J. Radiat. Res. (Japan)* 1, 107-110 (1960).

Il'in, D. I., and I. Petrova, Accumulation of radioactive elements by some groups of aquatic organisms, *Soviet J. Atomic Energy* 3, 1013-1016 (1958).

Jacobs, D. G., Release of tritium in an expanding nuclear power economy and its long range behavior in the environment, Health Physics Division Annual Progress Report for the Period Ending July 31, 1968, ORNL-4316 pp. 50-51 (1968).

Jarvis, N. D., R. W. Clough, and E. D. Clark, Iodine content of Pacific Coast salmon, Univ. Washington Pub. Fish. 1, 109-140 (1925-1928), In Robertson and Chaey 1953.

Jinks, S. M. and M. Eisenbud, Concentration factors in the aquatic environment, Radiat. Data. Rep. 13, 243-247 (1972).

Kahn, B., Radionuclides in the environment at nuclear power stations, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 30-36 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Kanazawa, T., K. Kanazawa, and J. A. Bassham, Tritium incorporation in the metabolism of Chlorella pyrenoidosa, Environ. Sci. Technol. 6, 638-642 (1972).

Kaye, N. W. and E. E. LeBourhis, Uptake and turnover of a single injected dose of iodine-131 in tadpoles of Rana clamitans, Zoologica 43, 73-76 (1958).

Kaye, S. V., Assessing potential radiological impacts to aquatic biota in response to the National Environmental Policy Act (NEPA) of 1969, IN Symposium on environmental behavior of radionuclides released in the nuclear industry, IAEA Symposium, Aix-en-Provence, France, 14-18 May 1973.

Kevern, N. R., Strontium and calcium uptake by the green algae, Oocystis eremosphaeria, Science 145, 1445-1446 (1964).

Kevern, N. R., Feeding rate of carp estimated by a radioisotopic method, Trans. Amer. Fisheries Soc. 95(4), 363-371 (1966).

Kevern, N. R., Effects of limnological factors on uptake of cesium-137, USAEC Technical Progress Report C00-1795-1 (1969).

Kevern, N. R. and N. A. Griffith, Cesium-134 distribution and transport in the carp, Health Physics Division Annual Progress Report for the Period Ending July 31, 1965, pp. 85-86, ORNL-3849 (1965).

Kevern, N. R. and N. A. Griffith, Effect of trophic level on radio-nuclide accumulation by fish, IN Radiation Ecology Section, Health Physics Division, Annual Progress Report, ORNL-4007 (1966).

Kevern, N. R. and S. A. Spigarelli, Effects of selected limnological factors on the accumulation of cesium-137 fallout by largemouth bass (Micropterus salmoides), IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 354-360 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Kevern, N. R., N. A. Griffith, and T. Grizzard, Biological half-life of cesium-134 in carp and two aquatic insects, Health Physics Division Annual Progress Report for the Period Ending July 31, 1964, Radiation Ecology Section, pp. 101-102, ORNL-3697 (1964).

Kevern, N. R., N. A. Griffith, G. M. Rosenthal, Jr., and D. J. Nelson, Uptake and accumulation of strontium and calcium by algae and snails, Health Physics Division Annual Progress Report for the Period Ending July 31, 1964, Radiation Ecology Section, pp. 96-101, ORNL-3697 (1964).

Kharkr, D. P., K. K. Turekian, and K. K. Bertine, Stream supply of dissolved silver, molybdenum, antimony, selenium, chromium, cobalt, ruddium and cesium to the oceans, *Geochim. Cosmochim. Acta* 32, 285-298 (1968).

King, S. F., Uptake and transfer of cesium-137 by Chlamydomonas, Daphnia, and bluegill fingerlings, *Ecology* 45, 852-859 (1964).

Kirchen, R. V., Radiopathology and dosimetry in the thyroid region of goitrous killfish fed iodine-131 under brinyhaline conditions--An Abstract, IN Biology of radioiodine, p. 261 (1964), Proceedings of the Hanford Symposium on the Biology of Radioiodine.

Klement, A. W., and V. Schultz, Russian radioecology, A bibliography of soviet publications, with citations of English translations and abstracts, TID-3915 (1972).

Knauss, H. J., and J. W. Porter, The absorption of inorganic ions by Chlorella pyrenoidosa, *Plant Physiol.* 29, 229-234 (1954).

Koch, A. L., A distributional basis for the variation in killing efficiencies by different tritiated compounds incorporated into Escherichia coli, *Radiat. Res.* 24, 389-411 (1965).

Kolehmainen, S. E., White Oak Lake studies, Health Physics Division Annual Progress Report for the Period Ending July 31, 1968, pp. 117-118 (1969).

Kolehmainen, S. E., Balances of cesium-137, stable cesium and potassium in bluegill, Lepomis macrochirus Raf., and other fish in White Oak Lake, *Health Phys.* 23, 301-315 (1972).

Kolehmainen, S. E. and D. J. Nelson, Balances of cesium-137, stable cesium, and the feeding rates of bluegill (Lepomis macrochirus Raf.) in White Oak Lake, ORNL-4445 (1969).

Kolehmainen, S., E. Hasanen, and J. K. Miettinen, Cesium-137 levels in fish of different limnological types of lakes in Finland during 1963, *Health Phys.* 12, 917-922 (1966).

Kolehmainen, S., E. Hasanen, and J. K. Miettinen, Cesium-137 in fish, plankton, and plants in Finnish lakes, IN Radioecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 913-919 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1966.

Kolehmainen, S., E. Hasanen, and J. K. Miettinen, Cesium-137 in the plants, plankton and fish of the Finnish lakes and factors affecting its accumulation, IN Proceedings of the First International Congress of Radiation Protection, W. S. Snyder (ed.), pp. 407-415 (1968), Pergamon Press, Oxford.

Kolehmainen, S., S. Takatalo, and J. K. Miettinen, A tracer experiment with iodine-131 in an oligotrophic lake, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 278-284 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Kolehmainen, S., Romantschuk, S. Takatalo, and J. K. Miettinen, Pollution experiments with Cs-137 in lakes of two limnologically different types, IN Congrès international sur la radio-protection du milieu devant le développement des utilisations pacifiques de l'énergie nucléaire, pp. 181-200, Toulouse, 14-16 Mars 1967, Société Française de Radio-protection, Paris (1968).

Koranda, J. J., Preliminary studies of the persistence of tritium and carbon-14 in the Pacific Proving Grounds, *Health Phys.* 11, 1445-1457 (1965).

Koranda, J. J. and J. R. Martin, The persistence of radionuclides at sites of nuclear detonations, USAEC Report UCRL-71867 (1969).

Kornegay, B. H., W. A. Vaughn, D. K. Jamison, and J. M. Morgan, Jr. (eds.), Transport of radionuclides in freshwater systems, USAEC Document, TID-7664 (1966).

Kouts, H. and J. Long, Tritium production in nuclear reactors, IN Tritium, A. A. Moghissi and M. W. Carter (eds.), pp. 38-45 (1973), Messinger Graphics, Phoenix, TID CONF-710809.

Krumholz, L. A., Observations on the fish population of a lake contaminated by radioactive wastes, *Bull. Am. Mus. Nat. His.* 110, 227-367 (1956).

Kulikov, N. V., Radioecology of freshwater plants and animals, IN Radioecology, V. M. Klechkovskii, G. G. Polikarpov, and R. M. Aleksakhin (eds.), pp. 323-337 (1973), Wiley, New York.

Lajtha, L. G., Some problems on the biological uses of tritiated thymidine, IN Chemical stability of tritiated pyrimidine nucleosides and some problems raised by their use in biology, *Actes de Colloque* tena a MOL/Belgium, March 1965.

Lambert, B. E. and R. J. Clifton, Radiation dose resulting from the ingestion of tritiated thymidine by the rat, *Health Phys.* 15, 3-9 (1968).

Lang, A. R. G. and S. G. Mason, Tritium exchange between cellulose and water: Accessibility measurements and effects of cyclic drying, *Can. J. Chem.* 38, 373-387 (1960).

Lebedeva, G. D., Influence of various salt contents of water on the accumulation and excretion of cesium-137 by freshwater fish, *Radio-biology* (Translated from Russian) 6, 118-124, AEC-tr-6773 (1966).

Lebedeva, G. D., Effects of various saline compositions of water on the buildup of strontium-90 and cesium-137 in freshwater fish, *Radiobiology* (Translated from Russian) 6, 227-234, AEC-tr-6944 (1966).

Leloup, J. and M. Fontain, Iodine metabolism in lower vertebrates, Part I. Intrathyroidal biochemical reaction, *Ann. N.Y. Acad. Sci.* 86, 316-353 (1960).

Lentsch, J. W., T. J. Kneip, M. E. Wrenn, G. P. Howells, and M. Eisenbud, Stable manganese and manganese-54 distributions in the physical and biological component of the Hudson River Estuary, IN *Radionuclides in Ecosystems*, D. J. Nelson (ed.), pp. 752-768, Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971 (1973).

Livingston, D. A., Chemical composition of rivers and lakes, *Geological Survey Professional Paper* 440-G (1963).

Lomenick, T. and D. A. Gardiner, The occurrence and retention of radionuclides in the sediments of White Oak Lake, *Health Phys.* 11, 567-577 (1965).

Lowman, F. G., Iron and cobalt in ecology, IN *Radioecology*, V. Schultz and A. W. Klement, Jr. (eds.), pp. 561-567 (1963), Proceedings of the First National Symposium on Radioecology, Fort Collins, Colorado, September 10-15, 1961.

Lucas, H. F., D. N. Edington, and P. J. Colby, Concentrations of trace elements in Great Lakes fishes, *J. Fish. Res. Bd. Can.* 27, 677-684 (1970).

Lunde, G., Über die Geochemie und Biochemie des Jods mit besonderer Berücksichtigung der norwegischen Kropfprophy Taxe, *Wien. klin. Wschr.* 40, 1559 (1928), IN Vinogradov 1953.

Lunde, G., J. Boe, and K. Closs, Iodine content of American marine animals, *J. Cons. Perm. Int. Explor. Mer.* 5, 216 (1930), IN Vinogradov 1953.

Lunde, G., K. Closs, H. Haaland, and S. O. Madsen, Jodinnholdet i norsk fisk og fiskerprodukter, *Aarsberetn. Norg. Fisk.*, No. 4, 1-41 (1928), IN Vinogradov 1953.

McKown, D. A., Health Physics biography on tritium, LAMS-2946 (1964).

Mahadevan, E. G., and E. V. Monse, "Equilibrium constant for the deuterium exchange between methylacetylene and water," *J. Chem. Phys.* 43, 2556 (1965).

Mahmoud, K. A., Y. A. Yousef, M. M. Ishak, W. E. Y. Abel-Malik, S. R. Khalil, A. Shafik, and R. M. K. El-Shinawy, Transfer mechanisms and concentrating processes of certain radionuclides in the aquatic ecology of Ismailia Canal, IAEA Symposium on Environmental Behavior of Radionuclides Released in the Nuclear Industry, France, May 14-18, 1973, IAEA/SM-172/51.

Maletskos, C. J., Intercalibration and intercomparison study of radio-nuclide concentrations in Lake Michigan samples, New England Deaconess Hospital, Progress Report October-March, 1972, COO-3371-1, USAEC Contract AT(11-1)-3371.

Marey, A. N. and M. M. Saurov, Material for assessment of the role played by food chains in strontium-90 migration from freshwater reservoirs into the diet of man, IN Radioecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 135-141 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1966.

Marshall, J. S. and J. H. LeRoy, Iron, manganese, cobalt, and zinc cycles in a South Carolina reservoir, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 465-473 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Martin, J. R. and J. J. Koranda, Biological half-life studies of tritium in chronically exposed kangaroo rats, *Radiat. Res.* 50, 426-440 (1972).

Marx, D. (Summarized by), "Etude des Constantes des Equilibres Isotopiques du Deuterium entre l'Eau et les Hydrures des Metalloides de la Deuxieme Famille," C.E.A. Report 1382, Saclay, France, 1960.

Mason, B. J., E. J. Wipper, and V. D. Leavitt, Tritium uptake following a thermonuclear test, IN Environmental contamination by radioactive materials, pp. 167-173, IAEA Symposium, Vienna, March 1969.

Mathis, B. J. and T. F. Cummings, Selected metals in sediments, water, and biota in the Illinois River, *J. Water Poll. Contr.* 45, 1573-1583 (1973).

Mazzocco, P., Iodine in food in salta, *Sem. Med. B. Aires* 37, 364 (1930).

Merlini, M., The freshwater clam as a biological indicator of radiomanganese, IN Radioecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 977-982 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1966.

Merlini, M., F. Giradi, and G. Possi, Activation analysis in studies of an aquatic ecosystem, IN Nuclear activation techniques in the life sciences, pp. 615-629 (1967), Proceedings of the Symposium held by the IAEA in Amsterdam, May 8-12, 1967, also STI/PUB/155, SM-91/7.

Merlini, M., C. Biglicocca, A. Berg, and G. Pozzi, Trends in the concentration of heavy metals in organisms of a mesotrophic lake as determined by activation analysis, IN Nuclear techniques in environmental pollution, pp. 447-458 (1971), IAEA Symposium on the Use of Nuclear Techniques in the Measurement and Control of Environmental Pollution, Salsburg, Austria, October 26-November 3, 1970.

Merlini, M., F. Girardi, R. Pietra, and A. Brozgelli, The stable manganese content of molluscs from Lake Maggiore determined by activation analysis, Limnol. Oceanogr. 10, 371-378 (1965).

Minckley, W. L., J. E. Craddock, and L. A. Krumholz, Natural radioactivity in the food web of the banded sculpin Cottus carolinus (Gill.), IN Radioecology, V. Schultz and A. W. Klement, Jr. (eds.), pp. 229-236 (1963), Proceedings of the First National Symposium on Radioecology, Fort Collins, Colorado, September 10-15, 1961.

Moghissi, A. A., M. W. Carter, and R. Lieberman, Long term evaluation of the biological half-life of tritium, Health Phys. 21, 57 (1971).

Money, W. L., V. Lucas, and R. W. Rawson, The turnover of radio-iodine (I-131) by the Rana pipiens tadpole, J. Exp. Biol. 128, 411-421 (1955).

Monier-Williams, G. W., Trace elements in food, John Wiley and Sons, New York, 1950.

Morton, R. J., Status report no. 5 on Clinch River study, ORNL-3721, Oak Ridge, Tennessee, 1965.

Moul, J. W. and P. F. Nace, Radioiodine metabolism in amphiuma after its intraperitoneal administration, Can. J. Physiol. Pharm. 42, 623-626 (1964).

Nelson, D. J., Clams as indicators of strontium-90, Science 137, 38-39 (1962).

Nelson, D. J., Strontium, strontium-90, and calcium analyses of Clinch and Tennessee River clams, ORNL-TM-270 (1962).

Nelson, D. J., Role of bottom organisms, IN Transport of radio-nuclides in freshwater systems, Kornegay et al. (eds.), pp. 193-207 (1963), TID-7664.

Nelson, D. J., The strontium and calcium relationships in Clinch and Tennessee River molluscs, IN Radioecology, V. Schultz and A. W. Klement, Jr. (eds.), pp. 203-211 (1963), Proceedings of the First National Symposium on Radioecology, Fort Collins, Colorado, September 10-15, 1961.

Nelson, D. J., Strontium in white crappie (Pomoxis annularis) flesh and bone, IN Health Physics Division Annual Progress Report for Period Ending June 30, 1963, Part 2, Radiation Ecology Section, p. 107, ORNL-3492 (1963).

Nelson, D. J., Biological vectors and reservoirs of strontium-90, Nature 203, 420 (1964).

Nelson, D. J., Interpretation of radionuclide uptake from aquatic environments, Nuclear Safety 5, 196-199 (1964).

Nelson, D. J., Deposition of strontium in relation to morphology of clam (Unionidae) shells, Verh. Internat. Verein. Limnol. 15, 893-902 (1964).

Nelson, D. J., Cesium and potassium in white crappie and other Clinch River fish, ORNL-4007, pp. 84-87 (1966).

Nelson, D. J., The prediction of strontium-90 uptake in fish using data on specific activities and biological half-lives, IN Radio-ecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 843-858 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1966.

Nelson, D. J., Ecological behavior of radionuclides in the Clinch River and Tennessee River, IN Proceedings of Reservoir Fishery Resources Symposium, pp. 169-187, Athens, Georgia, April 5-7, 1967.

Nelson, D. J., Cesium, cesium-137, and potassium concentrations in white crappie and other Clinch River fish, IN Symposium on radio-ecology, D. J. Nelson and F. C. Evans (eds.), pp. 240-248 (1969), Proceedings of the Second National Symposium on Radioecology, Ann Arbor, Michigan, May 15-17, 1967.

Nelson, D. J., Biological concentration of radioisotopes from aquatic environments, Nuclear Safety 2, 81-83 (1969).

Nelson, D. J. and C. R. Malone, Feeding rates of snails determined with ^{60}Co , IN Radiation Ecology Section, Health Physics Division, Annual Progress Report, July 31, 1968, ORNL-4316 (1969).

Nelson, D. J., N. A. Griffith, and R. C. Early, Calcium and strontium in white crappie flesh and bone, IN Health Physics Division Annual Progress Report for Period Ending July 31, 1964, Radiation Ecology Section, p. 95, ORNL-3697 (1964).

Nelson, D. J., N. A. Griffith, and S. A. Rucker, Radionuclide excretion studies, IN Health Physics Division Annual Progress Report for Period Ending July 31, 1969, pp. 119-120, ORNL-4446 (1970).

Nelson, D. J., T. C. Rains, and J. A. Norris, High purity calcium carbonate in freshwater clam shell, Science 152, 1368-1370 (1966).

Nelson, D. J., N. A. Griffith, J. W. Gooch, and S. A. Rucker, White Oak Lake studies, IN Radionuclide cycling in aquatic systems, ORNL-2634 (1970).

Nelson, D. J., N. A. Griffith, J. W. Gooch, and S. A. Rucker, White Oak Lake Studies, IN Ecological Sciences Division Annual Report, July 31, 1970, ORNL-4634 (1971).

Nelson, W. C. and F. W. Whicker, Cesium-137 in some Colorado game fish, 1965-1966, IN Symposium on Radioecology, D. J. Nelson and F. C. Evans (eds.), Proceedings of the Second National Symposium on Radioecology, Ann Arbor, Michigan, May 15-17, 1967.

Nilson, H. W. and E. J. Coulson, The mineral content of the edible portions of some American fisheries products, Invest. Rep. U.S. Bur. Fish. 41, 1-7 (1939), IN Vinogradov 1953.

Odum, H. T., Biogeochemical deposition of strontium, Pub. Inst. Mar. Sci. 4, 38-114 (1957).

Odum, E. P., and F. B. Golley, Radioactive tracers as an aid to the measurement of energy flow at the population level in nature, In Radioecology, V. Schultz and A. W. Klement (eds.), p. 403 (1963), Proceedings of the National Symposium on Radioecology, Fort Collins, Colorado, September 10-15, 1961.

Ohmomo, Y., H. Suzuki, M. Sumiya, and M. Saiki, Studies on accumulation excretion and distribution of iodine-131 and cadmium-115m in freshwater fish and marine fish, J. Rad. Res. 13, 3 (1972).

Ophel, I. L., The fate of radiostrontium in a freshwater community, IN Radioecology, V. Schultz and A. W. Klement, Jr. (eds.), pp. 213-216 (1963), Proceedings of the First National Symposium on Radioecology, Fort Collins, Colorado, September 10-15, 1961.

Ophel, I. L., The environmental capacity of freshwaters for waste radionuclides, IN Symposium on Environmental Behavior of Radionuclides Released in the Nuclear Industry, France, May 14-18, 1973, IAEA/SM-172/45.

Ophel, I. L. and C. D. Fraser, Calcium and strontium discrimination by aquatic plants, Ecology 51, 324-327 (1970).

Ophel, I. L., and E. D. Fraser, The fate of cobalt-60 in a natural freshwater ecosystem, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 323-327 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Ophel, I. L. and J. M. Judd, Absorption of radiostrontium by gills of freshwater fish, Nature 194, 1187-1188 (1962).

Ophel, I. L. and J. M. Judd, Experimental studies of radiostrontium accumulation by freshwater fish from food and water, IN Radioecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 859-865 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1966.

Ophel, I. L. and J. M. Judd, Strontium-calcium relationships in aquatic food chains, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 221-225 (1973), Proceedings of the Third National Symposium in Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Ophel, I. L., C. D. Fraser, and J. M. Judd, Strontium concentration factors in biota and bottom sediments of a freshwater lake, EUR-4800 (1972).

Orr, J. B. and I. Leitch, Iodine in nutrition, A review of existing information up to 1927, United Kingdom Medical Research Council, Special Report Series No. 123, His Majesty's Stationery Office, London (1929).

Osborne, R. V., Permissible levels of tritium in man and the environment, Radiat. Res. 50, 197-211 (1972).

Pahl, G., Radioactive and stable strontium analysis of upper Mississippi River clam shells, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 234-239 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Palumbo, R. F., P. R. Olson, J. R. Donaldson, F. G. Lowman, T. Gudjonsson, and Z. Short, Uptake of I-131 by freshwater organisms, An Abstract, IN Biology of radioiodine, L. K. Bustad (ed.), p. 133 (1964), Proceedings of the Hanford Symposium on the Biology of Radioiodine, Richland, Washington, July 17-19, 1962.

Parker, F. L., Radioactive wastes from fusion reactors, Science 159, 83-84 (1968).

Parker, F. S. and K. R. Bhaskar, Infrared studies of hydrogen-deuterium exchange in biological molecules, Appl. Spectr. Rev. 3, 91-142 (1969).

Patel, B., and G. R. Doshi, Lamellidens marginalis L. as indicator of manganese-54, Archo Oceanogr. Limnol. 17, 27-42 (1972).

Patten, B. C. and R. L. Iverson, Photosynthesis and uptake of strontium-85 in freshwater plankton, *Nature* 211, 96-97 (1966).

Patten, B. C., N. A. Griffith, S. A. Rucker, and D. J. Nelson, Aquarium experiments, IN Radiation Ecology Section, Health Physics Division, Annual Progress Report, July 31, 1968, ORNL-4316 (1969).

Patzer, R. G., A. A. Moghissi, and D. N. McNelis, Accumulation of tritium in various species of fish reared in tritiated water, IN Proceedings of IAEA/NEA/WHO Symposium on Environmental Behavior of Radionuclides Released in the Nuclear Industry, Aix-en-Provence, France, May 14-18, 1973, IAEA/SM-172/29, Vienna, 1973.

Pendleton, R. C., Absorption of cesium-137 by aquatic community, AEC Report HW-53500, p. 42 (1957).

Pendleton, R. C., Effects of some environmental factors on bioaccumulation of cesium-137 in an aquatic community, IN Hanford Biology Research Annual Report for 1958, J. J. Davis (ed.), pp. 42-46 (1958).

Pendleton, R. C. and W. C. Hanson, Absorption of cesium-137 by components of an aquatic community, IN Proceedings of the Second International Conference on Peaceful Uses of Atomic Energy, Vol. 18, 419-422 (1958), U.N. Geneva, September 1-13, 1958.

Pendleton, R. C., C. W. Mays, R. D. Lloyd, and B. W. Church, A trophic level effect on cesium-137 concentration, *Health Phys.* 11, 1503-1510 (1965).

Perkins, R. W. and J. M. Nielsen, Deposition of a wide spectrum of radionuclides by nuclear production and testing operations, BNWL-SA-993 (1967), also CONF-670313-1 (1967).

Peterson, H. I., Jr., Stable element and radionuclide reconcentration in aquatic systems. I. Stable element effects on equilibrium situation, IN Proceedings of the Fifth Annual Health Physics Society Mid-year Topical Symposium, Health Physics Aspects of Nuclear Siting, Vol. 2, pp. 289-328, AEC CONF-701106 (1970).

Pinson, E. A., The body absorption distribution and excretion of tritium in man and animals, USAEC Report LA-1218 (1951).

Pinson, E. A., Incorporation of tritium into the organic components of various tissue of the mouse, LA-1467 (1952).

Pinson, E. A. and W. H. Langham, Physiology and toxicology of tritium in man, *J. Appl. Physiol.* 10, 108-126 (1957).

Polikarpov, G. G., Radioecology of aquatic organisms, Reinhold Book Division, New York, 1966.

Porter, J. W., Incorporation of hydrogen isotopes into various fractions of rapidly growing algae, IN Biology Research Annual Report 1953, pp. 77-78, Hanford Atomic Products Operation, Richland, Washington, HW-30437 (1954).

Porter, J. W. and H. J. Knauss, The effect of tritium oxide on some synthetic processes of Chlorella pyrenoidosa, IN Biology Research, Annual Report 1953, pp. 73-76, Hanford Atomic Products Operation, Richland, Washington, HW-30437 (1954).

Porter, J. W. and M. S. Waston, Gross effects of growth-inhibiting levels of tritium oxide on Chlorella pyrenoidosa, IN Biology Research, Annual Report 1953, pp. 71-72, Hanford Atomic Products Operation, Richland, Washington, HW-30437 (1954).

Preston, A., D. F. Jefferies, and J. W. R. Dutton, The concentration of cesium-137 and strontium-90 in the flesh of brown trout taken from rivers and lakes in the British Isles between 1961-1966. The variables determining the concentrations and their use in radiological assessment, Water Res. 1, 475-496 (1967).

Price, J. W., A study of the food habits of some Lake Erie fish, Bull. Ohio Biological Survey, New Series, Vol. II, No. 1, Ohio State Univ., Columbus (1963).

Rambeck, W. A. and J. A. Bassham, Tritium incorporation and retention in photosynthesizing algae, Biochem. Biophys. Acta 304, 725-735 (1973).

Raney, F. and Y. Vaadia, Movement and distribution of THO in tissue water and vapor transpired by shoots of helianthus and nicotiana, Plant Phys. 40, 383-388 (1965).

Ravera, O., Radioactivity in freshwater organisms of some lakes of northern Italy, IN Biological problems in water pollution, pp. 195-201 (1965), Third Seminar, Research Branch, Robert A. Taft Sanitary Engineering Center, U.S. Department of Health, Education and Welfare, Madison, Wisconsin, 1962.

Ravera, O., Strontium-85 and cesium-137 uptake by freshwater seston, Rev. Int. Oceanogr. Med. 21, 109-124 (1971).

Ravera, O. and P. Gaglione, Concentrations di Mn-54 da fallout in relazione all'eta e all'organo di *Unio Mancus elongatus* (Molluschi, Lamellibranchi), Rend Ist Lombardo Sci. Lettere 96, 157 (1962).

Ravera, O., M. Merlini, and S. Musso, Assunzione ed Eliminazione de Radiocobalto da Parte di un Mollusco d'acqua Dolce (*Unio mancus* var. *elongatus*), Communicazione Presentata al XIV Congresso Nazionale dell'Associazione Italiane de Fisica Sanitaria e Protezione contro le Radiazioni, Formia, 29-31, Maggio 1968, also CONF-680535-29.

Rediske, J. H. and J. F. Cline, Comparative metabolism of protium, deuterium and tritium in the bean plant, IN Biology Research, Annual Report 1952, pp. 72-74, Hanford Atomic Products Operation, Richland, Washington, HW-28636 (1953).

Reed, J. R., Uptake and excretion of cobalt-60 by black bullheads Ictalurus melas (Raf.), Health Phys. 21, 835-844 (1971).

Reed, J. R. and D. J. Nelson, Radiostrontium uptake in blood and flesh in bluegills (Lepomis macrochirus), IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 226-233 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Reynolds, T. D. and E. F. Gloyna, Uptake and release of radionuclides by stream sediments, IN Proceedings of the Second International Conference on Advances in Water Pollution Research, Vol. 1, O. Jaag (ed.), pp. 151-172 (1964), Pergamon Press, New York.

Rickard, W. H. and L. L. Eberhardt, Fallout radiocesium in sedges and trout of a cascade mountain bog, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 349-353 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-11, 1971.

Robertson, O. H. and Albert L. Chaney, Thyroid hyperplasia and tissue iodine content in spawning rainbow trout: A comparative study of Lake Michigan and California sea-run trout, Physiol. Zool. 26, 328-340 (1953).

Rose, D. J., Hazards of tritium from controlled fusion, Health Phys. 18, 439-441 (1970).

Rosenthal, G. M., Jr. and M. L. Stewart, Tritium incorporation in algae and transfer in simple aquatic food chains, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 440-444 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Rosenthal, H. L., Uptake of calcium-45 and strontium-90 from water by freshwater fishes, Science 126, 699-700 (1957).

Sabo, J. and P. H. Bedrosian, Studies on the fate of certain radionuclides in estuarine and other aquatic environments, U.S. Department of Health, Education and Welfare, Public Health Service Publication No. 999-R-3 (1963).

Schroeder, H. A., J. J. Balassa, and I. H. Tipton, Essential trace metals in man: Manganese, J. Chron. Dis. 19, 545-571 (1966).

Schurr, J. M. and M. N. Stampler, Model for the accumulation of strontium and calcium by recently molted crayfish (Cambarus longulus longerostris Ort.), Limnol. Oceanogr. 7, 474-477 (1962).

Scott, A. G. and J. E. Guthrie, The accumulation of radioactive cesium, phosphorus, strontium, and yttrium by mosquito larvae and its relevance to radiation dose, *Can. J. Zool.* 50, 767-770 (1972).

Shellabarger, C. J., A. Gorbman, F. C. Schatzlein, and D. McGill, Some quantitative and qualitative aspects of iodine-131 metabolism in turtles, *Endocrinology* 59, 331-339 (1956).

Sherman, H. C., Essentials of nutrition, MacMillan Co., New York (1957).

Shiffman, R. H., The uptake of strontium from diet and water by rainbow trout, IN *Biology Research, Annual Report 1958*, pp. 16-19, Hanford Atomic Products Operation, Richland, Washington, HW-59500 (1959).

Shore, B., L. Anspaugh, R. Chertok, J. Gofman, F. Harrison, R. Heft, J. Koranda, Y. Ng, P. Phelps, G. Potter, and A. Tamplin, The fate and importance of radionuclides produced in nuclear events, IN *Proceedings of Symposium on Public Health Aspects of Peaceful Uses of Nuclear Explosives*, pp. 595-651, Las Vegas, Nevada, April 7-11, 1969.

Short, Z. F., R. F. Palumbo, P. R. Olson, and J. R. Donaldson, The uptake of iodine-131 by the biota of Fern Lake, Washington, in a laboratory and a field experiment, *Ecology* 50, 979-989 (1969).

Chia-Shun, S. and E. F. Gloyne, Radioactive transport in water-transport of strontium-85 and cesium-137 in an aquatic environment, EHE-01-6602, CRWR-10 (1966).

Siri, W. and J. Evers, Tritium exchange in biological systems, IN *IAEA Proceedings of the Symposium on the Detection and Use of Tritium in the Physical and Biological Sciences*, Vol. 2, pp. 71-84 (1961).

Skauen, D. M., Tritium in ecology--A review, IN *Radioecology*, V. Schultz and A. W. Klement, Jr. (eds.), pp. 603-609 (1963), *Proceedings of the First National Symposium on Radioecology*, Fort Collins, Colorado, September 10-15, 1961.

Skauen, D. M., The effects of tritium oxide on aquatic organisms, NYO-3039-1 (1964).

Slater, J. V., Radiocobalt accumulation in Tetrahymena, *Biol. Bull.* 122, 390-399 (1957).

Smith, G. N., O. L. Hollis, and R. C. Thompson, Distribution of bound tritium in the rat following exposure of tritium, IN *Biology Research, Annual Report 1953*, pp. 77-78, Hanford Atomic Products Operation, Richland, Washington, HW-30437 (1954).

Smith, T. E. and R. T. Taylor, Incorporation of tritium from tritiated water into carbohydrates, lipids, and nucleic acids, USAEC Report UCRL-50781 (1969).

Spector, W. S., Handbook of Biological Data, W. B. Saunders Co., Philadelphia, 1956.

Spigarelli, S. A., Ecological factors affecting the accumulation of cesium-137 fallout by a natural population of largemouth bass (Micropterus salmoides), IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 328-333 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Stewart, M. L., G. M. Rosenthal, and J. R. Kline, Tritium discrimination and concentration in freshwater microcosms, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 452-459 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 11-12, 1971.

Strand, J. A., W. L. Templeton, and E. G. Tangen, Accumulation and retention of tritium (tritiated water) in embryonic and larval fish, and radiation effect, IN Radionuclides in ecosystems, D. J. Nelson (ed.), pp. 445-451 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Struxness, E. G., P. H. Carrigan, Jr., M. A. Churchill, K. E. Cowser, R. J. Morton, D. J. Nelson, and F. L. Parker, Comprehensive report of the Clinch River study, USAEC Report ORNL-4035, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Stumm, W. and J. J. Morgan, Aquatic chemistry, John Wiley and Sons, New York, 1970.

Suzuki, Y., R. Nakamuyra, and T. Ueda, Accumulation of strontium and calcium in freshwater fishes of Japan, Japan Radiat. Res. Soc. 13, 199-207 (1972).

Takatalo, S., S. Kolehmainen, and J. K. Miettinen, Excretion of iodine from perch (Perca fluviatilis), IN Biological enrichment chains of radionuclides in Finnish Lakes, pp. 104-111 (1968), Annual Report for the period Sept. 1, 1967 to August 31, 1968, Department of Radiochemistry, Univ. Helsinki, Finland.

Templeton, W. L. and V. M. Brown, The relationship between concentrations of calcium, strontium and strontium-90 in wild brown trout, Salmo trutta L., and the concentrations of the stable elements in some waters of the United Kingdom, and the implications in radiological health studies, Int. J. Air Water Poll. 8, 49-75 (1964).

Templeton, W. L., J. M. Dean, D. G. Watson, and L. A. Rancitelli, Freshwater ecological studies in Panama and Columbia, Bioscience 19, 804-808 (1969).

Thompson, R. C., Studies of metabolic turnover with tritium as a tracer. I. Gross studies on the mouse, J. Biol. Chem. 197, 81-88 (1952).

Thompson, R. C., Studies of metabolic turnover with tritium as a tracer. II. Gross studies on the rat, *J. Biol. Chem.* 200, 731-739 (1953).

Thompson, R. C., Effect of body distribution and retention of tritium on the hazard of exposure to tritium oxide, IN Biology Research, Annual Report 1953, pp. 77-78, Hanford Atomic Products Operation, Richland, Washington, HW-30437 (1954).

Thompson, R. C. and J. E. Ballou, Comparative metabolism of deuterium and tritium in the rat, IN Biology Research, Annual Report 1952, pp. 72-74, Hanford Atomic Products Operation, Richland, Washington, HW-28636 (1953).

Thompson, R. C. and J. E. Ballou, Distribution of bound tritium in the rat following chronic exposure to tritium oxide, IN Biology Research, Annual Report 1953, pp. 77-78, Hanford Atomic Products Operation, Richland, Washington, HW-30437 (1954).

Thompson, R. C. and J. E. Ballou, Studies of metabolic turnover with tritium as a tracer. V. The predominantly nondynamic state of body constituents in the rat, *J. Biol. Chem.* 223, 795-809 (1956).

Thompson, S. E., C. A. Burton, D. J. Quinn, and Y. C. Ng, Concentration factors of chemical elements in edible aquatic organisms, Lawrence Livermore Laboratory, Univ. California, Livermore, 1972.

Timofeev-Resovskii, N. V., E. A. Timofeeva-Resovskaya, G. A. Milyutina, and A. B. Getsova, Coefficients of accumulation of radioisotopes of sixteen different elements by freshwater organisms and the effect of EDTA on some of them, *Dokl. Akad. Nauk SSSR* 132, 369-372 (1960).

Timofeeva, N. A., and N. V. Kulikov, The role of freshwater plants in accumulation of strontium-90 and its distribution over the components, IN Radioecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 835-841 (1967), Proceedings of the International Symposium, Stockholm, April 25-29 (1966).

Timofeyeva-Resovskaya, Ye. A., Distribution of radioisotopes in the main components of freshwater bodies (Monograph), *Tr. Inst. Biol. Akad. Nauk SSSR, Ural'skiy filia* 30, 1-78 (JPRS 21.816) 1963.

Timofeeva-Resovskaya, E. A., N. V. Timofeev-Resovskii, and E. A. Gileva, Specific accumulators of individual radioisotopes among freshwater organisms, *Dokl. Akad. Nauk SSSR* 140, 1437-1440 (1961).

Tong, W. and I. L. Chaikoff, Iodine-131 utilization by the aquarium snail and the cockroach, *Biochem. Biophys. Acta* 48, 347-351 (1961).

Townsley, S. J., The concentration of strontium-90 and other nuclides of strontium in fish, IN Radioecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 867-878 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1966.

Tressler, D. K. and A. W. Wells, Iodine content of sea foods, Docu. U.S. Bur. Fish., No. 967 (1924).

Underwood, E. J., Elements in human and animal nutrition, Academic Press, New York (1956).

Van der Borgh, O. and S. Van Puymbroeck, Ionic regulation and radiocontamination: Active transport of alkaline earth ions in freshwater gastropods (Mollusca), IN Radioecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 925-930 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1966.

Vanderploeg, H. A., R. S. Booth, and F. H. Clark, A specific activity and concentration model applied to cesium movement in an oligotrophic lake, IN Mineral Cycling in Southeastern Ecosystems, F. G. Howell (ed.), Symposium held 1-3 May 1974, Augusta, Georgia (in press).

Van Hook, R. I., Jr. and D. A. Crossley, Jr., Assimilation and biological turnover of cesium-134, iodine-131, and chromium-51 in brown crickets Acheta domesticus (L.), Health Phys. 16, 463-467 (1969).

Vaughn, B. E. and Staff of Ecosystems Department, Environmental and Life Sciences Division, Effects of tritium on aquatic environments, BNWL-1650 PT2 UC (1972).

Vennart, J., Radiotoxicity of tritium and carbon-14 compounds, Health Phys. 16, 429-440 (1969).

Vinogradov, A. P., Manganese in insects, Compt. Rend. Acad. Sci. USSR A1929, 277-320 (1929).

Vinogradov, A. P. and M. Nvestrueva, Manganese in insects, II. Compt. Rend. Acad. Sci. USSR A1930, 127-132 (1930).

Vinogradov, A. P., Manganese in insects (Formicidae), III, Compt. Rend. Acad. Sci. USSR 14, 357-359 (1937).

Vinogradov, A. P., The elementary chemical composition of marine organisms, Sears Foundation for Marine Research, Yale Univ., New Haven, 1953.

Volkova, G. A., The buildup factor of the radioisotopes of some chemical elements in aquatic insects, Entomologicheskoye Obozreniye, Vol. 17, 516-519 (1963), Trans. 63-41234 (1963).

Wangersky, P. J., Manganese in ecology, IN Radioecology, V. Schultz and A. W. Klement, Jr. (eds.), pp. 499-508, First National Symposium on Radioecology, Sept. 10-15, 1961 (1963).

Watts, J. R., and R. S. Harvey, Uptake and retention of Cs 137 by a blue-green alga in continuous flow and batch culture systems, *Limnol. Oceanogr.* 8, 45-49 (1963).

Weaver, C. L., E. D. Harward, and H. T. Peterson, Jr., Tritium in the environment from nuclear power plants, *Public Health Reports (US)* 84, 363-371 (1969).

Weinberger, D., Metabolism of hydrogen isotopes by algae, IN *Biology Research, Annual Report 1952*, pp. 72-74, Hanford Atomic Products Operation, Richland, Washington, HW-28636 (1953).

Weinberger, D. and J. W. Porter, Incorporation of tritium oxide into growing Chlorella pyrenoidosa cells, *Science* 117, 636-638 (1953).

Weston, R. E., Jr., Kinetic and equilibrium isotope effects of tritium substitution, IN *Tritium*, A. A. Moghissi and M. W. Carter (eds.), pp. 289-300 (1973), Messinger Graphics, Phoenix, TID CONF-710809.

Whilm, J. L., Transfer of radioisotopes between detritus and benthic macroinvertebrates in laboratory microecosystems, *Health Phys.* 18, 277-284 (1970).

Whitten, B. K. and C. J. Goodnight, The accumulation of strontium-89 and calcium-45 by an aquatic oligochaete, *Physiol. Zool.* 40, 371-385 (1967).

Wilke-Dorfurt, E., Über den Judgehalf der Schalen von Muscheltieren, Sum *Chemischen Teil des Kropfproblems*, II, *Biochem. Zeit.* 192, 74-82 (1928).

Williams, L. G., Uptake of cesium-137 by cells and detritus of Euglena and Chlorella, *Limnol. Oceanogr.* 5, 301-311 (1960).

Williams, L. G., Concentration of strontium-85 and cesium-137 from water solutions by Cladophora and Pithophora, *J. Phycol.* 6, 314-316 (1970).

Williams, L. G. and Q. Pickering, Direct and food-chain uptake of cesium-137 and strontium-85 in bluegill fingerlings, *Ecology* 42, 205-206 (1961).

Williams, L. G. and H. D. Swanson, Concentration of cesium-137 by algae, *Science* 127, 187-188 (1958).

Willis, D. L. and B. B. Valett, Metabolism of iodine-131 in two amphibian species (Taricha granulosa and Rana pipiens), IN *Radio-nuclides in ecosystems*, D. J. Nelson (ed.), pp. 394-400 (1973), *Proceedings of the Third National Symposium on Radioecology*, Oak Ridge, Tennessee, May 10-12, 1971.

Winchester, J. W., Chemical equilibria of iodine in Great Lakes and waters, IN Proceedings of the Thirteenth Conference on Great Lakes Research, pp. 137-140, 1970, International Association for Great Lakes Research.

Wiser, C. W. and D. J. Nelson, Uptake and elimination of cobalt-60 by crayfish, Am. Midl. Nat. 72, 181-202 (1964).

Wlodek, S., The behavior of cesium-137 in freshwater reservoirs with particular reference to the reactions of plant communities to contamination, IN Radioecological concentration processes, B. Aberg and F. P. Hungate (eds.), pp. 897-912 (1967), Proceedings of the International Symposium, Stockholm, April 25-29, 1971.

Wlodek, S., M. Bysiek, and D. Grzyboska, Autochthonous caesium and behavior of allochthonous caesium in a fish pond, Ecologia Polska, Seria A, Vol. 18(30), 1-11 (1970).

Wolff, J., Transport of iodine and other anions in the thyroid gland, Phys. Rev. 44, 45-90 (1964).

Wood, J. M., Biological cycles for toxic elements in the environment, Science 183, 1049-1051 (1974).

Woodard, H. Q., The biological effects of tritium, USAEC Report HASL-229, Health and Safety TID-4100 (1970).

Wrenn, M. E., J. W. Lentsch, M. Eisenbud, G. J. Lauer, and G. P. Howells, Radiocesium distribution in water, sediment, and biota in the Hudson River Estuary from 1964 through 1970, IN Radionuclides in Ecosystems, D. J. Nelson (ed.), pp. 334-343 (1973), Proceedings of the Third National Symposium on Radioecology, Oak Ridge, Tennessee, May 10-12, 1971.

Wyerman, T. A., R. K. Farnsworth, and G. L. Stewart, Tritium in streams in the United States, 1961-1968, Radiat. Health Data Reports 11, 421-439 (1970).

Wylie, K. F., W. A. Bigler, and G. R. Grove, Biological half-life of tritium, Health Phys. 9, 911-914 (1963).

Young, E. G., The occurrence of inorganic elements in marine algae of the Atlantic provinces of Canada, Can. J. Bot. 36, 301-310 (1958).

APPENDIX

EFFECT OF CARRIER CONCENTRATION ON
THE DISCRIMINATION COEFFICIENT

The discrimination coefficient, q_i , is usually treated as a constant. The purpose of this appendix is to show that the coefficient can vary with concentration of the carrier in the water and in other sources. Treating i as a single compartment, the time rate of change in amount, R_i , of radionuclide R in organism or tissue i is:

$$\frac{dR_i}{dt} = I_T - R_i k \quad (I-1)$$

and the steady-state amount, C_i^* , of carrier element C^* is given by

$$C_i^* = \frac{I_T^*}{k} \quad (I-2)$$

where

I_T = rate of input of radionuclide from all sources ($\mu\text{Ci/day}$)

I_T^* = rate of input of carrier element from all sources ($\mu\text{g/day}$),

k = excretion rate coefficient for the radionuclide in i (day^{-1}),

k^* = excretion rate coefficient for the carrier element in i at steady-state concentration (day^{-1}).

At steady state, Eqs. (I-1) and (I-2) imply that

$$(R/C^*)_i \equiv (R_i/C_i) = \frac{I_T k^*}{I_T^* k} \quad (I-3)$$

For the case where uptake is from a single prey j and from water:

$$I_T = I_w + I_j, \text{ and} \quad (I-4)$$

$$I_T^* = I_w^* + I_j^*, \quad (I-5)$$

where

I_w and I_j are uptake rates ($\mu\text{Ci/day}$) of a radionuclide from water and j , respectively; and I_w^* and I_j^* are uptake rates ($\mu\text{g/day}$) of carrier element from water and j , respectively.

For uptake from water:

$$I_w = a[R]_w \text{ and,} \quad (I-6)$$

$$I_w^* = a^*[C^*]_w \quad (I-7)$$

where

a = uptake rate coefficient for radionuclide from water (g/day), and

a^* = uptake rate coefficient for carrier element from water (g/day).

For uptake from food:

$$I_j = b[R]_j Q \text{ and,} \quad (I-8)$$

$$I_j^* = b^*[C^*]_j Q \quad (I-9)$$

where

b = absorbtion efficiency of radionuclide uptake from food (unitless),

b^* = absorbtion efficiency of carrier element uptake from food (unitless), and

Q = feeding rate (g/day).

Thus,

$$(R/C^*)_i = \frac{k^*}{k} \left[\frac{a[R]_w + b[R]_j Q}{a^*[C^*]_w + b^*[C^*]_j Q} \right] \quad (I-10)$$

$$= \frac{k^*}{k} \left[\frac{a[C^*]_w \left(\frac{R}{C^*} \right)_w + b[C^*]_j \left(\frac{R}{C^*} \right)_j Q}{a^*[C^*]_w + b^*[C^*]_j Q} \right] \quad (I-11)$$

Dividing Eq. (I-11) through by $(R/C^*)_w$ and incorporating the elimination coefficient of the prey, q_j , gives

$$q_i \equiv \frac{(R/C^*)_i}{(R/C^*)_w} = \frac{k^*}{k} \left[\frac{a[C^*]_w + b[C^*]_j q_j Q}{a^*[C^*]_w + b^*[C^*]_j Q} \right] , \quad (I-12)$$

Equation (I-12) may be rewritten as

$$q_i = \frac{k^*}{k} \left[\frac{a^* \left(\frac{a}{a^*} \right) [C^*]_w + b^* \left(\frac{b}{b^*} \right) [C^*]_j q_j Q}{I_T^*} \right] \quad (I-13)$$

$$= \frac{k^*}{k} \left[\frac{I_w^* \left(\frac{a}{a^*} \right) + I_j^* \left(\frac{b}{b^*} \right) q_j}{I_T^*} \right] \quad (I-14)$$

Because of homeostatic control, $[C^*]_j$ is constant. Thus, as can be seen from Equation (I-14), q_i should vary with $[C^*]_w$ since uptake from water is assumed proportional to $[C^*]_w$.

Equation (I-14) may be generalized for any set of sources:

$$q_i = \frac{k^*}{k} \left[\sum_{j \neq i} \frac{I_{ij}^*}{I_T^*} \left(\frac{a_{jj}}{a_{ij}^*} \right) q_j \right] \quad (I-15)$$

where

I_{ij}^* = intake rate of carrier element by i from j ($\mu\text{g/day}$),

and

a_{ij} = absorption efficiency of uptake of radionuclide from source j at surfaces of i (unitless), and

a_{ij}^* = absorption efficiency of uptake of carrier element from source j at surfaces of i (unitless).

If water is a source, where $j = 1$ for water, a_{i1} is absorption of the radionuclide brought to the body surfaces and $q_1 \equiv 1$.

ORNL-5002
UC-11 — Environmental
and Earth Sciences

INTERNAL DISTRIBUTION

1. S. I. Auerbach	42. F. R. O'Donnell
2. J. A. Auxier	43. R. V. O'Neill
3. C. J. Barton	44-58. D. C. Parzyck
4. B. G. Blaylock	59. H. Postma
5. R. S. Booth	60. D. E. Reichle
6. R. L. Burgess	61. C. R. Richmond
7. C. C. Coutant	62. P. S. Rohwer
8. F. L. Culler	63. E. G. Struxness
9. R. C. Dahlman	64. J. E. Till
10. J. W. Elwood	65. R. I. Van Hook
11. L. D. Eyman	66. J. P. Witherspoon
12. W. F. Harris, III	67-69. Central Research Library
13. R. F. Hibbs	70. ORNL-Y-12 Technical Library
14. S. G. Hildebrand	Document Reference Section
15. G. S. Hill	71-75. Laboratory Records Department
16-39. S. V. Kaye	76. Laboratory Records, ORNL R.C.
40. G. G. Killough, Jr.	77. ORNL Patent Office
41. R. E. Moore	

EXTERNAL DISTRIBUTION

78. L. R. Anspaugh, University of California, Lawrence Livermore Laboratory, Livermore, California 94550.
79. David S. Ballantine, Division of Biological and Environmental Research, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
80. R. F. Barker, Product Standards Branch, Directorate of Regulatory Standards, U.S. Nuclear Regulatory Commission, Washington, D. C. 20545.
81. Nathaniel F. Barr, Division of Biomedical and Environmental Research, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
82. D. S. Barth, National Environmental Research Center, EPA, P. O. Box 15027, Las Vegas, Nevada 89114.
83. Richard L. Blanchard, Radiochemistry & Nuclear Engineering Facility, U.S. Environmental Protection Agency, Cincinnati, Ohio 45268.
84. Andre Bouville, United Nations, Scientific Committee on the Effects of Atomic Radiation, New York, New York 10016.
85. J. A. Broadway, Eastern Environmental Research Facility, P. O. Box 61, Montgomery, Alabama 36101.
86. Carter D. Broyles, Department Manager, Test Sciences (7110), Sandia Corp., P. O. Box 5800, Albuquerque, New Mexico 87115.

87. W. W. Burr, Jr., Division of Biology and Medicine, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
88. M. W. Carter, Interdisciplinary Sciences, Georgia Institute of Technology, Atlanta, Georgia 20332.
89. Roger J. Cloutier, Special Training Division, Oak Ridge Associated Universities, Oak Ridge, Tennessee 37830.
90. R. G. Cochrell, Manager, Licensing Division, Westinghouse Electric Corporation, P. O. Box 158, Madison, Pennsylvania 15663.
91. Jerry Cohen, Lawrence Livermore Laboratory, P. O. Box 808, Livermore, California 94551.
92. Raymond D. Cooper, Division of Biomedical and Environmental Research, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
93. T. V. Crawford, Savannah River Laboratory, E. I. du Pont de Nemours & Company, Aiken, South Carolina 29801.
94. R. G. Cuddihy, Inhalation Toxicology Research Institute, 5200 Gibson Blvd. SE, Albuquerque, New Mexico 87108.
95. Adrian H. Dahl, U.S. Energy Research and Development Administration, Idaho Operations Office, 550 2nd St., Idaho Falls, Idaho 83401.
- 96-98. Jared J. Davis, Office of Nuclear Regulatory Research, U.S. Nuclear Regulatory Commission, Washington, D. C. 20545.
99. P. B. Dunaway, Office of Effects Evaluation, Nevada Operations Office, U.S. Energy Research and Development Administration, Las Vegas, Nevada 87114.
100. G. G. Eichholz, School of Nuclear Engineering, Georgia Institute of Technology, Atlanta, Georgia 20332.
101. Rudolf J. Engleman, Division of Biomedical and Environmental Research, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
102. Edward H. Fleming, University of California Lawrence Livermore Laboratory, Mail Stop, L-1, Box 808 Livermore, California 94550.
103. R. Franklin, Division of Biomedical and Environmental Research, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
104. Carl Gamertsfelder, Directorate of Licensing, U.S. Nuclear Regulatory Commission, Washington, D. C. 20545.
105. F. A. Gifford, Jr., Air Resources Atmospheric Turbulence and Diffusion Laboratory, NOAA, Oak Ridge, Tennessee 37830.
106. Abraham S. Goldin, Kresge Center for Environmental Health, Harvard University, School of Public Health, Boston, Massachusetts 02100.
107. R. S. Goor, National Science Foundation, 2101 Constitution Ave., Washington, D. C. 20418.
108. R. L. Gotchy, Directorate of Licensing, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
109. Douglas Grahn, Argonne National Laboratory, 9700 South Cass Avenue, Argonne, Illinois 60439.

110. Harold J. Groh, Jr., Separations Chemistry, Savannah River Laboratory, E. I. duPont de Nemours and Company, P. O. Box "A", Aiken, South Carolina 29802.
111. D. F. Harmon, Office of Regulation, U.S. Nuclear Regulatory Commission, Bethesda, Maryland 20014.
112. Monte Hawkins, Head, Safety and Environmental Control, Allied General Nuclear Services, P. O. Box 847, Barnwell, South Carolina 29812.
113. J. W. Healy, Health Physics Division, Los Alamos Scientific Laboratory, Box 1663, Los Alamos, New Mexico 87544.
114. John Heslep, Chief, Environmental Safety and Consumer Protection Program, State Department of Public Health, 2151 Berkeley Way, Berkeley, California 97404.
115. George Hinman, Gulf Central Atomic, Box 608, San Diego, California 92112.
116. I. A. Hobbs, Director, Technical and Production Division, Savannah River Operations Office, P. O. Box "A", Aiken, South Carolina 29802.
117. John B. Hursh, Department of Radiation Biology and Biophysics, University of Rochester, Rochester, New York 14620.
118. Stephen Jinks, Institute of Environmental Medicine, New York University Medical Center, Long Meadow Road, Tuxedo, New York 10987.
119. B. Kahn, Interdisciplinary Sciences, Georgia Institute of Technology, Atlanta, Georgia 20332.
- 120-169. Jacob Kastner, Radiological Assessment Branch, Directorate of Licensing, U.S. Nuclear Regulatory Commission, Washington, D. C. 20545.
- 170-171. J. R. Kercher, University of California, Lawrence Livermore Laboratory, Livermore, California 94550.
172. J. Kline, Argonne National Laboratory, Radiological Physics Division Ecological Studies, Argonne, Illinois 60439.
173. J. Koranda, Biomedical Division, Lawrence Radiation Laboratory, P. O. Box 808 (L-13), Livermore, California 94551.
174. Paul Kruger, Professor of Nuclear Chemistry, Department of Civil Engineering, Stanford University, Stanford, California.
175. William Martin, Battelle Memorial Institute, 505 King Avenue, Columbus, Ohio 43201.
176. P. J. Mellinger, EXXON Nuclear Company, 2955 George Washington Way, Richland, Washington 99352.
177. W. A. Mills, U.S. Environmental Protection Agency, Criteria and Standards Division, 401 M St., SW, Washington, D. C. 20460.
178. A. A. Moghissi, Office of Interdisciplinary Studies, Georgia Institute of Technology, Atlanta, Georgia 20332.
179. Daniel M. Montgomery, Radiochemistry and Nuclear Engineering Facility, U.S. Environmental Protection Agency, Cincinnati, Ohio 45268.
180. K. Z. Morgan, School of Nuclear Engineering, Georgia Institute of Technology, Atlanta, Georgia 20332.
181. Henry W. Morton, Nuclear Fuel Services, Inc., 6000 Executive Blvd., Suite 600, Rockville, Maryland 20852.

182. D. R. Nelson, Office of Radiation Programs, U.S. Environmental Protection Agency, 401 M Street, S. W., Washington, D. C. 20460.
183. Randolph R. Newton, Division of Biological and Environmental Research, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
184. T. R. Noonan, UT-AEC Agricultural Research Laboratory, 1299 Bethel Valley Road, Oak Ridge, Tennessee 37830.
185. H. T. Odum, Department of Environmental Engineering, University of Florida, Gainesville, Florida 32601.
186. J. S. Olson, Division of Biomedical and Environmental Research, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
187. W. S. Osburn, Division of Biomedical and Environmental Research, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
188. Charles Osterberg, Division of Biomedical and Environmental Research, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
189. Aaron Padgett, Carolina Power and Light, P. O. Box 1551, Raleigh, North Carolina 27602.
190. C. C. Palmiter, Office of Radiation Programs, Environmental Protection Agency, 401 M Street, SW, Washington, D. C. 20460.
191. F. L. Parker, Department of Civil Engineering, Vanderbilt University, Nashville, Tennessee 37203.
192. Michael A. Parsons, Radiological Assessment Branch, Directorate of Licensing, U.S. Nuclear Regulatory Commission, Washington, D. C. 20545.
193. H. R. Payne, U.S. Environmental Protection Agency Region IV, 1421 Peachtree St., Atlanta, Georgia 30309.
194. W. H. Pennington, Division of Biomedical and Environmental Research, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
195. J. F. Phillip, Director, Special Projects Division, San Francisco Operations Office, U.S. Energy Research and Development Administration, 1333 Broadway, Oakland, California 94612.
196. K. Purushothamam, University of Missouri at Rolla, Rolla Missouri 65401.
197. W. H. Ray, Office of Regulation, U.S. Nuclear Regulatory Commission, Bethesda, Maryland 20014.
198. Ronald L. Ritschard, University of California, Lawrence Livermore Laboratory, Livermore, California 94550.
199. L. Roberts, School of Nuclear Engineering, Georgia Institute of Technology, Atlanta, Georgia 20332.
200. J. N. Rogers, Division 8321, Sandia Laboratories, P. O. Box 969, Livermore, California 94550.
201. H. C. Rothschild, Radiation Protection Bureau, Brookfield Road, Ottawa K1A1C1, Ontario, Canada.
202. Keith J. Schiager, University of Pittsburgh, Radiation Center, Pittsburgh, Pennsylvania 15213.
203. G. A. Selmel, Division of Biology and Medicine, Battelle Pacific Northwest Laboratories, Richland, Washington 99352.

204. Seymour Shwiller, Staff Member, Joint Committee on Atomic Energy, Congress of the United States, Washington, D. C. 20510.
205. David S. Smith, Office of Radiation Programs, U.S. Environmental Protection Agency, 401 M Street, SW, Washington, D.C. 20460.
206. J. K. Soldat, Pacific Northwest Laboratories, Battelle Memorial Institute, Richland, Washington.
207. Jeff Swinebroad, Division of Biomedical and Environmental Research, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
208. Lauriston Taylor, National Academy of Sciences, Advisory Committee on Emergency Planning, 2101 Constitution Avenue, Washington, D. C. 20418.
209. Howard Tewes, Lawrence Livermore Laboratory, P. O. Box 808, Livermore, California 94551.
210. Paul C. Tompkins, 407 Clayton Road, Chapel Hill, North Carolina 27514.
211. J. R. Trabalka, KMS Fusion Inc., 3941 Research Park Drive, Ann Arbor, Michigan 48106.
212. B. H. Van Domelen, Governor's Science Adviser, Sandia Laboratory, 1820, Albuquerque, New Mexico 87115.
- 213-232. Henry A. Vanderploeg, Great Lakes Environmental Research Laboratory, NOAA, Ann Arbor, Michigan 48104.
233. James E. Watson, Jr., Radiological Hygiene Branch, Tennessee Valley Authority, River Oaks Building, Muscle Shoals, Alabama 35630.
234. R. Watters, Division of Biomedical and Environmental Research, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
235. Alvin M. Weinberg, Institute for Energy Analysis, Oak Ridge Associated Universities, Oak Ridge, Tennessee 37830.
236. Glenn C. Werth, Associate Director for Plowshare, Lawrence Livermore Laboratory, P. O. Box 808 (L-13), Livermore, California 94551.
237. F. W. Whicker, Department of Radiology and Radiation Biology, Colorado State University, Ft. Collins, Colorado 80521.
- 238-239. W. H. Wilcox, Environmental Quality Laboratory, 590-D N.W. Olean Boulevard, Port Charlotte, Florida 33952.
240. B. G. Wixson, University of Missouri at Rolla, Rolla, Missouri 65401.
241. R. W. Wood, Division of Biology and Medicine, U.S. Energy Research and Development Administration, Washington, D. C. 20545.
242. S. Yaniv, Office of Nuclear Regulatory Research, Mail Stop A1-3002, U.S. Nuclear Regulatory Commission, Washington, D. C. 20555.

243. Research and Technical Support Division, ERDA-ORO
244-439. Given distribution as shown in TID-4500 under category for
Environmental and Earth Sciences (25 copies - NTIS)