

CONF-9606243--23

BROOKHAVEN NATIONAL LABORATORY

---

22 November 1996

BNL - 63704

---

**Signals from Flavor Changing Scalar Currents at the Future  
Colliders\***

**David Atwood**

Theory Group, Continuous Electron Beam Accelerator Facility,  
Newport News, VA 23606d

and

**MASTER**

**L. Reina and Amarjit Soni**

Physics Department, Brookhaven National Laboratory, Upton, NY 11973

---

\* To be published in the Proceedings of the 1996 DPF/DPB Summer Study on New Directions for High Energy Physics, June 25-July 12, 1996.

This manuscript has been authored under contract number DE-AC02-76CH00016 with the U.S. Department of Energy. Accordingly, the U.S. Government retains a non-exclusive, royalty-free license to publish or reproduce the published form of this contribution, or allow others to do so, for U.S. Government purposes.

**DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED**

*LM*

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

## **DISCLAIMER**

**Portions of this document may be illegible  
in electronic image products. Images are  
produced from the best available original  
document.**

# Signals from Flavor Changing Scalar Currents at the Future Colliders \*

D. Atwood

Theory Group, Continuous Electron Beam Accelerator Facility, Newport News, VA 23606

L. Reina and A. Soni

Physics Department, Brookhaven National Laboratory, Upton, NY 11973

## ABSTRACT

We present a general phenomenological analysis of a class of Two Higgs Doublet Models with Flavor Changing Neutral Currents arising at the tree level. The existing constraints mainly affect the couplings of the first two generations of quarks, leaving the possibility for non negligible Flavor Changing couplings of the top quark open. The next generation of lepton and hadron colliders will offer the right environment to study the physics of the top quark and to unravel the presence of new physics beyond the Standard Model. In this context we discuss some interesting signals from Flavor Changing Scalar Neutral Currents.

## I. GENERAL FRAMEWORK

The next generation of lepton and hadron colliders will play a fundamental role in the study of new physics beyond the Standard Model (SM). Higher energies will allow a careful study of the physics of the top quark (its couplings in particular) and of the scalar and gauge sector of the fundamental theory of elementary particles.

In this context, we have analyzed the possibility of having a Two Higgs Doublet Model (2HDM) with Flavor Changing Neutral Currents (FCNC's) allowed at the tree level [1]-[3]. This Model constitutes a simple extension of the scalar sector of the Standard Model and closely mimics the Higgs sector of a SuperSymmetric Theory (SUSY). However, the possibility of having flavor changing (FC) tree level couplings in the neutral scalar sector definitely distinguishes it from both the SM and SUSY. Moreover, the discovery and study of extra scalar or pseudoscalar, neutral and charged particles with not too heavy masses will be in the reach of the future machines. From here our interest.

Although there is no *a priori* veto to the existence of FCNC at the tree level, the low energy phenomenology of the K- and of the B-meson as well as the existing precision measurements of the SM impose strong constraints on the possibility of having sizable effects from FCNC. However, under suitable assumptions, the FC couplings of the top quark partially escape these constraints and can be predicted to give non negligible signals as we will illustrate in the following.

### A. The Model

A mild extension of the SM with one additional scalar SU(2) doublet opens up the possibility of flavor changing scalar currents (FCSC's) at the tree level. In fact, when the up-type quarks and the down-type quarks are allowed simultaneously to couple

to more than one scalar doublet, the diagonalization of the up-type and down-type mass matrices does not automatically ensure the diagonalization of the couplings with each single scalar doublet. For this reason, the 2HDM scalar potential and Yukawa Lagrangian are usually constrained by an *ad hoc* discrete symmetry [4], whose only role is to protect the model from FCSC's at the tree level. Let us consider a Yukawa Lagrangian of the form

$$\mathcal{L}_Y = \eta_{ij}^U \bar{Q}_{i,L} \tilde{\phi}_1 U_{j,R} + \eta_{ij}^D \bar{Q}_{i,L} \phi_1 D_{j,R} + \xi_{ij}^U \bar{Q}_{i,L} \tilde{\phi}_2 U_{j,R} + \xi_{ij}^D \bar{Q}_{i,L} \phi_2 D_{j,R} + h.c. \quad (1)$$

where  $\phi_i$ , for  $i = 1, 2$ , are the two scalar doublets of a 2HDM, while  $\eta_{ij}^{U,D}$  and  $\xi_{ij}^{U,D}$  are the non diagonal matrices of the Yukawa couplings. Imposing the following *ad hoc* discrete symmetry

$$\begin{aligned} \phi_1 &\rightarrow -\phi_1 & \text{and} & \phi_2 \rightarrow \phi_2 \\ D_i &\rightarrow -D_i & \text{and} & U_i \rightarrow \mp U_i \end{aligned} \quad (2)$$

some of the terms in  $\mathcal{L}_Y$  have to be dropped and one obtains the so called Model I and Model II, depending on whether the up-type and down-type quarks are coupled to the same or to two different scalar doublets respectively [5].

In contrast we will consider the case in which no discrete symmetry is imposed and both up-type and down-type quarks then have FC couplings. For this type of 2HDM, which we will call Model III, the Yukawa Lagrangian for the quark fields is as in Eq. (1) and no term can be dropped *a priori*, see also refs. [6, 7].

For convenience we can choose to express  $\phi_1$  and  $\phi_2$  in a suitable basis such that only the  $\eta_{ij}^{U,D}$  couplings generate the fermion masses, i.e. such that

$$\langle \phi_1 \rangle = \begin{pmatrix} 0 \\ v/\sqrt{2} \end{pmatrix}, \quad \langle \phi_2 \rangle = 0. \quad (3)$$

The two doublets are in this case of the form

$$\begin{aligned} \phi_1 &= \frac{1}{\sqrt{2}} \left[ \begin{pmatrix} 0 \\ v+H^0 \end{pmatrix} + \begin{pmatrix} \sqrt{2}\chi^+ \\ i\chi^0 \end{pmatrix} \right] \\ \phi_2 &= \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2}H^+ \\ H^1 + iH^2 \end{pmatrix}. \end{aligned} \quad (4)$$

The scalar Lagrangian in the  $(H^0, H^1, H^2, H^\pm)$  basis is such that [8, 5]: the doublet  $\phi_1$  corresponds to the scalar doublet of the SM and  $H^0$  to the SM Higgs field (same couplings and no interactions with  $H^1$  and  $H^2$ ); all the new scalar fields belong

\* Work supported by U.S. Department of Energy contracts DC-AC05-84ER40150 (CEBAF) and DE-AC-76CH0016 (BNL).

described by introducing the parameter  $\rho_0$  [15, 16], defined as

$$\rho_0 = \frac{M_W^2}{\rho M_Z^2 \cos^2 \theta_W} \quad (12)$$

where the  $\rho$  parameter absorbs all the SM corrections to the gauge boson self energies. In the presence of new physics

$$\rho_0 = 1 + \Delta\rho_0^{\text{NEW}} \quad (13)$$

From the recent global fits of the electroweak data, which include the input for  $m_t$  from Ref. [17] and the new experimental results on  $R_b$ ,  $\rho_0$  turns out to be very close to unity [16, 9, 10]. This impose severe constraints on many extension of the SM, especially on the mass range of the new particles.

As is the case in 2HDM's with no FCNC's, it is very difficult to reconcile the measured values of the previous three observables in the presence of an extended scalar sector. Taking into account also the constraints from the  $F^0 - \tilde{F}^0$  mixings, two main scenarios emerge depending on the choice of enforcing or not  $R_b^{\text{exp}}$  [10].

1. If we **enforce the constraint from  $R_b^{\text{exp}}$**  (see Eq. (11)); then we can accommodate the present measurement of the  $Br(B \rightarrow X_s \gamma)$  (see Eq. (9)) and of the  $\Delta F = 2$  mixings (see Eq. (7)) and at the same time satisfy the global fit result for the  $\rho$  parameter [16] provided the following conditions are satisfied.

- i) The neutral scalar  $h^0$  and the pseudoscalar  $A^0$  are very light, i.e.

$$50 \text{ GeV} \leq M_h \sim M_A < 70 \text{ GeV}. \quad (14)$$

- ii) The charged scalar  $H^+$  is heavier than  $h^0$  and  $A^0$ , but not too heavy to be in conflict with the constraints from the  $\rho$  parameter. Thus

$$150 \text{ GeV} \leq m_c \leq 200 \text{ GeV}. \quad (15)$$

- iii) The  $\xi_{ij}^D$  couplings are enhanced with respect to the  $\xi_{ij}^U$  ones

$$\begin{aligned} \lambda_{bb} &\gg 1 \text{ and } \lambda_{tt} \ll 1 \\ \lambda_{sb} &\gg 1 \text{ and } \lambda_{ct} \ll 1. \end{aligned} \quad (16)$$

The choice of the phase  $\alpha$  is not as crucial as the above conditions and therefore we do not make any assumption on it.

2. If we **disregard the constraint from  $R_b^{\text{exp}}$**  there is no need to impose the bounds of Eqs. (14)-(16) and we can safely work in the scenario in which only the first generation FC couplings are suppressed

$$\lambda_{ui}, \lambda_{dj} \ll 1 \quad \text{for } i, j = 1, 2, 3 \quad (17)$$

in order to satisfy the experimental constraints on the  $F^0 - \tilde{F}^0$  mixings. We will assume the FC couplings of the second and third generations to be given by Eq. (6) with

$$\lambda_{ct} \simeq O(1) \text{ and } \lambda_{sb} \simeq O(1). \quad (18)$$

The value of the mixing angle  $\alpha$  is not relevant, while the masses are mainly dictated by the fit to  $Br(B \rightarrow X_s \gamma)$  and  $\Delta\rho_0$  [9]

$$M_H, M_h \leq M_c \leq M_A \text{ and } M_A \leq M_c \leq M_H, M_h. \quad (19)$$

We can see that, except in a very narrow window of the parameter space, it is in general very difficult to accommodate the present value of  $R_b^{\text{exp}}$  in Model III. Due to the present unclear experimental situation for  $R_b$ , we will mainly concentrate on the second scenario<sup>2</sup>. This scenario has the very interesting characteristics of providing sizable FC couplings for the top quark, in a way that will certainly be testable at the next generation of lepton and hadron colliders. We will discuss some of these phenomenological issues in the next section.

## II. SIGNALS OF TOP-CHARM PRODUCTION

If we assume  $\lambda_{sb} \simeq O(1)$  and  $\lambda_{ct} \simeq O(1)$  as in Eq. (18),  $\xi_{ct}^U$  becomes the most relevant FC coupling. The presence of a  $\xi_{ct}^U$  flavor changing coupling can be tested by looking at both top decays and top production (see ref. [10] and references therein). We want to concentrate here on top-charm production at lepton colliders, both  $e^+e^-$  and  $\mu^+\mu^-$ , because, as we have emphasized before [7, 18], in this environment the top-charm production has a particularly clean and distinctive signature. The SM prediction for this process is extremely suppressed and any signal would be a clear evidence of new physics with large FC couplings in the third family. Moreover it has a very distinctive signature, with a very massive jet recoiling against an almost massless one (very different from a  $b\bar{s}$  signal, for instance). This characteristic is enhanced even more in the experimental environment of a lepton collider.

In principle, the production of top-charm pairs arises both at the tree level, via the  $s$  channel exchange of a scalar field with FC couplings, and at the one loop level, via corrections to the  $Ztc$  and  $\gamma tc$  vertices. The  $s$  channel top-charm production is one of the new interesting possibilities offered by a  $\mu^+\mu^-$  collider in studying the physics of standard and non standard scalar fields. However, it is not relevant for an  $e^+e^-$  collider, because the coupling of the scalar fields to the electron is likely to be very suppressed (see Eq. (6)). Therefore we will consider these two cases separately.

In the case of an  $e^+e^-$  collider, top-charm production arises via  $\gamma$  and  $Z$  boson exchange, i.e. the process  $e^+e^- \rightarrow \gamma^*, Z^* \rightarrow$

<sup>2</sup>See ref. [10] for a discussion of the scenario which accommodates  $R_b^{\text{exp}}$ .

in which  $\mathcal{H} = h^0$ , and  $\alpha = 0$  (case 1) or  $\alpha = \pi/4$  (case 2). We estimate that for a Higgs particle of  $m_{\mathcal{H}} = 300$  GeV, a luminosity of  $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$  and a year of  $10^7 \text{ s}$  (1/3 efficiency), a sample of  $tc$  events ranging from almost one hundred to few thousands can be produced [18]. Given the distinctive nature of the final state and the lack of a Standard Model background, the predicted luminosity should allow the observation of such events. Therefore many properties of the Higgs- $tc$  coupling could be studied in detail.

Finally we want to consider the impact that a tree level  $\xi_{ct}^U$  coupling could have on the present scenario of the Higgs discovery. As was already pointed out in the literature [21], if  $M_{\mathcal{H}} > m_t$  (for  $\mathcal{H} = \bar{H}^0, h^0$  or  $A^0$ ) Model III allows the new decay channel

$$\mathcal{H} \rightarrow c\bar{t} + \bar{c}t \quad (23)$$

which should also be considered in the search for a non standard Higgs particle. In the mass range  $m_t < M_{\mathcal{H}} < 2m_t$ , this single top production is of particular interest because its rate can be greater than the rate for  $\mathcal{H} \rightarrow b\bar{b}$  while the decay  $\mathcal{H} \rightarrow t\bar{t}$  is not yet possible. Assuming Eq. (6), the rate for  $\mathcal{H} \rightarrow c\bar{t} + \bar{c}t$  is given by

$$\begin{aligned} \Gamma(\mathcal{H} \rightarrow c\bar{t} + \bar{c}t) &= N_c \frac{G_F}{4\sqrt{2}\pi} M_{\mathcal{H}} \lambda_{ct}^2 m_c m_t \cdot \\ &\left[ 1 - \frac{(m_t + m_c)^2}{M_{\mathcal{H}}^2} \right]^{3/2} \left[ 1 - \frac{(m_t - m_c)^2}{M_{\mathcal{H}}^2} \right]^{1/2} \end{aligned} \quad (24)$$

to be compared with the rate for  $\mathcal{H} \rightarrow q\bar{q}$ , i.e.

$$\Gamma(\mathcal{H} \rightarrow q\bar{q}) = N_c \frac{G_F}{4\sqrt{2}\pi} M_{\mathcal{H}} m_q^2 \left[ 1 - 4 \frac{m_q^2}{M_{\mathcal{H}}^2} \right]^{3/2} \quad (25)$$

We see for instance that for  $M_{\mathcal{H}} \simeq 300$  GeV,  $\Gamma(\mathcal{H} \rightarrow c\bar{t} + \bar{c}t) \sim 6\lambda_{ct}^2 \Gamma(\mathcal{H} \rightarrow b\bar{b})$ . Therefore, depending on  $\lambda_{ct}$ , there are cases in which in the range  $m_t < M_{\mathcal{H}} < 2m_t$  we could predict a distinctive signal, both with respect to the SM and to SUSY. When  $\mathcal{H} = h^0, \bar{H}^0$  then  $\mathcal{H} \rightarrow c\bar{t} + \bar{c}t$  competes only with the decays  $\mathcal{H} \rightarrow ZZ$  or  $WW$ , depending on the value of the phase  $\alpha$ . In the case  $\mathcal{H} = A^0$  the decays into gauge boson pairs are absent.

When the phase  $\alpha$  is chosen in such a way that the couplings  $\mathcal{H}ZZ$  and  $\mathcal{H}WW$  (for  $\mathcal{H} = h^0, \bar{H}^0$ ) are suppressed, the decay we are interested in can be produced for instance via  $e^+e^- \rightarrow h^0A^0, h^0\bar{H}^0 \rightarrow (t\bar{c} + c\bar{t})q\bar{q}$  (lepton collider) or  $gg \rightarrow \mathcal{H} \rightarrow t\bar{c} + c\bar{t}$  (hadron collider). Therefore both NLC and LHC should be able to look for it: the first one would offer the possibility of a much cleaner signal while the second one would provide a much higher statistics. As is the case of many other decays, a good b-tagging is clearly necessary. However the kinematic constraints of the  $\mathcal{H} \rightarrow t\bar{c} + c\bar{t}$  decay should be so distinctive to limit the size of the background. We think that dedicated simulations and systematic studies of the background will be useful in understanding the real potentiality of this decay channel.

In conclusion, we think that Model III offers a simple but interesting example in which some important topics of the physics at the future colliders can be investigated. With a few assumptions we are able to propose some distinctive processes, the existence of which would be clear evidence of some very new physics beyond the Standard Model.

### III. REFERENCES

- [1] T.P. Cheng and M. Sher, Phys. Rev. **D35**, 3484 (1987); **D44**, 1461 (1991); See also Ref. [2].
- [2] A. Antaramian, L.J. Hall, and A. Rasin, Phys. Rev. Lett. **69**, 1871 (1992).
- [3] L.J. Hall and S. Weinberg, Phys. Rev. **D48**, R979 (1993).
- [4] S. Glashow and S. Weinberg, Phys. Rev. **D15**, 1958 (1977).
- [5] For a review see J. Gunion, H. Haber, G. Kane, and S. Dawson, *The Higgs Hunter's Guide*, (Addison-Wesley, New York, 1990).
- [6] M. Luke and M.J. Savage, Phys. Lett. **B307**, 387 (1993).
- [7] D. Atwood, L. Reina and A. Soni, Phys. Rev. **D53**, 1199 (1996);
- [8] C.D. Froggatt, R.G. Moorhouse and I.G. Knowles, Nucl. Phys. **B386**, 63 (1992).
- [9] D. Atwood, L. Reina and A. Soni, hep-ph/9603210, to appear in Phys. Rev. D, and references therein.
- [10] D. Atwood, L. Reina and A. Soni, JL-TH-96-15 and hep-ph/9609280, and references therein.
- [11] A. Das and C. Kao, Phys. Lett. **B372**, 106 (1996).
- [12] R. Ammar *et al.*, CLEO Collaboration, Phys. Rev. Lett. **71**, 674 (1993); M.S. Alam *et al.*, CLEO Collaboration, Phys. Rev. Lett. **74**, 2885 (1995).
- [13] Results presented at the *XXVIII International Conference on High Energy Physics*, Warsaw, July 1996 and at the APS-DPF96 Meeting, Minneapolis, August 1996.
- [14] P. Bamert, C.P. Burgess, J.M. Cline, D. London and E. Nardi, hep-ph/9602438, to appear in Phys. Rev. D.
- [15] *Review of Particles Properties*, Phys. Rev. **D50** (1994).
- [16] P. Langacker, hep-ph/9412361. To be published in *"Precision Tests of the Standard Electroweak Model"*, ed. by P. Langacker, (World Scientific 1994).
- [17] F. Abe *et al.*, [CDF], Phys. Rev. Lett. **74**, 2626 (1995); S. Abachi *et al.*, [D0], Phys. Rev. Lett. **74**, 2632 (1995).
- [18] D. Atwood, L. Reina and A. Soni, Phys. Rev. Lett. **75**, 3800 (1995).
- [19] R.B. Palmer, preprint, "High Frequency  $\mu^+ \mu^-$  Collider Design," SLAC-AAS-Note81, 1993.
- [20] V. Barger, M. Berger, J. Gunion, and T. Han, Phys. Rev. Lett. **75**, 1462 (1995).
- [21] W.S. Hou, Phys. Lett. **B296**, 179 (1992); D. Chang, W.S. Hou and W.Y. Keung, Phys. Rev. **D48**, 217 (1993).