

CANONICAL HAMILTONIAN GUIDING CENTER VARIABLES

Roscoe B. White

Princeton Plasma Physics Laboratory

Princeton, New Jersey 08543

Abstract

A simplification of the canonical Hamiltonian variables for the guiding center motion of a charged particle in a general toroidal field is obtained using the Lagrangian formalism of Littlejohn.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Hamiltonian guiding center calculation of particle motion in general toroidal fields is an effective way to examine long-time particle confinement and induced particle losses due to static magnetic perturbations or magnetohydrodynamic fluctuations.^{1,2} The expansion parameter for the derivation of equations of motion is the ratio of gyroradius to system size, ρ . Recently, the derivation of the canonical variables for a given magnetic field representation has been considerably simplified by an elegant Lagrangian formalism due to Littlejohn.³⁻⁵ However, for a general magnetic field the canonical variables are algebraically complicated due to nonorthogonality of the magnetic coordinates. In this note we show that this complexity can be removed by a modification of the parallel velocity which is of second order in ρ . This simplification of the canonical variables, of use in the study of guiding center motion, is also of interest for the formulation of gyrokinetics in general toroidal fields.⁶

First consider magnetic configurations with toroidal symmetry and exact magnetic surfaces. Cases of broken symmetry are treated using perturbation theory.

In a general magnetic configuration possessing nested toroidal magnetic surfaces the field can be written,² in contravariant and covariant form, as

$$\mathbf{B} = \nabla\psi \times \nabla\theta - \nabla\psi_p \times \nabla\zeta, \quad (1)$$

$$\mathbf{B} = g(\psi_p)\nabla\zeta + I(\psi_p)\nabla\theta + \delta(\psi_p, \theta)\nabla\psi_p. \quad (2)$$

The magnetic coordinates $\psi, \psi_p, \zeta, \theta$ are, respectively, toroidal and poloidal flux and toroidal and poloidal angles. The functions g and I are related to the

poloidal and toroidal current, and δ describes the degree of nonorthogonality of the coordinate system. The field lines are straight in the ζ, θ plane, i.e., $d\psi/d\psi_p = q(\psi_p)$. The fact that I is a function of ψ_p only is the result of a special choice for the Jacobian $J = \nabla\psi_p \cdot (\nabla\theta \times \nabla\zeta)$, with $B^2 J = I + gq$.

Derivation of canonical variables is facilitated by the use of the guiding center Lagrangian due to Littlejohn,^{3,4} which has the form

$$L = \mathbf{A}^* \cdot \mathbf{v} + \mu \dot{\xi} - H \quad (3)$$

with \mathbf{v} the guiding center velocity, $\mathbf{A}^* = \mathbf{A} + \rho_{\parallel} \mathbf{B}$ the modified vector potential, $\rho_{\parallel} = v_{\parallel}/B$ the normalized parallel velocity, μ the magnetic moment, ξ the gyrophase and

$$H = \frac{\rho_{\parallel}^2 B^2}{2} + \mu B + \Phi, \quad (4)$$

the Hamiltonian. The field magnitude B is a function of ψ_p and θ only, with ζ an ignorable coordinate. We use units defined by the on-axis gyrofrequency (time) and the major radius (distance). Originally, canonical Hamiltonian variables were found^{1,2} by the rather laborious procedure of first finding the time variation, following a particle orbit, of the variables $\psi, \theta, \zeta, \rho_{\parallel}$ by using the guiding center motion \mathbf{v} , e.g., $\dot{\psi} = \mathbf{v} \cdot \nabla\psi$. Here and in the following all quantities are functions of the location, i.e., of ψ_p, θ, ζ only. The time derivative refers only to the change following a particle orbit. A convenient form for the guiding center motion, due to Morozov and Soloviev,⁷ is

$$\mathbf{v} = \rho_{\parallel} (\mathbf{B} + \nabla \times \rho_{\parallel} \mathbf{B}) / (1 + \rho_{\parallel} \hat{\mathbf{b}} \cdot \nabla \times \hat{\mathbf{b}}) \quad (5)$$

with $\hat{b} = \mathbf{B}/B$, where in this equation ρ_{\parallel} is considered a function of position through Eq. (4) with H replaced by the energy, and the denominator is a correction to the parallel drift discussed by several authors.⁸⁻¹¹ This second order parallel drift correction is introduced strictly to preserve the Hamiltonian character of the motion. There are other second order terms to \mathbf{v} , neglected to preserve Hamiltonian character order by order. Canonical variables were then found by writing the obtained expressions for $\dot{\psi}, \dot{\theta}, \dot{\zeta}, \dot{\rho}_{\parallel}$ in terms of partial derivatives of the Hamiltonian, and finding the Darboux transformation to put the equations in canonical form. Littlejohn⁵ subsequently pointed out that a Lagrangian formalism made this procedure much less arduous, and significantly more transparent. We will see that the Lagrangian formalism also allows for the simplification of the canonical variables through a redefinition of the guiding center.

Canonical variables are obtained by substituting expressions for \mathbf{A} and \mathbf{B} in terms of magnetic coordinates into Eq. (3), and rewriting it in the form

$$L = \sum_i P_i \dot{q}_i - H. \quad (6)$$

If this form can be obtained, the canonical momenta and coordinates are immediately identified.

Rewriting Eq. (1) in the form $\mathbf{B} = \nabla \times (\psi \nabla \theta - \psi_p \nabla \zeta) = \nabla \times \mathbf{A}$ and substituting \mathbf{A} and \mathbf{B} into the Lagrangian, we find

$$L = (\psi + \rho_{\parallel} J) \dot{\theta} + (\rho_{\parallel} g - \psi_p) \dot{\zeta} + \mu \dot{\xi} - H + \delta \rho_{\parallel} \dot{\psi}_p. \quad (7)$$

Except for the last term, this has the form given by Eq. (6). However, it can be put in canonical form. The appropriate modification, as found in Ref. 2, is given by using the poloidal variable $\theta_c = \theta - \rho_{\parallel} \delta / q$. Substituting $\dot{\theta} = \dot{\theta}_c + (d/dt)(\rho_{\parallel} \delta / q)$ into Eq. (7) and discarding the exact differential $(d/dt)(\rho_{\parallel} \delta \psi / q)$ as well as a term of order ρ^3 , we are left with $L = P_{\theta} \dot{\theta}_c + P_{\zeta} \dot{\zeta} + \mu \dot{\zeta} - H$, with

$$\begin{aligned} P_{\theta} &= \psi + \rho_{\parallel} I \\ P_{\zeta} &= \rho_{\parallel} g - \psi_p. \end{aligned} \quad (8)$$

Note that these expressions are immediately recognizable as the covariant components of the sum of particle momentum and vector potential, just as in classical electrodynamics. Explicit equations of motion are readily obtained. First note from Eq. (8) that ψ_p and ρ_{\parallel} are functions of P_{θ} , P_{ζ} only, and that

$$\begin{aligned} \frac{\partial \psi_p}{\partial P_{\theta}} &= \frac{g}{D} & \frac{\partial \psi_p}{\partial P_{\zeta}} &= \frac{-I}{D} \\ \frac{\partial \rho_{\parallel}}{\partial P_{\theta}} &= \frac{1 - \rho_{\parallel} g'}{D} & \frac{\partial \rho_{\parallel}}{\partial P_{\zeta}} &= \frac{q + \rho_{\parallel} I'}{D} \end{aligned} \quad (9)$$

with $D = gq + I + \rho_{\parallel}[gI' - Ig']$, and primes indicate differentiation with respect to ψ_p . It is then straightforward to find the explicit form of the equations of motion from $\dot{P}_{\zeta} = -\partial H / \partial \zeta$, $\dot{P}_{\theta} = -\partial H / \partial \theta$, $\dot{\theta}_c = \partial H / \partial P_{\theta}$, $\dot{\zeta} = \partial H / \partial P_{\zeta}$. For simplicity we set the electrostatic potential Φ equal to zero. The equations of motion take the form, to second order in ρ

$$\dot{\zeta} = \frac{\rho_{\parallel} B^2 (q + \rho_{\parallel} I')}{D} - (\mu + \rho_{\parallel}^2 B) \frac{\partial B}{\partial \psi_p} \frac{I}{D} + (\mu + \rho_{\parallel}^2 B) \frac{\partial B}{\partial \theta} \frac{\delta}{D}, \quad (10)$$

$$\dot{\theta}_c = \frac{\rho_{\parallel} B^2 (1 - \rho_{\parallel} g')}{D} + (\mu + \rho_{\parallel}^2 B) \frac{\partial B}{\partial \psi_p} \frac{g}{D} + (\mu + \rho_{\parallel}^2 B) \frac{\partial B}{\partial \theta} \frac{\delta}{Dq}, \quad (11)$$

$$\dot{\psi}_p = \frac{-g}{D} (\mu + \rho_{\parallel}^2 B) \frac{\partial B}{\partial \theta}, \quad (12)$$

$$\dot{\rho}_{\parallel} = -\frac{(\mu + \rho_{\parallel}^2 B)}{D} (1 - \rho_{\parallel} g') \frac{\partial B}{\partial \theta}, \quad (13)$$

where we have used the definition of θ_c to find that within corrections of higher order in ρ

$$\left. \frac{\partial \theta}{\partial P_{\theta}} \right|_{\theta_c, P_{\zeta}, \zeta} = \frac{\delta}{qD}, \quad \left. \frac{\partial \theta}{\partial P_{\zeta}} \right|_{\theta_c, P_{\theta}, \zeta} = \frac{\delta}{D}. \quad (14)$$

Equations (10–13) clearly describe Hamiltonian motion. Directly calculating $\mathbf{v} \cdot \nabla \theta$, $\mathbf{v} \cdot \nabla \zeta$ by substituting Eqs. (1,2) into Eq. (5) and using

$$\hat{\mathbf{b}} \cdot \nabla \times \hat{\mathbf{b}} = \frac{gI' - Ig' - g \partial \delta / \partial \theta}{gq + I}, \quad (15)$$

we find that \mathbf{v} given by Eq. (5) differs from the Hamiltonian motion by a second order term. Second order terms in the velocity can be modified by the addition of exact derivatives. For example, the term

$$\frac{d}{dt}(\rho_{\parallel} \delta) \approx \frac{\rho_{\parallel}^2 B^2}{B} \frac{\partial \delta}{\partial \theta} - \frac{\delta}{D} (\mu + \rho_{\parallel}^2 B) \frac{\partial B}{\partial \theta}, \quad (16)$$

where terms of higher order in ρ have been discarded, if added to Eq. (10) eliminates the term in δ and adds a $\partial \delta / \partial \theta$ term almost equal to that given by Eq. (15). Within the guiding center description such a modification of the velocity is not significant. It produces nonsecular, periodic motion of the magnitude of the gyroradius. This suggests that simplification of the Hamiltonian variables can be obtained by the addition of such terms.

The Lagrangian formalism can be used to retain the variables θ and ζ as canonical, eliminating the term containing δ in Eq. (7) through a modification of the second order parallel guiding center drift. Return to Eq. (3) and replace the guiding center velocity \mathbf{v} by $\mathbf{v} + \mathbf{w}$, with \mathbf{v} describing the guiding center motion, i.e., $\mathbf{v} \cdot \nabla\theta = \dot{\theta}$. This adds a new term to Eq. (7) of the form $\mathbf{A}^* \cdot \mathbf{w}$, and we choose \mathbf{w} so that

$$\mathbf{A}^* \cdot \mathbf{w} = -\delta \rho_{\parallel} \dot{\psi}_p. \quad (17)$$

First note that a change in gauge $\mathbf{A} \rightarrow \mathbf{A} + \nabla G$ adds an exact derivative to L . Without loss of generality we choose the gauge $\mathbf{B} \cdot \mathbf{A} = 0$. Noting from Eqs. (12) and (13) that to leading order $\dot{\psi}_p = g \dot{\rho}_{\parallel}$, we find

$$\mathbf{w} = -\frac{\delta \dot{\rho}_{\parallel} g \mathbf{B}}{B^2}. \quad (18)$$

With this change the canonical variables become $\theta, P_{\theta}, \zeta, P_{\zeta}$ and the terms proportional to δ are eliminated from Eqs. (10) and (11). Directly from the Hamiltonian we find

$$\dot{\zeta} = \frac{\rho_{\parallel} B^2}{D} (q + \rho_{\parallel} I') - (\mu + \rho_{\parallel}^2 B) \frac{I}{D} \frac{\partial B}{\partial \psi_p}, \quad (19)$$

$$\dot{\theta} = \frac{\rho_{\parallel} B^2}{D} (1 - \rho_{\parallel} g') + (\mu + \rho_{\parallel}^2 B) \frac{g}{D} \frac{\partial B}{\partial \psi_p}, \quad (20)$$

$$\dot{\psi}_p = \frac{-g}{D} (\mu + \rho_{\parallel}^2 B) \frac{\partial B}{\partial \theta}, \quad (21)$$

$$\dot{\rho}_{\parallel} = -\frac{(\mu + \rho_{\parallel}^2 B)}{D} (1 - \rho_{\parallel} g') \frac{\partial B}{\partial \theta}. \quad (22)$$

Discarding an exact derivative, we find

$$\mathbf{w} \cdot \nabla \theta = \rho_{\parallel} \frac{d}{dt} \left(\frac{g\delta}{D} \right) \approx \frac{\rho_{\parallel}^2 B^2}{D^2} g \frac{\partial \delta}{\partial \theta}. \quad (23)$$

Similarly discarding an exact derivative in $\mathbf{w} \cdot \nabla \zeta$, we find $(\mathbf{v} - \mathbf{w}) \cdot \nabla \zeta = \dot{\zeta}$, $(\mathbf{v} - \mathbf{w}) \cdot \nabla \theta = \dot{\theta}$. Thus we have obtained simplified expressions for the motion, with the magnetic coordinates θ, ζ as canonical Hamiltonian variables. The velocity differs from the usual expression, Eq. (5), only by the second order parallel velocity \mathbf{w} , which is zero if the magnetic coordinates are orthogonal. Since the second order parallel velocity is introduced only to maintain the Hamiltonian form, this is not a significant difference. Note that the function $\delta(\psi_p, \theta)$ has disappeared entirely from the equations of motion. All of the dependence of the description of the motion appears in the functions g, q, I , and B .

Now consider the addition of a general magnetic perturbation, described by the vector potential

$$\tilde{\mathbf{A}} = \alpha \nabla \zeta + \beta \nabla \theta + \gamma \nabla \psi_p \quad (24)$$

with α, β, γ general functions of position. The Lagrangian then becomes $L = (P_\theta + \beta)\dot{\theta} + (P_\zeta + \alpha)\dot{\zeta} + (\delta\rho_{\parallel} + \gamma)\dot{\psi}_p + \mu\dot{\xi} - H$ with P_θ, P_ζ given by Eq. (8). Thus we identify the canonical variables as

$$\begin{aligned} \theta & \quad P'_\theta = P_\theta + \beta(\psi_p, \theta, \zeta) \\ \zeta & \quad P'_\zeta = P_\zeta + \alpha(\psi_p, \theta, \zeta) \end{aligned} \quad (25)$$

and again eliminate the term $(\delta\rho_{\parallel} + \gamma)\dot{\psi}_p$ from the Lagrangian by a modification of the second order parallel velocity \mathbf{w} with $\mathbf{A}^* \cdot \mathbf{w} = -(\delta\rho_{\parallel} + \gamma)\dot{\psi}_p$. We then

find

$$w = -\frac{(\delta\rho_{\parallel} + \gamma)\dot{\psi}_p \mathbf{B}}{B^2}. \quad (26)$$

To find an expression for w , it is necessary to find the equations of motion to leading order. Note that, in general, in order to preserve the advantages arising from using a Hamiltonian formalism in numerical calculations of particle orbits, all orders in ρ must be retained in the equations of motion. Truncation of these equations is useful only to calculate leading order effects.

The exact equations of motion in the perturbed field are readily found. We simply sketch the procedure here, since the results are algebraically very cumbersome, and for most applications a perturbed field of the restricted form $\tilde{\mathbf{A}} = \alpha\mathbf{B}$ is sufficient, and much easier to implement.¹ From Eq. (25), using $\psi_p = \psi_p(P_\theta, P_\zeta)$ and Eq. (9) all of the partial derivatives of the transformation to the new canonical variables can be calculated. Finding the equations of motion is then an exercise in the use of the chain rule. We readily find, for example,

$$\begin{aligned} \dot{\psi}_p = & \frac{-g}{D}(\rho_{\parallel}^2 B + \mu) \frac{\partial B}{\partial \theta} + \left[\frac{I}{D} \frac{\partial \alpha}{\partial \theta} - \frac{g}{D} \frac{\partial \beta}{\partial \theta} \right] \frac{\rho_{\parallel} B^2}{D} \\ & + \left[\frac{I}{D} \frac{\partial \alpha}{\partial \zeta} - \frac{g}{D} \frac{\partial \beta}{\partial \zeta} \right] \frac{\rho_{\parallel} B^2 q}{D} \end{aligned} \quad (27)$$

plus terms of order ρ^3 .

Exact time derivatives of order ρ^2 discarded at various stages in these calculations integrate to zero by virtue of orbits closing in the ψ_p, θ plane, a property depending on the existence of the ignorable coordinate ζ . This is no longer

the case in the presence of field perturbations. If the perturbations are small, i.e., the orbits close except for small amounts $\Delta\theta, \Delta\psi_p$, then additional small changes are produced by these terms of order $\rho\Delta\theta, \rho\Delta\psi_p$. Motion across the drift surfaces produced by the neglected terms is smaller than that produced by the field perturbation by a factor ρ . If, however, the field is stochastic, then the initial guiding center coordinates do not uniquely determine the orbit. The motion is sensitive to the exact particle location, and depends for example on the initial gyrophase as well as the guiding center position. The inaccuracies given by discarding order ρ^2 exact derivatives are no larger than these initial condition inaccuracies. In such a system it makes little sense to discuss precise individual particle orbits. Only statistical quantities obtained using a distribution of initial values are relevant, and thus these terms can also be safely ignored in this case.

ACKNOWLEDGMENTS

The author would like to thank Allen Boozer and Harold Weitzner for useful discussions. This work is supported by U.S. Department of Energy Contract No. DE-AC02-76-CHO3073.

References

- ¹R.B. White, A.H. Boozer, and Ralph Hay, *Phys. Fluids* **25**, 575 (1982).
- ²R.B. White and M.S. Chance, *Phys. Fluids* **27**, 2455 (1984).
- ³R.G. Littlejohn, *J. Plasma Physics* **29**, 111 (1983).
- ⁴R.G. Littlejohn, *Physica Scripta* **T2/1**, 119 (1982).
- ⁵R.G. Littlejohn, *Phys. Fluids* **28**, 2015 (1985).
- ⁶P.J. Catto, W. M. Tang, and D.E. Baldwin, *Plasma Phys.* **23**, 639 (1981).
- ⁷A.I. Morozov and L.S. Soloviev, in *Reviews of Plasma Physics* (Consultants Bureau, New York, 1968), Vol. 2, translated from Russian; *Voprosy Teorii Plazmy* (Atomizdat, Moscow, 1963).
- ⁸R.G. Littlejohn, *Phys. Fluids* **24**, 1730 (1981).
- ⁹A.H. Boozer, *Phys. Fluids* **23**, 904 (1980).
- ¹⁰T.G. Northrop and J.A. Rome, *Phys. Fluids* **21**, 384 (1978).
- ¹¹R.D. Hazeltine, *Plasma Phys.* **15**, 77 (1973).