

2
RECEIVED BY TIC JAN 7 1980

NUREG/CR-1125
PNL-2940

Population Dose Commitments Due to Radioactive Releases From Nuclear Power Plant Sites in 1976

MASTER

Prepared by D. A. Baker

Pacific Northwest Laboratory

Prepared for
U. S. Nuclear Regulatory
Commission

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED
N

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, or any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for any third party's use, or the results of such use, of any information, apparatus product or process disclosed in this report, or represents that its use by such third party would not infringe privately owned rights.

Available from

 GPO Sales Program
 Division of Technical Information and Document Control
 U.S. Nuclear Regulatory Commission
 Washington, D.C. 20555

and

 National Technical Information Service
 Springfield, Virginia 22161

Population Dose Commitments Due to Radioactive Releases From Nuclear Power Plant Sites in 1976

Manuscript Completed: August 1979
Date Published: December 1979

Prepared by
D. A. Baker

Pacific Northwest Laboratory
Battelle Boulevard, P.O. Box 999
Richland, WA 99352

Prepared for
Division of Technical Support
Office of Management and Program Analysis
U.S. Nuclear Regulatory Commission
Washington, D.C. 20555
NRC FIN No. B2243-8

DISCLAIMER
This book was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.


DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

SUMMARY

Population radiation dose commitments have been estimated from reported radionuclide releases from commercial power reactors operating during 1976. Fifty-year dose commitments from a one-year exposure were calculated from both liquid and atmospheric releases for four population groups - infant, child, teenager and adult - residing between 2 and 80 km from each site.

Results are given in the form of tables giving the dose commitments for both liquid and airborne pathways for each age group and organ. Also included for each site is a histogram showing the fraction of the total population within the 2 to 80-km region around each site receiving various average dose commitments from the airborne pathways.

The total dose commitment from both liquid and airborne pathways ranged from a high of 160 person-rem to a low of 0.0002 person-rem with an arithmetic mean of 11 person-rem. The total population dose for all sites was estimated at 470 person-rem for the 90 million people considered at risk.



CONTENTS

SUMMARY	i
INTRODUCTION	1
SITE-DEPENDENT PARAMETERS	5
RESULTS	6
DISCUSSION OF RESULTS	6
SITE SUMMARIES	11
Arkansas	12
Beaver Valley	14
Big Rock Point	16
Browns Ferry 1, 2, & 3	18
Brunswick 1 & 2	20
Calvert Cliffs	22
Cook	24
Cooper Station	26
Dresden 1, 2, & 3	28
Duane Arnold	30
J. A. Fitzpatrick	32
Fort Calhoun	34
R. E. Ginna	36
Haddam Neck (Connecticut Yankee)	38
Edwin I. Hatch	40
Humboldt Bay	42
Indian Point	44
Kewaunee	46
LaCrosse	48
Maine Yankee	50
Millstone Point 1 & 2	52
Monticello	54
Nine Mile Point	56
Oconee 1, 2, & 3	58

Oyster Creek	60
Palisades	62
Peach Bottom 2 & 3	64
Pilgrim	66
Point Beach 1 & 2	68
Prairie Island 1 & 2	70
Quad Cities 1 & 2	72
Rancho Seco	74
H. B. Robinson	76
St. Lucie	78
Salem	80
San Onofre	82
Surry 1 & 2	84
Three Mile Island	86
Trojan	88
Turkey Point 3 & 4	90
Vermont Yankee	92
Yankee Rowe	94
Zion 1 & 2	96
REFERENCES	98
APPENDIX	A-1

INTRODUCTION

The Pacific Northwest Laboratory (PNL), operated by Battelle Memorial Institute for the Department of Energy (DOE), has conducted a study sponsored by the Nuclear Regulatory Commission (NRC) to estimate radiation dose commitments from reported radionuclide releases from 58 commercial power reactors operating during 1976. This work is a continuation of the study done for 1975 releases (Baker et al., 1977).

In this study, as in the last, we estimated the dose commitment* from both the liquid and gaseous releases to four age groups making up the population residing in the region of the site: infant, child, teenager, and adult.

The particular organs of reference in this study are listed in Table 1. The major pathways by which radionuclides travel from the reactor to the individual receptors are shown in Table 2. Other possible liquid pathways such as direct exposure from waterborne activities (swimming, boating, shoreline recreation) and internal exposure through ingestion of food products produced using contaminated irrigation water were not included. This was because we have found from past experience that the dose from these pathways is generally much smaller when compared to the pathways considered.

TABLE 1. Organs Considered

<u>Airborne</u>	<u>Waterborne</u>
• Total Body	• Total Body
• Thyroid	• Thyroid
• Bone	• Bone
• GI-Tract	• GI-Tract
• Liver	• Liver
• Lung	

*As used in this report the dose commitment describes the dose equivalent received over 50 years from one year of intake (1976).

TABLE 2. Pathways Considered

Airborne

- Air Submersion
- Contaminated Ground
- Inhalation
- Ingestion of Food Crops and Animal Products

Waterborne

- Ingestion of Drinking Water
- Ingestion of Fish and Invertebrates

The regional population for which we estimated doses included those persons estimated to be living in a region between 2 and 80 kilometers around the reactor sites during 1976. We estimated the numbers of persons by extrapolating Bureau of Census data for 1970 to 1976.

Atmospheric transport factors (annual average dilution and annual average deposition) were calculated for the region around each site using appropriate meteorological data supplied by the NRC, Office of Nuclear Reactor Regulation.

Models used to calculate the doses were those approved by the NRC. We incorporated these models into two small computer codes to expedite the dose calculations involved for each site.

Site specific parameters other than releases, meteorology, and population were obtained from Environmental Statements for the various reactors when available (Table 3). Such parameter values include the total population drinking contaminated water, river flow, dilution flow from the reactors (for non-river sites), fish and invertebrate harvest for region, and dilution factors for drinking water and aquatic foods.

The reactors included in this study, their type, licensed thermal power rating, and net electrical output for 1976 are listed in Table 4. Those reactors which had an operating augmented gaseous radioactive-waste system in 1976 are identified in the table along with populations at risk and the results of the study.

TABLE 3. Environmental Statements for Power Plants Included in This Study

<u>Site No.</u>		<u>Docket No.</u>	<u>Date</u>	<u>Remarks</u>
1	Big Rock Pt			ES not available
2	Brown's Ferry 1, 2, 3		Jul 71	Tennessee Valley Authority
3	Cooper Station	50-298	Feb 73	Draft Environmental Statement
4	Dresden 1			ES of Dresden 2, 3 used
4	Dresden 2, 3	50-237, 50-249	Nov 73	
* 5	Beaver Valley 1	50-334	Jul 73	
6	Humboldt Bay			ES not available
7	LaCrosse	50-409	Jun 76	Draft
8	Millstone Pt 1, 2	50-245, 50-336	Jun 73	
9	Monticello	50-263	Nov 72	
10	Nine Mile Pt 1	50-220	Jan 74	
11	Oyster Creek	50-219	Dec 74	
12	Peach Bottom 2, 3	50-277, 50-278	Apr 73	
13	Pilgrim 1	50-293	May 72	
14	Quad Cities 1, 2	50-254, 50-265	Sep 72	
15	Vermont Yankee	50-271	Jul 72	
* 16	St. Lucie	50-335	Jun 73	
17	Brunswick 1, 2	50-324, 50-325	Jun 73	Draft
18	Duane Arnold	50-331	Mar 73	
19	Fitzpatrick	50-333	Mar 73	
20	Hatch	50-321	Oct 72	
21	Arkansas 1	50-313	Feb 73	
22	Connecticut Yankee	50-213	Oct 73	(Haddam Neck)
23	Fort Calhoun	50-285	Aug 72	
24	H. B. Robinson	50-261	Apr 74	
25	Indian Pt 1, 2	50-247	Sep 72	ES of Indian Pt 2 used
* 26	Salem	50-272, 50-311	Apr 73	
27	Kewaunee	50-305	Dec 72	
28	Maine Yankee	50-309	Jul 72	
29	Oconee 1, 2, 3	50-269, 50-270 50-287	Mar 72	
30	Palisades	50-255	Jun 72	
31	Point Beach 1, 2	50-266, 50-301	May 72	
32	Prairie Island 1, 2	50-282, 50-306	May 73	
33	R. E. Ginna	50-244	Dec 73	
34	San Onofre 1	50-206	Oct 73	
35	Surry 1, 2	50-281	Jun 72	Unit 2
36	Three Mile Island 1	50-289	Dec 72	
37	Turkey Point 3, 4	50-250, 50-251	Feb 72	Draft
38	Yankee Rowe			ES not available
39	Zion 1, 2	50-295, 50-304	Dec 72	
40	Calvert Cliffs 1	50-317	Apr 73	
41	Cook 1	50-315	Aug 73	
* 42	Trojan	50-344	Jan 73	Draft
43	Rancho Seco	50-312	Mar 73	

* Added for 1976

TABLE 4. Reactor Characteristics and Population
Total-Body Dose Commitments 1976

Site	Unit	Type	Licensed	Electric	BWR	Population			Avg. Individual	
			Thermal Power (MW)	Generation 1976 (TW-hr)	Augmented Rad-Waste System (1976)	Dose Commitment (Person-Rem)	Liquid	Air	Total	Population At Risk
Arkansas	1	PWR	2568	3.89	-	4.5	0.038	4.5	1.6E5	2.8E-2
Beaver Valley	1	PWR	2652	0.442	-	0.0024	0.037	0.039	3.7E6	1.1E-5
Big Rock Point	1	BWR	240	0.244	No	7.3	0.41	7.7	1.3E5	6.0E-2
Browns Ferry	1	BWR	3293	1.30	Yes					
	2	BWR	3293	1.57	Yes					
	3	BWR	3293	1.42	Yes					
Browns Ferry	TOTAL		9879	4.29		0.50	0.57	1.1	6.6E5	1.6E-3
Brunswick	1	BWR	2436	2.49	No					
	2	BWR	2436	0	No					
Brunswick	TOTAL		5872	2.49		0.072	0.46	0.53	1.9E5	2.9E-3
Calvert Cliffs	1	PWR	2560	6.30	-	0.43	0.31	0.74	2.4E6	3.0E-4
Cook	1	PWR	3250	6.80	-	5.0	0.016	5.0	1.1E6	4.6E-3
Cooper	1	BWR	2381	3.64	Yes	--(a)	0.39	0.39	1.8E5	2.2E-3
Dresden	1	BWR	700	0.953	No					
	2	BWR	2527	4.37	No					
	3	BWR	2527	4.03	No					
Dresden	TOTAL		5754	9.35		--	120	120	6.4E6	1.9E-2
Duane Arnold	1	BWR	1658	2.49	Yes	0.0047	0.32	0.32	5.7E5	5.7E-4
J. A. Fitzpatrick	1	BWR	2436	4.16	Yes	0.27	1.1	1.4	8.3E5	1.6E-3
Fort Calhoun	1	PWR	1420	2.20	-	0.24	0.018	0.26	7.4E5	3.5E-4
R. E. Ginna	1	PWR	1520	2.06	-	0.43	0.084	0.51	1.2E6	4.4E-4
Haddam Neck	1	PWR	1825	4.03	-	0.057	3.6	3.7	3.4E6	1.1E-3
E. I. Hatch	1	BWR	2436	4.13	Yes	0.0095	0.024	0.034	2.7E5	1.2E-4
Humboldt Bay	3	BWR	220	0.197	No	0.0022	5.8	5.8	1.1E5	5.3E-2
Indian Point	1	PWR	2758	2.27	-	1.4	7.7	9.1	1.6E7	5.6E-4
Keweenaw	1	PWR	1650	3.38	-	5.1	0.019	5.1	5.9E5	8.6E-3
LaCrosse	1	BWR	165	0.173	No	9.4	2.7	12	3.3E5	3.7E-2
Maine Yankee	1	PWR	2440	5.93	-	0.037	0.024	0.061	5.6E5	1.1E-4
Millstone Point	1	BWR	2011	3.75	No					
	2	PWR	2560	4.54	-					
Millstone Point	TOTAL		4571	8.29		0.024	160	160	2.5E6	6.4E-2
Monticello	1	BWR	1670	3.99	Yes	0	0.25	0.25	2.1E6	1.2E-4
Nine Mile Point	1	BWR	1850	4.11	No	2.1	6.6	8.7	8.3E5	1.0E-2
Oconee	1	PWR	2568	3.99	-					
	2	PWR	2568	4.23	-					
	3	PWR	2568	4.76	-					
Oconee	TOTAL		7704	12.98		19	0.95	20	7.3E5	2.7E-2
Oyster Creek	1	BWR	1930	3.86	No	0.025	37	37	3.3E6	1.1E-2
Palisades	1	PWR	2200	2.85	-	0.64	0.0012	0.64	1.0E6	6.3E-4
Peach Bottom	2	BWR	3293	5.57	Yes					
	3	BWR	3293	6.05	Yes					
Peach Bottom	TOTAL		6586	11.6		3.9	13	17	4.1E6	4.1E-3
Pilgrim	1	BWR	1998	2.42	Yes	0.072	14	14	4.4E6	3.2E-3
Point Beach	1	PWR	1518	3.40	-					
	2	PWR	1518	3.76	-					
Point Beach	TOTAL		3036	7.16		0.19	0.14	0.33	5.9E5	5.6E-4
Prairie Island	1	PWR	1650	3.27	-					
	2	PWR	1650	2.66	-					
Prairie Island	TOTAL		3300	5.93		0.26	1.1	1.4	2.1E6	6.4E-4
Quad Cities	1	BWR	2511	3.39	Yes					
	2	BWR	2511	4.30	Yes					
Quad Cities	TOTAL		5022	7.69		9.3	1.4	11	6.7E5	1.6E-2
Rancho Seco	1	PWR	2772	2.21	-	0	0.0076	0.0076	1.5E6	5.2E-6
H. B. Robinson	1	PWR	2200	4.87	-	0.23	0.049	0.28	6.3E5	4.4E-4
St. Lucie	1	PWR	2560	0.099	-	0.0027	0.029	0.032	2.8E5	1.1E-4
Salem	1	PWR	3338	0	-	--	--	--	4.9E6	3.3E-8
San Onofre	1	PWR	1347	2.47	-	1.4	0.031	1.4	3.8E6	3.8E-4
Surry	1	PWR	2441	4.40	-					
	2	PWR	2441	3.34	-					
Surry	TOTAL		4882	7.74		2.3	1.1	3.4	1.7E6	2.0E-3
Three Mile Island	1	PWR	2535	4.34	-	0.082	1.3	1.4	1.8E6	7.6E-4
Trojan	1	PWR	3411	2.10	-	0.010	0.010	0.020	1.3E6	1.6E-5
Turkey Point	3	PWR	2200	4.32	-					
	4	PWR	2200	3.77	-					
Turkey Point	TOTAL		4400	8.09		0.017	0.26	0.28	2.0E6	1.4E-4
Vermont Yankee	1	BWR	1593	3.26	Yes	0.019	0.092	0.11	1.4E6	8.0E-5
Yankee Rowe	1	PWR	600	1.25	-	0.064	0.0072	0.071	1.6E6	4.5E-5
Zion	1	PWR	3250	4.76	-					
	2	PWR	3250	4.64	-					
Zion	TOTAL		6500	9.40		7.3	10	17	7.0E6	2.5E-3
TOTAL FOR ALL SITES				185		82	390	470	9.0E7	--
Arithmetic Mean				4.3		1.9	9.1	11	2.1E6	8.5E-3
Geometric Mean				2.9		0.19	0.24	1.0	1.1E6	9.0E-4

(a) -- indicates dose commitment <0.001 person-rem.

1 TW-hr = 3.6E15 joules

SITE-DEPENDENT PARAMETERS

For each reactor site included in the section on Site Summaries, the site name is given along with its location (including latitude and longitude). Next the estimated 1976 population within the 2 to 80-km region around the site is given. This population is derived from the 1970 census by extrapolation to 1976. In addition, the location of major metropolitan centers within 80 km are listed along with their 1976 extrapolated populations. The populations of the Standard Metropolitan Statistical Areas (SMSA) are given where applicable.

Next, the average production rates of vegetable crops and animal products are given for the area within a circle of 80-km radius based upon the state-wide average. This production has been reduced for sites on lakes and sea-coasts to account for the presence of the body of water. An animal grazing factor is estimated for each site location. This factor accounts for the fraction of the year grazing animals such as milk cows and beef cattle graze on fresh pasture in the region around the site.

The location of the meteorological station used in the acquisition of diffusion climatology data is indicated along with the percent data recovery and period of record.

Various site-dependent factors associated with the waterborne pathways are presented next. For lake and ocean sites the average dilution of plant effluents for the year 1976 used are those specified in Decker (1978). For river sites, the average annual river flow is tabulated. This flow was used in place of a dilution flow from the plant to account for dilutions of liquid releases at the locations of probable intake of drinking water and aquatic food catch. Any exceptions to this scheme have been footnoted. Next, the estimated 1976 population utilizing drinking water drawn from water supplies containing diluted effluents from the site are shown along with an estimated dilution factor where applicable. Fish and invertebrate catch data taken from the respective plant Environmental Statement when available (see Table 3) is listed next along with estimated dilution factors for the lake and ocean sites.

RESULTS

This report consists of a summary of values used for site-specific parameters at each site as explained above and the results of population dose commitment calculations. The population dose commitments are presented in two tables facing the page summarizing site-specific parameters for that site. These tables include both liquid and airborne pathway dose commitments for each of the several organs of reference for each age group investigated as well as the dose to the whole population which includes all age groups. The airborne population dose commitments for each of 160 segments partitioning the region around the site* were divided by the population residing within that segment to derive an average individual dose for that segment. These doses are summarized as a histogram showing percent of the population receiving a given dose level for each site. The fractional population dose from the liquid pathway was not determined in this manner, since the NRC does not at present take into account the location of individuals exposed via this pathway except for drinking water.

DISCUSSION OF RESULTS

Population dose commitments estimated for both the liquid pathway and airborne pathway varied widely over the 43 sites (58 reactors) studied. The total dose commitments (from both pathways) varied from a high of 160 to a low of 0.0002 person-rem. The arithmetic mean for the dose from liquid pathways was 2 person-rem and the mean for the dose from airborne pathways was 9 person-rem (see Table 4).

Monticello, Rancho Seco, and Dresden releases resulted in the smallest doses from liquid pathways (zero, zero, and near zero, respectively). This was because no liquid releases were reported for Monticello and Rancho Seco; and the receiving waters for the Dresden site are contaminated with Chicago sewage to such an extent as to severely limit use of this water for drinking

* See Appendix for definition of segments.

or fishing. The largest liquid pathway doses to the total body calculated were Oconee: 20 person-rem; and LaCrosse and Quad Cities: 9 person-rem. These doses resulted primarily from the radioactive cesium (Cs-134 and Cs-137) released by these plants.

The lowest total airborne pathway dose to the total body was estimated for Salem (1×10^{-8} person-rem)*; while the highest was at Millstone Point (160 person-rem) and Dresden (120 person-rem). The major contributors to these doses were the noble gases: Xe-133, Xe-135, and Kr-88. The total population dose commitments from all sites for 1976 were estimated to be 82 person-rem via liquid pathways and 390 person-rem via the airborne pathways (Table 4). Compared to 1975, the "liquid dose" is only slightly higher (82 vs. 76); however, the "air dose" is much smaller (390 vs. 1300).

We should point out here, however, that the doses estimated in this study are extremely low compared with an average annual background dose of 0.1 rem and well within all limits. Even the highest average individual dose commitment of 0.06 millirem (Table 4) from two sites (Millstone Point and Big Rock Point) is well within any national limits.

Figure 1 shows graphically the wide range of the airborne population dose commitments for the reactor sites (except Salem). The median, upper and lower quartiles, and upper and lower octiles for the distribution of doses calculated for each of 160 segments are indicated for each site.

Figure 2 is a histogram for all 43 sites taken together. We can see from this plot that about 23% of the total population at risk (90 million) would each receive a dose commitment of between 0.0003 and 0.001 millirem. We can see further that 6% receive a dose which is less than 1×10^{-6} millirem. Although not discernable from the plot, less than 0.002% received a dose of between 3 and 10 millirem.

It should be noted that we have been comparing these dose commitments with annual background, which is not quite exact. The dose commitments as stated above are those total-body doses received from the year's (1976) effluent release, over 50 years of the person's lifetime. However, most of the dose is delivered in the first year so that this comparison is not too greatly in error.

*However, Salem was only operating less than one month.




FIGURE 1. Airborne Population Dose Commitments for the Reactor Sites, 1976




FIGURE 2. Fraction of Total Population Receiving Various Individual Dose Commitments for All Sites, 1976

SITE SUMMARIES

1976

Site ARKANSAS POPE COUNTY, ARKANSAS
Location N 35.3075° W 93.2308°
Total Population Within 2-80-km Region 1.6E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Russellville</u>	<u>13,000</u>	<u>10 km</u> <u>E</u>
<u>Conway</u>	<u>17,000</u>	<u>77 km</u> <u>ESE</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 5.8E6 kg Milk 4.8E7 l Meat 7.2E7 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.7

Location of Meteorological Station Site Recovery 97%

Period of Record 1 JAN 75 - 31 DEC 75

Average Arkansas River Flow at Site 36,000 ft³/s

Drinking Water

Exposed Population None Dilution Factor -

Fish

Edible Harvest 1.4 (a) kg/yr Dilution Factor 1

(a) Average individual consumption rates as given in the FES (1973) were used in lieu of catch data.

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
ARKANSAS

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total</u>	<u>Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	1.9E-01	2.2E-02	7.3E-02	1.1E+00	1.1E+00	
Teen	3.7E-01	4.5E-02	5.3E-02	6.3E-01	9.6E-01	
Adult	4.0E+00	3.8E-01	3.5E-01	3.6E+00	5.7E+00	
TOTAL	4.5E+00	4.5E-01	4.7E-01	5.3E+00	7.8E+00	

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total</u>	<u>Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	5.3E-04	5.1E-04	6.0E-03	7.4E-04	7.9E-04	5.8E-04	
Child	6.0E-03	5.7E-03	3.8E-02	7.8E-03	7.8E-03	6.5E-03	
Teen	4.4E-03	4.2E-03	1.7E-02	4.8E-03	5.1E-03	5.0E-03	
Adult	2.7E-02	2.5E-02	7.1E-02	2.7E-02	2.8E-02	2.8E-02	
TOTAL	3.8E-02	3.6E-02	1.3E-01	4.0E-02	4.2E-02	4.0E-02	


Production / consumption factors:

Produce: <1

Milk: 2.3

Meat: 5.5

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site BEAVER VALLEY SHIPPINGPORT, PENNSYLVANIA
 Location N 40.6214° W 80.4347°
 Total Population Within 2-80-km Region 3.7E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Pittsburg SMSA	2.41E6	32 km SE
Youngstown-Warren SMSA	539,000	56 km NNW
Steubenville	31,000	31 km SW
Wheeling	50,000	69 km SSW

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 5.3E7 kg Milk 5.3E8 l Meat 5.4E7 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 92%

Period of Record 1 JAN 77 - 31 DEC 77

Average Ohio River Flow at Site 30,000 ft³/s

Drinking Water

Exposed Population 6,400 Dilution Factor 12(a)

Fish

Edible Harvest 410 kg/yr Dilution Factor 1

(a) This factor accounts for the incomplete dilution of plant effluent by river at point of drinking water intake at Midland.

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
BEAVER VALLEY

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	4.7E-05	3.2E-05	4.1E-05	6.8E-05	2.6E-05
Child	5.8E-04	4.8E-04	3.7E-04	1.0E-03	2.7E-04
Teen	2.0E-04	3.2E-04	1.3E-04	3.6E-04	1.1E-04
Adult	1.6E-03	3.0E-03	1.0E-03	2.8E-03	8.5E-04
TOTAL	2.4E-03	3.8E-03	1.6E-03	4.2E-03	1.3E-03

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	6.4E-04	6.4E-04	6.4E-04	1.5E-06	6.4E-04	6.4E-04
Child	7.6E-03	7.6E-03	7.6E-03	1.8E-05	7.6E-03	7.6E-03
Teen	4.7E-03	4.7E-03	4.7E-03	1.3E-05	4.7E-03	4.7E-03
Adult	2.4E-02	2.4E-02	2.4E-02	7.6E-05	2.4E-02	2.4E-02
TOTAL	3.7E-02	3.7E-02	3.7E-02	1.1E-04	3.7E-02	3.7E-02


Production / consumption factors:

Produce: <1

Milk: 1.1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site BIG ROCK POINT CHARLEVOIX COUNTY, MICHIGAN
 Location N 45.3589° W 85.1958°
 Total Population Within 2-80-km Region 1.3E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Traverse City</u>	<u>18,000</u>	<u>70 km</u> <u>SSW</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 6.9E7 kg Milk 2.9E8 l Meat 4.5E7 kg

Regional Productivity Factor 0.5

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 85%

Period of Record 9 FEB 61 - 8 FEB 63

Average Water Dilution Flow from Plant 83 ft³/s

Drinking Water

Exposed Population 24,000^(a) Dilution Factor 4.9E-5^(b)

Fish

Edible Harvest (c) kg/yr Dilution Factor 0.01

(a) Population exposed to contaminated drinking water derived from information obtained from J. Hennigan, Division of Radiation Health, Bureau of Environmental and Health, Michigan Department of Public Health.

(b) Drinking water dilution factor estimated by averaging dilution factor derived from Figure 6B-5, Vol. 1 of WASH-1258 (July 1973) suitably weighted for population

(c) Generic consumption rate used (Table A1).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
BIG ROCK POINT

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	1.2E-06	2.5E-07	2.9E-08	8.8E-06	1.1E-05
Child	3.0E-01	1.3E-02	3.0E-06	1.6E+00	1.8E+00
Teen	5.9E-01	2.6E-02	2.6E-06	9.8E-01	1.5E+00
Adult	6.4E+00	2.2E-01	2.0E-05	5.7E+00	9.0E+00
TOTAL	7.3E+00	2.6E-01	2.6E-05	8.3E+00	1.2E+01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	5.8E-03	5.8E-03	6.2E-03	5.9E-03	6.0E-03	6.0E-03
Child	6.5E-02	6.5E-02	6.9E-02	6.7E-02	6.7E-02	6.7E-02
Teen	4.8E-02	4.7E-02	4.9E-02	4.8E-02	4.8E-02	5.0E-02
Adult	2.9E-01	2.9E-01	3.0E-01	2.9E-01	2.9E-01	3.0E-01
TOTAL	4.1E-01	4.1E-01	4.2E-01	4.1E-01	4.1E-01	4.2E-01


Production / consumption factors:

Produce: 1.3

Milk: 8.5

Meat: 2.2

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site BROWNS FERRY DECATUR, ALABAMA
Location N 34.7036° W 87.1211°
Total Population Within 2-80-km Region 6.6E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Decatur	40,000	16 km SE
Huntsville	150,000	48 km E

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 1.7E7 kg Milk 5.7E7 l Meat 8.6E7 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.7

Location of Meteorological Station Site Recovery 94%

Period of Record 1 JAN 74 - 31 DEC 75

Average Tennessee River Flow at Site 45,000 ft³/s

Drinking Water

Exposed Population 23,000 Dilution Factor 1

Fish

Edible Harvest 1.6E6 kg/yr Dilution Factor 1

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
BROWNS FERRY 1, 2 & 3

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	3.1E-05	1.8E-05	8.2E-04	1.2E-04	7.4E-05
Child	2.9E-02	2.3E-02	1.1E-02	9.6E-02	1.2E-01
Teen	4.3E-02	4.8E-02	5.4E-03	5.8E-02	1.0E-01
Adult	4.3E-01	4.2E-01	3.6E-02	3.5E-01	6.1E-01
TOTAL	5.0E-01	4.9E-01	5.3E-02	5.0E-01	8.3E-01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	8.2E-03	8.1E-03	1.4E-02	8.4E-03	8.5E-03	1.1E-02
Child	9.1E-02	9.1E-02	1.3E-01	9.3E-02	9.3E-02	1.3E-01
Teen	6.7E-02	6.7E-02	8.1E-02	6.7E-02	6.7E-02	1.1E-01
Adult	4.0E-01	4.1E-01	4.6E-01	4.1E-01	4.1E-01	5.4E-01
TOTAL	5.7E-01	5.7E-01	6.8E-01	5.7E-01	5.7E-01	7.9E-01


Production / consumption factors:

Produce: <1

Milk: <1

Meat: 1.6

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site BRUNSWICK BRUNSWICK COUNTY, NORTH CAROLINA
Location N 33.9586° W 78.0208°
Total Population Within 2-80-km Region 1.9E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Wilmington</u>	<u>50,000</u>	<u>32 km</u> <u>N</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 2.6E7 kg Milk 1.0E8 l Meat 5.8E7 kg

Regional Productivity Factor 0.3

Animal Grazing Factor 0.7

Location of Meteorological Station Site Recovery 93%

Period of Record 1 JAN 75 - 31 DEC 75

Average Dilution Flow from Plant 250 ft³/s

Fish
Edible Harvest 2.1E5^(a) kg/yr Dilution Factor 0.001

Invertebrates
Edible Harvest 1.1E5^(a) kg/yr Dilution Factor 0.002

(a) Fish and invertebrate harvests together total harvest given in FES (1973).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
BRUNSWICK 1 & 2

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	1.2E-02	8.9E-03	6.6E-03	8.4E-03	2.0E-02
Teen	8.5E-03	1.8E-02	4.6E-03	5.6E-03	1.8E-02
Adult	5.1E-02	1.6E-01	3.0E-02	3.7E-02	1.1E-01
TOTAL	7.2E-02	1.9E-01	4.2E-02	5.1E-02	1.5E-01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	6.6E-03	6.5E-03	5.6E-02	6.7E-03	6.7E-03	6.7E-03
Child	7.3E-02	7.3E-02	3.7E-01	7.4E-02	7.4E-02	7.5E-02
Teen	5.3E-02	5.3E-02	1.7E-01	5.3E-02	5.4E-02	5.7E-02
Adult	3.2E-01	3.2E-01	7.4E-01	3.2E-01	3.2E-01	3.3E-01
TOTAL	4.6E-01	4.6E-01	1.3E+00	4.6E-01	4.6E-01	4.7E-01


Production / consumption factors:

Produce: <1

Milk: 1.3

Meat: 1.2

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site CALVERT CLIFFS LUSBY, MARYLAND
Location N 38.4272° W 76.4261°
Total Population Within 2-80-km Region 2.4E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Washington, DC SMSA (3/4)</u>	<u>2,300,000</u>	<u>75 km</u> <u>NW</u>
<u>Cambridge</u>	<u>12,000</u>	<u>32 km</u> <u>ENE</u>
<u>Annapolis</u>	<u>31,000</u>	<u>61 km</u> <u>N</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 4.5E7 kg Milk 5.0E8 l Meat 6.2E7 kg

Regional Productivity Factor 0.6

Animal Grazing Factor 0.6

Location of Meteorological Station Site Recovery 96%

Period of Record 1 JAN 75 - 31 DEC 75

Average Dilution Flow from Plant 2,400 ft³/s

Fish

Edible Harvest 1.0E7 kg/yr Dilution Factor 0.062^(a)

Invertebrates

Edible Harvest 7.4E6 kg/yr Dilution Factor 0.062^(a)

(a) Dilutions given in FES (1973).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
CALVERT CLIFFS

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	2.7E-02	1.9E-01	3.5E-01	9.4E-02	9.9E-02
Teen	3.7E-02	4.0E-01	2.5E-01	5.7E-02	8.3E-02
Adult	3.7E-01	3.4E+00	1.6E+00	3.3E-01	4.9E-01
TOTAL	4.3E-01	4.0E+00	2.2E+00	4.8E-01	6.8E-01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	4.7E-03	4.5E-03	1.6E-01	4.4E-03	5.0E-03	4.9E-03
Child	5.1E-02	5.0E-02	8.7E-01	4.7E-02	5.2E-02	5.5E-02
Teen	3.6E-02	3.6E-02	3.6E-01	3.3E-02	3.7E-02	4.3E-02
Adult	2.1E-01	2.1E-01	1.2E+00	2.0E-01	2.2E-01	2.4E-01
TOTAL	3.1E-01	3.0E-01	2.6E+00	2.8E-01	3.1E-01	3.4E-01


Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site COOK BENTON HARBOR, MICHIGAN
Location N 41.9742° W 86.5667°
Total Population Within 2-80-km Region 1.1E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Gary SMSA	650,000	80 km SW
South Bend SMSA	290,000	42 km SE
Benton Harbor	17,000	18 km NE

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 1.1E8 kg Milk 2.3E8 l Meat 1.9E8 kg

Regional Productivity Factor 0.6

Animal Grazing Factor _____ 0.5

Location of Meteorological Station Site Recovery 95%

Period of Record 1 MAY 75 - 31 APR 76

Average Water Dilution Flow from Plant _____ 78 ft³/s

Drinking Water

Exposed Population 270,000 Dilution Factor 0.025^(a)

Fish

Edible Harvest _____ 1.5E6 kg/yr Dilution Factor _____ 0.01

(a) Drinking water dilution factor estimated by averaging dilution factors derived from FES (1973) suitably weighted for population.

POPULATION DOSE-COMMITMENT ESTIMATES AND
 AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
 COOK

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	1.7E-02	1.5E-02	1.2E-01	1.3E-02	3.1E-02
Child	3.6E-01	1.9E-01	9.2E-01	9.1E-01	1.2E+00
Teen	4.0E-01	1.0E-01	3.0E-01	5.0E-01	9.0E-01
Adult	4.2E+00	8.6E-01	2.2E+00	2.9E+00	5.5E+00
TOTAL	5.0E+00	1.2E+00	3.6E+00	4.4E+00	7.7E+00

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	2.3E-04	2.3E-04	8.6E-04	2.3E-04	2.3E-04	2.5E-04
Child	2.6E-03	2.6E-03	6.8E-03	2.6E-03	2.6E-03	2.8E-03
Teen	1.9E-03	1.9E-03	3.5E-03	1.9E-03	1.9E-03	2.1E-02
Adult	1.1E-02	1.1E-02	1.7E-02	1.1E-02	1.1E-02	1.2E-02
TOTAL	1.6E-02	1.6E-02	2.8E-02	1.6E-02	1.6E-02	1.7E-02


Production / consumption factors:

Produce: <1

Milk: <1

Meat: 1.3

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
 TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site COOPER STATION NEMAHA COUNTY, NEBRASKA
Location N 40.3614° W 95.6378°
Total Population Within 2-80-km Region 1.8E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Shenandoah</u>	<u>6,000</u>	<u>58 km</u> <u>NNE</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 9.7E7 kg Milk 7.2E7 l Meat 2.0E8 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.6

Location of Meteorological Station Site Recovery 89%

Period of Record 1 MAR 70 - 31 DEC 75

Average Missouri River Flow at Site 31,000 ft³/s

Drinking Water

Exposed Population None Dilution Factor -

Fish

Edible Harvest 5.0 E3^(a) kg/yr Dilution Factor 0.5^(a)

(a) Assumes 1/2 fish harvest caught below plant.

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
COOPER STATION

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	2.6E-05	1.0E-05	2.6E-05	1.3E-04	1.3E-04
Teen	4.6E-05	2.2E-05	1.9E-05	7.7E-05	1.1E-04
Adult	4.9E-04	1.9E-04	1.2E-04	4.5E-04	6.8E-04
TOTAL	5.6E-04	2.2E-04	1.7E-04	6.5E-04	9.3E-04

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	5.6E-03	5.6E-03	9.1E-03	5.6E-03	5.7E-03	5.8E-03
Child	6.4E-02	6.4E-02	9.6E-02	6.2E-02	6.4E-02	6.6E-02
Teen	4.6E-02	4.6E-02	5.9E-02	4.6E-02	4.6E-02	5.0E-02
Adult	2.8E-01	2.8E-01	3.2E-01	2.8E-01	2.8E-01	2.9E-01
TOTAL	3.9E-01	4.0E-01	4.9E-01	3.9E-01	3.9E-01	4.1E-01


Production / consumption factors:

Produce: 2.8

Milk: 3.1

Meat: 14

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site DRESDEN GRUNDY COUNTY, ILLINOIS
 Location N 41.3897° W 88.2714°
 Total Population Within 2-80-km Region 6.4E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Chicago SMSA	7,000,000 ^(a)	72 km NE
Gary-Hammond-E. Chicago-SMSA	640,000 ^(a)	80 km ENE
Joliet	80,000	24 km NE
Aurora	75,000	43 km N
Elgin	56,000	74 km N

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 1.1E8 kg Milk 1.8E8 l Meat 1.9E8 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 77%

Period of Record 1 JAN 74 - 31 JAN 75

Average Illinois River Flow at Site 12,000 ft³/s

Drinking Water

Exposed Population (b) Dilution Factor -

Fish

Edible Harvest (b) kg/yr Dilution Factor -

(a) Population of total SMSA given, however population of SMSA fraction within 80 km of site would be somewhat smaller.

(b) River water used for sewage disposal for Chicago, so population doses from liquid pathways assumed to be near zero.

POPULATION DOSE COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
DRESDEN 1, 2, & 3

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
--	-------------------	---------------	----------------	-------------	--------------

TOTAL	~0	~0	~0	~0	~0
-------	----	----	----	----	----

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.7E+00	1.6E+00	3.0E+00	1.7E+00	1.7E+00	1.8E+00
Child	1.9E+01	1.8E+01	3.0E+01	2.1E+01	1.8E+01	2.0E+01
Teen	1.3E+01	1.4E+01	1.9E+01	1.4E+01	1.3E+01	1.5E+01
Adult	8.2E+01	8.2E+01	1.0E+02	8.5E+01	8.1E+01	8.7E+01
TOTAL	1.2E+02	1.2E+02	1.6E+02	1.2E+02	1.1E+02	1.2E+02


Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site DUANE ARNOLD CEDAR RAPIDS, IOWA
Location N 42.0981° W 91.7725°
Total Population Within 2-80-km Region 5.7E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Cedar Rapids	110,000	16 km SE
Waterloo	77,000	66 km NW
Iowa City	48,000	53 km SSE

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 9.8E7 kg Milk 2.6E8 l Meat 4.2E8 kg

Regional Productivity Factor 1

Animal Grazing Factor _____ 0.5

Location of Meteorological Station Site Recovery 57%

Period of Record 1 FEB 71 - 31 DEC 75

Average Cedar River Flow at Site 3,000 ft³/s

Drinking Water

Exposed Population 110,000 Dilution Factor 1

Fish

Edible Harvest _____ (a) _____ kg/yr Dilution Factor _____ 1

(a) No fish catch data given in FES (1973), thus population assumed to eat fish at generic consumption rates (Table A1).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
DUANE ARNOLD

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total</u>	<u>Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	2.1E-05	1.8E-05	1.0E-05	2.1E-05	1.5E-05	
Child	6.1E-04	1.4E-03	1.2E-04	8.8E-04	7.9E-04	
Teen	4.4E-04	2.6E-03	4.8E-05	5.3E-04	6.4E-04	
Adult	3.6E-03	2.3E-02	4.1E-04	3.7E-03	4.0E-03	
TOTAL	4.7E-03	2.7E-02	5.9E-04	5.1E-03	5.5E-03	

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total</u>	<u>Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	4.6E-03	4.6E-03	2.7E-02	4.6E-03	4.7E-03	4.7E-03	
Child	5.2E-02	5.2E-02	2.7E-01	5.2E-02	5.3E-02	5.3E-02	
Teen	3.8E-02	3.8E-02	1.2E-01	3.7E-02	3.8E-02	4.0E-02	
Adult	2.3E-01	2.3E-01	5.5E-01	2.3E-01	2.3E-01	2.3E-01	2.4E-01
TOTAL	3.2E-01	3.2E-01	9.7E-01	3.2E-01	3.2E-01	3.2E-01	3.3E-01


Production / consumption factors:

Produce: <1

Milk: 3.5

Meat: 9.2

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site J.A. FITZPATRICK OSWEGO, NEW YORK
Location N 43.5217° W 76.3980°
Total Population Within 2-80-km Region 8.3E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Syracuse SMSA	630,000	54 km SSE
Rome	50,000	80 km ESE
Auburn	34,000	66 km SSW

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 7.6E7 kg Milk 7.0E8 l Meat 3.3E7 kg

Regional Productivity Factor 0.7

Animal Grazing Factor 0.5

Location of Meteorological Station Nine Mile Point Site Recovery 97%

Period of Record 1 JAN 74 - 31 DEC 75

Average Water Dilution Flow from Plant 610 ft³/s

Drinking Water

Exposed Population 550,000^(a) Dilution Factor 0.003^(b)

Fish

Edible Harvest 7.3E5 kg/yr Dilution Factor 0.005^(b)

(a) Population exposed to drinking water derived from Nine Mile Point FES (1974).

(b) Dilution factors derived from FES (1973).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
J. A. FITZPATRICK

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	6.7E-04	5.8E-04	2.3E-02	1.5E-03	1.3E-03
Child	2.2E-02	2.5E-02	1.7E-01	5.3E-02	6.4E-02
Teen	2.3E-02	4.2E-02	5.3E-02	2.7E-02	5.0E-02
Adult	2.2E-01	3.7E-01	3.7E-01	1.7E-01	3.1E-01
TOTAL	2.7E-01	4.4E-01	6.1E-01	2.5E-01	4.2E-01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.7E-02	1.6E-02	3.9E-01	1.7E-02	1.7E-02	1.7E-02
Child	1.8E-01	1.8E-01	2.7E+00	1.9E-01	1.9E-01	1.9E-01
Teen	1.3E-01	1.3E-01	1.1E+00	1.3E-01	1.3E-01	1.4E-01
Adult	8.0E-01	7.9E-01	3.8E+00	8.0E-01	8.0E-01	8.2E-01
TOTAL	1.1E+00	1.1E+00	7.9E+00	1.1E+00	1.1E+00	1.2E+00


Production / consumption factors:

Produce: <1

Milk: 4.5

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site FORT CALHOUN WASHINGTON COUNTY, NEBRASKA
 Location N 41.5200° W 96.0780°
 Total Population Within 2-80-km Region 7.4E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Omaha SMSA</u>	<u>570,000</u>	<u>32 km SSE</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 9.7E7 kg Milk 7.2E7 l Meat 2.0E8 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 98%

Period of Record 1 JAN 74 - 31 DEC 74

Average Missouri River Flow at Site 27,000 ft³/s

Drinking Water

Exposed Population 570,000^(a) Dilution Factor 1

Fish

Edible Harvest 1.0E4 kg/yr Dilution Factor 1

(a) Drinking water population assumed to be Omaha SMSA (FES, 1972).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
FORT CALHOUN

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	2.9E-03	2.4E-03	7.1E-02	4.5E-03	8.2E-03
Child	3.6E-02	2.9E-02	5.1E-01	5.0E-02	8.0E-02
Teen	1.8E-02	1.3E-02	1.6E-01	1.4E-02	3.1E-02
Adult	1.8E-01	1.1E-01	1.1E+00	8.6E-02	2.1E-01
TOTAL	2.4E-01	1.6E-01	1.8E+00	1.5E-01	3.3E-01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	2.6E-04	2.5E-04	7.7E-03	2.6E-04	2.7E-04	2.7E-04
Child	2.9E-03	2.8E-03	7.9E-02	2.9E-03	3.0E-03	3.1E-03
Teen	2.1E-03	2.1E-03	3.2E-02	2.0E-03	2.1E-03	2.5E-03
Adult	1.2E-02	1.2E-02	1.2E-01	1.2E-02	1.3E-02	1.4E-02
TOTAL	1.8E-02	1.7E-02	2.4E-01	1.7E-02	1.8E-02	1.9E-02


Production / consumption factors:

Produce: <1

Milk: <1

Meat: 3.3

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site R. E. GINNA ONTARIO, NEW YORK
 Location N 43.2772° W 77.3081°
 Total Population Within 2-80-km Region 1.2E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Rochester SMSA	950,000	34 km WSW
Auburn	34,000	64 km SE
Oswego	24,000	56 km ENE

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 7.6E7 kg Milk 7.0E8 l Meat 3.3E7 kg

Regional Productivity Factor 0.6

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 89%

Period of Record 1 JAN 66 - 31 DEC 67 1 JAN 73 - 31 DEC 74

Average Water Dilution Flow from Plant 710 ft³/s

Drinking Water

Exposed Population 550,000 Dilution Factor 0.01^(a)

Fish

Edible Harvest 7.3E5 kg/yr Dilution Factor 0.01^(a)

(a) Dilution factors from FES (1973).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
R. E. GINNA

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	1.9E-03	1.7E-03	9.2E-02	1.9E-03	4.0E-03
Child	3.5E-02	2.5E-02	6.6E-01	8.6E-02	1.1E-01
Teen	3.4E-02	1.8E-02	2.1E-01	4.5E-02	7.7E-02
Adult	3.6E-01	1.6E-01	1.5E+00	2.6E-01	4.8E-01
TOTAL	4.3E-01	2.0E-01	2.4E+00	3.9E-01	6.7E-01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.2E-03	1.2E-03	1.8E-02	1.2E-03	1.3E-03	1.3E-03
Child	1.4E-02	1.4E-02	1.2E-01	1.3E-02	1.4E-02	1.5E-02
Teen	9.9E-03	9.8E-03	5.0E-02	9.2E-03	9.9E-03	1.2E-02
Adult	5.9E-02	5.9E-02	1.9E-01	5.5E-02	5.9E-02	6.5E-02
TOTAL	8.4E-02	8.4E-02	3.7E-01	7.8E-02	8.4E-02	9.2E-02


Production / consumption factors:

Produce: <1

Milk: 2.8

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site HADDAM NECK (CONN. YANKEE) HADDAM NECK, CONNECTICUT
 Location N 41.4822° W 72.4989°
 Total Population Within 2-80-km Region 3.4E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Hartford SMSA	830,000	37 km NW
New Haven SMSA	760,000	40 km WSW
Springfield SMSA	600,000	72 km N
Waterbury	110,000	47 km W

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 3.2E7 kg Milk 4.4E8 l Meat 2.0E7 kg

Regional Productivity Factor 0.7

Animal Grazing Factor 0.6

Location of Meteorological Station Site Recovery 95%

Period of Record 1 JAN 75 - 31 DEC 75

Average Discharge Canal Flow 780 ft³/s

Drinking Water

Exposed Population None Dilution Factor -

Fish

Edible Harvest 9.1E3^(a) kg/yr Dilution Factor 1

(a) Caught in discharge canal (FES, 1973).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
HADDAM NECK

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	2.8E-03	8.8E-04	7.3E-04	1.3E-02	1.4E-02
Teen	4.7E-03	1.0E-03	6.6E-04	7.7E-03	1.2E-02
Adult	4.9E-02	8.3E-03	5.2E-03	4.4E-02	7.0E-02
TOTAL	5.7E-02	1.0E-02	6.6E-03	6.5E-02	9.5E-02

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.8E-01	1.8E-01	1.8E-01	7.6E-01	1.8E-01	1.8E-01
Child	1.2E+00	1.2E+00	1.2E+00	5.0E+00	1.2E+00	1.2E+00
Teen	4.6E-01	4.6E-01	4.6E-01	1.5E+00	4.6E-01	4.6E-01
Adult	1.7E+00	1.7E+00	1.7E+00	4.3E+00	1.7E+00	1.7E+00
TOTAL	3.6E+00	3.6E+00	3.6E+00	1.2E+01	3.6E+00	3.6E+00


Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site E. I. HATCH BAXLEY, GEORGIA
Location N 31.9344° W 82.3444°
Total Population Within 2-80-km Region 2.7E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Waycross	21,000	80 km S

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 8.8E6 kg Milk 7.0E7 l Meat 8.1E7 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.8

Location of Meteorological Station Site Recovery 87%

Period of Record 1 JUN 70 - 31 AUG 74

Average Altamaha River Flow at Site 13,000 ft³/s

Drinking Water

Exposed Population None Dilution Factor -

Fish

Edible Harvest 6.3E5 (a) kg/yr Dilution Factor 1

(a) Commercial catch plus 3 pounds of game fish per year taken from river by average person (FES, 1972).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
E. I. HATCH

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	9.1E-04	5.1E-04	2.9E-04	1.7E-03	2.3E-03
Teen	9.3E-04	1.1E-03	2.1E-04	1.1E-03	2.1E-03
Adult	7.7E-03	9.7E-03	1.4E-03	7.1E-03	1.2E-02
TOTAL	9.5E-03	1.1E-02	1.9E-03	1.0E-02	1.7E-02

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	3.5E-04	3.4E-04	1.2E-03	3.5E-04	3.5E-04	3.6E-04
Child	3.9E-03	3.8E-03	8.5E-03	4.1E-03	3.9E-03	4.0E-03
Teen	2.8E-03	2.8E-03	4.6E-03	2.9E-03	2.8E-03	3.1E-03
Adult	1.7E-02	1.7E-02	2.3E-02	1.8E-02	1.7E-02	1.8E-02
TOTAL	2.4E-02	2.4E-02	3.7E-02	2.5E-02	2.4E-02	2.5E-02


Production / consumption factors:

Produce: <1

Milk: 2.0

Meat: 3.7

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site HUMBOLDT BAY EUREKA, CALIFORNIA
Location N 40.7417° W 124.2081°
Total Population Within 2-80-km Region 1.1E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Eureka	26,000	6.4 km N

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 4.8E7 kg Milk 2.3E8 l Meat 5.0E7 kg

Regional Productivity Factor 0.5

Animal Grazing Factor 1

Location of Meteorological Station Site Recovery 96%

Period of Record 1 JAN 66 - 31 DEC 67

Average Dilution Flow from Plant 150 ft³/s

Fish

Edible Harvest 4.E4^(a) kg/yr Dilution Factor 0.001

Invertebrates

Edible Harvest 5E3^(a) kg/yr Dilution Factor 0.002

(a) Edible seafood harvest taken from Noskin, et al., (1976).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
HUMBOLDT BAY

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	2.9E-04	1.2E-04	1.0E-05	3.1E-04	5.6E-04
Teen	2.3E-04	2.7E-04	7.2E-06	2.0E-04	5.0E-04
Adult	1.6E-03	2.4E-03	4.7E-05	1.3E-03	3.1E-03
TOTAL	2.2E-03	2.8E-03	6.5E-05	1.8E-03	4.2E-03

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	8.3E-02	8.3E-02	9.4E-02	8.3E-02	8.3E-02	8.5E-02
Child	9.2E-01	9.2E-01	1.1E+00	9.2E-01	9.2E-01	9.5E-01
Teen	6.7E-01	6.7E-01	7.5E-01	6.7E-01	6.7E-01	7.2E-01
Adult	4.1E+00	4.1E+00	4.5E+00	4.1E+00	4.1E+00	4.2E+00
TOTAL	5.8E+00	5.8E+00	6.4E+00	5.8E+00	5.8E+00	6.0E+00


Production / consumption factors:

Produce: 1.1

Milk: 8.1

Meat: 2.9

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site INDIAN POINT BUCHANAN, NEW YORK
 Location N 41.2656° W 73.9350°
 Total Population Within 2-80-km Region 1.6E7

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
New York SMSA	9,900,000	60 km S
Newark SMSA	2,100,000	64 km SSW
Stamford	110,000	42 km SE
Norwalk	78,000	48 km ESE
Bridgeport	160,000	64 km E
Poughkeepsie	32,000	48 km N
White Plains	50,000	30 km SSE

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 7.6E7 kg Milk 7.0E8 l Meat 3.3E7 kg

Regional Productivity Factor 0.8

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 96%

Period of Record 1 JAN 75 - 31 DEC 75

Average Hudson River Flow at Site 20,000 ft³/s

Drinking Water

Exposed Population None Dilution Factor -

Fish

Edible Harvest (a) kg/yr Dilution Factor 0.001^(b)

(a) No fish catch data given in FES so generic consumption rates used (Table A1).

(b) One percent of population obtain 10% of their fish from river (FES, 1972).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
INDIAN POINT

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	5.9E-02	3.9E-03	3.1E-02	2.5E-01	3.1E-01
Teen	1.1E-01	7.9E-03	2.2E-02	1.5E-01	2.7E-01
Adult	1.2E+00	6.7E-02	1.5E-01	8.8E-01	1.6E+00
TOTAL	1.4E+00	7.9E-02	2.0E-01	1.3E+00	2.2E+00

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.1E-01	1.1E-01	6.9E-01	1.1E-01	1.1E-01	1.2E-01
Child	1.2E+00	1.2E+00	5.3E+00	1.2E+00	1.3E+00	1.4E+00
Teen	9.0E-01	9.0E-01	3.0E+00	8.9E-01	9.1E-01	1.1E+00
Adult	5.5E+00	5.4E+00	1.4E+01	5.4E+00	5.5E+00	6.0E+00
TOTAL	7.7E+00	7.7E+00	2.3E+01	7.6E+00	7.8E+00	8.5E+00


Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site KEWAUNEE CARLTON, WISCONSIN
 Location N 44.3433° W 87.5350°
 Total Population Within 2-80-km Region 5.9E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Sheboygan	51,000	64 km SSW
Manitowoc	35,000	30 km SSW
Greenbay	92,000	40 km W
Appleton	60,000	72 km WSW

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 7.2E7 kg Milk 1.2E9 l Meat 1.0E8 kg

Regional Productivity Factor 0.5

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 76%

Period of Record 2 JAN 69 - 31 DEC 69

Average Water Dilution Flow from Plant 140 ft³/s

Drinking Water

Exposed Population 190,000 Dilution Factor 8.2E-3^(a)

Fish

Edible Harvest 1.1^(b) kg/yr Dilution Factor 0.01^(c)

(a) Drinking water dilution factor estimated by averaging dilution factors derived from FES (1972) suitably weighted for populations.

(b) Average individual consumption rate as given in the FES (1972) used in lieu of catch data.

(c) Dilution factor reduced 1/10 from that used in FES in consideration of lake mixing.

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
KEWAUNEE

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	3.4E-03	2.5E-03	1.5E-02	9.0E-03	1.3E-02
Child	2.5E-01	6.8E-02	1.2E-01	1.1E+00	1.3E+00
Teen	4.1E-01	9.1E-02	3.9E-02	6.4E-01	1.0E+00
Adult	4.5E+00	7.9E-01	2.8E-01	3.7E+00	6.2E+00
TOTAL	5.1E+00	9.6E-01	4.5E-01	5.5E+00	8.5E+00

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	2.7E-04	2.7E-04	6.5E-04	2.9E-04	2.7E-04	2.9E-04
Child	3.1E-03	3.0E-03	5.8E-03	3.8E-03	3.0E-03	3.4E-03
Teen	2.2E-03	2.2E-03	3.5E-03	2.6E-03	2.2E-03	2.6E-03
Adult	1.3E-02	1.3E-02	1.8E-02	1.6E-02	1.3E-02	1.5E-02
TOTAL	1.9E-02	1.9E-02	2.8E-02	2.2E-02	1.9E-02	2.1E-02


Production / consumption factors:

Produce: <1

Milk: 7.5

Meat: 1.1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site LA CROSSE GENOA, WISCONSIN
Location N 43.5597° W 91.2281°
Total Population Within 2-80-km Region 3.3E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>La Crosse</u>	<u>53,000</u>	<u>32 km</u>
		<u>N</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 7.2E7 kg Milk 1.2E9 l Meat 1.0E8 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 97%

Period of Record 1 JAN 75 - 31 DEC 75

Average Mississippi River Flow at Site 28,000 ft³/s

Drinking Water

Exposed Population None Dilution Factor -

Fish

Edible Harvest (a) kg/yr Dilution Factor 0.5^(a)

(a) No fish catch data given in FES so 1/2 population assumed to consume river fish at generic consumption rates (Table A1).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
LA CROSSE

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	4.0E-01	2.0E-02	3.1E-02	1.9E+00	2.2E+00
Teen	7.6E-01	4.0E-02	2.2E-02	1.1E+00	1.9E+00
Adult	8.2E+00	3.4E-01	1.5E-01	6.7E+00	1.1E+01
TOTAL	9.4E+00	4.0E-01	2.0E-01	9.7E+00	1.5E+01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	3.9E-02	3.9E-02	4.7E-02	3.9E-02	3.9E-02	4.0E-02
Child	4.4E-01	4.4E-01	5.2E-01	4.6E-01	4.4E-01	4.5E-01
Teen	3.2E-01	3.2E-01	3.5E-01	3.3E-01	3.2E-01	3.3E-01
Adult	1.9E+00	1.9E+00	2.1E+00	2.0E+00	1.9E+00	2.0E+00
TOTAL	2.7E+00	2.7E+00	3.0E+00	2.8E+00	2.7E+00	2.8E+00


Production / consumption factors:

Produce: 1.1

Milk: 27

Meat: 3.9

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site MAINE YANKEE LINCOLN COUNTY, MAINE
Location N 43.9503° W 69.6964°
Total Population Within 2-80-km Region 5.6E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Portland	70,000	51 km SW
Lewiston	45,000	40 km NW

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 2.4E8 kg Milk 6.6E7 l Meat 4.3E6 kg

Regional Productivity Factor 0.6

Animal Grazing Factor 0.5

Location of Meteorological Station Site _____ Recovery _____ 98%

Period of Record 1 APR 75 - 31 MAR 76

Average Dilution Flow from Plant _____ 791 ft³/s

Fish

Edible Harvest (a) kg/yr Dilution Factor 0.001

Invertebrates

Edible Harvest (a) kg/yr Dilution Factor 0.002

(a) No seafood harvest data given in FES (1972), thus generic population consumption rates used (Table A1).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
MAINE YANKEE

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	1.8E-03	7.1E-04	3.7E-03	7.8E-03	8.7E-03
Teen	3.1E-03	1.5E-03	2.6E-03	4.7E-03	7.3E-03
Adult	3.2E-02	1.3E-02	1.7E-02	2.7E-02	4.3E-02
TOTAL	3.7E-02	1.5E-02	2.3E-02	4.0E-02	5.9E-02

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	3.3E-04	3.3E-04	6.2E-04	3.3E-04	3.5E-04	3.7E-04
Child	4.0E-03	3.9E-03	8.2E-03	3.9E-03	4.3E-03	4.5E-03
Teen	2.9E-03	2.8E-03	4.6E-03	2.7E-03	3.0E-03	3.5E-03
Adult	1.7E-02	1.7E-02	2.4E-02	1.6E-02	1.7E-02	1.9E-02
TOTAL	2.4E-02	2.4E-02	3.7E-02	2.3E-02	2.5E-02	2.7E-02


Production / consumption factors:

Produce: 1.4

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site MILLSTONE POINT WATERFORD, CONNECTICUT
 Location N 41.3086° W 72.1675°
 Total Population Within 2-80-km Region 2.5E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Hartford SMSA	840,000	72 km NW
New Haven SMSA	760,000	64 km W
Warwick	86,000	80 km NE
Waterbury	110,000	70 km WNW

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 3.2E7 kg Milk 4.4E8 l Meat 2.0E7 kg

Regional Productivity Factor 0.6

Animal Grazing Factor 0.6

Location of Meteorological Station Site Recovery 95%

Period of Record 1 JAN 74 - 31 DEC 74

Average Dilution Flow from Plant 300 ft³/s

Fish

Edible Harvest 9.1E4 kg/yr Dilution Factor 0.001

Invertebrates

Edible Harvest 9.1E4 kg/yr Dilution Factor 0.002

POPULATION DOSE-COMMITMENT ESTIMATES AND
AND AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
MILLSTONE POINT 1 & 2

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	2.2E-03	3.1E-03	3.1E-03	5.9E-03	6.2E-03
Teen	2.3E-03	6.2E-03	2.1E-03	3.5E-03	5.3E-03
Adult	2.0E-02	5.3E-02	1.4E-02	2.1E-02	3.2E-02
TOTAL	2.4E-02	6.2E-02	1.9E-02	3.0E-02	4.3E-02

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	2.3E+00	2.3E+00	5.5E+00	2.3E+00	2.3E+00	2.4E+00
Child	2.6E+01	2.6E+01	4.3E+01	2.6E+01	2.6E+01	2.6E+01
Teen	1.9E+01	1.9E+01	2.6E+01	1.9E+01	1.9E+01	2.0E+01
Adult	1.1E+02	1.1E+02	1.4E+02	1.1E+02	1.1E+02	1.2E+02
TOTAL	1.6E+02	1.6E+02	2.1E+02	1.6E+02	1.6E+02	1.7E+02


Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site MONTICELLO MONTICELLO, MINNESOTA
 Location N 45.3339° W 93.8486°
 Total Population Within 2-80-km Region 2.1E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Minneapolis-St. Paul SMSA</u>	<u>2,000,000</u>	<u>64 km</u> <u>SE</u>
<u>St. Cloud</u>	<u>41,000</u>	<u>37 km</u> <u>NW</u>
<u>Anoka</u>	<u>14,000</u>	<u>35 km</u> <u>ESE</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 1.2E8 kg Milk 4.0E8 l Meat 1.1E8 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 92%

Period of Record 1 JAN 74 - 31 DEC 74

Average Mississippi River Flow at Site 4,600 ft³/s

Drinking Water

Exposed Population (a) Dilution Factor -

Fish

Edible Harvest (a) kg/yr Dilution Factor -

(a) No radionuclides released in liquid effluents reported (Decker, 1978).

POPULATION DOSE-COMMITMENT ESTIMATES AND
 AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
 MONTICELLO

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
--	-------------------	---------------	----------------	-------------	--------------

TOTAL	0	0	0	0	0
-------	---	---	---	---	---


Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	3.6E-03	3.3E-03	1.6E-01	4.1E-03	4.5E-03	3.6E-03
Child	4.5E-02	4.0E-02	1.1E+00	6.1E-02	4.8E-02	4.2E-02
Teen	3.0E-02	2.9E-02	4.4E-01	3.4E-02	3.1E-02	3.1E-02
Adult	1.7E-01	1.7E-01	1.6E+00	1.8E-01	1.7E-01	1.7E-01
TOTAL	2.5E-01	2.4E-01	3.3E+00	2.8E-01	2.6E-01	2.5E-01

Production / consumption factors:

Produce: <1 Milk: 1.5 Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
 TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site NINE MILE POINT OSWEGO, NEW YORK
 Location N 43.5217° W 76.3980°
 Total Population Within 2-80-km Region 8.3E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Syracuse SMSA</u>	<u>630,000</u>	<u>54 km SSE</u>
<u>Auburn</u>	<u>34,000</u>	<u>66 km SSW</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 7.6E7 kg Milk 7.0E8 l Meat 3.3E7 kg

Regional Productivity Factor 0.7

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 97%

Period of Record 1 JAN 74 - 31 DEC 75

Average Water Dilution Flow from Plant 270 ft³/s

Drinking Water

Exposed Population 550,000 Dilution Factor 0.01^(a)

Fish

Edible Harvest 7.3E5 kg/yr Dilution Factor 0.33^(b)

(a) Drinking water dilution factor estimated by averaging dilution factor derived from FES (1974) suitably weighted for population.

(b) Fish catch dilution factor (FES, 1974).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
NINE MILE POINT

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total</u>	<u>Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	5.7E-03	1.9E-03	1.2E-02	3.9E-02	3.0E-02	
Child	1.5E-01	3.7E-02	8.3E-02	7.3E-01	6.0E-01	
Teen	1.6E-01	3.4E-02	2.6E-02	3.0E-01	3.8E-01	
Adult	1.8E+00	3.1E-01	1.8E-01	1.8E+00	2.3E+00	
TOTAL	2.1E+00	3.8E-01	3.0E-01	2.9E+00	3.3E+00	

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total</u>	<u>Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	9.6E-02	9.4E-02	1.3E+00	9.8E-02	9.9E-02	9.6E-02	
Child	1.1E+00	1.1E+00	9.1E+00	1.1E+00	1.1E+00	1.1E+00	1.1E+00
Teen	7.7E-01	7.7E-01	3.8E+00	7.8E-01	7.8E-01	8.1E-01	
Adult	4.7E+00	4.7E+00	1.4E+01	4.7E+00	4.7E+00	4.7E+00	4.8E+00
TOTAL	6.6E+00	6.6E+00	2.9E+01	6.6E+00	6.6E+00	6.6E+00	6.8E+00


Production / consumption factors:

Produce: <1

Milk: 4.5

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site OCONEE OCONEE COUNTY, SOUTH CAROLINA
 Location N 34.7500° W 83.0458°
 Total Population Within 2-80-km Region 7.3E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Anderson</u>	<u>30,000</u>	<u>39 km</u> <u>SE</u>
<u>Greenville</u>	<u>67,000</u>	<u>47 km</u> <u>E</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 7.4E6 kg Milk 5.7E7 l Meat 5.0E7 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.7

Location of Meteorological Station Site Recovery 86%

Period of Record 1 JAN 75 - 31 DEC 75

Average Keowee River Flow at Site 1,100 ft³/s

Drinking Water

Exposed Population 44,000 Dilution Factor 1

Fish

Edible Harvest (a) kg/yr Dilution Factor 0.01^(b)

(a) No fish catch data given in FES so generic consumption rates used (Table A1).

(b) Ten percent of population obtain 10% of their fish diet from Hartwell Reservoir (FES, 1972).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
OCONEE 1, 2 & 3

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	8.8E-02	7.5E-02	2.7E+00	9.6E-02	2.0E-01
Child	1.6E+00	9.0E-01	1.9E+01	3.5E+00	4.9E+00
Teen	1.5E+00	4.3E-01	6.0E+00	1.8E+00	3.3E+00
Adult	1.6E+01	3.7E+00	4.3E+01	1.0E+01	2.0E+01
TOTAL	1.9E+01	5.1E+00	7.0E+01	1.6E+01	2.9E+01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.3E-02	1.3E-02	9.1E-02	1.2E-02	1.4E-02	1.4E-02
Child	1.5E-01	1.5E-01	6.1E-01	1.4E-01	1.6E-01	1.7E-01
Teen	1.1E-01	1.1E-01	3.1E-01	9.8E-02	1.1E-01	1.3E-01
Adult	6.7E-01	6.7E-01	1.4E+00	5.9E-01	6.8E-01	7.3E-01
TOTAL	9.5E-01	9.5E-01	2.4E+00	8.4E-01	9.6E-01	1.0E+00


Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site OYSTER CREEK OYSTER CREEK, NEW JERSEY
Location N 39.8139° W 74.2114°
Total Population Within 2-80-km Region 3.3E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Atlantic City	49,000	56 km SSW
Trenton	110,000	64 km NW

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 7.4E7 kg Milk 2.7E8 l Meat 2.4E7 kg

Regional Productivity Factor 0.5

Animal Grazing Factor 0.6

Location of Meteorological Station Site _____ Recovery _____ 63%

Period of Record 15 FEB 66 - 31 DEC 68

Average Dilution Flow from Plant _____ 1800 ft³/s

Fish

Edible Harvest 2.1 (a) kg/yr Dilution Factor 0.01^(b)

Invertebrates

Edible Harvest 0.96^(a) kg/yr Dilution Factor 0.01^(b)

- (a) Average individual consumption rate as given in the FES (1974) used in lieu of catch data.
- (b) 10% of seafood eaten assumed caught in bay waters diluted to 10% of that of discharge canal (FES, 1974).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
OYSTER CREEK

Dose Commitments (person-rem) from Liquid Pathways

	Total Body	GI-LLI	Thyroid	Bone	Liver
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	2.9E-03	3.8E-03	2.6E-04	3.8E-03	5.1E-03
Teen	2.5E-03	7.7E-03	1.8E-04	2.4E-03	4.4E-03
Adult	1.9E-02	6.7E-02	1.2E-03	1.5E-02	2.7E-02
TOTAL	2.5E-02	7.8E-02	1.7E-03	2.1E-02	3.6E-02

Dose Commitments (person-rem) from Airborne Pathways

	Total Body	GI-LLI	Thyroid	Bone	Liver	Lung
Infant	5.4E-01	5.4E-01	3.7E+00	5.5E-01	5.5E-01	5.5E-01
Child	6.0E+00	6.0E+00	2.8E+01	6.1E+00	6.0E+00	6.1E+00
Teen	4.4E+00	4.4E+00	1.5E+01	4.4E+00	4.4E+00	4.6E+00
Adult	2.6E+01	2.6E+01	6.7E+01	2.7E+01	2.7E+01	2.7E+01
TOTAL	3.7E+01	3.7E+01	1.1E+02	3.8E+01	3.8E+01	3.8E+01


Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site PALISADES COVERT TOWNSHIP, MICHIGAN
 Location N 42.3233° W 86.3142°
 Total Population Within 2-80-km Region 1.0E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Kalamazoo	88,000	56 km E
South Bend SMSA	290,000	72 km S
Michigan City	40,000	66 km SW

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 6.8E7 kg Milk 2.9E8 l Meat 4.5E7 kg

Regional Productivity Factor 0.6

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 67%

Period of Record 1 SEP 73 - 31 AUG 74

Average Water Dilution Flow from Plant 140 ft³/s

Drinking Water

Exposed Population 48,000 Dilution Factor 3.5E-3^(a)

Fish

Edible Harvest (b) kg/yr Dilution Factor 0.001^(b)

(a) Drinking water dilution factor estimated by averaging dilution factor derived from FES (1972) suitably weighted for population.

(b) Since the average individual consumption rate of 20 g/d as assumed in FES (1972) seemed unreasonably large, generic rates were used (Table A1).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
PALISADES

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Lung</u>
Infant	3.5E-05	2.1E-05	1.6E-04	1.6E-04	1.9E-04
Child	2.7E-02	6.6E-03	1.7E-03	1.4E-01	1.6E-01
Teen	5.2E-02	1.3E-02	7.4E-04	8.4E-02	1.3E-01
Adult	5.6E-01	1.1E-01	5.1E-03	4.9E-01	7.8E-01
TOTAL	6.4E-01	1.3E-01	7.7E-03	7.1E-01	1.1E+00

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	2.7E-05	1.4E-05	1.1E-02	4.1E-05	4.6E-05	1.4E-05
Child	2.6E-04	1.7E-04	6.5E-02	3.5E-04	3.5E-04	1.6E-04
Teen	1.5E-04	1.3E-04	2.5E-02	1.7E-04	2.0E-04	1.2E-04
Adult	7.9E-04	7.3E-04	8.3E-02	8.3E-04	9.1E-04	7.1E-04
TOTAL	1.2E-03	1.0E-03	1.8E-01	1.4E-03	1.5E-03	1.0E-03


Production / consumption factors:

Produce: <1

Milk: 1.3

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site PEACH BOTTOM YORK COUNTY, PENNSYLVANIA
 Location N 39.7592° W 76.2689°
 Total Population Within 2-80-km Region 4.1E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Lancaster SMSA</u>	<u>320,000</u>	<u>31 km</u> <u>N</u>
<u>Baltimore SMSA</u>	<u>2,100,000</u>	<u>58 km</u> <u>SW</u>
<u>Wilmington</u>	<u>500,000</u>	<u>64 km</u> <u>E</u>
<u>Reading</u>	<u>300,000</u>	<u>72 km</u> <u>NNE</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 5.3E7 kg Milk 5.3E8 l Meat 5.4E7 kg

Regional Productivity Factor 0.95

Animal Grazing Factor 0.6

Location of Meteorological Station Site Recovery 72%

Period of Record 1 AUG 67 - 31 JUL 71

Average Susquehanna River Flow at Site 36,000 ft³/s

Drinking Water

Exposed Population 2.1E6 Dilution Factor 1

Fish

Edible Harvest (a) kg/yr Dilution Factor 0.001^(b)

(a) No fish catch data given in FES (1974), thus generic consumption rates used (Table A1).

(b) One percent of people obtain 10% of their fish diet from river downstream from plant (FES, 1973).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
PEACH BOTTOM 2 & 3

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	2.2E-02	8.4E-03	4.4E-01	1.2E-01	1.9E-01
Child	3.8E-01	8.5E-02	3.1E+00	1.4E+00	1.7E+00
Teen	2.9E-01	4.9E-02	9.3E-01	3.7E-01	6.6E-01
Adult	3.3E+00	4.1E-01	6.5E+00	2.3E+00	4.3E+00
TOTAL	3.9E+00	5.6E-01	1.1E+01	4.1E+00	6.9E+00

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.9E-01	1.9E-01	1.9E+00	2.0E-01	2.0E-01	2.1E-01
Child	2.1E+00	2.1E+00	1.2E+01	2.2E+00	2.2E+00	2.3E+00
Teen	1.6E+00	1.5E+00	5.8E+00	1.6E+00	1.6E+00	1.8E+00
Adult	9.4E+00	9.3E+00	2.4E+01	9.4E+00	9.4E+00	1.0E+01
TOTAL	1.3E+01	1.3E+01	4.4E+01	1.3E+01	1.3E+01	1.5E+01


Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site PILGRIM PLYMOUTH, MASSACHUSETTS
 Location N 41.9444° W 70.5778°
 Total Population Within 2-80-km Region 4.4E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Boston SMSA	3,500,000	64 km NW
Providence SMSA	780,000	70 km WSW
Brockton	91,000	40 km WNW
New Bedford	100,000	45 km SW

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 2.0E7 kg Milk 2.6E8 l Meat 1.6E7 kg

Regional Productivity Factor 0.3

Animal Grazing Factor 0.6

Location of Meteorological Station Site Recovery 93%

Period of Record 1 MAY 74 - 30 APR 75

Average Dilution Flow from Plant 21 ft³/s

Fish

Edible Harvest 2.6E4 kg/yr Dilution Factor 0.001

Invertebrates

Edible Harvest 3.1E4 kg/yr Dilution Factor 0.002

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
PILGRIM

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	1.2E-02	1.1E-02	4.4E-04	1.3E-02	1.9E-02
Teen	8.3E-03	2.3E-02	3.1E-04	8.0E-03	1.7E-02
Adult	5.2E-02	2.0E-01	2.0E-03	5.1E-02	1.0E-01
TOTAL	7.2E-02	2.3E-01	2.8E-03	7.2E-02	1.4E-01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	2.1E-01	2.1E-01	3.5E-01	2.1E-01	2.1E-01	2.1E-01
Child	2.3E+00	2.3E+00	3.2E+00	2.3E+00	2.3E+00	2.4E+00
Teen	1.7E+00	1.7E+00	2.1E+00	1.7E+00	1.7E+00	1.8E+00
Adult	1.0E+01	1.0E+01	1.2E+01	1.0E+01	1.0E+01	1.1E+01
TOTAL	1.4E+01	1.4E+01	1.7E+01	1.4E+01	1.4E+01	1.5E+01


Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site POINT BEACH MANITOWOC COUNTY, WISCONSIN
 Location N 44.2764° W 87.5186°
 Total Population Within 2-80-km Region 5.9E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Sheboygan	51,000	56 km SSW
Manitowoc	35,000	19 km SSW
Greenbay	92,000	48 km NW
Appleton	60,000	72 km W

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 7.2E7 kg Milk 1.2E9 l Meat 1.0E8 kg

Regional Productivity Factor 0.5

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 83%

Period of Record 19 APR 67 - 18 APR 69

Average Water Dilution Flow from Plant 590 ft³/s

Drinking Water

Exposed Population 190,000 Dilution Factor 2.6E-3^(a)

Fish

Edible Harvest 6.7E4^(b) kg/yr Dilution Factor 0.013^(c)

(a) Drinking water dilution factor estimated by averaging dilution factor derived from FES (1972), suitably weighted for population.

(b) Includes both sport and commercial fish catch (FES, 1972).

(c) Dilution factor estimated by averaging sport and commercial dilution factor derived from FES (1972).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
POINT BEACH 1 & 2

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	5.9E-04	6.1E-04	6.1E-03	5.8E-04	1.3E-03
Child	1.3E-02	8.4E-03	4.6E-02	3.5E-02	4.8E-02
Teen	1.5E-02	8.5E-03	1.4E-02	1.9E-02	3.4E-02
Adult	1.6E-01	4.7E-02	1.0E-01	1.1E-01	2.1E-01
TOTAL	1.9E-01	6.1E-02	1.7E-01	1.7E-01	3.0E-01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.8E-03	1.8E-03	2.4E-03	1.1E-03	2.2E-03	1.9E-03
Child	2.6E-02	2.5E-02	2.9E-02	1.2E-02	2.8E-02	2.6E-02
Teen	1.7E-02	1.6E-02	1.8E-02	7.2E-03	1.7E-02	1.7E-02
Adult	9.4E-02	9.1E-02	9.7E-02	4.1E-02	9.5E-02	9.3E-02
TOTAL	1.4E-01	1.3E-01	1.5E-01	6.0E-02	1.4E-01	1.4E-01


Production / consumption factors:

Produce: <1

Milk: 7.6

Meat: 1.1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site PRAIRIE ISLAND RED WING, MINNESOTA
Location N 44.6217° W 92.6330°
Total Population Within 2-80-km Region 2.1E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Minneapolis-St Paul SMSA	2.0E6	50 km NW
Rochester	60,000	64 km SSE
Hastings	13,000	19 km NW

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 1.2E8 kg Milk 4.0E8 l Meat 1.0E8 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.5

Location of Meteorological Station Site _____ Recovery _____ 65%

Period of Record 22 MAR 74 - 21 MAR 75

Average Mississippi River Flow at Site 15,000 ft³/s

Drinking Water

Exposed Population None Dilution Factor 1

Fish

Edible Harvest 6.8E5 kg/yr Dilution Factor 1

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
PRAIRIE ISLAND 1 & 2

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	1.5E-02	1.6E-02	2.2E-03	4.4E-02	5.7E-02
Teen	2.2E-02	3.1E-02	1.8E-03	2.7E-02	4.9E-02
Adult	2.2E-01	2.7E-01	1.3E-02	1.6E-01	3.0E-01
TOTAL	2.6E-01	3.2E-01	1.7E-02	2.3E-01	4.1E-01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.6E-02	1.6E-02	3.2E-02	1.6E-02	1.6E-02	1.7E-02
Child	1.8E-01	1.8E-01	2.8E-01	1.8E-01	1.8E-01	1.9E-01
Teen	1.3E-01	1.3E-01	1.7E-01	1.3E-01	1.3E-01	1.4E-01
Adult	8.0E-01	8.0E-01	9.3E-01	7.9E-01	8.0E-01	8.2E-01
TOTAL	1.1E+00	1.1E+00	1.4E+00	1.1E+00	1.1E+00	1.2E+00


Production / consumption factors:

Produce: <1

Milk: 1.4

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site QUAD CITIES ROCK ISLAND, ILLINOIS
 Location N 41.7272° W 90.3417°
 Total Population Within 2-80-km Region 6.7E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Davenport SMSA</u>	<u>360,000</u>	<u>40 km SW</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 1.1E8 kg Milk 1.8E8 l Meat 1.9E8 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 88%

Period of Record 1 JAN 74 - 31 DEC 75

Average Mississippi River Flow at Site 47,000 ft³/s

Drinking Water

Exposed Population 370,000^(a) Dilution Factor 1

Fish

Edible Harvest 2.1E6^(b) kg/yr Dilution Factor 0.5^(b)

(a) All people in Davenport SMSA assumed to drink river water.

(b) Assumes 1/2 fish harvest caught below plant.

POPULATION DOSE-COMMITMENT ESTIMATES AND
 AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
 QUAD CITIES 1 & 2

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	3.7E-03	1.1E-03	7.0E-02	2.5E-02	2.9E-02
Child	4.3E-01	3.4E-02	5.1E-01	2.1E+00	2.4E+00
Teen	7.5E-01	4.6E-02	1.6E-01	1.2E+00	1.9E+00
Adult	8.1E+00	3.9E-01	1.1E+00	6.8E+00	1.1E+01
TOTAL	9.3E+00	4.7E-01	1.9E+00	1.0E+01	1.5E+01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	2.0E-02	1.9E-02	2.9E-01	2.0E-02	2.1E-02	2.0E-02
Child	2.3E-01	2.2E-01	2.9E+00	2.5E-01	2.4E-01	2.3E-01
Teen	1.6E-01	1.6E-01	1.2E+00	1.6E-01	1.7E-01	1.7E-01
Adult	9.7E-01	9.7E-01	5.0E+00	9.6E-01	9.8E-01	1.0E+00
TOTAL	1.4E+00	1.4E+00	9.4E+00	1.4E+00	1.4E+00	1.4E+00


Production / consumption factors:

Produce: <1

Milk: 2.0

Meat: 3.5

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
 TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site RANCHO SECO SACRAMENTO COUNTY, CALIFORNIA
Location N 38.3461° W 121.1186°
Total Population Within 2-80-km Region 1.5E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Sacramento	860,000	35 km NNW
Stockton	120,000	45 km SW

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 4.8E7 kg Milk 2.3E8 l Meat 5.0E7 kg

Regional Productivity Factor

Animal Grazing Factor 0.9
Location of Meteorological Station Site Recovery 98%
Period of Record 1 FEB 75 - 31 JAN 76

Average Water Dilution Flow from Plant _____ (a) _____ ft³/s

Drinking Water

Fish

Edible Harvest (a) kg/yr Dilution Factor -

(a) No radionuclides released in liquid effluents reported (Decker, 1978).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
RANCHO SECO

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total</u>	<u>Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
--	--------------	-------------	---------------	----------------	-------------	--------------

TOTAL	0	0	0	0	0
-------	---	---	---	---	---

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total</u>	<u>Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.1E-04	1.1E-04	1.3E-03	6.2E-05	1.1E-04	1.1E-04	1.1E-04
Child	1.4E-03	1.3E-03	8.0E-03	7.0E-04	1.4E-03	1.4E-03	1.4E-03
Teen	9.2E-04	9.1E-04	3.5E-03	4.9E-04	9.2E-04	9.2E-04	9.8E-04
Adult	5.2E-03	5.2E-03	1.3E-02	2.9E-03	5.2E-03	5.2E-03	5.4E-03
TOTAL	7.6E-03	7.5E-03	2.6E-02	4.2E-03	7.6E-03	7.6E-03	7.9E-03


Production / consumption factors:

Produce: <1

Milk: 1.2

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site H. B. ROBINSON HARTSVILLE, SOUTH CAROLINA
 Location N 34.4014° W 80.1567°
 Total Population Within 2-80-km Region 6.2E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Florence</u>	<u>29,000</u>	<u>40 km</u> <u>SE</u>
<u>Sumter</u>	<u>27,000</u>	<u>53 km</u> <u>SSW</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 7.4E6 kg Milk 5.7E7 l Meat 5.0E7 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.8

Location of Meteorological Station Site Recovery 94%

Period of Record 1 JAN 75 - 31 DEC 75

Average Water Dilution Flow from Plant 720 ft³/s

Drinking Water

Exposed Population None Dilution Factor -

Fish

Edible Harvest 1.8(a) kg/yr Dilution Factor 0.02(b)

(a) Average individual consumption rate as given in the RES (1975) used in lieu of catch data.

(b) Ten percent of population consumes fish taken from water diluted by a factor of 0.2 (FES, 1975).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
H. B. ROBINSON

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	1.1E-02	2.6E-03	5.6E-04	4.3E-02	5.3E-02
Teen	1.9E-02	4.9E-03	4.8E-04	2.6E-02	4.5E-02
Adult	2.0E-01	4.3E-02	3.6E-03	1.5E-01	2.7E-01
TOTAL	2.3E-01	5.0E-02	4.7E-03	2.2E-01	3.7E-01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	6.6E-04	6.1E-04	3.9E-02	2.9E-04	7.3E-04	6.2E-04
Child	8.7E-03	8.4E-03	2.3E-01	2.7E-03	9.0E-03	8.5E-03
Teen	5.9E-03	5.8E-03	9.5E-02	1.7E-03	6.0E-03	6.0E-03
Adult	3.4E-02	3.4E-02	3.5E-01	9.6E-03	3.5E-02	3.4E-02
TOTAL	4.9E-02	4.9E-02	7.1E-01	1.4E-02	5.0E-02	4.9E-02


Production / consumption factors:

Produce: <1

Milk: <1

Meat: 1.0

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site ST. LUCIE FORT PIERCE, FLORIDA

Location N 27.4466° W 80.3266°

Total Population Within 2-80-km Region 2.8E5

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
West Palm Beach	71,000	80 km SSE
Ft. Pierce	37,000	13 km NW
Vero Beach	15,000	40 km NNW

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 2.8E7 kg Milk 1.1E8 l Meat 7.2E7 kg

Regional Productivity Factor 0.5

Animal Grazing Factor 1

Location of Meteorological Station Site Recovery 90%

Period of Record 1 JAN 76 - 31 DEC 76

Average Dilution Flow from Plant 41 ft³/s

Fish

Edible Harvest 2.6E5 kg/yr Dilution Factor 0.005

Invertebrates

Edible Harvest 2.7E4 kg/yr Dilution Factor 0.005

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE COMMITMENT HISTOGRAM FOR
ST. LUCIE

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	4.7E-04	2.7E-03	1.9E-04	3.8E-04	6.8E-04
Teen	3.2E-04	5.6E-03	1.3E-04	2.3E-04	6.0E-04
Adult	1.9E-03	4.8E-02	8.8E-04	1.4E-03	3.6E-03
TOTAL	2.7E-03	5.6E-02	1.2E-03	2.0E-03	4.9E-03

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	4.1E-04	4.1E-04	9.3E-04	4.1E-04	4.1E-04	4.5E-04
Child	4.6E-03	4.6E-03	7.8E-03	4.6E-03	4.6E-03	5.1E-03
Teen	3.4E-03	3.4E-03	4.7E-03	3.3E-03	3.4E-03	4.0E-03
Adult	2.0E-02	2.0E-02	2.5E-02	2.0E-02	2.0E-02	2.2E-02
TOTAL	2.9E-02	2.9E-02	3.9E-02	2.8E-02	2.9E-02	3.2E-02


Production / consumption factors:

Produce: <1

Milk: 1.5

Meat: 1.6

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site SALEM SALEM, NEW JERSEY
 Location N 39.4628° W 75.5356°
 Total Population Within 2-80-km Region 4.9E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Philadelphia SMSA	4,850,000	64 Km NE
Wilmington	501,000	29 km N
Chester	56,600	45 km NNE

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 7.4E7 kg Milk 2.7E8 Meat 2.4E7 kg

Regional Productivity Factor 0.9

Animal Grazing Factor 0.6

Location of Meteorological Station Site Recovery 95%

Period of Record 1 JUN 70 - 31 MAY 71

Average Delaware River Flow at Site 16,000 ft³/s

Fish

Edible Harvest 3.6E5 kg/yr Dilution Factor 1

Invertebrates^(a)

Edible Harvest 1.6E5 kg/yr Dilution Factor 1

(a) Environment primarily salt water so invertebrates considered in lieu of drinking water.

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE COMMITMENT HISTOGRAM FOR
SALEM

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	2.2E-05	1.2E-06	1.5E-08	8.5E-05	1.5E-08
Teen	1.7E-05	2.0E-06	1.3E-08	6.9E-05	1.3E-08
Adult	1.3E-04	1.5E-05	1.1E-07	5.1E-04	1.1E-07
TOTAL	1.6E-04	1.8E-05	1.3E-07	6.7E-04	1.3E-07

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.9E-10	1.9E-10	1.9E-10	6.0E-11	1.9E-10	1.9E-10
Child	2.5E-09	2.5E-09	2.5E-09	6.6E-10	2.5E-09	2.6E-09
Teen	1.8E-09	1.8E-09	1.8E-09	4.8E-10	1.8E-09	1.9E-09
Adult	1.0E-08	1.0E-08	1.0E-08	2.9E-09	1.0E-08	1.0E-08
TOTAL	1.4E-08	1.4E-08	1.4E-08	4.1E-09	1.4E-08	1.5E-08

Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS

100% of population received an average
individual dose commitment of <1 x 10⁻⁶ mrem

Site SAN ONOFRE CAMP PENDLETON, CALIFORNIA
 Location N 33.3686° W 117.5544°
 Total Population Within 2-80-km Region 3.8E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
San Diego SMSA	1,400,000	68 km SSE
Riverside	150,000	69 km NE
Anaheim	180,000	69 km NNW
Santa Ana	170,000	55 km NNW

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 4.8E7 kg Milk 2.3E8 l Meat 5.0E7 kg

Regional Productivity Factor 0.6

Animal Grazing Factor 1

Location of Meteorological Station Site Recovery 88%

Period of Record 25 JAN 73 - 24 JAN 76

Average Dilution Flow from Plant 510 ft³/s

Fish

Edible Harvest 2.9E4^(a) kg/yr Dilution Factor 1^(a)

Invertebrates

Edible Harvest 2.9E3^(a) kg/yr Dilution Factor 1^(a)

(a) Seafood caught in undiluted effluent (FES, 1973).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
SAN ONOFRE

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	6.8E-02	3.5E-02	6.4E-03	2.8E-01	3.3E-01
Teen	1.1E-01	6.9E-02	5.0E-03	1.7E-01	2.8E-01
Adult	1.2E+00	6.0E-01	3.6E-02	9.9E-01	1.7E+00
TOTAL	1.4E+00	7.1E-01	4.8E-02	1.4E+00	2.3E+00

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	4.1E-04	4.0E-04	5.3E-03	2.0E-04	4.2E-04	4.2E-04
Child	5.3E-03	5.3E-03	3.1E-02	2.1E-03	5.4E-03	5.5E-03
Teen	3.8E-03	3.8E-03	1.4E-02	1.5E-03	3.8E-03	4.1E-03
Adult	2.2E-02	2.2E-02	5.4E-02	9.2E-03	2.2E-02	2.3E-02
TOTAL	3.1E-02	3.1E-02	1.1E-01	1.3E-02	3.2E-02	3.3E-02


Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site SURRY SURRY COUNTY, VIRGINIA
 Location N 37.1664° W 76.6972°
 Total Population Within 2-80-km Region 1.7E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Newport News SMSA	360,000	16 km ESE
Norfolk SMSA	790,000	50 km SE
Richmond SMSA	580,000	78 km NW

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 3.5E7 kg Milk 1.5E8 l Meat 7.4E7 kg

Regional Productivity Factor 0.8

Animal Grazing Factor 0.7

Location of Meteorological Station Site Recovery 91%

Period of Record 3 MAR 74 - 2 MAR 75

Average James River Flow at Site 25,000 ft³/s ^(a)

Fish

Edible Harvest 6.0E5 kg/yr Dilution Factor 1

Invertebrates ^(b)

Edible Harvest 1.1E6 kg/yr Dilution Factor 1

(a) Flow includes fresh water river flow and saline "mixing flow" of estuary (FES, 1972).

(b) Environment primarily salt water so invertebrates considered in lieu of drinking water.

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
SURRY 1 & 2

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	2.6E-01	4.0E-01	3.3E+00	2.4E-01	3.5E-01
Teen	2.3E-01	8.5E-01	2.1E+00	1.4E-01	2.9E-01
Adult	1.8E+00	7.6E+00	1.5E+01	8.3E-01	1.8E+00
TOTAL	2.3E+00	8.8E+00	2.1E+01	1.2E+00	2.4E+00

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.5E-02	1.5E-02	2.0E-01	1.3E-02	1.6E-02	1.7E-02
Child	1.8E-01	1.8E-01	1.3E+00	1.4E-01	1.8E-01	2.0E-01
Teen	1.3E-01	1.3E-01	6.1E-01	1.0E-01	1.3E-01	1.5E-01
Adult	7.6E-01	7.6E-01	2.5E+00	6.0E-01	7.6E-01	8.5E-01
TOTAL	1.1E+00	1.1E+00	4.7E+00	8.6E-01	1.1E+00	1.2E+00


Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site THREE MILE ISLAND THREE MILE ISLAND, PENNSYLVANIA
 Location N 40.1533° W 76.7269°
 Total Population Within 2-80-km Region 1.8E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Lancaster SMSA</u>	<u>320,000</u>	<u>40 km</u> <u>ESE</u>
<u>Harrisburg SMSA</u>	<u>410,000</u>	<u>16 km</u> <u>NW</u>
<u>Reading SMSA</u>	<u>300,000</u>	<u>71 km</u> <u>ENE</u>
<u>York SMSA</u>	<u>330,000</u>	<u>23 km</u> <u>S</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 5.3E7 kg Milk 5.3E8 l Meat 5.4E7 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 80%

Period of Record 1 OCT 72 - 30 SEP 73

Average Susquehanna River Flow at Site 34,000 ft³/s

Drinking Water

Exposed Population 200,000 Dilution Factor 1

Fish

Edible Harvest (a) kg/yr Dilution Factor 0.025^(b)

(a) No fish catch data given in FES so generic consumption rates used (Table A1).

(b) Ten percent of population consumes 25% of their fish from river (FES, 1972).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
THREE MILE ISLAND

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	9.8E-04	9.6E-04	1.4E-03	1.7E-04	1.1E-03
Child	1.2E-02	1.1E-02	1.4E-02	7.8E-03	1.9E-02
Teen	6.6E-03	4.7E-03	5.1E-03	4.0E-03	1.0E-02
Adult	6.2E-02	4.0E-02	4.2E-02	2.4E-02	7.2E-02
TOTAL	8.2E-02	5.7E-02	6.2E-02	3.6E-02	1.0E-01

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.7E-02	1.7E-02	2.3E-02	3.8E-03	1.7E-02	1.7E-02
Child	2.3E-01	2.3E-01	2.7E-01	4.2E-02	2.3E-01	2.3E-01
Teen	1.6E-01	1.6E-01	1.7E-01	3.1E-02	1.6E-01	1.6E-01
Adult	8.7E-01	8.7E-01	9.5E-01	1.8E-01	8.7E-01	8.9E-01
TOTAL	1.3E+00	1.3E+00	1.4E+00	2.6E-01	1.3E+00	1.3E+00


Production / consumption factors:

Produce: <1

Milk: 2.2

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site TROJAN PRESCOTT, OREGON
 Location N 46.0372° W 122.8847°
 Total Population Within 2-80-km Region 1.3E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Portland SMSA	1.1E6	63 km SSE
Longview	30,000	14 km NNW
Astoria	11,000	72 km WNW

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 6.4E7 kg Milk 3.7E7 l Meat 2.6E7 kg

Regional Productivity Factor 0.9

Animal Grazing Factor 0.75

Location of Meteorological Station Site Recovery 90%

Period of Record 1 SEP 71 - 31 AUG 74

Average Columbia River Flow at Site 2.3E5 ft³/s

Drinking Water

Exposed Population 480^(a) kg/yr Dilution Factor 1

Fish

Edible Harvest 1.0E6 kg/yr Dilution Factor 1

(a) Population of Rainier divided by 4, since residents only there for 25% of the year.

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
TROJAN

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	7.2E-07	2.6E-07	1.1E-05	3.2E-06	1.8E-07
Child	1.1E-03	1.8E-03	2.0E-03	4.7E-03	1.1E-03
Teen	9.9E-04	3.7E-03	1.4E-03	3.4E-03	9.4E-04
Adult	8.1E-03	3.2E-02	9.1E-03	2.3E-02	5.6E-03
TOTAL	1.0E-02	3.8E-02	1.2E-02	3.1E-02	7.6E-03

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	9.4E-05	8.5E-05	1.4E-03	2.0E-04	8.6E-05	1.6E-04
Child	2.2E-03	1.2E-03	1.4E-02	8.2E-03	1.0E-03	1.9E-03
Teen	1.2E-03	9.2E-04	6.8E-03	4.4E-03	7.3E-04	1.5E-03
Adult	6.7E-03	5.4E-03	3.0E-02	2.2E-02	4.3E-03	7.2E-03
TOTAL	1.0E-02	7.5E-03	5.2E-02	3.5E-02	6.1E-03	1.1E-02


Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site TURKEY POINT DADE COUNTY, FLORIDA
 Location N 25.4336° W 80.3317°
 Total Population Within 2-80-km Region 2.0E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Miami SMSA</u>	<u>1,500,000</u>	<u>48 km</u> <u>NNE</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 2.8E7 kg Milk 1.1E8 l Meat 7.2E7 kg

Regional Productivity Factor 0.4

Animal Grazing Factor 1

Location of Meteorological Station Site Recovery 98%

Period of Record 1 JAN 73 - 31 DEC 73

Average Dilution Flow from Plant 3,310 ft³/s

Fish

Edible Harvest (a) kg/yr Dilution Factor 0.001

Invertebrates

Edible Harvest (a) kg/yr Dilution Factor 0.002

(a) No catch data given in DES (1972) so generic consumptions rates used (Table A1).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
TURKEY POINT 3 & 4

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	1.1E-03	3.1E-03	4.7E-02	3.2E-03	3.7E-03
Teen	1.5E-03	6.5E-03	3.3E-02	1.9E-03	3.1E-03
Adult	1.4E-02	5.6E-02	2.2E-01	1.1E-02	1.9E-02
TOTAL	1.7E-02	6.6E-02	3.0E-01	1.7E-02	2.6E-02

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	3.9E-03	3.7E-03	1.8E-01	4.4E-03	4.6E-03	4.0E-03
Child	4.3E-02	4.1E-02	9.9E-01	4.6E-02	4.7E-02	4.6E-02
Teen	3.1E-02	3.1E-02	4.0E-01	3.1E-02	3.2E-02	3.6E-02
Adult	1.9E-01	1.9E-01	1.4E+00	1.8E-01	1.9E-01	2.0E-01
TOTAL	2.6E-01	2.6E-01	3.0E+00	2.7E-01	2.7E-01	2.9E-01


Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site VERMONT YANKEE VERNON, VERMONT
 Location N 42.7803° W 72.5158°
 Total Population Within 2-80-km Region 1.4E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Pittsfield</u>	<u>58,000</u>	<u>70 km</u> <u>SW</u>
<u>Springfield SMSA</u>	<u>620,000</u>	<u>72 km</u> <u>S</u>
<u>Worcester SMSA</u>	<u>680,000</u>	<u>80 km</u> <u>SE</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 4.4E6 kg Milk 7.3E8 l Meat 2.7E7 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.4

Location of Meteorological Station Site Recovery 97%

Period of Record 1 APR 75 - 31 MAR 76

Average Connecticut River Flow at Site 10,000 ft³/s

Drinking Water

Exposed Population None Dilution Factor -

Fish

Edible Harvest (a) kg/yr Dilution Factor 0.5^(a)

(a) No fish catch data given in FES (1972). Thus 1/2 population assumed to eat fish at level of generic consumption rates (Table A1).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
VERMONT YANKEE

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	8.8E-04	6.6E-05	7.8E-06	4.3E-03	4.5E-03
Teen	1.5E-03	1.3E-04	7.0E-06	2.6E-03	3.8E-03
Adult	1.6E-02	1.1E-03	5.6E-05	1.5E-02	2.3E-02
TOTAL	1.9E-02	1.3E-03	7.1E-05	2.2E-02	3.1E-02

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.4E-03	1.4E-03	2.8E-03	1.1E-03	1.4E-03	1.5E-03
Child	1.5E-02	1.5E-02	2.6E-02	1.2E-02	1.5E-02	1.8E-02
Teen	1.1E-02	1.1E-02	1.6E-02	8.9E-03	1.1E-02	1.4E-02
Adult	6.5E-02	6.5E-02	8.6E-02	5.4E-02	6.5E-02	7.5E-02
TOTAL	9.2E-02	9.3E-02	1.3E-01	7.6E-02	9.3E-02	1.1E-01


Production / consumption factors:

Produce: <1

Milk: 4.0

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site YANKEE ROWE ROWE, MASSACHUSETTS

Location N 42.7281° W 72.9247°

Total Population Within 2-80-km Region 1.6E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
Springfield SMSA	600,000	64 km SSE
Pittsfield	58,000	34 km SW
Albany SMSA	800,000	72 km W

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 2.0E7 kg Milk 2.6E8 l Meat 1.6E7 kg

Regional Productivity Factor 1

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 94%

Period of Record 1 OCT 71 - 30 SEP 72

Average Deerfield River Flow at Site 570 ft³/s

Drinking Water

Exposed Population None Dilution Factor -

Fish

Edible Harvest (a) kg/yr Dilution Factor 0.025^(b)

(a) No catch data available so generic consumption rates used (Table A1).

(b) Ten percent of population obtain 25% of their fish from river.

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
YANKEE ROWE

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	0.0E+00	0.0E+00	0.0E+00	0.0E+00	0.0E+00
Child	4.6E-03	2.9E-03	4.2E-03	2.0E-02	1.4E-02
Teen	5.6E-03	2.3E-03	2.9E-03	1.2E-02	1.2E-02
Adult	5.3E-02	1.6E-02	1.9E-02	6.6E-02	7.0E-02
TOTAL	6.4E-02	2.1E-02	2.6E-02	9.8E-02	9.5E-02

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	3.5E-04	3.5E-04	3.7E-04	1.5E-03	3.5E-04	3.5E-04
Child	2.5E-03	2.5E-03	2.6E-03	1.0E-02	2.5E-03	2.5E-03
Teen	9.2E-04	9.2E-04	9.7E-04	3.2E-03	9.2E-04	9.4E-04
Adult	3.5E-03	3.5E-03	3.7E-03	9.6E-03	3.5E-03	3.6E-03
TOTAL	7.2E-03	7.2E-03	7.6E-03	2.4E-02	7.2E-03	7.4E-03


Production / consumption factors:

Produce: <1

Milk: 1.2

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



Site ZION ZION, ILLINOIS
 Location N 42.4594° W 87.8061°
 Total Population Within 2-80-km Region 7.0E6

Major Metropolitan Centers Within Region

<u>Center</u>	<u>Population</u>	<u>Location</u>
<u>Chicago SMSA (2/3)</u>	<u>4,600,000</u>	<u>64 km</u> <u>SSE</u>
<u>Milwaukee</u>	<u>1,400,000</u>	<u>64 km</u> <u>N</u>
<u>Kenosha</u>	<u>80,000</u>	<u>14 km</u> <u>NNW</u>

Average Annual State Production of Crops and Animal Products in 80-km Radius Circle

Veg 1.1E8 kg Milk 1.8E8 l Meat 1.9E8 kg

Regional Productivity Factor 0.5

Animal Grazing Factor 0.5

Location of Meteorological Station Site Recovery 88%

Period of Record 1 JAN 74 - 31 DEC 75

Average Water Dilution Flow from Plant 1100 ft³/s

Drinking Water

Exposed Population 6,700,000 Dilution Factor 0.037^(a)

Fish

Edible Harvest 5.0E6 kg/yr Dilution Factor 0.01^(b)

(a) Drinking water dilution factor estimated by averaging dilution factors derived from FES (1972) suitably weighted for population.

(b) Dilution factor derived from FES (1972).

POPULATION DOSE-COMMITMENT ESTIMATES AND
AVERAGE INDIVIDUAL DOSE-COMMITMENT HISTOGRAM FOR
ZION 1 & 2

Dose Commitments (person-rem) from Liquid Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>
Infant	1.4E-01	1.4E-01	1.4E-01	0.0E+00	1.4E-01
Child	1.5E+00	1.5E+00	1.5E+00	0.0E+00	1.5E+00
Teen	5.9E-01	5.9E-01	5.9E-01	0.0E+00	5.9E-01
Adult	5.0E+00	5.0E+00	5.0E+00	0.0E+00	5.0E+00
TOTAL	7.3E+00	7.3E+00	7.3E+00	0.0E+00	7.3E+00

Dose Commitments (person-rem) from Airborne Pathways

	<u>Total Body</u>	<u>GI-LLI</u>	<u>Thyroid</u>	<u>Bone</u>	<u>Liver</u>	<u>Lung</u>
Infant	1.5E-01	1.5E-01	1.8E-01	1.5E-01	1.5E-01	1.6E-01
Child	1.7E+00	1.7E+00	2.0E+00	1.7E+00	1.7E+00	1.8E+00
Teen	1.2E+00	1.2E+00	1.4E+00	1.2E+00	1.2E+00	1.4E+00
Adult	7.4E+00	7.4E+00	8.0E+00	7.4E+00	7.4E+00	8.0E+00
TOTAL	1.0E+01	1.0E+01	1.1E+01	1.0E+01	1.0E+01	1.1E+01


Production / consumption factors:

Produce: <1

Milk: <1

Meat: <1

FRACTION OF POPULATION RECEIVING AN INDICATED AVERAGE
TOTAL-BODY DOSE COMMITMENT FROM AIRBORNE PATHWAYS



APPENDIX

APPENDIX

MODELS

The calculational models used were primarily those given in the Nuclear Regulatory Commission's Regulatory Guide 1.109 (1977). Computer programs were written to use these models to generate population dose commitments for four age groups. The fractions of the population comprising the four age groups were 1.44% infant (0 to 1 yr); 16.0% child (1 to 11 yr); 11.7% teen (11 to 17 yr); and 70.9% adult (17 and older) (Population Estimates and Projections, 1975). The site-dependent parameters were taken where possible from the Environmental Statements issued for each reactor (Table 3). The generic parameters used for this study such as consumption rates, occupancy factors and holdup times are given in Tables A1 and A2 below. It should be noted that generic consumption rates for aquatic foods and inhalation rates are now taken from the revised Regulatory Guide 1.109 (1977).

Bioaccumulation factors and terrestrial food transfer factors were taken from Regulatory Guide 1.109 (1977). Dose commitment factors for the four age groups were taken from the NRC document NUREG-0172 (Hoenes et al., 1977).

TABLE A1. Generic Consumption Rates and Occupancy Factors Used for the Study for Average Members of the Population(a)

Pathway	Infant	Child	Teen	Adult
Fruits, vegetables and grain (kg/yr)	0	200	240	190
Milk (l/yr)	170	170	200	110
Meat and poultry kg/yr)	0	37	59	95
Fish (kg/yr)(b)	0	2.2	5.2	6.9
Invertebrates (kg/yr)	0	0.33	0.75	1.0
Drinking water (l/yr)	170(c)	260	260	370
Inhalation (m ³ /yr)	1400(d)	3700	8000	8000
Air submersion and ground irradiation occupancy factor	0.5	0.5	0.5	0.5

(a) Regulatory Guide 1.109 (1977)

(b) Both fresh and saltwater

(c) Assumed to be equal to milk consumption

(d) Same as for maximum individual

TABLE A2. Holdup Times Between Harvest
and Consumption of Foods(a)

Food	Holdup Time (day)
Fruits, Grains, and Vegetables	14
Milk(b)	4
Meat(b)	20
Aquatic Foods (fish and invertebrates)	7
Drinking Water	1

(a) Regulatory Guide 1.109 (1977)

(b) Value given is time after milking or slaughter. For the portion of the time animals were fed stored feed, an additional 90 days was added to the holdup time.

RELEASES

The doses were estimated using the measured releases as reported by the site operators for 1976 (Decker, 1978).^(a) These releases include all radionuclides specified by the NRC to be measured and reported by the operators of all commercial nuclear power plants. Radionuclides given as a combination of parent/daughter isotopes such as Y/Sr-90, Zr/Nb-95, Ba/La-140, I/Xe-133, and Pr/Ce-144 were divided evenly between the parent and daughter.

The radionuclides used in this study along with their half-lives are given in Table A3. Note that the "+D" after some of the nuclides indicates that the decay energy of the daughter is included with the parent. Thus, whenever a parent nuclide release is specified, the result of the dose calculation will be as though an additional equilibrium amount of the daughter nuclide is specified. The daughter nuclide itself will be included separately if it can be released independently of the parent and/or if it has a relatively long half-life.

(a) Very short-lived isotopes such as Kr-90, 91, 93, 94, Xe-139, 140, 141, 143, and RB-88M; those not likely to be produced; and those which were daughters whose decay energies were accounted for in the dose factor for the parent were not included in the dose.

TABLE A3. Radionuclides Considered in This Study

No.	Nuclide	Decay Const. (1/sec)	No.	Nuclide	Decay Const. (1/sec)
1	H3	1.78E-09	43	NB97	1.57E-04
2	BE10	1.37E-14	44	M099+D	2.92E-06
3	C14	3.83E-12	45	TC99M	3.19E-05
4	N13	1.16E-03	46	RU103+D	2.02E-07
5	F18	1.05E-04	47	RU106+D	2.17E-08
6	NA22	8.44E-09	48	AG110M+D	3.19E-08
7	NA24	1.28E-05	49	CD115M	1.80E-07
8	AR41	1.05E-04	50	CD115	3.60E-06
9	SC46	9.58E-08	51	SN125+D	8.31E-07
10	CR51	2.89E-07	52	SB124	1.33E-07
11	MN54	2.57E-08	53	SB125+D	8.06E-09
12	MN56	7.47E-05	54	TE132+D	2.47E-06
13	FE55	8.14E-09	55	TE133M+D	2.09E-04
14	FE59	1.80E-07	56	I131+D	9.97E-07
15	C057	2.97E-08	57	I132	8.42E-05
16	C058	1.12E-07	58	I133+D	9.25E-06
17	C060	4.17E-09	59	I134	2.20E-04
18	NI57	5.35E-06	60	I135+D	2.92E-05
19	NI63	2.20E-10	61	XE131M	6.69E-07
20	NI65	7.64E-05	62	XE133M	3.61E-06
21	CU64	1.52E-05	63	XE133	1.52E-06
22	ZN65	3.31E-08	64	XE135M	7.56E-04
23	ZN69M+D	1.39E-05	65	XE135	2.10E-05
24	AS76	7.32E-06	66	XE137	3.01E-03
25	BR82	5.44E-06	67	XE138+D	8.14E-04
26	KR83M	1.04E-04	68	CS134	1.07E-08
27	KR85M	4.31E-05	69	CS136	6.17E-07
28	KR85	2.05E-09	70	CS137+D	7.31E-10
29	KR87	1.52E-04	71	CS138	3.58E-04
30	KR88+D	6.89E-05	72	CS139+D	1.24E-03
31	KR89	3.64E-03	73	BA139	1.39E-04
32	RB88	6.53E-04	74	BA140+D	6.28E-07
33	RB89+D	7.61E-04	75	LA140	4.78E-06
34	SR89+D	1.59E-07	76	LA141	4.97E-05
35	SR90+D	7.58E-10	77	CE141	2.47E-07
36	SR91+D	2.03E-05	78	CE144+D	2.83E-08
37	SR92+D	7.11E-05	79	EU152	1.69E-09
38	Y90	3.01E-06	80	EU154	2.55E-09
39	Y91M+D	2.32E-04	81	W187	8.06E-06
40	ZR95+D	1.22E-07	82	TH232+D	1.57E-18
41	ZR97+D	1.14E-05	83	NP239	3.42E-06
42	NB95	2.29E-07			

METEOROLOGY

In instances where more than one set of meteorological (joint frequency) data were available for a site, the one which appeared to be the most reliable was used to generate atmospheric transport factors. Factors were calculated for 16 compass points, and ten radii from 2 to 80 km (see Table A4) using the NRC computer program X0QDOQ (Sagendorf, 1977).

TABLE A4. Radii Intervals and Midpoints for Airborne Dose Calculations (km)

<u>Interval</u>	<u>Midpoint</u>
2 - 3	2.5
3 - 4	3.5
4 - 6	5
6 - 9	7.5
9 - 14	11.5
14 - 20	17
20 - 30	25
30 - 40	35
40 - 60	50
60 - 80	70

The X0QDOQ program generates four sets of atmospheric transport factors:

- Average annual atmospheric dilution factors which are not corrected for cloud depletion or radioactive decay;
- Dilution factors which are only corrected for decay assuming a 2.26-day half-life;
- Dilution factors which are corrected for depletion and for decay assuming an 8-day half-life;
- Relative deposition per unit area.

These factors were used to estimate the dose from airborne releases using methods similar to the NRC GASPAR program (Eckerman, 1976). Except for the four additional sites, the transport factors used this year were the same as those used for the 1975 estimates.

- The assumptions used in the calculation of these transport factors were: 50-meter source height with no correction for plume rise or building wake effects.

- semi-infinite cloud model with sector average, Gaussian plume dispersion
- no correction for terrain height variation.

Since information as to height and locations at each site for the releases given in Decker (1978) was unavailable, a single generic height of 50 meters was used at each site for the release point. Because of these uncertainties as to heights and locations of releases, estimates of dose to persons living within 2 km of the site could be in serious error; only persons living between 2 to 80 km from the site were included in the dose estimates.

POPULATION

The determination of population distributions around each site from 2 to 80 km was accomplished using a computer program and data base derived from the 1970 census. The program and data base were developed by the Department of Commerce and subsequently adopted for use with population exposure problems by the Environmental Protection Agency (Athey, 74; Hill, 77).

The population data base used was an edited and compressed version of the 1970 Census Bureau's "Master Enumeration District List with Coordinates." It contains housing and population counts for each census enumeration district (CED) and the geographic coordinates of the population centroid for each district. Using a modified version of the EPA program and the data base, an estimate of the population in each of the 160 segments around each site was made from a distance of 2 to 80 km.

The populations for 1976 were estimated using the "net increase" factors by state over the population values for 1970 as given in Statistical Abstract of the U. S., 1977 (Table No. 12).

FOOD PRODUCTION VS. FOOD CONSUMPTION

The total food production for the region within 80 km around each site was the product of the NRC statewide productivity figure for each state and a site productivity factor. At some sites this total production may be more or less than the total consumption; i.e., population times average individual consumption. (See Table A1 for generic consumption rates.) When production was more than consumption for a site, it was assumed that all persons in the

2 to 80-km region ate contaminated food; when production was less than consumption, it was assumed that dilution would occur, with uncontaminated food shipped into the area from outside. Thus, the calculated doses for a particular food type were reduced in proportion to the ratio of production ÷ consumption (production/consumption <1).

The dose to persons outside the 80-km limit from food shipped out of the region, in the case of production more than consumption, was not included in this report since only the dose within the 80-km radius was desired. These production/consumption factors are given as footnotes to the airborne dose-commitment tables in the Site Summary section for reference.

DRINKING WATER

The population between 2 and 80 km of the site exposed to drinking water contaminated with radionuclides released from the plants at each site was obtained generally from the Environmental Statement (ES) for the plant. For all sites on salt water it was assumed no drinking-water dose was received. The consumption rates used are given in Table A1 for drinking water.

The radionuclide concentration in the drinking water consumed by a population downstream from a site was estimated in most cases assuming 100% mixing of the plant effluent with the river. For lakes, an overall dilution factor was estimated from dilution factors given in the ES for each population center along the shore (within 80 km) which consume the contaminated lake water. These individual factors were weighted by population and averaged to obtain an effective dilution factor for the total population exposed to contaminated drinking water.

AQUATIC FOOD

Wherever possible, the fish-catch data from the plant ES was used to estimate aquatic food consumption rates for the population in the region. When this data was not found in the ES or considered unrealistic, the generic values of Table A1 were used.

The average radionuclide concentration of the waters in which this food was harvested was estimated assuming an additional dilution over the effluent flow from the reactor. For rivers, it was assumed the fish were caught in waters in which the plant effluent was completely diluted. For lakes, an additional dilution factor as given in the ES was used, but when absent, a generic value of 0.01 was used. For ocean and bay sites, a generic value of 0.001 and 0.002 was used for fish and invertebrates, respectively, if the ES yielded no values for these parameters. Invertebrates were not assumed to be caught in sufficient quantity at freshwater sites (river and lake) to affect the population dose and therefore were not included in the dose calculation. Any exceptions to these general guidelines are explained in the footnotes to the individual site summaries.

APPENDIX REFERENCES

Athey, T. W., Tell, R. A. and Janis, D. E., The Use of an Automated Population Date Base in Population Exposure Calculations, CONF. 741 018, Proceedings of Symposium on Population Exposures, 8th Midyear Topical Symposium, Health Physics Society, Knoxville, TN, Oct. 21-24, 1974. Published by the East Tennessee Chapter of the Health Physics Society.

Decker, T. R., Radioactive Materials Released from Nuclear Power Plants, (1976), NUREG-0367, U.S. Nuclear Regulatory Commission, March 1978.

Eckerman, K., GASPAR DOSE CODE, (Computer Listing) U.S. Nuclear Regulatory Commission, November 5, 1976.

Final Environmental Statement Concerning Proposed Rule Making Action: Numerical Guides for Design Objectives and Limiting Conditions for Operation to Meet the Criterion 'As Low as Practicable' for Radioactive Material in Light-Water-Cooled Nuclear Power Reactor Effluents, Directorate of Regulatory Standards, U.S. Atomic Energy Commission, WASH-1258, Vol. 1, July 1973.

Hill, E. R. Calculation of Population Distribution Using a Computer Data Base, BNWL-2395, Pacific Northwest Laboratory, August 1977.

Hoenes, G. R. and Soldat, J. K., Age-Specific Radiation Dose Commitment Factors for a One-Year Chronic Intake. NUREG-0172, U.S. Nuclear Regulatory Commission, November 1977.

Population Estimates and Projections, Current Population Reports, U.S. Dept. of Commerce, Social and Economic Statistics Administration, Bureau of the Census, Series P-25, No. 541, February 1975, p. 8.

Regulatory Guide 1.109, Calculation of Annual Doses to Man from Routine Releases of Reactor Effluents for the Purpose of Evaluating Compliance with 10 CFR Part 50, Appendix I, U.S. Nuclear Regulatory Commission, October 1977, Revision 1.

Sagendorf, J. F. and Goll, J. T., X0QDOQ - Program for the Meteorological Evaluation of Routine Effluent Releases at Nuclear Power Stations, NUREG-0324, U.S. Nuclear Regulatory Commission, September 1977.

Statistical Abstract of the United States: 1977, (98th edition), U.S. Dept. of Commerce, Social and Economic Statistics Administration, Bureau of the Census, Washington, DC, 1977.

U.S. NUCLEAR REGULATORY COMMISSION
BIBLIOGRAPHIC DATA SHEET

LE AND SUBTITLE (Add Volume No. if appropriate)
Population Dose Commitments Due to Radioactive Releases
from Nuclear Power Plant Sites in 1976

1. REPORT NUMBER (Assigned by DDC)

NUREG/CR-1125

2. (Leave blank)

3. RECIPIENT'S ACCESSION NO.

7. AUTHOR(S)

David A. Baker

5. DATE REPORT COMPLETED

MONTH August YEAR 1979

9. PERFORMING ORGANIZATION NAME AND MAILING ADDRESS (Include Zip Code)

Pacific Northwest Laboratories
• Battelle Boulevard
Richland, Washington 99352

DATE REPORT ISSUED

MONTH December YEAR 1979

6. (Leave blank)

8. (Leave blank)

10. PROJECT/TASK/WORK UNIT NO.

11. CONTRACT NO. NRC Fin No.
B2243-8

13. TYPE OF REPORT

Technical

PERIOD COVERED (Inclusive dates)

1976

15. SUPPLEMENTARY NOTES

14. (Leave blank)

16. ABSTRACT (200 words or less)

Population radiation dose commitments have been estimated from reported radionuclide releases from commercial power reactors operating during 1976. Fifty-year dose commitments from a one-year exposure were calculated from both liquid and atmospheric releases for four population groups - infant, child, teenager and adult - residing between 2 and 80 km from each site.

Results are given in the form of tables giving the dose commitments for both liquid and airborne pathways for each age group and organ. Also included for each site is a histogram showing the fraction of the total population within the 2 to 80-km region around each site receiving various average dose commitments from the airborne pathways.

The total dose commitment from both liquid and airborne pathways ranged from a high of 160 person-rem to a low of 0.0002 person-rem with an arithmetic mean of 11 person-rem. The total population dose for all sites was estimated at 470 person-rem for the 90 million people considered at risk.

17. KEY WORDS AND DOCUMENT ANALYSIS

17a. DESCRIPTORS

17b. IDENTIFIERS/OPEN-ENDED TERMS

18. AVAILABILITY STATEMENT

Unlimited

19. SECURITY CLASS (This report)

Unclassified

21. NO. OF PAGES

108

20. SECURITY CLASS (This page)

Unclassified

22. PRICE

S