

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

ORNL/TM--8831

DE83 016395

Health and Safety Research Division

POLONIUM-210 AND LEAD-210 IN FOOD AND TOBACCO PRODUCTS: A REVIEW OF PARAMETERS AND AN ESTIMATE OF POTENTIAL EXPOSURE AND DOSE

A. P. Watson

Date Published: July 1983

**Prepared for the
FLORIDA INSTITUTE OF PHOSPHATE RESEARCH
1855 West Main Street
Bartow, Florida 33830
under Interagency Agreement DFP#81002/DOEERD-81-163**

by

**OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee
operated by
UNION CARBIDE CORPORATION
for the
U. S. DEPARTMENT OF ENERGY
under Contract No. W-7405-eng-26**

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

PL4

CONTENTS

	<u>Page</u>
LIST OF TABLES	v
ABSTRACT	vii
1. INTRODUCTION	1
2. Pb-210 AND Po-210 SOURCES.	1
3. ROUTES OF UPTAKE	2
3.1 FOOD.	2
3.1.1 Distribution and Uptake in Edible Crops.	6
3.1.2 Distribution and Uptake in Animal Products	11
3.1.3 Estimated Dietary Exposure	13
3.2 TOBACCO	17
3.2.1 Origin	17
3.2.2 Estimated Inhalation Exposure.	18
4. ESTIMATED DOSE FROM DIETARY AND TOBACCO SMOKE EXPOSURES. . .	22
4.1 DIETARY INGESTION	22
4.2 TOBACCO SMOKE INHALATION.	25
5. RECOMMENDATIONS AND CONCLUSIONS.	27
REFERENCES	29

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1 Average concentrations (pCi/kg fresh weight) of naturally occurring Pb-210 and Po-210 in some common food categories.	4
2 Parameters representing soil-plant transfer of Pb-210. . .	8
3 Parameters representing soil-plant transfer of Po-210. . .	9
4 Parameters representing transfer of Pb-210 and Po-210 to meat and milk	12
5 Daily intake of naturally occurring Pb-210 (pCi/d) and Po-210 (pCi/d) in the diet (excluding water and beverages)	14
6 Pb-210 and Po-210 content of commercial cigarettes	19
7 Estimated daily inhalation exposure of Pb-210 and Po-210 to smokers.	23
8 Estimated range of adult 50-year dose commitment (rem) for one year's dietary intake of Pb-210 and Po-210 from natural sources.	24
9 Estimated range of committed dose equivalent (rem) for one year's inhalation exposure tobacco smoke containing natural concentrations of Pb-210 and Po-210.	26

ABSTRACT

Food-chain transport of Pb-210 and Po-210 from soil to edible plant parts and from animal feed to meat and milk were evaluated from a review of literature. The degree of transfer was characterized by estimating concentration factors (unweighted arithmetic means) as well as the transfer coefficients B_v , B_r (unweighted geometric means), f_m and f_f (unweighted arithmetic means). Global dietary intake of Pb-210 and Po-210 was also summarized, and 50-year dose estimates to target organs calculated. The greatest estimated ingestion doses were those to populations with large dietary complements of animal protein in the form of seafood (Japan) or caribou/reindeer muscle and organ meats (Arctic Eskimos and Lapps). The magnitude of this latter source illustrates the importance of simple food chains in generating significant exposures to populations dependent upon them.

The origin and magnitude of inhalation exposure and dose from tobacco products was also assessed. For the majority of internal organs evaluated, the dose resulting from smoking commercially available tobacco products is comparable to or greater than the dose estimates for ingestion of naturally occurring dietary Pb-210 and Po-210.

1. INTRODUCTION

The mining and milling of phosphate ores in Southwest Florida has produced large volumes of processed ore and tailings materials that contain quantities of the naturally occurring isotope U-233 ($T_{1/2} = 4.5 \times 10^9$ y) (Kocher, 1977). This isotope and the principal members of its decay series, Ra-226 ($T_{1/2} = 1600$ y), Pb-210 ($T_{1/2} = 22.3$ y), and Po-210 [$T_{1/2} = 138.4$ d (Kocher, 1977)] have been identified as potential sources of radiation exposure to local residents via food ingestion, and have therefore been chosen as candidates for evaluation. The transport and presence of Ra-226 in drinking water and food has been assessed in a previous report by the author and her colleagues (Watson, Etnier, and McDowell-Boyer, 1983); Pb-210 and Po-210 will be addressed in the current evaluation.

2. Pb-210 AND Po-210 SOURCES

Lead-210 and Po-210 are directly produced during radioactive decay of the parent radionuclide Ra-226; which is, in turn, a daughter of U-238 decay. Lead-210 and Po-210 are found at elevated concentrations in association with a Ra-226 source, but are often detected at some distance from supposed sites of release due to the extreme mobility of the noble gas Rn-222 [$T_{1/2} = 3.8$ d (Kocher, 1977)], an intermediate product of Ra-226 decay. The rapid decay of Rn-222 in the atmosphere generates Pb-210 and Po-210, which are adsorbed by aerosols and returned to earth as surficial deposition or rainout. The degree of potential long-range atmospheric transport in North America has been previously demonstrated for Rn-222 (Travis et al., 1979). Other intermediates in the decay

series between Ra-226 and Po-210 exist, but are all relatively short-lived and are thus not important to food-chain transport.

Radium-226 is found in the presence of uranium-bearing ores such as in phosphate and uranium mines. It and its daughters also occur naturally in other minerals (shale, coal, granite, and lead), soil, geothermal sites, and well water (particularly those draining granitic aquifers). Technologically enhanced sources of these isotopes, i.e., from an "activity not expressly designed to produce radiation" (Gesell and Prichard, 1975) include operation of coal-fired power plants, combustion of gasoline containing anti-knock compounds composed of lead, and application of phosphate fertilizers to agricultural soils (Moore, Martell, and Poet, 1976; Jaworowski and Grzybowska, 1977; Jaworowski and Kownacka, 1976; Eisenbud and Petrow, 1964; Breslin and Glauberman, 1970; Jaworowski, Bilkiewicz, and Zyliez, 1971; Moore and Poet, 1976). Human intake of Pb-210 and Po-210 from these sources via food and tobacco products will be the focus of this report.

3. ROUTES OF UPTAKE

3.1 FOOD

Ingestion of food is considered the single greatest source of naturally occurring lead among individuals consuming a modern, Western-style diet in a major metropolitan area. In a study of New York City residents, as much as 85.7% of total daily Pb-210 intake was supplied by food when compared with intake via air (12.9%) and water (2.9%) (Bogen, Welford, and Morse, 1976). Inhalation of cigarette smoke was not considered in their survey. Similar results were obtained in a controlled

diet study of 12 institutionalized adult males in Illinois (Spencer et al., 1977). When ingestion of food/water, inhalation of air/cigarette smoke and decay of inhaled Rn-222 were all taken into consideration, 78.1% of the total daily Pb-210 intake was estimated to be supplied in food, 69% in water, 5.0% in air, and 0.4% by Rn-222 decay. Cigarette smoke inhalation supplied the remaining 9.4% (Spencer et al., 1977).

Food ingestion also represents the major source of naturally occurring Po-210 intake. Additional data collected by Spencer and her colleagues (1977) indicates that 77.3% of an adult male's daily Po-210 intake is supplied by food, 4.7% by water and 0.6% from air. Inhalation of cigarette smoke provides more Po-210 (17.1%) than drinking water and air combined. A similar distribution was observed in a market basket survey of typical food items comprising an adult diet in Bombay, India (Khandekar, 1977). With no correction for cigarette smoking sources, food in Bombay was estimated to supply 86.4% of the total daily adult Po-210 intake, while 9.3% was supplied by water, and 4.3% by inhalation.

Concentrations of Pb-210 and Po-210 in food categories of mixed diets consumed in cities of the U.S.A., U.S.S.R., southern India, and Japan are summarized in Table 1. Lowest values were observed in milk and dairy products, which do not comprise a major portion of the non-Western diet. The greatest values were determined for Pb-210 and Po-210 in the meat/fish category of Finland, Japan, and Kalpakkam, India (Kauranen and Miettinen, 1969; Okabayashi et al., 1975; and Iyengar et al., 1980). Animal protein utilized in these regions is comprised of caribou and seafood, respectively. Maximal values such as these are illustrative of the role that local availability and ethnic/cultural preferences can

Table 1. Average concentrations (pCi/kg fresh weight) of naturally occurring Pb-210 and Po-210 in some common food categories^a

Country/City	Cereals/		Leafy		Other		Potatoes		Grain		Milk/milk		Meat/		Citation
	Pb-210	Po-210	Pb-210	Po-210	Pb-210	Po-210	Pb-210	Po-210	Pb-210	Po-210	Pb-210	Po-210	Pb-210	Po-210	
U.S.A.															
New York City	0.8	-	-	-	0.7	-	1.5	-	1.6	-	0.3	-	1.2	-	Morse and Helford, 1971
U.S.S.R.															
Rostov-on-Don	5.7	4.0	-	-	3.5	1.5	6.2	3.1	2.2	1.7	0.6	0.2	4.7	5.0	Ladinskaya et al., 1973
India															
Bombay	-	1.4	-	5.7	-	1.6	-	0.4	-	-	-	0.4	-	3.0	Khandekar, 1977
Bombay	3.5	-	-	-	-	-	-	-	-	-	-	-	-	-	Lalit, Ramachandran, and Dajan, 1990
Kalpakkam	6.3	9.1	44.5	6.3	3.3	4.5	3.0	3.2	-	-	-	-	10.8	53.7	Ivenkar et al., 1990
Finland															
Inari, Lapland	-	-	-	-	-	-	-	-	-	-	-	-	5.9 to 574	122.7 to 2773	Kauranen and Viettinen, 1969
Japan															
National composite	1.5	-	17.0	-	1.6	-	0.7	-	4.2	-	0.3	-	91.1	-	Takata, Matanabe, and Ichikawa, 1969
Chiba and Okayama Prefecture	-	1.1	-	3.5	-	-	-	0.8	-	3.5	-	-	-	242.4	Okabayashi et al., 1975
Tokyo	5.0	-	10.8	-	3.0	-	2.7	-	-	-	0	-	40.1	-	Kanetani et al., 1981

^aUnweighted means.

make in determining the quantity of radionuclides ingested; the degree of dietary exposure is a direct consequence of the consumption rates for each category in Table 1.

In regions of harsh climate, where the food supply is governed by the productivity of simple food chains, the concentration of Pb-210 and Po-210 available for consumption may exceed "normal" U.S. rates by more than two orders of magnitude. Such is the case in the Arctic, where the lichen-caribou foodchain is exploited by native hunters and their families, who consume muscle and organ meats supplied by their herds (see the Pb-210 and Po-210 values for meat consumed by Lapp herders, Table 1). Polonium-210 in muscle from winter-killed caribou has averaged 248.3 pCi/kg; muscle from summer-killed caribou has contained 85.3 pCi (Holtzman, 1966a). These values illustrate a second point; human dietary content may be highly seasonal. In this case, lichens, upon which Pb-210 and Po-210 are deposited as natural fallout, possess a roughened surface from which deposited particles are not readily removed by wind and rain. This feature, coupled with the fact that lichens are slow-growing perennials, permits these species to passively accumulate elevated concentrations of dust. Caribou and reindeer graze almost exclusively on lichens during the winter; summer pastures provide a mixed forage of annual grasses and sedges as well as lichen. Annual species are less effective accumulators than lichens.

Kalpakkam is a coastal town on the Bay of Bengal, where fishing is a commercial activity. It is also an area of high natural radiation background, as the beach sands contain monazite. This factor, plus the understanding that unsupported Po-210 (i.e., Po-210 in the absence of

its parent Pb-210) is preferentially taken up by marine organisms, is thought to account for the elevated fish muscle values found at this site (Iyengar et al., 1980). The possibility of surface deposition from the monazite sands is also a likely explanation for the high levels of Pb-210 found associated with leafy vegetables at this site.

3.1.1 Distribution and Uptake in Edible Crops

These two isotopes enter the human food chain via two mechanisms: (1) plant uptake from soil and/or water, and (2) particle deposition onto plant surfaces. The Pb-210 and Po-210 content of edible foliage and reproductive structures (seeds, berries, fruits, etc.) may be the result of either one or both mechanisms. Careful studies which discriminate between each source are rare. Actual root uptake and translocation into edible portions of plants used as human food has been observed to equal <1%. The dominant mechanism is considered to be surface deposition of particulate matter on which Pb-210 and Po-210, originating as decay daughters of Rn-222 gas, are adsorbed (Hill, 1960; Berger, Erhardt, and Francis, 1965). The overwhelming source of plant Po-210 is the deposition of its precursor, Pb-210, on plant surfaces during rainfall events (Francis, Chesters, and Erhardt, 1968).

The degree of soil-to-plant uptake of any nuclide may be characterized by mathematical expressions involving several parameters. The soil-to-plant concentration factor (CF_{sp}) is defined as the unitless ratio of fresh weight specific activity in plants to dry weight specific activity in soil at harvest or equilibrium (McDowell-Boyer, Watson, and Travis, 1980). Further discrimination into edible plant parts permits evaluation of the ratio for vegetative and reproductive organs, i.e., B_v

(ratio of soil-to-plant transfer for vegetative portions of food crops and feed plants) and B_r (ratio of soil-to-plant transfer for reproductive and storage organs of food crops and feed plants) (Baes et al., in review). Reasonable values for these parameters are needed for operation of equilibrium food-chain models to estimate potential exposure when site-specific transfer factors are unknown (Ng, 1982). Since only a very few published studies of Pb-210 and Po-210 contain data appropriate for estimating transfer coefficients for these specific isotopes, the CF_{sp} , B_v and B_r values reported here (Tables 2 and 3) are element-specific and include data on several isotopes.

These parameters are thought to be lognormally distributed (Ng, 1982); thus, geometric means are an appropriate statistic by which to summarize results. Both arithmetic and geometric means are presented in Tables 2 and 3. These estimates also represent an average over a range of soil and plant conditions; soil pH and organic/phosphorus content, plant species, and the physiologic condition of the plant are all known to govern uptake and translocation (Dedolph et al., 1970; John and Van Laerhoven, 1972a; Menzel, 1965; MacLean, Halstead, and Finn, 1969; Rains, 1971; Cox and Rains, 1972; John, 1972; Zimdahl and Foster, 1976; Koeppen, 1977). To determine a more accurate estimate of soil-plant transfer, Tables 2 and 3 include values derived from only those studies in which aerosol contributions were considered insignificant. Values are presented by main crop type as recommended by Ng (1982), who considers a single soil-to-plant transfer factor for all plant material as unnecessarily adding to the uncertainty already inherent in the summarization process.

Table 2. Parameters representing soil-plant transfer of Pb-210

Transport mode	Mean CF _{sp} ^a ($\times 10^{-3}$)	Number of derived values	Range ($\times 10^{-3}$)	Mean R _v ^b ($\times 10^{-3}$)	Mean R _p ^b ($\times 10^{-3}$)	Reference
<u>Leafy vegetables (fresh weight)</u>	2.9	21	0.6 to 9.7	2.3	-	
Beet greens	1.1	3	0.8 to 1.5			Tracy, Prantl, and Quinn, 1983
Cabbage	4.9	1	-			Ter Haar, 1970
Lettuce	3.6	13	0.8 to 9.7			Ter Haar, 1970; John and Van Laerhoven, 1972(a); Rabinowitz, 1972
Spinach	1.7	2	0.6 to 2.9			John and Van Laerhoven, 1972(a)
Swiss chard	1.5	2	0.9 to 2.0			Tracy, Prantl, and Quinn, 1983
<u>Fruits and storage organs (fresh weight)</u>	2.2	30	0.1 to 15	-	1.0	
Beans (green)	3.1	3	0.2 to 7.8			Ter Haar, 1970; Tracy, Prantl, and Quinn, 1983
Beets	0.5	2	0.4 to 0.5			Tracy, Prantl, and Quinn, 1983
Carrut	3.5	7	0.1 to 15			Ter Haar, 1970; John and Van Laerhoven, 1972(a); Tracy, Prantl, and Quinn, 1983
Onion	1.2	3	0.7 to 2.5			Tracy, Prantl, and Quinn, 1983
Potato	1.9	3	0.2 to 4.8			Ter Haar, 1970; Tracy, Prantl, and Quinn, 1983
Pumpkin	0.2	1	-			Tracy, Prantl, and Quinn, 1983
Radish	3.4	4	2.2 to 4.7			John and Van Laerhoven, 1972(a); Dedolph et al., 1970
Raspberries	0.8	2	0.2 to 1.3			Tracy, Prantl, and Quinn, 1983
Strawberries	0.5	1	-			Tracy, Prantl, and Quinn, 1983
Sweet corn	4.1	1	-			Ter Haar, 1970
Tomato	0.9	3	0.1 to 2.5			Ter Haar, 1970; Tracy, Prantl, and Quinn, 1983
<u>Grains (dry weight)</u>	12.1	3	4.9 to 2.2	-	10.0	
Oats	13.5	2	4.9 to 22			John and Van Laerhoven, 1972(a)
Wheat	9.1	1	-			Ter Haar, 1970
<u>Forage, hay, feed (dry weight)</u>	91.1	23	11.6 to 370	73.1	-	
Fodder	40.0	1	-			Ter Haar, 1970
Grass hay	84.7	11	11.6 to 188.2			Dedolph et al., 1970; Milosević et al., 1980; Rayno, Momani and Sabeu, 1980
Oat tops	102.3	11	77 to 370			John and Van Laerhoven, 1972(b); Rabinowitz, 1972

^aUnweighted arithmetic mean; concentration factor.^bUnweighted geometric mean; transfer coefficients.

Table 3. Parameters representing soil-plant transfer of Po-210^a

Transport mode	Mean CF ^b ($\times 10^{-4}$)	Number of derived values	Range ($\times 10^{-4}$)	Mean η_c ($\times 10^{-4}$)	Mean η_r ($\times 10^{-4}$)	Reference
<u>Leafy vegetables (fresh weight)</u>						
Celery (leaves)	3.6	5	0.4 to 1.0	2.1	-	
Spinach	1.0	2	0.4 to 1.2			Walters, Johnson, and Hansen, 1960
	5.3	3	2.3 to 8.0			Walters, Johnson, and Hansen, 1961
<u>Fruits and storage organs (fresh weight)</u>						
Onion	2.2	10	1.7 to 10 ² to 5.2	-	2.1	Walters, Johnson, and Hansen, 1960
Pea	0.4	2	0.4 to 0.5			Walters, Johnson, and Hansen, 1960
Potato (whole)	3.3	2	0.1 to 1.4			Walters, Johnson, and Hansen, 1960
Radish	4.4	2	2.5 to 6.2			Walters, Johnson, and Hansen, 1960
Tomato	0.1	2	5.7 to 5.3			Walters, Johnson, and Hansen, 1960
		2	3.7 to 10 ² to 1.6 to 10 ⁻¹			Walters, Johnson, and Hansen, 1960
<u>Grains (dry weight)</u>						
Barley	2.6	4	1.7 to 10 ² to 4	-	2.9	Walters, Johnson, and Hansen, 1960
Wheat	5.0	2	2.6 to 7.4			Walters, Johnson, and Hansen, 1963
Corn	0.2	1	-			Walters, Johnson, and Hansen, 1960
	0.2	1	-			Walters, Johnson, and Hansen, 1961
<u>Forage, hay, feed</u>						
Fodder	39.1	13	7.3 to 1.1	2.1	-	
Grain straw	5.4	1	0.4 to 1.1			Johnson, Hansen, and Hansen, 1960
Pasture hay	54	3	1.4 to 9.1			Johnson, Hansen, and Hansen, 1960
	57	7	1.7 to 1.9			Walters et al., 1961

^aValues are nuclide-specific for chloride, oxide and naturally occurring elements of Po-210 .^bUnweighted arithmetic mean; concentration factor.^cUnweighted geometric mean; transfer coefficients.

The CF_{sp} 's, B_v 's and B_r 's are given in dry weight concentrations for forage, hay, feed and grain; and in fresh weight concentrations for vegetables and fruits (i.e., moisture content as normally consumed). When necessary, conversion of literature data to fresh or dry weight was made with the use of information supplied in the text of each paper cited or by standard conversions documented in Spector (1956). The unweighted arithmetic average of mean CF_{sp} 's, geometric means of B_v 's and B_r 's, and range of individual values are given for each food category. The ranges indicate that much uncertainty is involved in determining a single value for any transfer parameter; much of this uncertainty is probably due to the variability in experimental conditions among the studies cited as well as the wide range of soil, climatic, and nutritional conditions evaluated.

Edible plant portions appear to take up less Po-210 from soil than Pb-210; the difference is an approximate order of magnitude for leafy vegetables, fruits and storage organs; and approximately two orders of magnitude for grains and animal hay or forage. For either nuclide, the degree of uptake by fruits and vegetables is nearly the same; among dry foodstuffs, uptake is greater in hay and animal forage than grains.

The actual fraction of total grain Pb-210 or Po-210 present in the grain hull versus that present within the grain kernel is not known; most of the available studies do not mention whether analyses were performed on whole or de-hulled grain. The one study that described grain processing analyzed Po-210 in wheat "chaff" (Watters, Johnson, and Hansen, 1969). A CF_{sp} of 1.1×10^{-3} was determined for chaff, as compared to a CF_{sp} of 1.7×10^{-5} for the kernel. On the basis of this

single determination, it appears that approximately 98% of the wheat Po-210 may be incorporated into the hull. It is assumed that all remaining data presented for grain Pb-210 and Po-210 are for whole grain.

3.1.2 Distribution and Uptake in Animal Products

Coefficients developed for evaluating transfer to meat and milk incorporate daily intake rather than concentration ratios and thus consider accumulation in muscle and milk according to ingestion rates. The transfer coefficient, f_m , represents the fraction of the element or nuclide ingested daily that is secreted per liter of milk at equilibrium (day per liter); f_f represents the fraction of the element ingested daily that appears in each kilogram of meat at the time of slaughter (day per kilogram).

Very little work specifically designed to measure transfer of naturally occurring Pb-210 and Po-210 from feed sources to meat and milk of domestic animals has been performed. Those studies that exist are the result of acute controlled feeding exposure (Stanley, Mullen, and Bretthauer, 1971), Po-210 introduction directly to the rumen of individual cows (Watters and McInroy, 1969; Watters, Johnson, and McInroy, 1971), or sample collection from animals feeding near sources of industrial emissions (uranium mines and/or mills) (Holtzman et al., 1979). As a result of the sparse record of nuclide-specific data, information collected during longer-term studies of stable isotope transfer to muscle and milk were also included in the summary presented as Table 4. Sources included lead mines and smelters (Donovan, Feeley, and Canavan, 1969; Kerin and Kerin, 1971; Hammond and Aronson, 1964; Djuric et al., 1971),

Table 4. Parameters representing transfer of Pb-210 and Po-210 to meat and milk

Transport mode	Unweighted mean	Range	Reference
Pb-210			
Milk, f_m^a	1.2×10^{-4} (n=14)	1.8×10^{-6} to 4×10^{-4}	Donovan, Feeley, and Canavan, 1969; Kerin and Kerin, 1971; Bovay, 1971; Nelmes et al., 1974; Stanley et al., 1971
Beef, f_f^b	1×10^{-3} (n=4)	2×10^{-4} to 2×10^{-3}	Nelmes et al., 1974; Holtzman et al., 1979
Po-210			
Milk, f_m	1.7×10^{-4} (n=3)	3.9×10^{-5} to 2.9×10^{-4}	Watters and McInvoy, 1959; Watters, Johnson, and McInvoy, 1971; Johnson and Watters, 1972
Beef, f_f^c	2.2×10^{-3} (n=2)	3.0×10^{-4} to 4.0×10^{-3}	McDowell-Boyer and Baes, 1980; Baes et al. (in review)

^a f_m expresses the estimated fraction of the element ingested daily that is secreted per liter of milk at equilibrium (d/L).

^b f_f expresses the estimated fraction of the element ingested daily that appears in each kilogram of muscle at time of slaughter (d/kg).

^cBased in part on data for sheep, caribou, and reindeer.

automobile exhaust (Bovay, 1971) and sewage sludge containing lead (Nelmes et al., 1974). Even fewer data are available for Po-210, and all that could be obtained are included in Table 4.

Very little ingested lead is found in muscle, as this element is a bone-seeker. The principal sink for Pb-210 is the skeleton. The skeleton, as well as the liver and kidneys, are sinks for Po-210 (Beasley and Palmer, 1966; Hill, 1967; Kauranen and Miettinen, 1969). The wider distribution of Po-210 in soft tissues is thought to be partly the result of skeletal Pb-210 decay giving rise to daughter Po-210 which is, in turn, re-allocated by the bloodstream. The majority of Pb-210 in animal tissues is considered to be of dietary origin; Po-210 arises from both diet and internal decay of Pb-210. Animals and humans that regularly feed on bones (for broth or soup) and internal organs of grazers exposed to dietary Pb-210 or Po-210 have been observed to accumulate greater body burdens than those individuals who consume only muscle (Holtzman, 1966a,b).

3.1.3. Estimated Dietary Exposure

Global diets have been extensively sampled for the presence of naturally occurring radionuclides. Cultural food preferences, availability and anomalies of geology or atmosphere are all factors in governing the variability in world dietary exposures to Pb-210 and Po-210 summarized in Table 5. In general the smallest ingestion exposures were obtained from diets in developed countries (Federal Republic of Germany, U.K., U.S.A. and U.S.S.R) and the high grain and vegetable diet of metropolitan India. The greatest exposures were estimated for diets of Japan and Arctic dwellers in Finland, Canada, and Alaska; all of which contain animal protein in the form of seafood (Japan) or large,

Table 5. Daily intake of naturally occurring Pb-210 (pCi/d) and Po-210 (nCi/d) in the diet (excluding water and beverages)

Country/City	Pb-210 (pCi/d)	Po-210 (pCi/d)	Reference
U.S.A.			
Illinois	1.8 ^b	-	Holtzman, 1963
Illinois (hospital)	-	1.6 ^c	Spencer et al., 1977
Boston	1.6 ^d	-	Magno, Groulx, and Apidianakis, 1970
Chicago	1.7 ^d	-	Magno, Groulx, and Apidianakis, 1970
Honolulu	1.5 ^d	-	Magno, Groulx, and Apidianakis, 1970
Los Angeles	1.7 ^d	-	Magno, Groulx, and Apidianakis, 1970
New Orleans	1.7 ^d	-	Magno, Groulx, and Apidianakis, 1970
Palmer, Alaska	1.6 ^d	-	Magno, Groulx, and Apidianakis, 1970
Chicago (hospital)	1.3 ^c	1.6 ^c	Holtzman et al., 1974
New York City	1.2	-	Morse and Helford, 1971
New York City	1.8		Cohen, Jaakkola, and Wrenn, 1973
San Francisco (middle-income)	2.7 ^e	-	Michelson et al., 1962
Chicago (middle-income)	3.7 ^e	-	Michelson et al., 1962
New York City (middle-income)	2.0 ^e	-	Michelson et al., 1962
San Francisco (low-income)	3.9 ^e	-	Michelson et al., 1962
Chicago (low-income)	3.0 ^e	-	Michelson et al., 1962
New York City (low-income)	nd ^{e,f}	-	Michelson et al., 1962
Nine-city composite	2.4	-	Michelson, 1961; Michelson et al., 1962

Table 5. (continued)

Country/City	Pb-210 (pCi/d)	Po-210 (pCi/d)	Reference
United Kingdom	3.2	-	Hill, 1965
United Kingdom	1 to 10	1 to 10	Hill, 1967
India			
Bombay	3.4	-	Lalit, Ramachandran, and Rajan, 1980
Bombay	-	1.5 ^g	Khandekar, 1977
Federal Republic of Germany	4.6	4.6	Glöbel, Muth, and Oberhausen, 1966
U.S.S.R.			51
Leningrad	4.0	3.0	Yermolaeva-Markovskaya et al., 1968
Rostov-on-Don	6.2	4.1	Ladinskaya et al., 1973
Bulgaria	-	1.6 to 2.1	Keslev et al., 1975
Japan			
Tokyo	5.9	-	Kametani et al., 1981
Range, 7 regional sites	13.8 to 22.5 ^h	-	Takata, Watanabe, and Ichikawa, 1958
Mean, 7 regional sites	17.0 ^h	-	Takata, Watanabe, and Ichikawa, 1968
Chiba Prefecture	-	9.0 to 13.0	Okabayashi et al., 1975

Table 5. (continued)

Country/City	Pb-210 (pCi/d)	Po-210 (pCi/d)	Reference
Arctic Dwellers			
Finland (Lapps)	8.6	6.9	Kauranen and Miettinen, 1969
Canada	-	100	Hill, 1967
Alaska	10	100	Beasley and Palmer, 1966
Alaska	-	60	Blanchard, 1966
Alaska	10	100	Holtzman, 1966(a,b); Holtzman and Ilcewicz, 1971
Alaska	40	40	Ramzaev et al., 1974

^aMany of these values taken as cited by Holtzman, 1980.

^bBased on fecal and urine analyses.

^cAdult patients on controlled hospital diet.

^dAssume 2 kg/d food ingestion; data collected in 1966 for children aged 9-12 years.

^eTeenage diet.

^fNot detected.

^gSixty percent of diet sample.

^hSeven regional sites comprise the entire country, including major population centers of Tokyo and Kyoto-Osaka.

native ruminants (Arctic reindeer and caribou). The reasons for elevated Pb-210 and Po-210 concentrations in these two groups of animals have been previously discussed in Sect. 3.1.1.

3.2 TOBACCO

Inhalation of tobacco smoke is ranked second to food in providing Pb-210 and Po-210 exposure to man (Spencer et al., 1977). Among smokers, a larger portion of daily Po-210 intake (17.1%) than Pb-210 intake (9.4%) is provided by smoke. Exposure to alpha-decay products from these and other Rn-222 daughters has been established as the major risk factor in the incidence of bronchial cancer among uranium miners. It follows that exposure to alpha radiation from smoke inhalation may also be a factor in cigarette-induced bronchial cancer (Radford, 1974). Nevertheless, the reader should understand that estimates for continuous background inhalation exposure to Rn-222 and daughters (200-2000 pCi/d) greatly exceed that value estimated for Rn-222 and daughter exposure from tobacco smoke (Martell, 1974; Hamrick and Walsh, 1974).

3.2.1 Origin

The two possible sources of Pb-210 and Po-210 associated with tobacco are uptake from soils containing these isotopes as daughter products of Ra-226 or leaf surface deposition of airborne aerosols to which naturally occurring Pb-210 and Po-210 have become adsorbed. Greenhouse experiments with artificially contaminated soils and field sampling of tobacco plantations indicate that leaf scavenging of aerosols, particularly during rainfall events is the dominant mode of Pb-210 and Po-210 entry into tobacco plants (Francis, Chesters, and Erhardt, 1968).

The time elapsed between leaf harvest and actual manufacture into a cigarette or cigar determines the Po-210 content of the consumer product, which is a function of initial Po-210 deposition plus Pb-210 decay. The activity of some domestic and imported tobacco products is summarized in Table 6.

Polonium is readily volatilized at the temperature of the burning tip of a cigarette or cigar (600 to 800°C) (Abe and Abe, 1969; Radford and Hunt, 1964) and is known to be adsorbed to submicron particles (Martell, 1974), such as are found in smoke. Lead is not sublimated at these temperatures, but is a component of the resulting smoke and ash; Pb-210 can be found electrically attached to particles <0.3 microns in diameter (Kawans and Nakatani, 1964, as quoted in Francis, Chesters, and Erhardt, 1968). Lead-210 and Po-210 aerosols attached to the sticky exudate of tobacco leaf surfaces can also be encapsulated by the exudate polymer produced as part of the curing process. During tobacco combustion, these insoluble, polymerized particles are released (Fleischer and Parungo, 1974; Martell, 1974) to be inhaled by the smoker or other persons in the vicinity. Regardless of the specific mechanism, there appears to be ample opportunity for Pb-210 and Po-210 exposure to bronchial epithelium and lung tissue during the smoking process.

3.2.2. Estimated Inhalation Exposure

Numerous variables govern the degree of exposure to tobacco smoke; geographic region where the tobacco is grown, fineness of the tobacco cut; presence or absence of a cigarette filter, size and composition of the filter, smoking habits, etc. (Black and Bretthauer, 1968; Ferri and Baratta, 1966; Marsden, 1964). Estimated ranges of daily smoke intake

Table 5. Pb-210 and Po-210 content of commercial cigarettes^a

Country of manufacture	pCi per cigarette	pCi per g	Reference
Pb-210			
U.S.A.	0.017 ^b	-	Holtzman and Ilcewicz, 1966
	0.018 ^b	-	Holtzman and Ilcewicz, 1966
	0.015 ^b	-	Holtzman and Ilcewicz, 1966
	0.005 ^b	-	Holtzman and Ilcewicz, 1966
	0.013 ^b	-	Holtzman and Ilcewicz, 1966
Po-210			
Australia	-	0.64	Hill, 1965
Brazil	0.46	0.47	Dos Santos, 1970
Bulgaria	0.38	-	Nikolova and Parfenov, 1971
Canada	0.21	0.26	Black and Bretthauer, 1968
Central and South America	-	0.67	Hill, 1965
Czechoslovakia	0.2-0.45	-	Solnicka and Bischof, 1970
Egypt	0.38	0.44	Black and Bretthauer, 1968
Federal Republic of Germany	0.41-0.60	0.41-0.50	Rajewsky and Stahlhofen, 1966
Finland	0.52	0.56	Black and Bretthauer, 1968
France	0.29	0.63	Black and Bretthauer, 1968
	0.63	0.56	Black and Bretthauer, 1968

Table 6. (continued)

Country of manufacture	pCi per cigarette	pCi per g	Reference
<u>Po-210 (continued)</u>			
India and Pakistan	-	0.41	Hill, 1965
India	0-0.4	-	Athalye et al., 1972
Indonesia		0.24	Hill, 1965
Italy	-	0.37-0.53	Carfi and Lonati, 1956
Japan	-	0.31-0.34	Okabayashi et al., 1975
Japan	-	0.54	Suzuki, 1967
Japan	0.61	0.60	Black and Bretthauer, 1968
Norway	0.23	0.22	Black and Bretthauer, 1968
Philippines	0.22	0.18	Black and Bretthauer, 1968
Rhodesia	-	0.65	Hill, 1965
Turkey and Greece	-	0.24	Hill, 1965
U.S.S.R.	0.38	0.60	Black and Bretthauer, 1968
United Kingdom	0.47	0.46	Black and Bretthauer, 1968
U.S.A.	0.39-0.48	-	Radford and Hunt, 1964
U.S.A.	0.32-0.34	0.32-0.34	Rajewsky and Stahlhoffen, 1966
U.S.A.	0.54	0.52	Black and Bretthauer, 1968
U.S.A.	-	0.51	Hill, 1965

Table 6. (continued)

Country of manufacture	pCi per cigarette	pCi per g	Reference
<u>Po-210 (continued)</u>			
U.S.A.	0.023 ^b	-	Holtzman and Ilcewicz, 1966
	0.027 ^b	-	Holtzman and Ilcewicz, 1966
	0.016 ^b	-	Holtzman and Ilcewicz, 1966
	0.018 ^b	-	Holtzman and Ilcewicz, 1966
	0.020 ^b	-	Holtzman and Ilcewicz, 1966
Yugoslavia	-	0.40-0.60	Kilibarda et al., 1966

^aPartly derived from Parfenov (1974).

^bIn cigarette smoke.

of Pb-210 and Po-210 are presented in Table 7. Many of these results indicate that inhalation exposure via smoking may exceed normal atmospheric inhalation exposure by at least a factor of two in the case of Pb-210 and two orders of magnitude for Po-210 (Spencer et al., 1977).

4. ESTIMATED DOSE FROM DIETARY AND TOBACCO SMOKE EXPOSURES

4.1 DIETARY INGESTION

The range of total exposure estimated from global dietary studies (Table 5) was converted to 50-year dose commitments by use of the dose conversion factors recommended by Dunning et al. (1981). These estimates are presented in Table 8.

A comparison of the estimated maximum doses indicates that adults ingesting an "ordinary," mixed Western-style diet in the U.S.A., U.K., and Federal Republic of Germany would receive between 10-50% the adsorbed radiation to target organs that adults ingesting observed diets in Japan and the Arctic would receive. Nevertheless, even the maximum estimated dietary dose (Po-210 dose to the spleen of Arctic dwellers; 0.23 to 0.58 annual rems) is less than 40% of the annual permissible dose of 1.5 rem to the public for non-blood-forming organs (U.S. DOE, 1981). To date, there are no available estimates of Pb-210 and Po-210 dietary exposures specific to residents of the phosphate-bearing areas of Southwest Florida. It would be particularly useful to compare dietary exposure between populations ingesting foods grown locally on reclaimed or mine-adjacent properties and those ingesting a mixed diet largely derived from non-local sources. It is expected that such exposures would not be comparable to the maximally exposed populations of Japan or the Arctic.

Table 7. Estimated daily inhalation exposure of Pb-210 and Po-210 to smokers

Radionuclide	Estimated exposure (pCi/d)	Reference
Pb-210	0.76	Nikolova and Parfenov, 1971
	2.1	Yermolaeva-Makovskaya, Pertsov and Popov, 1969
	0.52-1.1	Ferri and Christiansen, 1967
	0.12-0.36	Holtzman and Ilcewicz, 1966
	0.15	Spencer et al., 1977
Po-210	16.0	Turner and Radley, 1960
	1.6-7.2	Parfenov, 1974
	8.0	Marsden, 1964
	2.0	Rajewsky and Stahlhoffen, 1966
	1.4	Nikolova and Parfenov, 1971
	1.5	Little and McGandy, 1966
	0.36	Spencer et al., 1977

Table 3. Estimated range of adult 50-year dose commitments (rem) for one year's dietary intake of Pb-210 and Po-210 from natural sources^{a,b}

Target organ	Region							
	U.S.A.	U.K.	India	Federal Republic of Germany	U.S.S.R.	Taiwan	Arctic	
Pb-210								
Bone	9.2E-3 to 3.0E-2	7.7E-3 to 7.7E-2	2.6E-2	3.5E-2	3.1E-2 to 4.9E-2	4.5E-2	1E-1	6.6E-2 to 7.1E-1
Endosteal tissue	4.2E-3 to 1.4E-2	3.5E-3 to 3.5E-2	1.2E-2	1.6E-2	1.4E-2 to 2.2E-2	2.	2E-2	2.7E-2 to 1.4E-1
Red marrow	4.4E-4 to 1.4E-3	3.7E-4 to 3.7E-3	1.2E-3	1.7E-3	1.5E-3 to 2.7E-3	2.	-3	2.1E-3 to 1.5E-2
Kidney	4.1E-4 to 1.3E-3	3.4E-4 to 3.4E-3	1.2E-3	1.6E-3	1.4E-3 to 2.1E-3	2.	-3	1.0E-3 to 1.4E-2
Liver	6.1E-4 to 2.0E-3	5.1E-4 to 5.1E-3	1.7E-3	2.4E-3	2.0E-3 to 3.2E-3	3.0E-3 to 1.2E-2	4.4E-3 to 2.0E-2	
Whole body	7.4E-4 to 2.4E-3	6.2E-4 to 6.2E-3	2.1E-3	2.9E-3	2.5E-3 to 3.9E-3	3.7E-3 to 1.4E-2	5.3E-3 to 2.5E-2	
Po-210^c								
Spleen	9.3E-3	5.8E-3 to 5.8E-2	3.3E-3	2.7E-2	9.9E-3 to 2.4E-2	7.6E-2	2.3E-1 to 5.9E-1	
Kidney	5.4E-3	3.4E-3 to 3.4E-2	5.1E-3	1.6E-2	5.4E-3 to 1.4E-2	4.4E-2	1.4E-1 to 3.4E-1	
Liver	9.3E-4	5.9E-4 to 5.8E-3	3.3E-4	2.7E-3	9.3E-4 to 2.4E-3	7.6E-3	2.3E-2 to 5.9E-2	
Whole body	2.4E-4	1.5E-4 to 1.5E-3	2.2E-4	6.9E-4	2.4E-4 to 6.1E-4	2.0E-3	5.0E-3 to 1.5E-2	

^aUsing dose conversion factors developed by Dunning et al. (1931).

^bUsing range of exposures as presented in Table 5.

^cDirect ingestion only; does not include contribution of internal Pb-210 decay.

When compared to previous estimates for U.S. ingestion of the parent Ra-226 (Watson, Etnier, and McDowell-Boyer, 1983), Pb-210 appears to provide a comparable dose for bone, kidney, liver, and whole body. The upper values of the dietary dose ranges for Pb-210 exceed those for Ra-226 in the kidney and liver by approximately an order of magnitude. The liver doses for dietary Po-210 and Ra-226 are approximately equal; the estimated Po-210 dose in the kidney exceeds that for Ra-226 by an approximate order of magnitude; the whole body Po-210 dose is less than that for Ra-226 by an approximate order of magnitude.

4.2 TOBACCO SMOKE INHALATION

Exposures additional to those received from ingestion and unique to smokers inhaling Pb-210 and Po-210 (summarized in Table 7) were converted to internal dose commitment by applying the dose conversion factors recommended by Eckerman, Ford, and Watson (1981). An activity median aerodynamic diameter (AMAD) of 0.3 μm was assumed. Estimated annual dose commitments are summarized in Table 9.

Tobacco smoking appears to provide a dose equal to or greater than that provided by dietary ingestion for both Pb-210 and Po-210 in bone tissues, liver and kidneys; and for Po-210 in the spleen for the three Western-style diets addressed in Sect. 4.1. The smoking dose estimates are most comparable to those obtained for dietary intake by Arctic dwellers.

Table 9. Estimated range of committed dose equivalent (rem) for one year's inhalation exposure to tobacco smoke containing natural concentrations of Pb-210 and Po-210^{a,b}

Target organ	Dose commitment (rem)
<u>Pb-210</u>	
Bone surface	9.2E-3 to 1.6E-1
Liver	2.6E-3 to 4.5E-2
Kidneys	1.2E-3 to 2.1E-2
Lungs	5.3E-5 to 9.3E-4
<u>Po-210^c</u>	
Spleen	1.1E-2 to 4.8E-1
Kidneys	6.3E-3 to 2.8E-1
Liver	1.1E-3 to 4.9E-2
Lungs	4.9E-4 to 2.2E-2

^aUsing internal dose conversion factors developed by Eckerman, Ford, and Watson, 1981.

^bUsing range of values reported in Table 7.

^cDoes not include contribution of internal Pb-210 decay.

5. CONCLUSIONS AND RECOMMENDATIONS

During the present evaluation, the author has identified a number of research areas that should be addressed in order to obtain a more realistic quantification of food chain transport of Pb-210 and Po-210 than is currently possible in the absence of site-specific data:

- (1) For the majority of internal organs evaluated, the dose resulting from smoking commercially available tobacco products is comparable to or greater than the dose estimates for ingestion of naturally occurring dietary Pb-210 and Po-210.
- (2) No data are readily available to characterize dietary exposures from Pb-210 and Po-210 for the residents of southwestern Florida ingesting locally grown food. These data could be readily collected from market-basket or whole meal surveys of the local populace.
- (3) A principal problem area is the lack of experimentation specifically designed to determine milk and meat transfer coefficients in cattle. Not only should transfer coefficients for the chemical forms of Pb-210 and Po-210 specific to background, historical, and current phosphate mining areas of southwestern Florida be determined; but releases from tailings piles should also be simulated as nearly as possible in feeding studies by offering contaminated feed. In addition, sufficient numbers of animals should be investigated to allow statistical comparison of results.

- (4) The importance of ingested soil and water as a source of contamination for milk and meat has not been critically examined. Data from feeding studies are needed to determine the degree of Pb-210 and Po-210 solubilization from ingested mineral soil in the digestive tract, and the subsequent internal transport.
- (5) For most food items and tobacco, aerosol deposition seems to be the principal mode of Pb-210 and Po-210 entry. This feature is of particular concern for leafy vegetables. As a result, only fruit-bearing crops such as citrus, berries, and cane fruits should be grown on phosphate-reclaimed land.

REFERENCES

Abe, S. and M. Abe., 1969, "Volatility of ^{210}Po (RaF) in airborne dusts at various temperatures," Health Phys. 17(2):340-341.

Athalye, V. V., V. M. Shah, and K. B. Mistry, 1972, "Polonium-210 in cigarette smoke," Indian J. Environ. Health 19(1):54-62.

Baes, C. F., III, R. D. Sharp, A. L. Sjoreen, and R. W. Shor, A Review and Analysis of Parameters for Assessing Transport of Environmentally Released Radionuclides Through Agriculture, ORNL-5786, Oak Ridge National Laboratory, Oak Ridge, Tennessee (in review).

Beasley, T. M. and H. E. Palmer, 1966, "Lead-210 and ^{210}Po in biological samples from Alaska," Science 152:1062-1064.

Berger, K. C., W. H. Erhardt, and C. W. Francis, 1965, "Polonium-210 analyses of vegetables, cured and uncured tobacco and associated soils," Science 150: 1738-1739.

Diack, S. C. and E. W. Bretthauer, 1968, "Polonium-210 in tobacco," Radiat. Data Rep. 9:145-152.

Blanchard, R. L., 1966, "Relationship between ^{210}Po and ^{210}Pb in man and his environment," in Radioecological Concentration Processes (B. Aberg and F. P. Hungate, eds.), pp. 281-296, Pergamon Press, Oxford.

Bogen, D. C., G. A. Welford, and R. S. Morse, 1976, "General population exposure of stable lead and ^{210}Pb to residents of New York City," Health Phys. 30:359-362.

Bovay, E., 1971, "Lead deposition on vegetation growing along highways traveled by automobiles; milk cow feeding experiments with hay contaminated with lead," Bull. Eidgenoess. Gesundh. Beil B., No. 3: 101-117.

Breslin, A. J. and H. Glauberman, 1970, "Investigation of radioactive dust dispersed from uranium mill tailings piles," in Environmental Surveillance in the Vicinity of Nuclear Facilities (W. C. Reinig, ed.), pp. 249-253, Charles C. Thomas Publisher, Springfield, Illinois.

Carfi, N. and R. D. Lonati, 1966, "Polonium-210 in Italian tobacco," in Proc. First Inter'l Cong. Rad. Prot., Vol. 2 (W. S. Snyder, H. H. Abel, L. K. Burton, R. Maushart, A. Benco, F. Duhamel, and B. M. Wheatley, eds.), pp. 1097-1098, Pergamon Press, Oxford.

Cohen, N., T. Jaakkola, and M. E. Wrenn, 1973, "Lead-210 concentrations in the bone, blood and excreta of a former uranium miner," Health Phys. 24:601-609.

Cox, W. J. and D. W. Rains, 1972, "Effect of lime on lead uptake by five plant species," J. Environ. Qual. 1(2):167-169.

Dedolph, R., G. Ter Haar, R. Holtzman and H. Lucas, Jr., 1970, "Survey of lead in perennial ryegrass and radishes," Environ. Sci. Tech. 4(3):217-225.

Djuric, D., Z. Kerin, L. Graovac-Leposavić, L. Novak, and M. Kop, 1971, "Environmental contamination by lead from a mine and smelter," Arch. Environ. Health 23:275-279.

Donovan, P. P., D. T. Feeley, and P. O. Canavan, 1969, "Lead contamination in mining areas in Western Ireland II. Survey of animals, pastures, foods and waters," J. Sci. Food Agric. 20:43-45.

Dos Santos, D. L., 1970, Rev. Biol. Med. Nucl. 2:73 (as cited in Parfenov, Y. D., 1974).

Dunning, D. E., Jr., G. G. Killough, S. R. Bernard, J. C. Pleasant, and P. J. Walsh, 1981, Estimates of Internal Dose Equivalent to 22 Target Organs for Radionuclides Occurring in Routine Releases From Nuclear Fuel-Cycle Facilities, Vol. III, NUREG/CR-0150, Vol. 3, ORNL/NUREG/TM-190/V3, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Eckerman, K. F., M. R. Ford, and S. B. Watson, 1981, Internal Dosimetry Data and Methods of ICRP-Part 2, Vol. 1: Committed Dose Equivalent and Secondary Limits, NUREG/CR-1962, Vol. 1., ORNL/NUREG/TM-433/V1, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Eisenbud, M. and H. G. Petrow, 1964, "Radioactivity in the atmospheric effluents of power plants that use fossil fuels," Science 144:288-289.

Ferri, E. S. and E. J. Baratta, 1966, "Polonium-210 in tobacco, cigarette smoke, and selected human organs," Public Health Rep. 81:121-127.

Ferri, E. S. and H. Christiansen, 1967, "Lead-210 in tobacco and cigarette smoke," Public Health Rep. 82:828-832.

Fleischer, R. L. and F. P. Parungo, 1974, "Aerosol particles on tobacco trichomes," Nature 250:158-159.

Francis, C. W., G. Chesters, and W. H. Erhardt, 1968, "Polonium-210 entry into plants," Environ. Sci. Tech. 2:690-695.

Gesell, T. F. and H. F. Richard, 1975, "The technologically enhanced natural radiation environment," Health Phys. 28:361-366.

Glöbel, B., H. Muth, and E. Oberhausen, 1966, "Uptake and elimination of the natural radionuclides ^{210}Pb and ^{210}Po by humans," Strahlentherapie 133:218-226.

Hammond, P. B. and A. L. Aronson, 1964, "Lead poisoning in cattle and horses in the vicinity of a smelter," Ann. N.Y. Acad. Sci. 111:595-611.

Hamrick, P. A. and P. J. Walsh, 1974, "Environmental radioactivity and the lung," Env. Health Perspec. 9:33-52.

Hill, C. R., 1960, "Lead-210 and ^{210}Po in grass," Nature 187:211-212.

Hill, C. R., 1965, "Polonium-210 in man," Nature 208:423-428.

Hill, C. R., 1967, "Routes of uptake of ^{210}Po into human tissues," in Radioecological Concentration Processes (B. Aberg and F. P. Hungate, eds.), pp. 297-302, Pergamon Press, Oxford.

Holtzman, R. B., 1963, "Measurement of the natural contents of RaD (^{210}Pb) and RaF (^{210}Po) in human bone - estimates of whole-body burdens," Health Phys. 9:385-400.

Holtzman, R. B., 1966(a), "Radium-226 and the natural airborne nuclides ^{210}Pb and ^{210}Po in Arctic biota," in Proc. First Int'l Cong. Rad. Prot., Vol. 2 (W. S. Snyder, H. H. Abel, L. K. Burton, R. Maushart, A. Benco, F. Duhamel, and B. M. Wheatley, eds.), pp. 1087-1096, Pergamon Press, Oxford.

Holtzman, R. B., 1966(b), "Natural levels of ^{210}Pb , ^{210}Po and ^{226}Ra in humans and biota of the Arctic," Nature 210:1094-1097.

Holtzman, R. B. and F. H. Ilcewicz, 1966, "Lead-210 and ^{210}Po in tissues of cigarette smokers," Science 153:1259-1260.

Holtzman, R. B. and F. H. Ilcewicz, 1971, "Renal excretion rates of ^{137}Cs , ^{210}Pb , ^{210}Po and stable lead by northern Canadians," in Argonne National Laboratory, Radiological Physics Division, Annual Report, ANL-7860 (Pt II), Argonne National Laboratory, Argonne, Illinois.

Holtzman, R. B., H. Spencer, F. H. Ilcewicz, and L. Kramer, 1974, "Metabolic balances of ^{210}Pb and ^{210}Po in unexposed men," in Proc. Third Int'l Rad. Prot. Cong. (W. S. Snyder, ed.), CONF-730907, pp. 1406-1411, NTIS, Springfield, Virginia.

Holtzman, R. B., P. W. Urnezis, A. Padova, and C. M. Bobula III, 1979, Contamination of the Human Food Chain by Uranium Mill Tailings Piles, NUREG/CR-0758, ANL/ES-69, Argonne National Laboratory, Argonne, Illinois.

Holtzman, R. B., 1980, "Normal dietary levels of radium-226, radium-228, lead-210 and polonium-210 for man," in Natural Radiation Environment III, Vol. 1 (T. F. Gesell and W. M. Lowder, eds.), CONF-780422, pp. 755-782, Technical Information Center, U.S. DOE, Oak Ridge, Tennessee.

Iyengar, M.A.R., M. P. Rajan, S. Ganapathy, and P. R. Kamath, 1980, "Sources of natural radiation exposure in a low-monazite environment," in Natural Radiation Environment III, Vol. 2 (T. F. Gesell and W. M. Lowder, eds.), CONF-780422, pp. 1090-1106, Technical Information Center, U.S.DOE, Oak Ridge, Tennessee.

Jaworowski, Z., J. Bilkiewicz, and E. Zylicz, 1971, "Radium-226 in contemporary and fossil snow," Health Phys. 20:449-450.

Jaworowski, Z. and L. Kownacka, 1976, "Lead and radium in the lower stratosphere," Nature 263:303-304.

Jaworowski, Z. and D. Grzybowska, 1977, "Natural radionuclides in industrial and rural soils," Sci. Total Environ. 7:45-52.

John, M. K. and C. J. Van Laerhoven, 1972(a), "Lead distribution in plants grown on contaminated soil," Environ. Lett. 3(2):111-116.

John, M. K. and C. J. Van Laerhoven, 1972(b), "Lead uptake by lettuce and oats as affected by lime, nitrogen and sources of lead," J. Environ. Qual. 1(2):169-171.

John, M. K., 1972, "Lead availability related to soil properties and extractable lead," J. Environ. Qual. 1(3):295-298.

Johnson, J. E. and R. L. Watters, 1972, PoO₂-210 Metabolism in Ruminants, Final Report and Summary, 1969-1972 to U.S. AEC, COO-2044-5, U.S. AEC, Chicago Operations Office, Argonne, Illinois.

Kametani, K., H. Ikebuchi, T. Matsumara, and H. Kawakami, 1981, "Radium-226 and ²¹⁰Pb measurements in foodstuffs," Radioisotopes 30(12):681-683.

Kaurenens, P. and J. K. Miettinen, 1969, "Polonium-210 and ²¹⁰Pb in the Arctic food chain and the natural radiation exposure of Lapps," Health Phys. 16:287-295.

Kawans, M. and S. Nakatani, 1964, The Natural Radiation Environment, pp. 291-312, University of Chicago Press, Chicago, Illinois (as cited in Francis, C. W., G. Chesters, and W. H. Erhardt, 1968).

Kerin, D. and Z. Kerin, 1971, "Lead contamination of milk and honey caused by industrial lead aerosols," Prot. Vitae 16(2):61-62.

Keslev, D., E. Novakova, A. Boyadzhiev, and A. Kerteva, 1975, "Contents of ²¹⁰Po in food products of Bulgaria," Nauchni Tr. Nauchnoizsled Inst. Radiobiol. Radiats. Khig. 5:193-199.

Khandekar, R. N., 1977, "Polonium-210 in Bombay diet," Health Phys. 33(2):148-150.

Kilibarda, M., D. Petrovic, D. Panov, and D. Djuric, 1966, "Contamination with ^{210}Po , uranium and ^{226}Ra due to smoking," Health Phys. 12:1808.

Kocher, D. C., 1977, Nuclear Decay Data for Radionuclides Occurring in Routine Releases from Light-Water Cooled Nuclear Power Reactors, ORNL/NUREG/TM-102, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Koeppe, D. E., 1977, "The uptake, distribution and effect of cadmium and lead in plants," Sci. Total Environ. 7:197-206.

Ladinskaya, L. A., Y. D. Parfenov, D. K. Popov, and A. V. Fedorova, 1973, "Lead-210 and ^{210}Po content in air, water, foodstuffs and the human body," Arch. Environ. Health 27:254-258.

Lalit, B. Y., T. V. Ramachandran, and S. Rajan, 1980, "Lead-210 content of food samples in India," Radiat. Environ. Biophys. 18(1):13-18.

Little, J. B. and R. B. McGandy, 1966, "Measurement of ^{210}Po in human blood," Nature 211:842-843.

MacLean, A. J., R. L. Halstead, and B. J. Finn, 1969, "Extractability of added lead in soils and its concentrations in plants," Can. J. Soil Sci. 49:327-334.

McDowell-Boyer, L. M., A. P. Watson, and C. C. Travis, 1980, "A review of parameters describing terrestrial food chain transport of ^{210}Pb and ^{226}Ra ," Nucl. Safety 21(4):486-495.

McDowell-Boyer, L. M. and C. F. Baes III, 1980, "Terrestrial food chain transport," in Recommendations Concerning Models and Parameters Best Suited to Breeder Reactor Environmental Radiological Assessments, ORNL-5529, pp. 23-54, NTIS, Springfield, Virginia.

Magno, P. J., P. R. Groulx, and J. C. Apidianakis, 1970, "Lead-210 in air and total diets in the United States during 1966," Health Phys. 18:383-388.

Marsden, E., 1964, "Incidence and possible significance of inhaled or ingested polonium," Nature 203:230-233.

Martell, E. A., 1974, "Radioactivity of tobacco trichomes and insoluble cigarette smoke particles," Nature 249:215-217.

Menzel, R. G., 1965, "Soil-plant relationships of radioactive elements," Health Phys. 11:1325-1332.

Michelson, I., 1961, "Radium-226 in the total diet," Health Phys. 5:229-230.

Michelson, I., J. C. Thompson, Jr., B. W. Hess, and C. L. Comar, 1962, "Radioactivity in total diet," J. Nutrition 78:371-383.

Milosević, Z., E. Horšić, R. Kljajic, and A. Bauman, 1980, "Distribution of uranium, ^{226}Ra , ^{210}Pb and ^{210}Po in the ecological cycle in mountain regions of central Yugoslavia," in Proc. Fifth Int'l Conf. Int'l Rad. Prot. Assoc., Vol. 2, INIS-MF-5876, pp. 1123-1126, Pergamon Press, Oxford.

Moore, H. E. and S. E. Poet, 1976, "Background levels of ^{226}Ra in the lower troposphere," Atmos. Environ. 10:381-383.

Moore, H. E., E. A. Martell, and S. E. Poet, 1976, "Sources of ^{210}Po in the atmosphere," Environ. Sci. Tech. 10(6):586-591.

Morse, R. S. and G. A. Welford, 1971, "Dietary intake of ^{210}Pb ," Health Phys. 21:53-55.

Nelmes, A. J., R. St. J. Buxton, F. A. Fairweather, and J. E. Martin, 1974, "The implication of the transfer of trace metals from sewage sludge to man," in Trace Substances in Environmental Health, VIII (D. D. Hemphill, ed.), pp. 145-153, University of Missouri Press, Columbia, Missouri.

Ng, Y., 1982, "A review of transfer factors for assessing the dose from radionuclides in agricultural products," Nucl. Safety 23(1):57-71.

Nikolova, M. E. and Y. D. Parfenov, 1971, Rentgenol. Radiol. 10:163 (as cited in Parfenov, Y. D., 1974).

Okabayashi, H., M. Suzuki-Yasumoto, S. Hongo, and S. Watanabe, 1975, "On the evaluation of ²¹⁰Po bioassay for uranium mine workers in Japan for the personal exposure index to radon daughters," J. Radiat. Res. 16:142-151.

Parfenov, Y. D., 1974, "Polonium-210 in the environment and in the human organism," Atomic Energy Rev. 12:75-143.

Rabinowitz, M., 1972, "Plant uptake of soil and atmospheric lead in southern California," Chemosphere 1(4):175-180.

Radford, E. P., Jr. and V. R. Hunt, 1964, "Polonium-210: A volatile radioelement in cigarettes," Science 143:247-249.

Radford, E. P., Jr., 1974, "Polonium-210 alpha radiation as a cancer initiator in tobacco smoke," Proc. Fifth Int'l Cong. on Rad. Res., July 14-20, Seattle, Washington.

Rains, D. W., 1971, "Lead accumulation by wild oats (Avena fatua) in a contaminated area," Nature 233:210-211.

Rajewsky, B. and W. Stahlhofen, 1966, "Polonium-210 activity in the lungs of cigarette smokers," Nature 209:1312-1313.

Ramzaev, P. V., et al., 1974, Absorption of ^{210}Pb and ^{210}Po in the Gastrointestinal Tract of Rats and Man, UNSCEAR Document A/AC.82/G/1.1467, Pamphlet U.S.S.R. State Comm. on the Use of Atomic Energy. in USAEC Health and Safety Laboratory Translation, July, 1974.

Rayno, D. R., M. H. Momeni, and C. Sabau, 1980, "Forage uptake of uranium series radionuclides in the vicinity of the Anaconda uranium mill," in Uranium Mill Tailings Management, pp. 57-66, Proc. of the Third Symposium, November 24-25, 1980, Civil Engineering Department, Colorado State University, Fort Collins, Colorado.

Solnicka, H. and P. Bischof, 1970, Cas. Lek. Cesk. 109:28 (as cited in Parfenov, Y. D., 1974).

Spector, W. S. (Ed.), 1956, Handbook of Biological Data, Committee on the Handbook of Biological Data, Division of Biology and Agriculture, The National Academy of Sciences, The National Research Council, W. B. Saunders Co., Philadelphia and London.

Spencer, H., R. B. Holtzman, L. Kramer, and F. H. Ilcewicz, 1977, "Metabolic balances of ^{210}Pb and ^{210}Po at natural levels," Radiat. Res. 69(1):166-184.

Stanley, R. E., A. A. Mullen, and E. W. Bretthauer, 1971, "Transfer to milk of ingested radiolead," Health Phys. 21:211-215.

Suzuki, M., et al., 1967, Health Phys. Res. Abst. IAEA I:71 (as cited in Parfenov, Y.D., 1974).

Takata, N., H. Watanabe, and R. Ichikawa, 1968, "Lead-210 content in foodstuffs and its dietary intake in Japan," J. Rad. Res. 9:29-34.

Ter Haar, G., 1970, "Air as a source of lead in edible crops," Environ. Sci. Tech. 4(3):226-230.

Tracy, B. L., F. A. Prantl, and J. M. Quinn, 1983, "Transfer of ^{226}Ra , ^{210}Pb and uranium from soil to garden produce: assessment of risk," Health Phys. 44(5):469-477.

Travis, C. C., A. P. Watson, L. M. McDowell-Boyer, S. J. Cotter, M. L. Randolph, and D. E. Fields, 1979, A Radiological Assessment of ^{222}Rn Released from Uranium Mills and Other Natural and Technologically Enhanced Sources, NUREG/CR-0573, ORNL/NUREG-55, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Turner, R. C. and J. M. Radley, 1960, "Naturally occurring alpha activity of cigarette tobacco," Lancet 1:1197-1198.

U. S. Department of Energy (U. S. DOE), 1981, "Requirements for radiation protection," Chap. 11 in Environmental Protection, Safety and Health Protection Program for DOE Operations, U. S. DOE Order No. 54301, August 13, 1981.

Watson, A. P., E. L. Etnier, and L. M. McDowell-Boyer, 1983, Radium-226 in Drinking Water and Terrestrial Food Chains: A Review of Parameters and an Estimate of Potential Exposure and Dose, ORNL/TM-8597, Oak Ridge National Laboratory, Oak Ridge, Tennessee.

Watters, R. L. and J. F. McInroy, 1969, "The transfer of ^{210}Po from cattle feed to milk," Health Phys. 16:221-225.

Watters, R. L., J. E. Johnson, and W. R. Hansen, 1969, A Study of Unsup-ported ^{210}Po for Ion Exchange in Soil and Uptake in Vegetation, Second Technical Progress Report to U. S. AEC, Department of Radi-
ology and Radiation Bot., Colorado State University, Fort Collins, Colorado.

Watters, R. L., J. E. Johnson, and J. F. McInroy, 1971, $^{210}\text{PoO}_2$ Metabolism in Ruminants, Second Technical Progress Report, January 15, 1970 to January 25, 1971, COO-2044-3, U. S. AEC, Chicago Operations Office, Argonne, Illinois.

Yermolayeva-Markovskaya, A. P., L. A. Pertsov, and D. K. Popov, 1969, Polonium-210 in the Human Body and in the Environment, UNSCEAR Document A/AC.82/G./L.1260, pp. 163-170, in U. S. AEC Health and Safety Laboratory Translation, February, 1969.

Zimdahl, R. L. and J. M. Foster, 1976, "The influence of applied phosphorus, manure, or lime on uptake of lead from soil," J. Environ. Qual. 5:31-34.

INTERNAL DISTRIBUTION

1. C. F. Baes III	12. L. E. Stratton
2. B. A. Berven	13. T. Tamura
3. R. O. Chester	14. C. C. Travis
4. S. J. Cotter	15. P. J. Walsh
5. K. F. Eckerman	16-20. A. P. Watson
6. C. W. Gehrs	21. J. P. Witherspoon
7. S. G. Hildebrand	22-23. TA Section Library
8. G. S. Hill, Jr.	24-25. Central Research Library
9. S. V. Kaye	26-27. Laboratory Records, Dept.
10. F. F. Knapp, Jr.	28. Laboratory Records, ORNL-RC
11. P. S. Rohwer	29. ORNL Y-12 Technical Library
	30. ORNL Patent Office

EXTERNAL DISTRIBUTION

- 31-130. Dr. David P. Borris, Florida Institute of Phosphate Research, 1855 West Main Street, Bartow, Florida 33830.
- 131. Office of Assistant Manager for Energy Research and Development, Department of Energy, Oak Ridge Operations Office, Oak Ridge, Tennessee 37830.
- 132-158. Technical Information Center, Department of Energy, Oak Ridge, Tennessee 37830.