

LA--10929-MS

DE87 009465

Groundwater Chemistry at Yucca Mountain, Nevada, and Vicinity

Jerry F. Kerrisk

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

Los Alamos Los Alamos National Laboratory
Los Alamos, New Mexico 87545

CONTENTS

ABSTRACT.....	1
I. INTRODUCTION.....	2
II. PREVIOUS WORK.....	6
III. YUCCA MOUNTAIN WELLS.....	9
IV. SOURCES OF GROUNDWATER.....	11
V. ACTIVE CHEMICAL AND PHYSICAL PROCESSES.....	12
A. Transport with Water.....	12
B. Rock-Water Reactions.....	12
C. Ion Exchange.....	12
D. Gas Dissolution.....	13
E. Mixing of Different Water Compositions.....	13
F. Evaporation.....	13
VI. IMPORTANT GROUNDWATER COMPOSITIONAL VARIABLES AND RELATIONS AMONG THESE VARIABLES.....	14
A. Major Species and Variables.....	14
B. Isotopic Data.....	32
C. Other Species and Variables.....	41
D. Relations with Location or Depth.....	46
VII. CONTROLS ON GROUNDWATER COMPOSITION.....	52
A. Control of Cation Content, Aqueous Silica Content, and pH of Tuffaceous Water.....	52
B. Control of Carbonate Content of Tuffaceous Water.....	55
C. Control of Chloride and Sulfate Content of Tuffaceous Water.....	64
D. Control of the Redox State of the Water.....	65
E. Control of Other Compositional Variables of Tuffaceous Water.....	66
F. Controls on Unsaturated-Zone Water Composition.....	67
VIII. IMPLICATIONS FOR RADIONUCLIDE TRANSPORT.....	68
ACKNOWLEDGEMENTS.....	71
REFERENCES.....	72
APPENDIX A: WATER CHEMISTRY DATA.....	77
APPENDIX B: IMPORTANCE OF RADIONUCLIDE TRANSPORT BY PARTICULATES ENTRAINED IN FLOWING GROUNDWATERS.....	114

TABLES

I.	Wells at Yucca Mountain and Vicinity.....	10
II.	Water Sampling Locations.....	16
III.	Variation of Water Properties With Depth.....	50
IV.	Carbon Model Parameters Used in Figures 26 and 27.....	61

FIGURES

1. Area map of Yucca Mountain and vicinity with well locations.....	3
2. Sodium content as a function of total carbonate content.	
(a) All sampling locations (see Table II for meaning of legend abbreviations).....	15
(b) Yucca Mountain and vicinity.....	15
3. Calcium content as a function of total carbonate content.	
(a) All sampling locations (see Table II for meaning of legend abbreviations).....	18
(b) Yucca Mountain and vicinity.....	18
4. Ternary diagram of relative sodium, calcium, and potassium content.	
(a) All sampling locations (see Table II for meaning of legend abbreviations).....	19
(b) Yucca Mountain and vicinity.....	19
5. $\text{Na}/(\text{Na}+\text{Ca}+\text{K})$ as a function of pH at Yucca Mountain and vicinity. Solid lines represent calculated results from Kerrick (1983).....	21
6. The \log_{10} (calcium content) as a function of pH at Yucca Mountain and vicinity.....	21
7. The pH as a function of total carbonate content at Yucca Mountain and vicinity.....	22
8. The \log_{10} (carbon dioxide pressure) as a function of pH.	
(a) All sampling locations (see Table II for meaning of legend abbreviations).....	23
(b) Yucca Mountain and vicinity.....	23
9. The \log_{10} (carbon dioxide pressure) as a function of total carbonate content at Yucca Mountain and vicinity.....	25
10. Calcite $\log_{10}(Q/K)$ as a function of pH at Yucca Mountain and vicinity.....	25
11. The (chloride + sulfate) content as a function of total carbonate content.	
(a) All sampling locations (see Table II for meaning of legend abbreviations).....	27
(b) Yucca Mountain and vicinity.....	27
12. Sulfate and chloride contents as a function of total carbonate content at Yucca Mountain and vicinity.	
(a) Sulfate content as a function of total carbonate.....	29
(b) Chloride content as a function of total carbonate.....	29

13. Sulfate content as a function of chloride content (see Table II for meaning of legend abbreviations).	30
(a) All sampling locations.....	30
(b) Expanded plot of region near the origin.....	30
14. Percentage of modern carbon as a function of total carbonate content.	
(a) All sampling locations (see Table II for meaning of legend abbreviations).....	33
(b) Yucca Mountain and vicinity.....	33
15. $\delta^{13}\text{C}$ as a function of total carbonate content.	
(a) All sampling locations (see Table II for meaning of legend abbreviations).....	35
(b) Yucca Mountain and vicinity.....	35
16. The pH as a function of the percentage of modern carbon and $\delta^{13}\text{C}$ at Yucca Mountain and vicinity.	
(a) pH as a function of the percentage of modern carbon.....	37
(b) pH as a function of $\delta^{13}\text{C}$	37
17. δD as a function of $\delta^{18}\text{O}$.	
(a) All sampling locations (see Table II for meaning of legend abbreviations).....	38
(b) Yucca Mountain and vicinity.....	38
18. δD as a function of the percentage of modern carbon and total carbonate content at Yucca Mountain and vicinity.	
(a) δD as a function of the percentage of modern carbon.....	40
(b) δD as a function of total carbonate content.....	40
19. Nitrate content as a function of the percentage of modern carbon and total carbonate content at Yucca Mountain and vicinity.	
(a) Nitrate content as a function of the percentage of modern carbon.....	42
(b) Nitrate content as a function of total carbonate content.....	42
20. Fluoride content as a function of sodium content and total carbonate at Yucca Mountain and vicinity.	
(a) Fluoride content as a function of sodium content.....	44
(b) Fluoride content as a function of total carbonate content.....	44
21. Water temperature as a function of the percentage of modern carbon and total carbonate content at Yucca Mountain and vicinity.	
(a) Water temperature as a function of the percentage of modern carbon.....	47
(b) Water temperature as a function of total carbonate.....	47
22. $\text{Na}/(\text{Na}+\text{Ca}+\text{K})$ as a function of location for wells near Yucca Mountain.....	48

23. Calcium content as a function of sodium content at Yucca Mountain and vicinity.....	48
24. Total carbonate content as a function of location for wells near Yucca Mountain.....	49
25. Ternary diagrams of relative sodium, calcium, and potassium content at Rainier Mesa. (a) Surface water and pore water (Benson 1976).....	53
(b) Surface water and fracture water (Henne 1982).....	53
26. Percentage of modern carbon as a function of total carbonate content at Yucca Mountain with carbon-model prediction (see Table IV for solid and dashed curve definitions).....	60
27. $\delta^{13}\text{C}$ as a function of total carbonate content at Yucca Mountain with carbon-model prediction (see Table IV for solid and dashed curve definitions).....	60
28. δD as a function of apparent age and carbon-model age at Yucca Mountain. (a) δD as a function of apparent age.....	63
(b) δD as a function of carbon-model age.....	63

GROUNDWATER CHEMISTRY AT YUCCA MOUNTAIN,
NEVADA, AND VICINITY

by

Jerry F. Kerrisk

ABSTRACT

The chemistry of groundwater at Yucca Mountain and vicinity has been reviewed and compared with the chemistry of water from the Nevada Test Site and surrounding areas such as the Amargosa Desert and Oasis Valley. Sodium is the primary cation and carbonate is the primary anion in water from the saturated zone of the tuffaceous aquifer at Yucca Mountain. Other major cations present are calcium, potassium, and magnesium; other major anions are sulfate and chloride, with lesser quantities of fluoride and nitrate. Aqueous silica is also present. The primary purpose of this review was to survey water-composition data and look for relations among the compositional variables that could provide insight into the processes that control the composition and would ultimately affect radionuclide transport. The following conclusions were inferred from the review. Major cation concentrations are controlled by rock dissolution and mineral precipitation reactions as well as by cation exchange with existing minerals. Aqueous carbonate initially comes from atmospheric and soil-zone carbon dioxide, but there is evidence at Yucca Mountain that carbon dioxide in the gas phase of the unsaturated zone supplies additional carbonate to saturated-zone water in the tuffaceous aquifer as mineral dissolution and precipitation reactions raise the pH of the water. This combination is effectively mineral dissolution and precipitation in a system that is open with respect to carbon dioxide. A carbon model for this process is discussed; one conclusion of the model is that the true age of water that has obtained significant amounts of carbonate from the gas phase of the unsaturated zone is older than its apparent age. The primary source of aqueous chloride and sulfate

is probably from precipitation; there does not appear to be any mineralogical controls on these species at Yucca Mountain. There is some evidence that water in the deep saturated zone at Yucca Mountain may be reducing; this conclusion would be difficult to prove with existing data. Water near the static-water level is oxidizing. Water in the western part of Yucca Mountain is lower in calcium than water to the east. Carbonate and fluoride tend to be more concentrated in the water in the southwestern part of Yucca Mountain. The implications of these conclusions for radionuclide transport primarily reflect the range of water compositions that could be encountered at Yucca Mountain. Oxidizing conditions, and carbonate and fluoride complexes, will have the largest impact on solubility and speciation. There are not enough data available at this time to discuss the processes that control unsaturated-zone water compositions at Yucca Mountain.

I. INTRODUCTION

The Nevada Nuclear Waste Storage Investigations (NNWSI) Project is studying a site at Yucca Mountain in southern Nevada as a potential nuclear waste repository. The site is located on the southwestern edge of the Nevada Test Site (NTS) and on adjacent US Bureau of Land Management land and land controlled by the US Air Force (see Fig. 1). The Topopah Spring Member tuff in the unsaturated zone of Yucca Mountain has been selected as the candidate rock unit for the repository. Waste-element transport in water that moves through the waste-storage area and toward the environment is considered the most likely mechanism for release of most waste elements. To evaluate the importance of this mechanism, site-characterization information is required about water flow paths, water fluxes, and water chemistry. This report discusses water chemistry at Yucca Mountain and in nearby areas. Water chemistry is an important parameter for waste-element transport because the solubility, speciation, and sorption of waste elements on local minerals all depend on water chemistry. Water chemistry can also affect the composition and stability of the minerals in contact with the water. Questions concerning water fluxes and flow paths are being addressed by the US Geological Survey (USGS) in another part of the NNWSI Project (Montazer and Wilson 1984; Waddell et al. 1984).




Fig. 1. Area map of Yucca Mountain and vicinity with well locations.

The Yucca Mountain site lies within the southwest Nevada volcanic field in the southern Great Basin and is a remnant of the Timber Mountain-Oasis Valley caldera complex. Thick deposits of ash-flow tuffs overlying Paleozoic carbonate deposits predominate at Yucca Mountain. Alluvium fills some of the washes on Yucca Mountain as well as adjacent basins such as Jackass Flats and Crater Flat. The unsaturated zone is quite thick at Yucca Mountain (about 500 to 750 m). The repository would be located 300 m or more below the land surface in the densely welded, devitrified tuff of the Topopah Spring Member. In addition to the lower portions of the Topopah Spring tuff, tuffs of the Calico Hills and the Prow Pass Member of the Crater Flat tuff are unsaturated below the repository location. Two additional units of the Crater Flat tuff (the Bullfrog and Tram Members) are in the saturated zone below Yucca Mountain. To the east of Yucca Mountain, all the tuff units mentioned above are partially or completely below the static water level. A summary of the mineralogy of Yucca Mountain has recently been published (Bish and Vaniman 1985). Farther from Yucca Mountain, the sequence of alluvium and tuff overlying Paleozoic carbonates persists in areas such as Jackass Flats, Yucca Flat, and Rainier Mesa; some exposure of Paleozoic carbonates also occurs. Static water levels in alluvium, tuff, or carbonates have been found (Claassen 1973).

At Yucca Mountain and vicinity, sodium is the primary cation and carbonate (as H_2CO_3 , HCO_3^- , and CO_3^{--}) is the primary anion in water from the saturated zone (Benson et al. 1983; Ogard and Kerrisk 1984; Benson and McKinley 1985). Other major cations present are calcium, potassium, and magnesium; other major anions are sulfate and chloride, with lesser quantities of fluoride and nitrate. Sodium and total carbonate concentrations are generally in the range of 2 to 10 mmoles/l. Aqueous silica is also present at a concentration of about 1 mmoles/l. Most waters have a pH in the 7 to 8 range and a temperature in the 25 to 40 °C range. In the wider area covering NTS, Oasis Valley, and the Amargosa Desert, the relative proportions of dissolved species is generally similar to water from Yucca Mountain; waters in the higher range of total concentrations of dissolved species are more often found, however, particularly in areas where evaporation is important. Examples also exist of different water compositions in specialized environments.

Current USGS models indicate that, at Yucca Mountain, water would move generally downward (with some possibility of lateral transport) through the unsaturated zone and into groundwater in the saturated zone (Montazer and Wilson 1984). Thus, water chemistry in both the unsaturated and saturated zones is of interest. To date, numerous wells have sampled water from the saturated zone at Yucca Mountain, and other studies have examined water chemistry in nearby areas such as Oasis Valley, the Amargosa Desert, Pahute Mesa, and Rainier Mesa. Thus, information is available about saturated-zone water chemistry at Yucca Mountain and its relation to the chemistry of water in adjacent areas. However, no analyses of pore water from the unsaturated zone at Yucca Mountain have been reported. (Gas analyses from the unsaturated zone at Yucca Mountain have been published.) Some pore-water compositions from Rainier Mesa have been reported and can be used as a guide until data from Yucca Mountain are available. Because the distribution of available data is weighted heavily toward the saturated zone, this report will necessarily concentrate on chemistry of saturated-zone water. The importance of the chemistry of unsaturated-zone water at Yucca Mountain is recognized, however, and when sufficient data are available, a study emphasizing the unsaturated zone is planned.

The phrase "water chemistry" can have a variety of meanings. For this report, water chemistry relates to the identities and concentrations of dissolved species present in the water (including isotopic data where available), the origins of these species, and the chemical controls on their concentrations. An understanding of water chemistry at Yucca Mountain starts with compilations of local water compositions and mineralogy. However, Yucca Mountain cannot be viewed in isolation, so that similar data from surrounding areas are also useful. The similarity of water compositions and mineralogy in a larger area containing Yucca Mountain makes an area-wide comparison even more valuable. An understanding of the origins of dissolved species and chemical controls on their concentrations leads to knowledge of the chemical and physical processes that affect water compositions. This knowledge is important because Yucca Mountain is not a static system; future climatic changes and even the presence of a repository can change some of the parameters that control water composition, such as recharge rate, recharge mechanism, temperature, or materials that contact the water. Thus, in order to assess the effects of changes in boundary conditions, an understanding is

necessary of the origins of dissolved species in the water and the processes that control their concentrations. For quantitative predictions, a quantitative model of water chemistry is required. At this time, however, only a qualitative understanding is available of most aspects of Yucca Mountain water chemistry.

The following sections of this report (1) describe previous reports of water chemistry at Yucca Mountain and nearby areas, (2) discuss the wells at Yucca Mountain and vicinity that are the primary focus of this report, (3) discuss some possible sources of groundwater at Yucca Mountain, (4) outline chemical and physical processes that can affect water compositions in this area, (5) present data for important compositional variables of water from the area and some relations among the compositional variables, (6) discuss controls that appear to exist on water chemistry at Yucca Mountain, and (7) discuss the implications of the water chemistry for solubility and speciation of waste elements and for mineral stability.

II. PREVIOUS WORK

Water from Yucca Mountain and many nearby areas has been sampled and analyzed. The operation of NTS led to test wells and producing wells in areas such as Yucca Flat, Frenchman Flat, Jackass Flats, and Pahute Mesa. In addition, springs and seeps have been sampled in some of these same areas as well as at Rainier Mesa. The compositions of water from areas near NTS have also been reported; these include Oasis Valley to the west of Yucca Mountain and NTS, and the Amargosa Desert to the south of NTS. In recent years, a number of test wells have been dug and sampled at Yucca Mountain as part of the NNWSI Project. Data from all of these locations provide a base for the analyses done in this report.

The sources of water-composition data that were reviewed and included in this investigation of groundwater chemistry are listed below along with a few words about the water sampled.

1. Clebsch and Barker (1960), composition of water from wells and springs at Rainier Mesa,
2. Clebsch (1961), tritium analyses of water from NTS,
3. Blankenagel and Weir (1973), composition of water from wells at Pahute Mesa and vicinity,

4. Claassen (1973), composition of water from various water-supply wells around NTS,
5. Benson (1976), composition of pore water from Rainier Mesa,
6. White (1979), composition of water from Oasis Valley,
7. White et al. (1980), composition of fracture water from Rainier Mesa,
8. Henne (1982), composition of fracture water from Rainier Mesa,
9. Benson et al. (1983), composition of water from wells in the vicinity of Yucca Mountain,
10. Ogard and Kerrisk (1984), composition of water from wells in the vicinity of Yucca Mountain,
11. Benson and McKinley (1985), composition of water from wells in the vicinity of Yucca Mountain,
12. Claassen (1985), composition of water from wells and springs in the Amargosa Desert and at other locations at NTS and vicinity, and
13. White and Chuma (1986), composition of water from Oasis Valley and Pahute Mesa.

A tabulation of the various water compositions is given in Appendix A. In addition to the water-composition data outlined above, analyses of the gas phase from the unsaturated zone at Yucca Mountain have been reported by Yang et al. (1985). These data were taken from near-surface locations to about 368-m depth; they provide direct information about the gas phase contacting pore water in the unsaturated zone at Yucca Mountain, but only indirect information about the pore water.

A general review of groundwater chemistry in the south-central Great Basin was given by Winograd and Thordarson (1975). They identified five types of groundwater in the area:

1. a calcium-magnesium-bicarbonate water that moves only through the carbonate aquifer or through a valley-fill aquifer that is rich in carbonate detritus,
2. a sodium-potassium-bicarbonate water that moves only through the tuff or lava-flow terrane, or through a valley-fill aquifer that is rich in volcanic detritus,
3. a mixture of the first two types that results from downward flow of the sodium-potassium-bicarbonate water into the carbonate aquifer,
4. a sodium-sulfate-bicarbonate water in east-central Death Valley, and

5. a playa water that exists as shallow groundwater beneath saturated playas.

Winograd and Thordarson (1975) infer, on the basis of hydrochemical data, that groundwater beneath NTS moves toward the Ash Meadows area.

White (1979) reviewed the geochemistry of groundwater associated with tuffaceous rocks in Oasis Valley, which is west of NTS and Yucca Mountain. He concluded that most recharge entering Oasis Valley comes from the north and east, including Pahute Mesa. The water moves from the recharge areas through the fracture system in the tuffaceous rocks and into the alluvium in the valley floor. The close proximity of the water in the alluvium to the ground surface promotes evapotranspiration and an increase in concentration of the dissolved species. Some dissolved species such as sodium and chloride were not selectively added or removed from the water, but others such as calcium, fluoride, aqueous silica, and bicarbonate were affected by precipitation or dissolution reactions.

White, Claassen, and Benson (1980) described the effect of volcanic glass dissolution on water chemistry in a tuffaceous aquifer at Rainier Mesa (see also Claassen and White 1979). Water containing dissolved carbon dioxide reacts with vitric and crystalline tuffs. As dissolution proceeds, the water becomes saturated with respect to various minerals. The competition between species added by dissolution and removed by precipitation controls water composition. Experiments with vitric and crystalline tuff produced aqueous solutions of different composition, crystalline tuff giving more calcium-rich water and vitric tuff giving more sodium-rich water. The authors concluded that dissolution of vitric tuff was the predominant reaction affecting water composition; this predominance was probably related to different flow mechanisms in the two materials (fracture flow dominating in crystalline tuff and interstitial flow in vitric tuff). Cation compositions in fracture and pore water sampled at Rainier Mesa were similar, but pore water was higher in chloride and sulfate relative to bicarbonate than was fracture waters (see also Benson 1976). Kerrisk (1983) modeled glass dissolution and mineral precipitation reactions with reaction-path calculations. He was able to reproduce the general trends of major-species water composition and mineral precipitation by using glass dissolution data measured by White, Claassen, and Benson (1980).

Claassen (1985) reviewed data for groundwater in the west-central Amargosa Desert. He concluded that overland flow of snowmelt in or near present-day stream channels was the primary recharge mechanism rather than subsurface flow from highland recharge areas in the north. Two of these stream channels are the Amargosa River, which enters the Amargosa Desert in the west from Oasis Valley, and Fortymile Wash (Fortymile Canyon), which passes Yucca Mountain on its way to the Amargosa Desert. Recharge through alluvium composed of fragments of tuff or carbonate, or mixtures of the two, and through playa deposits results in different water compositions. He also saw evidence for upward leakage of water from a regional carbonate aquifer into alluvium and mixing with water recharged directly into the alluvium at the eastern edge of the study area.

White and Chuma (1986) reviewed data for water from Oasis Valley and Fortymile Wash. They used isotopic data to conclude that groundwater from Pahute Mesa discharges through tuffaceous aquifers into Oasis Valley, but not into Fortymile Wash. They also concluded that water in the alluvium in the upper north-central Amargosa Desert originated principally as groundwater in tuffs in Fortymile Wash; this disagrees with the conclusion of Claassen (1985) that overland flow was more important. White and Chuma (1986) infer that, during passage through Oasis Valley, dissolved carbonate in the water is exchanged with soil-zone carbon dioxide and carbonate in caliche; this exchange is facilitated by the proximity of the water to the ground surface in Oasis Valley.

III. YUCCA MOUNTAIN WELLS

A number of wells have been completed and sampled at Yucca Mountain and vicinity as part of the NNWSI Project or other programs at NTS. Compositions of water from these wells have been reported primarily by Benson et al. (1983), Ogard and Kerrisk (1984), and Benson and McKinley (1985). Table I lists the 15 wells from Yucca Mountain and vicinity that were reviewed as part of this report. Wells J-12 and J-13 are water-supply wells; the others are test wells. Table I also lists the well locations and Fig. 1 shows the locations around Yucca Mountain.

All but one of the wells (UE-25p#1) sample the tuffaceous aquifer around Yucca Mountain. Well UE-25p#1 encountered the carbonate aquifer at about 1300-m depth and continued to about 1800-m depth (Craig and Robison 1984).

TABLE I
WELLS AT YUCCA MOUNTAIN AND VICINITY

<u>Well</u>	<u>Location</u>	<u>Water Samples^a</u>
J-12	Fortymile Wash, east of Yucca Mountain	Integral
J-13	Fortymile Wash, east of Yucca Mountain	Integral
UE-25b#1	Yucca Mountain, Drill Hole Wash	Integral, 863-875 m
UE-25c#1	Yucca Mountain, Midway Valley	Integral
UE-25c#2	Yucca Mountain, Midway Valley	Integral
UE-25c#3	Yucca Mountain, Midway Valley	Integral
UE-25p#1	Yucca Mountain, Midway Valley	381-1179 m, 1279-1805 m
UE-29a#2	Fortymile Wash, northeast of Yucca Mountain	87-213 m, 247-354 m
USW G-4	Yucca Mountain, Drill Hole Wash	Integral
USW H-1	Yucca Mountain, Drill Hole Wash	572-687 m, 687-1829 m
USW H-3	Yucca Mountain crest	822-1220 m
USW H-4	Yucca Mountain, wash south of Drill Hole Wash	Integral
USW H-5	Yucca Mountain crest	Integral
USW H-6	Jet Ridge, across Solitario Canyon, west of Yucca Mountain	Integral, 608-646 m, 753-835 m
USW VH-1	Crater Flat, southwest of Yucca Mountain	Integral

^aLists depth interval sampled or whether the entire well bore was pumped (integral sample).

Water samples from two depths have been reported from UE-25p#1. A deep sample that appears to represent the carbonate aquifer in this location is called "carbonate water" in this report. A shallower sample that represents a mixture of tuffaceous and carbonate aquifer water is called "tuffaceous-carbonate water" in this report. These two samples have different compositions and are usually differentiated in discussions. Most samples from the other wells, which all sample the tuffaceous aquifer, have been integral samples in which no attempt was made to selectively pump particular zones. The exceptions are shown in Table I, where information under the heading WATER SAMPLES indicates whether an integral sample was taken or whether specific intervals were sampled. Discussions of water from the tuffaceous aquifer around Yucca Mountain have usually not differentiated between samples from different depth intervals.

IV. SOURCES OF GROUNDWATER

One of the boundary conditions that influences the composition of saturated-zone water at Yucca Mountain is the source of the water. Water entering the area carries dissolved species with it. The identities and concentrations of these dissolved species depend on the past history of the water. There are three potential sources of groundwater in the tuffaceous aquifer at Yucca Mountain and vicinity: (1) subsurface flow from other recharge areas, (2) direct recharge from local precipitation or runoff, and (3) flow from the carbonate aquifer that underlies the tuffaceous aquifer. A hydrologic model of Yucca Mountain that outlines sources of groundwater in the saturated zone is still being developed by the USGS. However, some ideas about the likelihood of these sources can be obtained from the preliminary models that have been developed (Waddell et al. 1984; Robison 1984; Czarnecki 1985) and from consideration of nearby areas.

The general similarity of the composition of the tuffaceous-aquifer minerals around NTS and vicinity makes it difficult to distinguish, from geochemical evidence alone, between subsurface flow through the tuffaceous aquifer from other recharge areas and direct recharge. Two different proposals for the source of groundwater in the southern part of Fortymile Wash have been made. Claassen (1985) concluded that groundwater was recharged primarily by overland flow and local recharge, but White and Chuma (1986) considered subsurface flow more likely. In either case, the water would contact tuffaceous material in the aquifers or valley fill and have a similar overall composition.

The general flow direction from north to south proposed by Winograd and Thordarson (1975) for groundwater beneath NTS makes highland areas to the north of Yucca Mountain potential recharge areas. White (1979) and White and Chuma (1986) have concluded that groundwater from Pahute Mesa flows into Oasis Valley, but that it is not a source for groundwater in Fortymile Wash. The relationship between water from Pahute Mesa and Yucca Mountain is uncertain at this time.

Claassen (1985) found evidence for upward flow of water from the carbonate aquifer and mixing with water from the tuffaceous aquifer in local areas in the eastern Amargosa Desert. Springs in the Ash Meadows area also discharge water from the carbonate aquifer. No geochemical evidence of upward flow of carbonate aquifer water around Yucca Mountain has been reported;

however, there are potentiometric data indicating this possibility (Robison 1984). Broxton et al. (1986) have proposed this process as the cause of a trend from more sodium- and potassium-rich zeolites in western Yucca Mountain to more calcium-rich zeolites in the eastern part.

V. ACTIVE CHEMICAL AND PHYSICAL PROCESSES

This section presents short discussions of chemical and physical processes that are active at Yucca Mountain and that could affect water compositions. Its purpose is to remind the reader of those processes that could control water composition or that could change water composition as conditions change at Yucca Mountain.

A. Transport with Water

As discussed in Section IV above, transport of dissolved species with water that enters or leaves the Yucca Mountain area is an important process affecting water compositions. Because of the similarity of mineralogy in the tuffaceous aquifers at NTS, it may be that water entering Yucca Mountain by subsurface flow through the tuffaceous aquifer has already achieved a steady-state composition in the major cations and anions. Water from the carbonate aquifer has its own characteristic composition. Water directly recharged through the unsaturated zone may also have a different composition, but this has not been determined as yet.

B. Rock-Water Reactions

Dissolution of volcanic glass and minerals provides a major source of alkali metal and alkaline earth cations, and aqueous silica to water in the tuffaceous aquifer (Hoover 1968; Claassen and White 1979; White et al. 1980). Dissolution and secondary mineral precipitation are responsible for the sorptive minerals found at Yucca Mountain. Water-mineral reactions also stabilize the pH of the water (Ogard and Kerrisk 1984). These processes are particularly important for water that is recharged through the unsaturated zone. Reactions between water and carbonate minerals in the Paleozoic (carbonate) aquifer strongly influence the chemistry of that water.

C. Ion Exchange

Cation exchange between local water and secondary minerals such as clays and zeolites at Yucca Mountain can also change or control alkali metal and alkaline earth concentrations in the water. Isotopic exchange between

dissolved carbonate species and carbonate minerals has also been reported (White and Chuma 1986).

D. Gas Dissolution

Various gases dissolved in water influence water chemistry. At Yucca Mountain, two important gases are oxygen and carbon dioxide. Most saturated-zone water from Yucca Mountain and vicinity is oxidizing, containing measurable quantities of dissolved oxygen (Ogard and Kerrisk 1984). This affects the chemistry of many of the waste elements, particularly the actinides.

Water containing dissolved carbonate species has an equilibrium carbon dioxide pressure that is a function of the amount of dissolved carbonate and the water composition, including pH. The atmosphere and the soil zone are two important sources of gaseous carbon dioxide. Precipitation and water passing through the soil zone pick up dissolved carbon dioxide. The carbon dioxide content of gas from the unsaturated zone and the isotopic composition of the gas has been measured at and near NTS (Henne 1982; White and Chuma 1986) and at Yucca Mountain (Yang et al. 1985). These data have not been related to the carbonate content of water from the saturated zone as yet.

E. Mixing of Different Water Compositions

There are two major aquifers in the vicinity of Yucca Mountain, the tuffaceous aquifer and the carbonate aquifer, which generally underlies the tuffaceous aquifer at NTS. The waters from these two aquifers have somewhat different compositions, and mixtures of the two extremes have been noted (Winograd and Thordarson 1975). This process could be important at Yucca Mountain if water from the carbonate aquifer enters the tuffaceous aquifer.

F. Evaporation


Evaporation is observed in areas such as Oasis Valley where the static water level approaches the land surface. At Yucca Mountain, however, the deep unsaturated zone would preclude evaporation from saturated-zone water. There are two situations where evaporation may be important at Yucca Mountain: in the unsaturated zone and during the thermal pulse in the repository. In the matrix of the unsaturated zone, where liquid water presents a large surface area to the gas phase, evaporation may affect water compositions. Because of the limited amount of data about unsaturated-zone water compositions, little can be said about evaporation in the unsaturated zone at this time.

VI. IMPORTANT GROUNDWATER COMPOSITIONAL VARIABLES AND RELATIONS AMONG THESE VARIABLES


The composition of groundwater represents direct evidence of the chemical and physical processes that are active in control of that composition. Water compositions and relations among compositional variables are often used to support interpretations or models about the origin or age of the water, or about its relation to the local area (Fritz and Fontes 1980; Drever 1982; Claassen 1985; White and Chuma 1986). This section presents information about relations among various compositional variables of water from Yucca Mountain, NTS, and vicinity. The object of these presentations is to show relations where they exist and, in this and the following sections, to interpret these relations in terms of the physical and chemical processes that control water composition.

A. Major Species and Variables

Sodium is the primary cation in essentially all the saturated-zone water in this area. Calcium and potassium are next in importance, with calcium predominating in waters from the carbonate aquifer. Magnesium is also present in smaller quantities in these waters; it is more prominent in waters from the carbonate aquifer. The dissolution of volcanic glass and minerals in the tuffaceous and carbonate aquifers is the major source of these cations (Hoover 1968; Winograd and Thordarson 1975; Claassen and White 1979; White et al. 1980). Figure 2 shows plots of total sodium content as a function of total carbonate (the primary anion) content of water from Yucca Mountain, NTS, and vicinity. The meanings of the abbreviations shown in the legends are described in Table II. In Fig. 2(a), data from 187 samples are plotted; the data are segregated into 10 sets, primarily based on location. Definite clustering of some sets of data can be seen. Above about 6 mmoles/l total carbonate, the data split into two groups; water from Oasis Valley continues to increase in sodium content with increasing carbonate, but other water does not. White (1979) concluded that the water in Oasis Valley is undergoing evaporation as it moves down the valley; this is demonstrated by the roughly linear variation of sodium and carbonate concentrations (as well as a number of other concentrations that will be discussed later) in that water. Three of the samples with high carbonate (9-10 mmoles/l) and intermediate sodium (5-7 mmoles/l) are from carbonate aquifer wells (UE-25p#1 carbonate water, Well C, and Well C-1). Some of the higher carbonate waters that fall in the group



(a)



(b)

Fig. 2. Sodium content as a function of total carbonate content.
 a. All sampling locations (see Table II for meaning of legend abbreviations).
 b. Yucca Mountain and vicinity.

TABLE II
WATER SAMPLING LOCATIONS

<u>Abbreviation</u>	<u>Location</u>
YM/FMW	Wells at Yucca Mountain and vicinity including Fortymile Wash (J-12, J-13, UE-25b#1, UE-25c#1, UE-25c#2, UE-25c#3, UE-25p#1, UE-29a#2, USW G-4, USW H-1, USW H-3, USW H-4, USW H-5, USW H-6, and USW VH-1).
AD/FMW	Amargosa Desert wells and springs in the Fortymile Wash stream channel - sites 3 to 19 from Claassen (1985).
AD/MISC	Miscellaneous wells and springs in the Amargosa Desert - from Claassen (1985).
AD/OV	Amargosa Desert wells and springs in the Oasis Valley drainage system above the Fortymile Wash stream channel - sites 45 to 52 from Claassen (1985).
OTHER	NTS wells and springs from Yucca Flat, Frenchman Flat, and the Calico Hills - Claassen (1973) and Claassen (1985).
OV	Wells and Springs from Oasis Valley - White (1979) and White and Chuma (1986).
PM	Wells from Pahute Mesa
RM/PW	Pore water from the unsaturated zone at Rainier Mesa - Benson (1976).
RM/FW	Fracture water from Rainier Mesa - White et al. (1980).
RM/MISC	Miscellaneous wells, springs, and fracture-water samples from Rainier Mesa.

between the carbonate wells and the main body of the data may result from mixing more dilute water from the tuffaceous aquifer with water from the carbonate aquifer; two of these wells are from the tuffaceous aquifer at Pahute Mesa (UE-19c and UE-19d). Winograd and Thordarson (1975) identify a mixed tuffaceous-carbonate water in terms of downward flow of water from the tuffaceous aquifer into the carbonate aquifer. It is also possible that upward flow of water from the carbonate aquifer into the tuffaceous aquifer (Waddell et al. 1984) could result in a similar mixing process and similar water compositions.

In Fig. 2(b), the shaded area represents the range of compositions covered in Fig. 2(a) by all the data, and only the compositions from Yucca Mountain and vicinity are plotted as specific points. Excluding data for the UE-25p#1 carbonate water (9.3 mmoles/l total carbonate), which is a carbonate aquifer sample, there is about a factor of 2 variation in the sodium and carbonate contents of water from the tuffaceous aquifer at Yucca Mountain. The trend of the compositions of the Yucca Mountain water with increasing carbonate content generally stays below the Oasis Valley data.

Figure 3 shows plots of total calcium content as a function of total carbonate content of water from Yucca Mountain, NTS, and vicinity. In Fig. 3(a), data from the same 187 sources that were shown in Fig. 2(a) are presented. As in Fig. 2(a), definite clustering of some sets of data can be seen in Fig. 3(a). The Oasis Valley data show a relatively constant calcium content with increasing carbonate, probably because of solubility constraints. In Fig. 3(b), the shaded area represents the range of compositions covered in Fig. 3(a) by all the data, and only the compositions from Yucca Mountain and vicinity are plotted as specific points. Except for both samples from UE-25p#1, the water from Yucca Mountain has relatively low calcium content (less than 0.5 mmoles/l). This is particularly true of water from wells USW H-3, H-6, H-5, and H-1, which have calcium contents of 0.15 mmoles/l or less. These wells are on the western part of Yucca Mountain or across Solitario Canyon. The Ghost Dance Fault is an approximate dividing line between these low calcium wells and those to the east with higher calcium content (see Section VI(D)).

Figure 4(a) shows a ternary plot of the relative sodium, calcium, and potassium contents of water from Yucca Mountain, NTS, and vicinity. Water from the carbonate aquifer has high calcium (and magnesium), up to about 45% in the data shown in Fig. 4(a). Most of the Yucca Mountain waters plot along a line with about 0-5% potassium, and sodium ranging from about 70% up to essentially 100%; this is more apparent in Fig. 4(b), which shows a shaded area representing all the data in Fig. 4(a) with only Yucca Mountain data plotted as specific points. Two of the wells in Fortymile Wash (J-12 and J-13) have slightly higher relative potassium contents. An attempt was made to correlate the relative sodium content ($Na/(Na+Ca+K)$) of the water with other compositional variables, but no significant correlations were found for





Fig. 3. Calcium content as a function of total carbonate content.

a. All sampling locations (see Table II for meaning of legend abbreviations).

b. Yucca Mountain and vicinity.




Fig. 4. Ternary diagram of relative sodium, calcium, and potassium content.
 a. All sampling locations (see Table II for meaning of legend abbreviations).
 b. Yucca Mountain and vicinity.

all the data. There is, however, a relation between relative sodium content and pH for water from Yucca Mountain and vicinity, and between relative sodium content and the location of Yucca Mountain wells. Figure 5 shows a plot of $\text{Na}/(\text{Na}+\text{Ca}+\text{K})$ as a function of pH for water from Yucca Mountain and vicinity. In addition to the water data plotted there, the two lines show variation in the same variables from reaction-path calculations of volcanic-glass dissolution and secondary-mineral precipitation at Rainier Mesa (Kerrick 1983). The general shape of the data, relatively flat at lower $\text{Na}/(\text{Na}+\text{Ca}+\text{K})$ values with increasing pH as $\text{Na}/(\text{Na}+\text{Ca}+\text{K})$ approaches 1, is also predicted by the calculations. The variation of $\text{Na}/(\text{Na}+\text{Ca}+\text{K})$ with location is discussed in Section VI(D).

The pH of water from Yucca Mountain, NTS, and vicinity is generally in the range of 7 to 9. Two extreme values are observed at Yucca Mountain; the carbonate-water sample from UE-25p#1 has a pH of 6.6 and water from H-3 has a pH of 9.2. The pH of all water samples as a group did not show any correlations with other compositional variables. However, at Yucca Mountain and vicinity, pH is related to a number of other compositional variables. The relation between pH and $\text{Na}/(\text{Na}+\text{Ca}+\text{K})$ was noted above and shown in Fig. 5. The pH is also inversely related to calcium content and, to some extent, to magnesium content. If the calcium content is related to pH through calcite solubility, a plot of \log_{10} (calcium content) as a function of pH should be linear with slope -2 when the carbon dioxide partial pressure is constant, or curved with the slope varying from -2 through 0 with increasing pH when the total carbonate content is constant. Figure 6 shows a plot of \log_{10} (calcium content) as a function of pH of water from Yucca Mountain and vicinity. The slope of a curve through the data would decrease from about -2 at pH 6-7 to about -1 at pH 8-9. This variation approximates the expected behavior for constant total carbonate content and would indicate that calcite solubility may be a factor in controlling the calcium content of water at Yucca Mountain. There is also a relation between pH and total carbonate content of tuffaceous-aquifer water from Yucca Mountain and vicinity (see Fig. 7). Although most of the tuffaceous-aquifer data are grouped together in one area, water with higher total carbonate content tends to have higher pH. Water from UE-25p#1 has high carbonate content but low pH; thus, it does not fit this trend.




Fig. 5. $\text{Na}/(\text{Na} + \text{Ca} + \text{K})$ as a function of pH at Yucca Mountain and vicinity. Solid lines represent calculated results from Kerrisk (1983).




Fig. 6. The \log_{10} (calcium content) as a function of pH at Yucca Mountain and vicinity.




Fig. 7. The pH as a function of total carbonate content at Yucca Mountain and vicinity.

The carbon dioxide partial pressure in equilibrium with water compositions was calculated using the EQ3 chemical equilibrium computer program (Wolery 1983) for about two-thirds of the samples reviewed here. Figure 8(a) shows a plot of \log_{10} (carbon dioxide pressure) as a function of pH for samples from Yucca Mountain, NTS, and vicinity. There is a trend toward decreasing carbon dioxide pressure with increasing pH. As expected, water from the carbonate aquifer tends to have higher equilibrium carbon dioxide pressures than water from the tuffaceous aquifer. The Rainier Mesa pore water shown in Fig. 8(a) has lower equilibrium carbon dioxide pressures than the fracture water. Figure 8(b) shows a similar plot for data just from Yucca Mountain and vicinity. There is much less scatter in these data. The slope of a line through the UE-25p#1 data is about -2; in the higher pH range (7.5 to 9), the slope is about -1. The maximum carbon dioxide pressure calculated for tuffaceous waters from Yucca Mountain ($\sim 10^{-2}$ atm) is similar to peak carbon dioxide pressures observed in the soil zone at Yucca Mountain (Yang et al. 1985). Thus, soil-zone carbon dioxide is a possible source of the aqueous




Fig. 8. The \log_{10} (carbon dioxide pressure) as a function of pH.

- All sampling locations (see Table II for meaning of legend abbreviations).
- Yucca Mountain and vicinity.

carbonate in the water. The calculated carbon dioxide pressure of water from Well H-3 at Yucca Mountain ($10^{-3.8}$ atm) is below the carbon dioxide pressure in the atmosphere ($10^{-3.5}$ atm) and well below carbon dioxide pressures measured in the gas phase of the unsaturated zone at Yucca Mountain (Yang et al. 1985). This water also has the highest total carbonate content of the tuffaceous aquifer wells at Yucca Mountain. The high carbonate content of this water could result from some process that increases the pH, thus driving down the equilibrium carbon dioxide pressure of the water to the level where it is below carbon dioxide partial pressures in the unsaturated-zone gas phase above the water, and allowing dissolution of gas-phase carbon dioxide in the water.

As could be implied from the relations between pH and total carbonate (Fig. 7), and carbon dioxide pressure and pH (Fig. 8(b)) for tuffaceous water at Yucca Mountain, there is also a relation between carbon dioxide pressure and total carbonate for these data (see Fig. 9). For the tuffaceous wells, those with higher total carbonate tend to have lower carbon dioxide pressures. This is consistent with the possibility that gas-phase carbon dioxide in the unsaturated zone could be the source of aqueous carbonate contents above about 2 mmoles/l in tuffaceous-aquifer water from Yucca Mountain. The two samples from UE-25p#1, which represent carbonate aquifer water and mixed tuffaceous-carbonate water, do not fit this trend (see Figs. 7 and 9).

The discussions about relations among pH, total carbonate, calcium content, and equilibrium carbon dioxide pressure in the previous two paragraphs alluded to the possibility that calcite or dolomite solubilities may also be involved in controlling water chemistry at Yucca Mountain. The state of water with respect to saturation with a mineral can be represented by the quantity $\log_{10}(Q/K)$, where Q is the ion activity product and K is the equilibrium constant for the solubility reaction (Stumm and Morgan 1981). The quantity $\log_{10}(Q/K)$ is negative for undersaturation, zero at saturation, and positive for oversaturation. Figure 10 shows a plot of calcite $\log_{10}(Q/K)$ as a function of pH for water from Yucca Mountain and vicinity. The $\log_{10}(Q/K)$ data were calculated from water compositions using the EQ3 chemical equilibrium computer program at the same time that carbon dioxide pressures were calculated (Walery 1983). Most of the waters with high pH are near saturation with respect to calcite; these are the same waters that showed low




Fig. 9. The log_{10} (carbon dioxide pressure) as a function of the carbonate content at Yucca Mountain and vicinity.




Fig. 10. Calcite $\text{log}_{10}(Q/K)$ as a function of pH at Yucca Mountain and vicinity.

calcium contents in Fig. 6. Thus, the low calcium content of these waters may be the result of calcite (or dolomite) precipitation as water pH increases. The low calcium content and the high carbonate content in the tuffaceous-aquifer water in the western part of Yucca Mountain may be connected phenomena.

Essentially all the aqueous silica contents of the waters from Yucca Mountain, NTS, and vicinity range from about 0.6 to 1.3 mmoles/l, with an average of 0.89 mmoles/l. This puts the water in the range of cristobalite to amorphous silica saturation. A few samples show higher and lower values. In particular, water sampled from the surface soil of Rainier Mesa show lower aqueous silica contents, 0.3 to 0.6 mmoles/l (Benson 1976; Henne 1982). Wells that tap the carbonate aquifer (Wells Army-1, C, and C-1) are also low in aqueous silica; Winograd and Thordarson (1975) noted the low aqueous silica content of water from the carbonate aquifer. The aqueous silica contents of waters from this area did not show significant correlations with any of the other compositional variables examined. Aqueous silica is supplied to these waters primarily by dissolution of tuffaceous rock and minerals; concentrations are probably controlled by precipitation of various solid silica polymorphs. Kerrisk (1983) has proposed that high aqueous silica activity (in equilibrium with cristobalite) is required for the presence of the zeolites found at Yucca Mountain.

Carbonate is the primary anion in essentially all the water from Yucca Mountain, NTS, and vicinity. After carbonate, chloride and sulfate are next in order of amount present; these two anions are generally present in about equimolar quantities. Carbonate, chloride, and sulfate usually represent 95% or more of the anion content of the water. Figure 11 shows plots of (chloride + sulfate) content as a function of total carbonate content of these waters. In Fig. 11(a), data from 185 sources are plotted. Three types of behavior can be seen moving away from the dilute region of the plot. Many samples of Rainier Mesa pore water show high (chloride + sulfate) content (1-2.5 mmoles/l) for the amount of carbonate present (0.4-1 mmoles/l) (the data cluster near the vertical axis); this is consistent with the low equilibrium carbon dioxide pressures calculated for this water (see Fig. 8(a)). Data from Oasis Valley show a roughly linear trend of increasing (chloride + sulfate)




Fig. 11. The (chloride + sulfate) content as a function of total carbonate content.

- All sampling locations (see Table II for meaning of legend abbreviations).
- Yucca Mountain and vicinity.

with carbonate; water from the Amargosa Desert in the Oasis Valley drainage system has somewhat higher (chloride + sulfate) content. These data tend to cluster in the middle of the plot. Water from carbonate aquifer wells and other wells around NTS shows lower (chloride + sulfate) content for the amount of carbonate present; these data cluster near the horizontal axis. In Fig. 11(b), the shaded area represents the range of compositions covered in Fig. 11(a) by all the data, and only compositions from Yucca Mountain and vicinity are plotted as specific points. Except for UE-25p#1 carbonate water, water from Yucca Mountain and vicinity is low in (chloride + sulfate), 0.3-0.8 mmoles/l. Figure 12(a) shows an expanded-scale plot of sulfate content alone as a function of carbonate content for water from Yucca Mountain and vicinity; Fig. 12(b) shows a similar plot for chloride. Neither plot includes the carbonate water from UE-25p#1 (sulfate content 1.67 mmoles/l, chloride content 0.79 mmoles/l, and carbonate content 9.3 mmoles/l), but they both include the mixed tuffaceous-carbonate water from this well. The sulfate content shows an almost linear relation with carbonate except for Well VH-1 (see Fig. 12(a)). There is enough scatter in the chloride plot so that there appears to be little or no relationship between chloride and carbonate for these wells (see Fig. 12(b)).

Figure 13 shows plots of sulfate content as a function of chloride content for water from Yucca Mountain, NTS, and vicinity. In Fig. 13(a), data from 185 sources are plotted. Most of the data cluster around the origin, with less than 0.5 mmoles/l sulfate and chloride contents. The straight line through the origin, with a slope of 0.82, represents a least squares fit of all the data. Claassen (1985) shows a similar plot with a line representing evaporation of modern precipitation; the slope of that line is approximately 1.65, that is, twice the slope of the line in Fig. 13(a). Claassen (1985) also noted that groundwater in the Amargosa Desert contains less sulfate relative to chloride than modern precipitation. EQ3 calculations of gypsum $\log_{10}(Q/K)$ for about two-thirds of the samples in Figs. 12 and 13 gave values from -5 up to about -1. This indicates that all these waters are undersaturated with respect to gypsum so that gypsum solubility should not limit sulfate concentrations. The Rainier Mesa pore-water data with high (chloride + sulfate) in Fig. 11(a) plot below the line in Fig. 13(a), with sulfate content about half the chloride content. Figure 13(b) shows an




Fig. 12. Sulfate and chloride contents as a function of total carbonate content at Yucca Mountain and vicinity.
 a. Sulfate content as a function of total carbonate.
 b. Chloride content as a function of total carbonate.




Fig. 13. Sulfate content as a function of chloride content (see Table II for meaning of legend abbreviations).
 a. All sampling locations.
 b. Expanded plot of region near the origin.

expanded view of the region near the origin; most of the data from Yucca Mountain wells appear in this region. The straight line is the same line shown in Fig. 13(a).

Measurements of the redox state of water have been made for some wells at Yucca Mountain and vicinity but have not been reported for any of the other waters reviewed here. Dissolved oxygen, Eh, and nitrite/nitrate ratios have been measured (Ogard and Kerrisk 1984). Most of the waters measured showed detectable dissolved oxygen (2 to 6 mg/l) and Eh values in the 200- to 400-mV range (against a H₂ electrode). Both measures are signs of oxidizing conditions. Two wells, UE-25b#1 and USW H-3, had dissolved oxygen contents below detection limits (0.1 mg/l) and negative values of Eh in pumped samples. During a 28-day pumping test from a packed-off interval of the Bullfrog Member tuff in Well UE-25b#1, reducing conditions at the start of the test gave way to oxidizing conditions at the end; the measured Eh of the water increased, dissolved oxygen increased during the latter third of the test, and the total iron, total manganese, and nitrite/nitrate ratio decreased (Daniels et al. 1983; Ogard et al. 1983; Rundberg et al. 1985). All these trends are indications that the water became more oxidizing as the test progressed. However, the three direct measures of the redox state, dissolved oxygen, Eh, and nitrite/nitrate ratio, gave conflicting quantitative information. The lack of equilibrium among various redox couples that can exist in solution is common (Lindberg and Runnels 1984). The other well to show reducing conditions during pumping tests, USW H-3, had detectable dissolved oxygen that decreased with time early in a three-month pumping test; however, the water stabilized at oxygen levels below detection and Eh in the range of -80 to -140 mV for most of the test (Crowe and Vaniman 1985). The oxidizing conditions observed early in the USW H-3 test were probably a result of contamination during drilling. Two other wells, USW H-1 and H-4, showed reducing conditions (negative values of Eh and detectable sulfide concentrations) in samples taken in evacuated bottles that were lowered into the well bores to various depths after pumping tests were completed and the pumps removed (Ogard and Kerrisk 1984). Well USW H-4 water had positive Eh and considerable dissolved oxygen in the sample taken during pumping. In addition to the redox measurements on water samples, measurements of the composition of the unsaturated-zone gas phase at Yucca Mountain have shown about 20% oxygen present even at 300-m

depth (Yang 1986). The presence of gaseous oxygen above the static water level is an indication that water at that level should be oxidizing. The existence of a wide range of redox conditions in water from Yucca Mountain and vicinity, and the variation of redox conditions in individual wells, have not been satisfactorily explained at this time.

B. Isotopic Data

There have been a number of measurements of carbon, hydrogen, and oxygen isotopic data on the waters reviewed here. These data often provide information about the origin of the water or the physical and chemical processes that the water has undergone (Fritz and Fontes 1980; Faure 1977; Dansgaard 1964; Craig 1961). Figure 14 shows plots of the percentage of modern carbon (PMC) in aqueous carbonate as a function of total carbonate content of water from Yucca Mountain, NTS, and vicinity. PMC is a measure of the ^{14}C content of carbon in terms of the ^{14}C content of a carbon standard, and it ranges from about 100% for carbon that is in equilibrium with the atmosphere to 0% for carbon that has been isolated from sources of ^{14}C for a long time (Fritz and Fontes 1980). Values of PMC greater than 100% are also found because the standard represents ^{14}C contents before atmospheric nuclear testing increased the ^{14}C content of the atmosphere. Carbonate derived from Paleozoic rocks has very low values of PMC. The apparent age (t_a) of the water is related to PMC by

$$t_a = -[\ln(\text{PMC}/100)]/\lambda, \quad (1)$$

where $\lambda = 1.24 \times 10^{-4}/\text{yr}$ is the decay constant of ^{14}C . In Fig. 14(a), data for 53 samples are plotted. There are two distinct trends apparent. The data from Oasis Valley show an increase in PMC with increasing carbonate content, and were explained by White and Chuma (1986) in terms of contact between the water in the shallow saturated zone of Oasis Valley and soil-zone carbon dioxide or caliche; this leads to increasing PMC as the water moves down the valley. Some of the increase is caused by an addition of carbonate to the water and some by isotopic exchange between aqueous carbonate and the soil-zone sources. The data from Yucca Mountain and vicinity, on the other hand, show a range of PMC (20% to 60% or apparent ages of 13000 to 4000 yr) at 2 mmoles/l total carbonate with a trend toward decreasing PMC with increasing carbonate content above 2 mmoles/l; this trend is more apparent in Fig. 14(b),




Fig. 14. Percentage of modern carbon as a function of total carbonate content.

a. All sampling locations (see Table II for meaning of legend abbreviations).

b. Yucca Mountain and vicinity.

where only the Yucca Mountain data are plotted. It is not clear whether there is a connection between the tuffaceous water data and UE-25p#1 data, or whether the tuffaceous water data level off at about 10% PMC (apparent age 18500 yr). A connection could indicate that the trend in Fig. 14(b) is caused by the addition of carbonate water to the tuffaceous aquifer (see Section VII(B)).

Another feature of the data in Fig. 14(b) is that those wells with the highest values of PMC or youngest apparent age (UE-29a#2, J-12, and J-13) are in Fortymile Wash where local recharge may be high (Claassen 1985). These two characteristics, high local recharge and young water, are consistent.

Figure 15 shows plots of the relative $^{13}\text{C}/^{12}\text{C}$ content ($\delta^{13}\text{C}$) of aqueous carbonate in these same waters as a function of the total carbonate content. The $\delta^{13}\text{C}$ data can provide some information about the origin of the carbonate (Faure 1977; Fritz and Fontes 1980; Haas et al. 1983; Thorstenson et al. 1983; Wigley et al. 1978). In particular, paleozoic carbonates have a $\delta^{13}\text{C}$ value of $\sim -2\text{‰}$, atmospheric carbon dioxide has a $\delta^{13}\text{C}$ value of ~ -7 to -9‰ , and soil-zone carbon dioxide that is derived from plants has $\delta^{13}\text{C}$ values of ~ -12 to -25‰ . Some care must be taken in comparing $\delta^{13}\text{C}$ values of gas, aqueous, and solid phases because of fractionation; thus, carbon dioxide gas will have a different $\delta^{13}\text{C}$ than the carbonate in an aqueous phase in equilibrium with it (Wigley et al. 1978). Figure 15(a) shows data for 55 samples from Yucca Mountain, NTS, and vicinity. As with the ^{14}C data (PMC) in Fig. 14, there are two trends apparent in Fig. 15(a). The data from Oasis Valley, although scattered, show a decrease in $\delta^{13}\text{C}$ with increasing carbonate. This trend is consistent with the variation of PMC with carbonate; soil-zone carbonate that is added to the aqueous phase or exchanges with aqueous carbonate is derived from plant respiration and has more negative $\delta^{13}\text{C}$ than the original carbonate in the water. The data from Yucca Mountain and vicinity (see also Fig. 15(b)) show a range of $\delta^{13}\text{C}$ values (-13‰ to -7‰) at about 2 mmoles/l total carbonate, with some tendency to more positive $\delta^{13}\text{C}$ with increasing carbonate. Only one tuffaceous aquifer well from Yucca Mountain (USW H-3) has a $\delta^{13}\text{C}$ value above about -7‰ . As with PMC, it is not clear whether there is a connection between the tuffaceous water data and the UE-25p#1 data. If there is a connection, the increase in $\delta^{13}\text{C}$ with increasing carbonate could be caused by the addition of carbonate aquifer water to the tuffaceous aquifer.




Fig. 15. $\delta^{13}\text{C}$ as a function of total carbonate content.
 a. All sampling locations (see Table II for meaning of legend abbreviations).
 b. Yucca Mountain and vicinity.

The relation between pH and total carbonate shown in Fig. 7, and the relations between PMC and total carbonate and between $\delta^{13}\text{C}$ and total carbonate shown in Figs. 14(b) and 15(b), imply a relation between pH and the isotopic variables. Figure 16(a) shows a plot of pH as a function of PMC and Fig. 16(b) shows a plot of pH as a function of $\delta^{13}\text{C}$ for water from Yucca Mountain and vicinity. Both plots show some trends in the data for the tuffaceous waters, but the UE-25p#1 samples do not follow these trends. This disparity between the UE-25p#1 data and the tuffaceous water data was also evident in plots of pH and carbon dioxide pressure as a function of total carbonate (Figs. 7 and 9), and calcite $\log_{10}(Q/K)$ as a function of pH (Fig. 10). This behavior contrasts with the apparent continuity between the UE-25p#1 data and tuffaceous data in plots of calcium content and carbon dioxide pressure as a function of pH (Figs. 6 and 8(b)) and of PMC and $\delta^{13}\text{C}$ as a function of total carbonate (Figs. 14(b) and 15(b)). Although some data point to a connection between the tuffaceous and carbonate waters at Yucca Mountain, it is far from clear that an actual physical or chemical connection exists (see Section VII(B)).

Stable hydrogen and oxygen isotopic data can also provide information about the origins of water or some of the physical and chemical processes that the water has undergone (Craig 1961; Faure 1977; Dansgaard 1964; Fritz and Fontes 1980). Figure 17(a) shows a plot of the relative deuterium/hydrogen content (δD) of water from Yucca Mountain, NTS, and vicinity as a function of the relative $^{18}\text{O}/^{16}\text{O}$ content ($\delta^{18}\text{O}$). The straight line represents a relation obtained by Craig (1961) for meteoric water. Claassen (1985) shows a similar plot (with only some of the data) and interprets the distribution of the data along the meteoric water line as a temperature effect, with more negative δD and $\delta^{18}\text{O}$ originating as precipitation at lower temperatures. Claassen (1985) also shows relations between δD and $\delta^{18}\text{O}$, and apparent age, with older waters (lower PMC) showing more negative δD and $\delta^{18}\text{O}$. He relates this to colder temperatures 10,000 to 15,000 years ago (see also the following paragraph). White and Chuma (1986) reported the data from Pahute Mesa and Oasis Valley in Fig. 17(a). They interpreted the spread in Oasis Valley as caused by a mixing of Pahute Mesa water ($\delta\text{D} = \sim -114\text{‰}$ and $\delta^{18}\text{O} = \sim -14.5\text{ to } -14.7\text{‰}$) with water from the Bullfrog Hills in Oasis Valley ($\delta\text{D} = \sim -102\text{‰}$ and $\delta^{18}\text{O} = \sim -13.4\text{‰}$). They also cite the difference between Pahute Mesa water and water from Fortymile Wash and the portion of the Amargosa Desert that drains




Fig. 16. The pH as a function of the percentage of modern carbon and $\delta^{13}\text{C}$ at Yucca Mountain and vicinity.
 a. pH as a function of the percentage of modern carbon.
 b. pH as a function of $\delta^{13}\text{C}$.




Fig. 17. δD as a function of $\delta^{18}\text{O}$.

a. All sampling locations (see Table II for meaning of legend abbreviations).

b. Yucca Mountain and vicinity.

Fortymile Wash as evidence that Pahute Mesa is not a recharge area for Fortymile Wash. Figure 17(b) shows a plot of δD as a function of $\delta^{18}O$ for water from Yucca Mountain and vicinity only. The wells from Fortymile Wash (J-12, J-13, and UE-29a#2) plot at more positive δD and $\delta^{18}O$ than the Yucca Mountain wells, consistent with that water originating as precipitation at higher temperatures or lower elevations (Dansgaard 1964; Fritz and Fontes 1980).

The relation between δD and apparent age reported by Claassen (1985) for water from the Amargosa Desert also holds for Yucca Mountain. Figure 18(a) shows a plot of δD as a function of PMC for water from Yucca Mountain and vicinity. Water with lower values of PMC (older apparent age) has more negative values of δD , indicating it originated as precipitation at lower temperatures or at higher elevations. Because PMC and the total carbonate content of water from Yucca Mountain are related (see Fig. 14(b)), there is also a relation between δD and total carbonate; this is shown in Fig. 18(b), where increasing carbonate content above about 2 mmoles/l leads to more negative values of δD . Both plots show scatter; however, the relationship between δD and PMC (Fig. 18(a)) seems to be better defined. This leaves two possible explanations for the range of δD values observed: that it is a temperature effect with older water originating as precipitation at lower temperatures (Claassen 1985) or that it is a mixing of waters with different values of δD . Some combination of these two limiting cases is also possible.

Tritium is a radioactive isotope of hydrogen that is sometimes useful for dating water (Fritz and Fontes 1980). Because of its short half-life (12.3 yr), water isolated from atmospheric sources of tritium would not contain measurable levels after about 100 yr. Before atmospheric testing, the natural tritium content of precipitation was less than about 60 pCi/l; atmospheric testing of nuclear weapons increased tritium contents of precipitation by several orders of magnitude (Fritz and Fontes 1980). The tritium contents of water from a number of wells at Yucca Mountain (Benson and McKinley 1985) and around NTS (Clebsch 1961; Claassen 1973) have been reported. Most of the results represent tritium contents below detection limits of the analysis, however some analyses have reported measurable tritium contents. Clebsch (1961) found detectable tritium in Whiterock Spring and in water from a tunnel at Rainier Mesa; tritium was not detected in water from a number of wells around NTS. Claassen (1973) reported detectable tritium




Fig. 18. δD as a function of the percentage of modern carbon and total carbonate content at Yucca Mountain and vicinity.
 a. δD as a function of the percentage of modern carbon.
 b. δD as a function of total carbonate content.

levels in water from a number of NTS water-supply wells during a 1964 survey; levels of 1000 to 2500 pCi/l (300 to 800 tritium units) were measured for some wells, but most wells had much lower levels. Benson and McKinley (1985) reported detectable tritium levels in six wells from Yucca Mountain and vicinity, UE-25b#1 (2 pCi/l), UE-25c#3 (2 pCi/l), UE-25p#1 carbonate water (10 pCi/l), UE-29a#2 (37 pCi/l), USW H-3 (2 pCi/l), and USW H-6 (1-4 pCi/l).* The highest tritium level around Yucca Mountain is from the shallow (29-m depth to static water level) Well UE-29a#2 in Fortymile Wash; this well also has the highest value of PMC (youngest apparent age) of these wells. The meaning of measurable tritium levels in water from deep wells (over 100-m depth to static water level) is uncertain. If they are representative of the water at depth, they imply a connection that allows fast transport (less than 100 yr) between an atmospheric source of tritium and the saturated zone, or a source of tritium at depth. Low but measurable levels of tritium may also result from sample contamination.

C. Other Species and Variables

In addition to the discussions of major species and isotopic data presented above, there are a number of minor species or other variables that are important or that show some interesting relationships. This section discusses nitrate, fluoride, organic and particulate concentrations, and water temperature.

Analyses for nitrate are often not done. Only about a quarter of the water samples reviewed here had nitrate contents reported; however, two-thirds of the samples from Yucca Mountain and vicinity included nitrate data. The nitrate content of all the waters as a group did not show any relationship with other compositional variables, but the data from Yucca Mountain and vicinity did. Figure 19(a) shows a plot of nitrate content of water from Yucca Mountain and vicinity as a function of PMC. Figure 19(b) shows nitrate content as a function of total carbonate. The behavior of nitrate content as a function of these variables shows the same general trends as that of δD (see Figs. 18). For these wells, low nitrate content is associated with low PMC (older apparent age) and, to some extent, with higher total carbonate

*In their report on UE-25p#1, Craig and Robison (1984) quoted tritium levels less than 10 pCi/l for both carbonate water and mixed tuffaceous-carbonate water; thus, the report of 10 pCi/l for UE-25p#1 carbonate water in Benson and McKinley (1985) may be a misprint.




Fig. 19. Nitrate content as a function of the percentage of modern carbon and total carbonate content at Yucca Mountain and vicinity.

a. Nitrate content as a function of the percentage of modern carbon.

b. Nitrate content as a function of total carbonate content.

content. High nitrate content in water is usually associated with surface contamination such as fertilizers or sewage, or with shallow wells (Feth 1966). The well with the highest nitrate content in Figs. 19 (UE-29a#2) is very shallow (29-m depth to water (Benson and McKinley 1985)), compared to the other wells around Yucca Mountain (180- to over 700-m depth to water (Robison 1984)). Data on nitrate contents were unavailable for most other waters examined here; data were reported for some wells from NTS (Claassen 1973) and Pahute Mesa (Blankenagel and Weir 1973). The data from Pahute Mesa showed low nitrate contents (less than 0.05 mmoles/l); the data from wells in Yucca Flat and Frenchman Flat showed higher nitrate contents (up to 0.2 mmoles/l). A survey of 950 groundwater analyses around the United States showed that two-thirds were in the 0- to 0.15-mmoles/l range (Feth 1966). This range covers all the water from around Yucca Mountain except UE-29a#2. Based on this comparison, the presence of nitrate in the water and the range of nitrate concentrations observed at Yucca Mountain are not unusual.

Data on fluoride contents are available for about half the water samples reviewed here; all of the samples from Yucca Mountain and vicinity included fluoride data. The fluoride content of all the water samples taken together did not seem to be strongly correlated with other compositional variables, however there was a tendency for increasing fluoride content with increasing sodium and carbonate contents and relative sodium content ($\text{Na}/(\text{Na}+\text{Ca}+\text{K})$). Considering only the data from Yucca Mountain and vicinity, there is a nearly linear relation between fluoride content and sodium and carbonate contents. Figure 20(a) shows a plot of fluoride content of water from Yucca Mountain and vicinity as a function of sodium content and Fig. 20(b) shows fluoride content as a function of total carbonate content. The wells with high sodium and carbonate contents also have high fluoride contents. The carbonate water from UE-25p#1 does not fit the fluoride-carbonate relation well (see Fig. 20(b)). Ogard and Kerrisk (1984) also showed a tendency for $\text{F}^-/(\text{F}^-+\text{Cl}^-)$ to increase with increasing $\text{Na}/(\text{Na}+\text{Ca}+\text{K})$ for water from Yucca Mountain. EQ3 calculations of fluorite $\log_{10}(\text{Q}/\text{K})$ for 56 water samples from Yucca Mountain, NTS, and vicinity gave values from -3 up to 0.5. Two wells from Yucca Mountain and vicinity (UE-25p#1 carbonate water and USW H-4) were supersaturated with respect to fluorite; other supersaturated waters were from Oasis Valley and Pahute Mesa. Waters with both phosphate and fluoride concentrations reported (Claassen 1973) were highly supersaturated with respect to fluorapatite.




Fig. 20. Fluoride content as a function of sodium content and total carbonate at Yucca Mountain and vicinity.

a. Fluoride content as a function of sodium content.

b. Fluoride content as a function of total carbonate content.

The organic contents of water from two wells at Yucca Mountain and vicinity were reported by Means et al. (1982). No other organic analyses were found for water from Yucca Mountain, NTS, and vicinity. The total organic carbon content of Well J-13 water was 0.15 mg/l, and of UE-25b#1 water was 0.55 mg/l. About 50% of the organic content of J-13 water and 33% of the organic content of UE-25b#1 water were high molecular-weight organics (molecular weight greater than 1000). The organic content of J-13 water is probably more representative of conditions in the saturated zone because it is a producing well and all drilling fluids have been removed by extensive pumping. At this time, there are not enough data on the organic contents of water from Yucca Mountain and vicinity to attempt to relate organic concentrations to other compositional or physical variables. The low levels of organics may make further investigation unnecessary.

Particulate material, although in suspension rather than in solution, can significantly affect concentrations of dissolved material. The particulate content of water from only one well near Yucca Mountain (Well J-13) has been examined. This well was chosen because it has been used for over 20 years and the particulate material produced during drilling should no longer contaminate the water. Water from Well J-13 was pumped through filters during a 14-day test. Two size fractions of solids were collected, $>0.4 \mu\text{m}$, and 5 nm to $0.4 \mu\text{m}$. Based on the quantity of water filtered and the mass of solids collected, the average concentration of the larger size fraction was approximately 3×10^{-5} g solids/l water; the smaller size fraction was estimated to be present as approximately 6×10^{-7} g solids/l water. Analysis of the cation content of the larger size fraction showed 60 wt% silicon, 20 wt% iron, 11 wt% calcium, and 4 wt% aluminum. The smaller size fraction contained 44 wt% sodium, 42 wt% silicon, 8 wt% calcium, and 4 wt% iron. Aside from the iron, these compositions could easily result from particulates of local minerals. The relatively large amount of iron in both groups may result from particulate material from the pumping and piping system of the well. A more complete description of this work, along with a discussion of the possible influence of particulate concentrations at this level on radionuclide transport, is contained in Appendix B of this report. There are no other data on particulate concentrations in water from Yucca Mountain, NTS, and vicinity available for comparison.

As expected, the temperature of water from wells at Yucca Mountain, NTS, and vicinity tends to increase with increasing depth of the water sample. In addition to this variation, the temperature of water from wells at Yucca Mountain and vicinity is correlated with PMC and total carbonate content of the water (see Figs. 21(a) and 21(b)). Part of this relation is from water sample depth. For example, the carbonate water from UE-25p#1 is the deepest sample, has the lowest PMC, and has the highest temperature; water from UE-29a#2 is the shallowest sample, has the highest PMC, and has the lowest temperature. However, for the intermediate samples, water temperature is not a function of sample depth, well depth, or depth to the static water level. Water from H-3 seems to have an anomalously low temperature (see Figs. 21). The trends of temperature with PMC and carbonate content seen in Figs. 21 are similar to the trends of $\delta^{13}\text{C}$, pH, and fluoride content seen in Figs. 15(b), 16(a), and 20(b).

D. Relations with Location or Depth

During a discussion of the relative sodium content ($\text{Na}/(\text{Na}+\text{Ca}+\text{K})$) of water from Yucca Mountain and vicinity (Section VI(A)), a relation between relative sodium content and well location was mentioned. Figure 22 shows a map of the Yucca Mountain area with locations of wells near the Exploration Block shown; associated with each well is a value or range of values of $\text{Na}/(\text{Na}+\text{Ca}+\text{K})$ for that well. There is a definite east to west trend in this variable. Wells west of the Ghost Dance Fault (USW H-3, H-5, and H-6) show high relative sodium content, 0.94 to 0.98, compared to wells directly to the east of the fault (USW G-4, H-4, and UE-25b#1), 0.69 to 0.79. This trend of low relative sodium content in the east and high in the west also includes the other Yucca Mountain wells. The high relative sodium content of water from wells west of the Ghost Dance Fault is more a result of the low calcium content of these waters that was noted in Section VI(A) than from a high sodium content (see Fig. 23). The reasons for the trends in calcium and sodium content are not evident from these data alone.

Figure 24 shows a map of the Yucca Mountain area in which the total carbonate contents are noted. Wells to the south and west (USW H-3, H-4, and H-6) tend to have higher carbonate contents than wells to the north and east. This trend is not as well defined as that noted above for relative sodium content. In particular, the Ghost Dance Fault no longer represents a dividing line for these data. Because of the relations between PMC and $\delta^{13}\text{C}$ and the




Fig. 21. Water temperature as a function of the percentage of modern carbon and total carbonate content at Yucca Mountain and vicinity.

- Water temperature as a function of the percentage of modern carbon.
- Water temperature as a function of total carbonate.




Fig. 22. $\text{Na}/(\text{Na} + \text{Ca} + \text{K})$ as a function of location for wells near Yucca Mountain.




Fig. 23. Calcium content as a function of sodium content at Yucca Mountain and vicinity.




Fig. 24. Total carbonate content as a function of location for wells near Yucca Mountain.

total carbonate content for wells at Yucca Mountain (see Figs. 14(b) and 15(b)), both $\delta^{13}\text{C}$ and δPMC show trends with location that are similar to that of total carbonate as seen in Fig. 24. The fluoride content of waters from Yucca Mountain also shows a trend with location that is similar to the total carbonate trend; that is, Wells USW H-3, H-4, and H-6 are high in fluoride compared to other nearby wells to the northeast. The relation between fluoride content and total carbonate was discussed in Section VI(C) and shown in Fig. 20(b). As with the trends in cation concentrations noted above, the reasons for the trends in carbonate content and associated variables are uncertain at this time.

Five wells at Yucca Mountain and vicinity have been pumped from two packed-off intervals or from one interval and the entire well bore (integral sample). Compositional variables from these five wells were examined to look for trends with depth. Table III lists the depth intervals sampled and the values of four variables (δD , δPMC , $\delta^{13}\text{C}$, and total carbonate) at these

TABLE III
VARIATION OF WATER PROPERTIES WITH DEPTH

<u>Parameter</u>	<u>Well</u>	<u>Integral^a Sample</u>	<u>Shallow Interval</u>	<u>Deep Interval</u>
Depth Interval	UE-25b#1	Yes	No	863-875
Sampled (m)	UE-25p#1	No	381-1197	1297-1805
	UE-29a#2	No	87-213	247-354
	USW H-1	No	572-687	687-1829
	USW H-6	Yes	608-646	753-835
δD ($^{\circ}/\text{‰}$)	UE-25b#1	-99.5 to -101	-	-99.5
	UE-25p#1	-	-106	-106
	UE-29a#2	-	-93	-93.5
	USW H-1	-	-103	-101
	USW H-6	-106	-107	-105
PMC (%)	UE-25b#1	16.7	-	18.9
	UE-25p#1	-	3.5	2.3
	UE-29a#2	-	60.0	62.3
	USW H-1	-	19.9	23.9
	USW H-6	16.3	12.4	10.0
$\delta^{13}\text{C}$ ($^{\circ}/\text{‰}$)	UE-25b#1	-10.4 to -10.7	-	-8.6
	UE-25p#1	-	-4.2	-2.3
	UE-29a#2	-	-13.1	-13.0
	USW H-1	-	-	-11.4
	USW H-6	-7.5	-7.1	-7.3
Total Carbonate (mmoles/l)	UE-25b#1	2.3-2.8	-	2.2
	UE-25p#1	-	4.6	9.3
	UE-29a#2	-	1.8	1.8
	USW H-1	-	1.9	2.0
	USW H-6	3.0	3.8	3.6

^aEntire well bore pumped.

intervals. The variation of some variables (PMC, $\delta^{13}\text{C}$, and total carbonate) for UE-25p#1 is understandable because the deep interval sampled the carbonate aquifer and the shallow interval sampled a mixture of carbonate and tuffaceous water (Craig and Robison 1984). However, there is no obvious trend in the data for the other four wells, which sample the tuffaceous aquifer at all the intervals sampled. The same conclusion holds for other compositional variables not listed in Table III.

During a 28-day pumping test of the 863-875 m packed-off interval (Bullfrog Member) of UE-25b#1, a significant variation in a number of the compositional variables was observed (Daniels et al. 1983; Rundberg et al. 1985). Dissolved oxygen and Eh increased and total iron, manganese, and nitrite decreased with time after the first few days; alkalinity also showed a slight drop. Major cation concentrations were relatively constant. One possible explanation for these results is that the water sampled early in the test is representative of the interval pumped, but with continued pumping, water was being drawn from other vertical locations (Daniels et al. 1983). If this is true, the significance of pumping packed-off intervals to investigate variation of water composition with depth at Yucca Mountain is uncertain. Similar pumping tests of two intervals from Well USW H-6, the 608- to 646-m interval (Bullfrog Member) and the 753- to 835-m interval (Tram Unit), gave water compositions that were essentially constant with time and very similar (Ogard and Vaniman 1985).

Another technique was used to sample Wells USW H-1 and H-4 (Ogard and Kerrisk 1984). Water samples were taken from static holes after the completion of pumping tests by lowering evacuated, stainless-steel bottles to selected depths. (These samples were already mentioned in Section VI(A) during the discussion of redox conditions.) The USW H-1 samples were taken at four depths, about a year after the pumping test was completed. Concentrations of some major species varied considerably from sample to sample (factors of 2 to 4 for calcium, potassium, silicon, chloride, and sulfate); the static-sample compositions also differed from the integral sample taken during the pumping test on this well. There was no consistent variation in the USW H-1 static samples with depth. The USW H-4 static samples were taken at eight depths, about a week after completion of the pumping test. Concentrations of the major species were similar at all depths and generally similar to the integral sample taken during the pumping test. The similarity of all the USW H-4 compositions may result from mixing during pumping that did not have time to equilibrate with local formation water in one week. Ogard and Kerrisk (1984) did not interpret the static sample compositions in terms of the water chemistry; they recommended further static sampling as a function of time after the completion of pumping tests before interpretation.

There are two reports of measurements of water compositions of near surface water (within a few meters of the surface) in conjunction with

measurements at depth. At Rainier Mesa, Benson (1976) measured compositions of two surface samples along with pore-water compositions at 130- to 530-m depth. Also at Rainier Mesa, Henne (1982) measured surface water compositions and fracture water compositions from tunnel samples. Figure 25(a) shows a ternary plot of the sodium-calcium-potassium composition of the water analyzed by Benson (1976), separated into surface samples and two depth intervals. Figure 25(b) shows a similar plot for the data of Henne (1982), separated into surface samples and samples at depth. In both cases there is a tendency toward increasing relative sodium content ($Na/(Na+Ca+K)$) with increasing depth. This is also a tendency toward increasing maturity of water in the process of glass and mineral dissolution and secondary mineral precipitation (White et al. 1980; Kerrisk 1983; Claassen 1985). As noted in Section VI(A) in the discussion of aqueous silica, the surface samples taken by Benson (1976) and Henne (1982) also have about half the aqueous silica content of waters at depth. This bias also indicates that dissolution continues with increasing depth. The two surface samples analyzed by Benson (1976) had much lower chloride and sulfate contents (0.05 to 0.08 mmoles/l) than the pore-water samples and than essentially all other waters reviewed in this report; this trend did not occur with the surface samples of Henne (1982). The total carbonate content and pH of the surface waters did not differ from other tuffaceous waters.


VII. CONTROLS ON GROUNDWATER COMPOSITION

In the previous section, relations were presented among many of the compositional variables of water from Yucca Mountain, NTS, and vicinity. These relations were primarily presented in the form of plots of one variable as a function of another. Very little effort was made in the previous section to connect the various relations observed with each other or with physical or chemical processes. This section describes a number of processes that appear to control water composition at Yucca Mountain and vicinity and relates these processes to the variations among compositional variables described in the previous section.

A. Control of Cation Content, Aqueous Silica Content, and pH of Tuffaceous Water

The primary cations in tuffaceous water at Yucca Mountain and vicinity are sodium, calcium, potassium, and magnesium. A number of studies in nearby

(a) Rainier Mesa Pore Water



(b) Rainier Mesa Fracture Water




Fig. 25. Ternary diagrams of relative sodium, calcium, and potassium content at Rainier Mesa.

- Surface water and pore water (Benson 1976).
- Surface water and fracture water (Henne 1982).

and similar systems have led to the general conclusion that glass and mineral dissolution, accompanied by secondary mineral precipitation, provides the primary control on cation composition of tuffaceous water (Hoover 1968; Winograd and Thordarson 1975; Claassen and White 1979; White et al. 1980; Moncure et al. 1981; Kerrisk 1983; Claassen 1985). The dissolution process represents an exchange of H^+ from the water for cations in the solid, as well as a breakup of the silicate structure of the solid. The reaction-path calculations reported by Kerrisk (1983) indicate that early in the dissolution process, dissolution rates control the relative concentrations of sodium, calcium, potassium, and magnesium. As dissolution proceeds, calcium, potassium, and magnesium are incorporated into various secondary mineral precipitates such as zeolites and clays, thus decreasing their concentrations. The presence of these secondary minerals at Yucca Mountain (Bish and Vaniman 1985; Broxton et al. 1986) gives additional evidence that this process has been active. The net result of the dissolution and precipitation processes is an increase in relative sodium content ($Na/(Na+Ca+K)$) of the water as well as an increase in pH. The variation of pH with relative sodium content of water from Yucca Mountain and vicinity (see Fig. 5) is consistent with this process. Concentrations of calcium may also be limited by calcite solubility in the high-pH water in the western part of Yucca Mountain (see Fig. 10) and may be influenced by other supplies of calcium in the eastern part (see below). Water-composition data show that aqueous silica concentrations are near saturation with respect to cristobalite. Reaction-path calculations indicate that these high aqueous-silica activities are necessary for the stability of the zeolites found at Yucca Mountain (Kerrisk 1983).

Surface samples of water taken at Rainier Mesa demonstrate that this dissolution process starts at the surface. However, the tendency of the surface waters to have higher relative calcium contents ($Ca/(Na+Ca+K)$) and lower aqueous silica contents than water at depth (see Figs. 25) suggests that this process is continuing as water moves down through the unsaturated zone. The moderate cation concentrations of water from the saturated zone of the tuffaceous aquifer at Yucca Mountain (see Figs. 2 and 3) indicate that evaporation is not an important process for controlling concentrations. Ogard and Kerrisk (1984) have proposed that the pH of water in the tuffaceous

aquifer at Yucca Mountain is buffered by aqueous carbonate and the minerals present.

There is an east to west variation in $\text{Na}/(\text{Na}+\text{Ca}+\text{K})$ of tuffaceous-aquifer water at Yucca Mountain (see Fig. 22) that is primarily caused by higher calcium content waters in the east than in the west. There is an opposite trend in total carbonate content; it is higher in the southwest than in the east (see Fig. 24). These two trends combine to result in water in the eastern part of Yucca Mountain (for example, Wells J-12, J-13, UE-25b#1, UE-25c#1, UE-25c#2, and UE-25c#3) with higher calcium and lower carbonate contents than water in the west (for example, USW H-3 and H-6). The variation in calcium content of Yucca Mountain water coincides with the east-west variation in calcium content of zeolites at Yucca Mountain described by Broxton et al. (1986). They state that the variation in calcium content of zeolites probably developed during initial stages of zeolite formation (11 to 14 million years ago) and that the enrichment of calcium in the eastern zeolites may have been caused by water from the carbonate aquifer mixing with tuffaceous water in that area. Based on a discussion of the carbonate content of water from Yucca Mountain (see section VII(B) below), mixing of tuffaceous- and carbonate-aquifer waters does not appear to be occurring in water recently sampled. Thus, carbonate aquifer water is not causing the higher calcium content of tuffaceous-aquifer water presently found in the eastern part of Yucca Mountain. However, the eastern zeolites provide a source of calcium and that source is independent of carbonate; thus, it is possible to have the eastern water higher in calcium but lower in total carbonate than water in the west if calcium is supplied to water presently in the eastern part of Yucca Mountain by cation exchange with the high calcium content zeolites. This mechanism provides an example of mineral compositions established in the past that influence present-day water compositions.

B. Control of Carbonate Content of Tuffaceous Water

Carbonate in water in the tuffaceous aquifer is normally considered to come from soil-zone carbon dioxide (Claassen 1985; White and Chuma 1986). This hypothesis is generally consistent with calculated carbon-dioxide pressures of tuffaceous water from Yucca Mountain and vicinity (see Fig. 8(b)), and with measured carbon-dioxide pressures in the gas phase of the unsaturated zone at Yucca Mountain (Yang et al. 1985). There is, however, a range of carbonate contents in the tuffaceous waters at Yucca Mountain (1.8

to 4.5 mmoles/l) that is larger than one might expect if passage through the soil zone early in the recharge cycle is the only source. There are also relations between carbonate content and a number of other compositional variables that point toward a more complex picture (see Figs. 2(b), 3(b), 7, 9, 12(a), 14(b), 15(b), 18(b), 19(b), 20(b), and 21(b)). In particular, tuffaceous-aquifer waters with higher carbonate contents tend to have higher pH (Fig. 7), lower carbon-dioxide pressures because of the shift in carbonate equilibria with pH (Fig. 9), and lower calcium contents (Fig. 3(b)). There are two sources of carbonate available to these waters: carbon dioxide (from the atmosphere, soil zone, or unsaturated-zone gas phase) and the carbonate aquifer. Some of the aqueous carbonate undoubtedly comes from carbon dioxide in the atmosphere and the soil zone as precipitation is recharged into the groundwater system. The relations between carbonate content and other compositional variables point to carbon dioxide as a more likely source of the additional carbonate in some waters at Yucca Mountain for three reasons. First, as pH increases, the equilibrium carbon-dioxide pressure drops, creating a driving force for dissolution of carbon dioxide from the gas phase. Second, carbonate in water from the carbonate aquifer would bring along calcium, which is quite low in concentration in the higher carbonate waters and in minerals in the western part of Yucca Mountain (Broxton et al. 1986), where the higher-carbonate waters are generally located. Third, it is unlikely that mixing low-pH water from the carbonate aquifer (UE-25p#1 carbonate water) with dilute, intermediate-pH water from the tuffaceous aquifer is the cause of high-pH water in the tuffaceous aquifer. The conclusion that gaseous carbon dioxide is the source of additional aqueous carbonate at Yucca Mountain wells does not necessarily extend to other waters examined in this report. In particular, there are tuffaceous-aquifer wells at Pahute Mesa (UE-19c and UE-19d) with high total carbonate, intermediate pH, and moderate to high calcium content that could result from mixing of tuffaceous and carbonate waters.

The reasoning of the previous paragraph relates an increase in carbonate content to an increase in pH of the water. As discussed above in Section VII(A), and indicated in Fig. 5, the higher pH waters from the tuffaceous aquifer at Yucca Mountain tend to have higher relative sodium content ($Na/(Na+Ca+K)$), consistent with the process of glass and mineral dissolution and precipitation of secondary minerals as the cause (Claassen 1985; Kerrisk

1983). This combination of glass and mineral dissolution, increasing water pH, and carbon-dioxide dissolution can be considered as mineral dissolution in a system that is open with respect to carbon dioxide. Mass transfer probably limits carbon-dioxide contents of deeper parts of the saturated zone, so that the entire saturated zone may only be partially open with respect to carbon dioxide. This may be the reason that calculated carbon-dioxide pressures of saturated zone water do not approach a limit with increasing pH (see Fig. 8(b)). Claassen (1985) mentions that mineral dissolution in a system that is open with respect to carbon dioxide is a likely process in surface or shallow saturated-zone conditions. However, at Yucca Mountain, most of the tuffaceous-aquifer wells have deep unsaturated zones (300 to 700 m). Although gaseous carbon dioxide has been observed in the deep unsaturated zone at Yucca Mountain (Yang et al. 1985), it is surprising that it could act as a source of aqueous carbonate.

Some additional information about this process can be obtained from the isotopic carbon data of Yucca Mountain waters (see Figs. 14(b) and 15(b)). In the Section VI(B) discussion of the variation of both ^{14}C (measured as percentage modern carbon, PMC) and ^{13}C (measured as $\delta^{13}\text{C}$) contents of Yucca Mountain water with total carbonate content, the possibility was mentioned that the observed variation could result from mixing tuffaceous and carbonate waters. This process was proposed because of the continuity of the tuffaceous and carbonate data seen in Figs. 14(b) and 15(b). However, based on the model described above of glass and mineral dissolution in a system open or partially open to carbon dioxide, mixing of carbonate and tuffaceous waters is not considered likely at Yucca Mountain. In an attempt to see if the model of carbon-dioxide addition to saturated-zone water could explain the variation of PMC and $\delta^{13}\text{C}$ with total carbonate content, a simplified carbon-balance model was developed. The model is based on the assumption that an initial charge of carbonate (probably from the soil zone) is present in the water and that carbonate from another source (carbon dioxide in the gas phase of the unsaturated zone) is added at some rate as time progresses. The rate of change of the isotope (^{14}C or ^{13}C) content of the water is the sum of three components, (1) addition with the added carbonate, (2) addition or loss by isotopic exchange between the water and the carbon-dioxide source, and (3) loss by radioactive decay (for ^{14}C only). If x is the isotope concentration

(PMC for ^{14}C , or $\delta^{13}\text{C}$ for ^{13}C) in the water and C is the total carbonate content of the water at time t , then

$$\frac{d(xC)}{dt} = x_s (dC/dt) - \gamma(x - x_s) - \lambda x C, \quad (2)$$

where x_s is the isotope concentration of the carbon-dioxide source (assumed constant), λ is the radioactive decay constant ($\lambda = 1.24 \times 10^{-4}/\text{yr}$ for ^{14}C and 0 for ^{13}C), and γ is a constant relating the rate of isotopic exchange to the concentration difference between the water and the carbon-dioxide source. By expanding the derivative on the left, Eq. (2) can be rewritten as

$$\frac{dx}{dt} = -(1/C)[\gamma + (dC/dt)](x - x_s) - \lambda x. \quad (3)$$

In Eqs. (2) and (3), the quantity (dC/dt) represents the addition rate of carbonate to the water, and C is related to t by

$$C = C_0 + \int_0^t (dC/d\tau)d\tau, \quad (4)$$

where C_0 is the initial carbonate content of the water ($t = 0$). Equation (3) is a linear, first-order differential equation with nonconstant coefficients. An initial condition, specifying the initial isotope concentration ($x = x_0$ at $t = 0$), is required in addition to the initial condition on total carbonate content.

The rate of addition of carbonate to the water that is needed to solve Eq. (3) should be a function of the difference between the gas-phase carbon-dioxide pressure and the equilibrium carbon-dioxide pressure over the water. No attempt was made to model this aspect of the process; instead, isotope concentrations calculated using a variety of different carbonate addition rates were compared with the observed variation of PMC or $\delta^{13}\text{C}$ with carbonate content. Even for the simplest case considered, dC/dt constant, a closed form solution to Eq. (3) was not found; the results discussed here were obtained from a numerical solution. Initial attempts to use a constant carbonate addition rate gave poor agreement between calculated and observed PMC values at Yucca Mountain; the calculated results did not drop as steeply as the observations at low carbonate content (see Fig. 14(b)). It was found

that a carbonate addition rate that increased with time gave much better results. For calculational purposes, a carbonate addition rate of

$$\frac{dC}{dt} = \alpha e^{(\beta t)}$$

was used, where α and β are positive constants. With this form, Eq (4) gives

$$C = C_0 + (\alpha/\beta)[e^{(\beta t)} - 1]. \quad (5)$$

For the ^{14}C data, reasonable comparisons between calculated and observed PMC values could be obtained with ($\gamma > 0$) or without ($\gamma = 0$) isotopic exchange. However, for the ^{13}C data, calculated results without isotopic exchange did not compare well with all the observations. Figures 26 and 27 show plots of PMC and $\delta^{13}\text{C}$ as a function of total carbonate for water from Yucca Mountain and vicinity; the data points plotted are the same as those in Figs. 14(b) and 15(b) and the lines represent calculated results from this model. Table IV lists the values of the parameters used to obtain the calculated results in these figures. The same definition of the carbonate addition rate (values of α and β) and initial carbonate content (C_0) were used for all the calculations. The model fits the shape of the PMC variation with carbonate content (Fig. 26) reasonably well. Acceptable values for x_s (the PMC value of the carbon dioxide in the gas phase of the unsaturated zone) decrease from about 75% to about 40% as isotopic exchange increases (γ increases). Yang et al. (1985) observed values of PMC of 60 to 80% for carbon dioxide in the gas phase of the unsaturated zone at 20- to 150-m depth in Yucca Mountain. Above 20-m depth the effects of modern ^{14}C were seen (PMC > 100%), and below 150-m depth there was more scatter, the data ranging from 50 to 100% PMC. Attempts to use very low values of x_s (such as might result from the carbonate aquifer being the source of carbon) did not produce a good comparison between the model and the data.

The relation between the model and the variation of $\delta^{13}\text{C}$ of Yucca Mountain water with carbonate content is not as clear (see Fig. 27). A group of wells with $\delta^{13}\text{C}$ of about -7‰ (J-13, UE-25c#1, UE-25c#2, USW 4-4, and H-6) span a range of carbonate contents from about 2 to 4 mmoles/l. A constant $\delta^{13}\text{C}$ with increasing carbonate content could be accommodated by the model if $x_0 = x_s = -7\text{‰}$ (a value of x_s of -7‰ would correspond to $\delta^{13}\text{C}$ of about -14 to




Fig. 26. Percentage of modern carbon as a function of total carbonate content at Yucca Mountain with carbon-model prediction (see Table IV for solid and dashed curve definitions).




Fig. 27. $\delta^{13}\text{C}$ as a function of total carbonate content at Yucca Mountain with carbon-model prediction (see Table IV for solid and dashed curve definitions).

TABLE IV
CARBON MODEL PARAMETERS USED IN FIGURES 26 AND 27

<u>Constant</u>	Figure 26		Figure 27	
	<u>Solid Curve</u>	<u>Dashed Curve</u>	<u>Solid Curve</u>	<u>Dashed Curve</u>
C_0 (mmoles/l)	1.65	1.65	1.65	1.65
α (mmoles/l.yr)	3.0×10^{-5}	3.0×10^{-5}	3.0×10^{-5}	3.0×10^{-5}
β (1/yr)	2.5×10^{-5}	2.5×10^{-5}	2.5×10^{-5}	2.5×10^{-5}
γ (mmoles/l.yr)	0.0	1.0×10^{-4}	0.0	1.0×10^{-4}
λ (1/yr)	1.24×10^{-4}	1.24×10^{-4}	0.0	0.0
x_s	75.0%	40.0%	$-4.0^{\circ}/\text{oo}$	$-4.0^{\circ}/\text{oo}$
x_0	100.0%	100.0%	$-13.0^{\circ}/\text{oo}$	$-15.0^{\circ}/\text{oo}$

$-15^{\circ}/\text{oo}$ for the carbon dioxide in the gas phase because of fractionation between the aqueous (pH 7 to 8) and gas phases). However, data from other wells are not consistent with this assumption. The curves in Fig. 27 show how $\delta^{13}\text{C}$ of the water would vary from a relatively negative value as carbonate with $\delta^{13}\text{C} = -4^{\circ}/\text{oo}$ (corresponding to carbon dioxide with $\delta^{13}\text{C}$ of about -11 to $-12^{\circ}/\text{oo}$) is added. Without isotopic exchange, the model does not fit the data well; the inclusion of isotopic exchange is needed to get a steep rise in $\delta^{13}\text{C}$ at low carbonate contents. Yang et al. (1985) observed $\delta^{13}\text{C}$ values for carbon dioxide in the gas phase of the unsaturated zone at Yucca Mountain of -10 to $-18^{\circ}/\text{oo}$ below about 60-m depth. The choice of x_s for ^{13}C is consistent with these observations.

This choice of parameters in Fig. 26 (particularly α and β) results in a time of approximately 49,000 yr to achieve 4.5 mmoles/l total carbonate content and 11% PMC, which are characteristic of USW H-3. The apparent age of H-3 water based on the measured PMC of 10.5% is 18,100 yr. The model requires 10,000 yr to achieve 2.0 mmoles/l total carbonate and 30% PMC, which are characteristic of J-13; the apparent age of J-13 water based on the measured PMC of 29.2% is 9,900 yr. The net result of this model is that for water with less than about 2 mmoles/l total carbonate, the age of the water is essentially the apparent age. However, as the carbonate content rises above 2

mmoles/l, the apparent age underestimates the true age. The continual addition of younger carbon to the water means that as the carbonate content increases, larger corrections are required to the apparent age. This situation contrasts with what would exist if the variation of PMC with carbonate content was caused by the addition of carbonate water or old carbon dioxide (PMC = ~0%) to the tuffaceous water; the apparent age of the mixture would then overestimate the true age.

The variation of δD with PMC and total carbonate content of water from Yucca Mountain and vicinity was shown in Figs. 18. Claassen (1985) interpreted a similar variation of δD with apparent age for water from the Amargosa Desert and Fortymile Wash to indicate that older water (apparent age, about 15,000 yr) precipitated under colder conditions (see Section VI(B)). The carbonate model proposed here predicts that the true ages of water with higher carbonate contents would be older than their apparent ages. Figure 28(a) shows a plot of δD as a function of apparent age for water from Yucca Mountain and vicinity. This plot uses the data plotted in Fig. 18(a) with apparent age calculated from Eq. (1); the data from UE-25p#1 were not included because carbonate derived from Paleozoic carbon would require corrections to be comparable to carbonate in tuffaceous waters. Figure 28(b) shows the same data plotted as a function of the carbon-model age, where the carbon-model age was calculated from the total carbonate content using Eq. (5) and the values of α and β shown in Table IV. In Fig. 28(a), the data bunch up in the 15,000- to 20,000-yr apparent age range; in Fig. 28(b), this range of apparent ages is spread out over 20,000- to 50,000-yr carbon-model age. Assuming the relation of δD with precipitation temperature, the application of the carbon model to the Yucca Mountain carbonate data indicates that colder temperatures (lower values of δD) existed 20,000 to 50,000 yr ago. Spaulding (1985) estimated that colder average-annual temperatures existed in the vicinity of NTS from 18,000 to 38,000 yr ago, with somewhat warmer temperatures (still below modern temperatures) from 38,000 to 45,000 yr ago. Thus, the carbon model proposed here is generally consistent with those estimates.

The results of applying the carbon model to Yucca Mountain data should probably not be interpreted quantitatively. The parameters in Table IV were chosen to show that the model was generally consistent with the variation of PMC and $\delta^{13}\text{C}$ with total carbonate content and with the proposal that colder





Fig. 28. δD as a function of apparent age and carbon-model age at Yucca Mountain.

a. δD as a function of apparent age.

b. δD as a function of carbon-model age.

temperatures were the cause of the variation of δD with PMC. Even though the carbon-model ages may not be quantitatively accurate, the concept that water with higher total carbonate contents (for example, USW H-3, H-4, and H-5) may be older than the apparent age of the water is quite plausible. Claassen (1985) observed that groundwaters with apparent ages older than about 17,000 yr were not present in the tuffaceous aquifers of the Amargosa Desert area he studied. Apparent ages of water from the tuffaceous aquifer at Yucca Mountain are also less than about 20,000 yr (see Fig. 28(a)). Claassen's preferred explanation for this observation was that snowfall earlier than about 20,000 yr ago was insufficient to result in recharge (Claassen 1985; Spaulding 1985). The discussion of aqueous carbonate in water from the tuffaceous aquifer at Yucca Mountain and vicinity that was presented here provides an alternate explanation for this observation. That explanation is that the apparent age of Yucca Mountain water with more than about 2 mmoles/l total carbonate tends to underestimate the true age. Thus, the model predicts that water older than 20,000 yr is present at Yucca Mountain, but corrections are required to the apparent age.

C. Control of Chloride and Sulfate Content of Tuffaceous Water

Precipitation is usually considered a primary source of chloride and sulfate in the water around Yucca Mountain, NTS, and vicinity. Under some conditions both chloride and sulfate are considered as conservative species, that is, having no sources or sinks in the groundwater system (Claassen 1985). In other situations, a source of sulfate has been proposed for some waters. Winograd and Thordarson (1975) proposed dissolution of gypsum in the basal strata of Tertiary rocks as a source for the lower carbonate aquifer and some particular wells. White (1979) proposed a hydrothermal source of sulfate in Oasis Valley. Young (1972) also proposed hydrothermal alteration as the source of high sulfate concentrations in water from one well in eastern Jackass Flats, near the Calico Hills. Most of the tuffaceous waters at Yucca Mountain and vicinity discussed in Section VI(A) have low chloride (0.16 to 0.31 mmoles/l) and sulfate (0.17 to 0.32 mmoles/l) concentrations (see Figs 11, 12, and 13). One well near Yucca Mountain, USW VH-1 in Crater Flat, has somewhat higher sulfate content (0.45 mmoles/l) than the other tuffaceous waters and may indicate a minor, local source of sulfate in Crater Flat. Compared to other areas such as Oasis Valley, evaporation does not appear to be important in the saturated zone of the tuffaceous aquifer at Yucca Mountain.

Both chlorine and sulfur are present in rocks at Yucca Mountain. Warren and Broxton (1986) have measured the chlorine content of a large number of core samples from NTS; an average chlorine content is about 500 ppm. They also observed that zeolitized tuff has little or no chlorine left. Vaniman (1986) has reported the sulfur content of Topopah Spring and Calico Hills tuff from drill hole USW G-4; values range from 46 to 137 ppm sulfur. Thus, the tuff may also represent a source of chloride and sulfate for water in the saturated zone. The sulfur/chlorine molar ratio of the tuff source is only about one-tenth the sulfate/chloride ratio in the water; for this reason, the tuff is probably only a minor source of these species at most.

Figure 12(a) showed an nearly linear relation between sulfate and total carbonate contents of water from Yucca Mountain and vicinity; chloride did not show a similar relation (see Fig. 12(b)). The carbon model described above (Section VII(B)) proposed gaseous carbon dioxide as the source of additional carbonate in this water. A source of sulfate added with the carbon dioxide does not seem likely. Other variables that may lead to this relationship are the higher pH and older age of water with added carbonate. A physical or chemical process that results in this relationship has not been identified at this time.

Based on the chloride and sulfate contents of saturated-zone water around Yucca Mountain, there seems no need of sources of these species other than precipitation. The difference between the average sulfate/chloride molar ratio of 0.82 for all the waters reviewed here (see Fig. 13(a)) and the ratio for modern precipitation of ~1.65 (Claassen 1985) may be that modern precipitation contains higher concentrations of sulfate and nitrate because of fossil fuel burning (Mayewski et al. 1986).

D. Control of the Redox State of the Water

Measurements of the redox state of water are available only for samples from Yucca Mountain and these data present a somewhat confusing picture. Most measurements indicate oxidizing conditions. However, measurements on some samples pumped from packed-off zones or taken from static wells show reducing conditions. This situation may be a sign of reducing conditions in deep saturated-zone water with oxidizing conditions near the static water level where interaction with oxygen in the unsaturated-zone gas phase is possible; vertical mixing during pumping may disturb this condition.

There is little information available about what reactions might contribute to reducing conditions in the deep saturated zone. Ogard and Kerrisk (1984) proposed that reducing conditions in the water were produced by reactions with organic matter as the water recharged through the soil zone. They further proposed that conditions in the past were suitable for these reactions to occur, but present conditions are not. Caporuscio and Vaniman (1985) have reported that iron is present in reduced form in rocks at Yucca Mountain, but that it appears to be generally inaccessible to the water. Thus, the iron is probably not a factor in controlling the redox state of the water.

It would be difficult to prove with the present data base that reducing conditions exist in deep saturated-zone water at Yucca Mountain. Regardless of the state of the deep saturated zone, water at the static water level appears to be oxidizing. Pore water or fracture water in the unsaturated zone will probably also be oxidizing. Thus, oxidizing conditions will be present in water over a significant portion of the transport path of radionuclides from the repository.

E. Control of Other Compositional Variables of Tuffaceous Water

Three other compositional variables discussed in Section VI(C) showed relations that may indicate controls on these variables. They are nitrate, fluoride, and water temperature.

The nitrate content of water from Yucca Mountain and vicinity decreases with decreasing PMC (increasing age) as seen in Fig. 19(a). The soil zone or precipitation are the most likely sources of nitrate in water recharged to the saturated zone. There are no aqueous-solid reactions that would provide a sink for nitrate in these waters. However, the reduction of nitrate in the saturated zone represents a possible sink. Some evidence for reducing conditions in water below the static water level at Yucca Mountain and for the variation of nitrate content with redox conditions in Well UE-25b#1 was given in Section VI(A). Although no proof is available, reduction of nitrate represents a plausible mechanism for decreasing nitrate content with age.

Figures 20 showed a tendency for fluoride content to increase with increasing sodium and carbonate content of water from Yucca Mountain and vicinity (see Section VI(C)). The fluoride content of these waters also showed a consistent variation with location similar to the total carbonate variation represented in Fig. 24 (see Section VI(D)). The $\log_{10}(Q/K)$ for

fluorite ranged from -1.9 (undersaturated) to 0.04 (just above saturation) for water from the tuffaceous aquifer; most waters had $\log_{10}(Q/K)$ of -0.6 or less. Fluorite has occasionally been observed in fractures at Yucca Mountain (Bish and Vaniman 1985), but not to the extent that it would represent a general mineralogical control on fluoride concentrations. A few analyses of the fluorine content of glass from NTS have been done; results range from 0 to 0.26 wt% fluorine with an average of 0.19 wt% (Warren and Broxton 1986). Compared with the 500 ppm chlorine content discussed above (Section VII(C)), there is considerably more fluorine than chlorine in the tuff (the fluorine/chlorine molar ratio is about 7). Thus, the tuff may act as a source of fluoride for the water. The increase of fluoride with increasing sodium and carbonate contents as well as $\text{Na}/(\text{Na}+\text{Ca}+\text{K})$ may be an indication that as glass dissolution and mineral precipitation reactions proceed, fluorine is also released to the water.

The temperature of water from Yucca Mountain and vicinity tended to increase with decreasing PMC (increasing age) and with increasing total carbonate content (see Figs. 21). The temperature of Well USW H-3 water did not fit this relationship. There appears to be some relation other than just depth that influences water temperature; however, it has not been identified at this time.

F. Controls on Unsaturated-Zone Water Composition

The only data on unsaturated-zone water compositions come from the work of Benson (1976) at Rainier Mesa. This water showed one unusual compositional feature, the chloride and sulfate contents were high relative to the total carbonate content (see Figs. 11). This led to lower calculated carbon-dioxide equilibrium pressures than most other waters in the same pH range (see Fig. 8(a)). The major cation composition was similar to other waters reviewed (see Fig. 4(a)). There were no ^{14}C , ^{13}C , ^{18}O , or deuterium contents available for these water samples; this limits the possible comparisons that can be made. The composition of the gas phase of the unsaturated zone at Yucca Mountain has also been measured. The overall chemical composition is similar to air (Yang 1986). Below the soil zone (top 20 to 30 m), carbon-dioxide pressures of $\sim 10^{-3}$ atm or less were measured (Yang et al. 1985).

For this discussion, the primary difference between processes that are important in the unsaturated and saturated zones is the presence of the gas phase in the unsaturated zone. Pore water in the unsaturated zone can react

with the rock or the gas-phase constituents and should present a relatively large surface area to both phases. The presence of the gas phase allows evaporation to occur more readily in the unsaturated zone, and the large gas-aqueous surface area means higher mass-transfer rates and closer approach to gas-aqueous equilibrium. Evaporation of pore water could be the mechanism for increasing chloride and sulfate concentrations. Also, carbon-dioxide equilibrium pressures in pore water should be closer to the gas phase carbon-dioxide pressure. The calculated carbon-dioxide equilibrium pressures of Rainier Mesa pore water (see Fig. 8(a)) are consistent with the carbon-dioxide pressures observed in the gas phase at Yucca Mountain. The pH of Rainier Mesa pore water is generally in the 7 to 8 range. The calcium content of this water is higher than most of the tuffaceous waters at Yucca Mountain, but consistent with other tuffaceous water such as in the Amargosa Desert (see Fig. 4(a)). The aqueous silica content is also similar to other tuffaceous waters. Thus, the pore water is probably undergoing the same glass and mineral dissolution and precipitation reactions as saturated-zone water in the tuffaceous aquifer. The extent of these reactions may be influenced by increased concentrations from evaporation.

The trends in unsaturated-zone compositions seen at Rainier Mesa may be the result of glass and mineral dissolution and precipitation reactions that are similar to those discussed for the saturated zone, accompanied by evaporation in a system that is open with respect to carbon dioxide. At present, this hypothesis is rather speculative. More compositional data, in particular some isotopic data, are needed to establish what processes control unsaturated-zone water compositions.

VIII. IMPLICATIONS FOR RADIONUCLIDE TRANSPORT

One of the primary reasons for studying water chemistry at Yucca Mountain and vicinity is to provide information for calculations of radionuclide transport. Water chemistry can influence waste-element solubility, speciation, and sorption, and through these processes, the rates at which waste elements are transported away from the repository. Water chemistry can also affect the stability and composition of minerals. This section discusses some implications of the water chemistry for solubility, speciation, and mineral stability.

The relative concentrations of major cations (sodium, calcium, and potassium) in water from the tuffaceous aquifer at Yucca Mountain and vicinity vary over a range of relative sodium and calcium contents at nearly constant relative potassium content (see Fig. 4(b)), with higher relative sodium content in the west than in the east (see Fig. 22). Broxton et al. (1986) see related variations in the sodium and calcium contents of zeolites. Sorption of some waste elements on zeolites is by ion exchange, in which the waste-element cation replaces an existing cation (such as sodium, calcium, or potassium) in the zeolite. At this time, no variations in zeolite sorptive behavior (measured values of the sorption coefficient) have been related to variations in cation contents of the zeolites (Daniels et al. 1982; Ogard and Vaniman 1985); a relation of this nature may be difficult to demonstrate because of the precision of the data. Solubilities of some waste elements may be affected to a minor extent by changes in cation content of the water if the solid controlling solubility contains one of the cations; an example of this is neptunium, where $\text{Na}_3\text{NpO}_2(\text{CO}_3)_2 \cdot n\text{H}_2\text{O}$ was identified as the solid that precipitated from neptunium solutions in Well J-13 water (Nitsche and Edelstein 1985). The range of major cation concentrations observed at Yucca Mountain and vicinity should not significantly affect solubility or sorption of waste elements; however, these effects are being examined by sorption and solubility experiments using various water compositions (Ogard and Vaniman 1985; Kerrisk 1985).

The process of glass and mineral dissolution in a system open or partially open with respect to carbon dioxide was proposed as an explanation for the variation in carbonate content of tuffaceous waters seen at Yucca Mountain. Based on this model, carbonate contents of saturated zone waters could vary depending on the extent of dissolution or age of the water. Because carbonate forms complexes and solids with some important waste elements such as americium, uranium, and neptunium (Ogard and Kerrisk 1984; Nitsche and Edelstein 1985), this variation may have significant effects. Solubility and sorption experiments using water compositions that cover a range of carbonate concentrations are being done by the NNWSI Project (Ogard and Vaniman 1985; Kerrisk 1985).

The chloride and sulfate contents of saturated-zone tuffaceous waters from Yucca Mountain and vicinity are relatively low compared to most other waters in this area (see Fig. 11(b)). Although chloride does not form strong

complexes with waste elements, sulfate does form complexes and solids that may affect speciation, sorption, or solubility (Ogard and Kerrisk 1984). Unsaturated-zone pore water from Rainier Mesa showed high chloride and sulfate content relative to the carbonate content of the water (see Fig. 11(a)). If this condition also occurs at Yucca Mountain, the effects of varying sulfate content on solubility, sorption, or speciation may have to be investigated.

The fluoride content of water from Yucca Mountain and vicinity varies over a relatively wide range (see Figs. 20), but fluoride is still a minor anion. Some waste elements form complexes with fluoride (Ogard and Kerrisk 1984), so that solubility and sorption experiments should also give consideration to this variation. The nitrate content also varies over a relatively wide range (see Figs. 19). However, nitrate does not readily form complexes or solids, so that variations of the nitrate content are not important for solubility or sorption.

The pH of water from the tuffaceous aquifer at Yucca Mountain and vicinity falls in the range of 7 to 9.2, with most samples in the 7 to 8 range. Both the carbonate content of the water and the local minerals buffer pH (Ogard and Kerrisk 1984). Variations in pH over this range can have a significant effect on solubility and speciation of some waste elements (Allard 1982; Apps et al. 1983). Solubility and sorption experiments using water compositions that cover a range of pH are being done by the NNWSI Project (Ogard and Vaniman 1985; Kerrisk 1985). Although there are some indications that water in deep regions of the saturated zone at Yucca Mountain may be reducing, this situation, which would lead to lower solubilities of many waste elements, could be difficult to prove. Water in the unsaturated zone and near the static water level is probably oxidizing. Solubility and sorption experiments in the NNWSI Project are being done under oxidizing conditions.

Except for a few samples of water taken near the ground surface, aqueous silica contents of water from Yucca Mountain, NTS, and vicinity are at or above cristobalite saturation. The relatively high aqueous silica content of these waters may affect solubilities of some waste elements (Mendel 1984). If waste-element silicates do control solubilities of any important waste elements under conditions at Yucca Mountain, this information will be obtained from the NNWSI solubility-experiments that are in progress (Kerrisk 1985). Kerrisk (1983) proposed that a high aqueous silica activity was needed for

stability of the zeolites found at Yucca Mountain. The stability of zeolites is also being investigated by the NNWSI Project (Wolfsberg and Vaniman 1984).

ACKNOWLEDGEMENTS

The author would like to thank Katherine Campbell of Los Alamos National Laboratory for her help in setting up the SAS® data base and discussing the variation of water composition with location. He would also like to thank Allen E. Ogard of Los Alamos National Laboratory and Art F. White of Lawrence Berkeley Laboratory for their helpful comments on an early draft of this report.

REFERENCES

Allard, B., "Solubilities of Actinides in Neutral or Basic Solutions," in Actinides in Perspective, N. M. Edelstein (Ed.), Pergamon Press, New York, 1982.

Apps, J. A., C. L. Carnahan, P. C. Lichtner, M. C. Michel, D. Perry, R. J. Silva, O. Weres, and A. F. White, "Status of Geochemical Problems Relating to the Burial of High-Level Radioactive Waste, 1982," Lawrence Berkeley Laboratory report LBL-15103 (NUREG/CR-3062), (March 1983).

Benson, L. V., "Mass Transport in Vitric Tuffs of Rainier Mesa, Nye County, Nevada," Water Resources Center, Desert Research Institute report NVO-1253-10 (October 1976).

Benson, L. V. and P. W. McKinley, "Chemical Composition of Ground Water in the Yucca Mountain Area, Nevada, 1971-84," U. S. Geological Survey Open-File Report 85-484 (1985).

Benson, L. V., J. H. Robison, R. K. Blankennagel, and A. E. Ogard, "Chemical Composition of Ground Water and the Location of Permeable Zones in the Yucca Mountain Area, Nevada," U. S. Geological Survey Open-File Report 83-854 (1983).

Bish, D. L. and D. T. Vaniman, "Mineralogic Summary of Yucca Mountain, Nevada," Los Alamos National Laboratory report LA-10543-MS (October 1985).

Blankennagel, R. K. and J. E. Weir, Jr., "Geohydrology of the eastern Part of Pahute Mesa, Nevada Test Site, Nye County, Nevada," U. S. Geological Survey Professional Paper 712-B (1973).

Broxton, D. E., D. L. Bish, and R. G. Warren, "Distribution and Chemistry of Diagenetic Minerals at Yucca Mountain, Nye County, Nevada," to be published in Clays and Clay Minerals (1986).

Caporuscio, F. A. and D. T. Vaniman, "Iron and Manganese in Oxide Minerals and in Glasses: Preliminary Considerations of Eh Buffering Potential at Yucca Mountain, Nevada," Los Alamos National Laboratory report LA-10369-MS (April 1985).

Claassen, H. C., "Water Quality and Physical Characteristics of Nevada Test Site Water-Supply Wells," U. S. Geological Survey report USGS-474-158 (January 1973).

Claassen, H. C., "Sources and Mechanisms of Recharge for Ground Water in the West-Central Amargosa Desert, Nevada - A Geochemical Interpretation," U. S. Geological Survey Professional Paper 712-F (1985).

Claassen H. C. and A. F. White, "Application of Geochemical Kinetic Data to Groundwater Systems," in Chemical Modeling - Speciation, Sorption, Solubility, and Kinetics in Aqueous Systems, E. A. Jenne (Ed.), Amer. Chemical Soc. Symposium Series 93, Amer. Chemical Soc. (1979).

Clebsch A., Jr., "Tritium-Age of Ground water at the Nevada Test Site, Nye County, Nevada," U. S. Geological Survey Professional Paper 424-C, Article 194 (1961).

Clebsch, A., Jr. and F. B. Barker, "Analysis of Ground Water from Rainier Mesa, Nevada Test Site, Nye County, Nevada," U. S. Geological Survey Trace Elements Investigations Report 763 (TEI-763) (May 1960).

Craig, H., "Isotopic Variations in Meteoric Waters," *Science*, 133, 1702-1703 (1961).

Craig, R. W. and J. H. Robison, "Geohydrology of Rocks Penetrated by Test Well UE-25p#1, Yucca Mountain Area, Nye County, Nevada," U. S. Geological Survey Water-Resources Investigations Report 84-4248 (1984).

Crowe, B. M. and D. T. Vaniman, "Research and Development Related to the Nevada Nuclear Waste Storage Investigations, January 1 - March 30, 1984," Los Alamos National Laboratory report LA-10154-PR (February 1985).

Czarnecki, J. B., "Simulated Effects of Increased Recharge on the Ground-Water Flow System of Yucca Mountain and Vicinity, Nevada-California," U. S. Geological Survey Water-Resources Investigations Report USGS/WRI-84-4344 (1985).

Daniels, W. R., K. Wolfsberg, R. S. Rundberg, A. E. Ogard, J. F. Kerrisk, C. J. Duffy, et al., "Summary Report on the Geochemistry of Yucca Mountain and Environs," Los Alamos National Laboratory report LA-9328-MS (December 1982).

Daniels, W. R., B. R. Erdal, and D. T. Vaniman, "Research and Development Related to the Nevada Nuclear Waste Storage Investigations, July 1 - September 30, 1982," Los Alamos National Laboratory report LA-9577-PR (March 1983).

Dansgaard, W., "Stable Isotopes in Precipitation," *Tellus*, XVI, 436-468 (1964).

Drever, J. I., The Geochemistry of Natural Waters, (Prentice-Hall, Inc., Englewood Cliffs, New Jersey, 1982).

Faure, G., Principles of Isotope Geology (John Wiley and Sons, New York, 1977), pp. 323-401.

Feth, J. H., "Nitrogen Compounds in Natural Water - A Review," *Water Resources Research*, 2, 41-58 (1966).

Fritz, P. and J. Ch. Fontes, Handbook of Environmental Isotope Geochemistry, Vol. 1, The Terrestrial Environment, A (Elsevier Scientific Publishing Co., New York, 1980).

Haas, H., D. W. Fisher, D. C. Thorstenson, and E. P. Weeks, " $^{13}\text{CO}_2$ and $^{14}\text{CO}_2$ Measurements on Soil Atmosphere Sampled in the Sub-Surface Unsaturated Zone in the Western Great Plains of the US," *Radiocarbon*, 25, 301-314 (1983).

Henne, M. S., "The Dissolution of Rainier Mesa Volcanic Tuffs, and Its Application to the Analysis of the Groundwater Environment," M. S. Thesis, University of Nevada, Reno (1982).

Hoover, D. L., "Genisis of Zeolites, Nevada Test Site," in Nevada Test Site, E. B. Eckel, Ed., Geological Society of America Memoir 110, 275-283 (1968).

Kerrisk, J. F., "Reaction-Path Calculations of Groundwater Chemistry and Mineral Formation at Rainier Mesa, Nevada," Los Alamos National Laboratory report LA-9912-MS (December 1983).

Kerrisk, J. F., "Solubility Experiments for the Nevada Nuclear Waste Storage Investigations Project," Los Alamos National Laboratory report LA-10560-MS (November 1985).

Lindberg, R. D. and D. D. Runnels, "Ground Water Redox Reactions: An Analysis of Equilibrium State Applied to Eh Measurements and Geochemical Modeling," Science, 225, 925-927 (1984).

Mayewski, P. A., W. B. Lyons, M. J. Spencer, M. Twickler, W. Dansgaard, B. Koci, C. I. Davidson, and R. E. Honrath, "Sulfate and Nitrate Concentrations for a South Greenland Ice Core," Science, 232, 975-977 (1986).

Means, J. L., A. S. Maest, and D. A. Crerar, "The Organic Geochemistry of Deep Groundwaters and Organo-Radionuclide Interactions at Hydrothermal Conditions," Office of Nuclear Waste Isolation report ONWI-448 (September 1982).

Mendel, J. E., (Comp.), "Final Report of the Defense High-Level Waste Leaching Mechanisms Program," Pacific Northwest Laboratory report PNL-5157 (August 1984).

Moncure, G. K., R. C. Surdam, and H. L. McKague, "Zeolite Diagenesis Below Pahute Mesa, Nevada Test Site," Clays and Clay Minerals, 29, 385-395 (1981).

Montazar, P. and W. E. Wilson, "Conceptual Hydrologic Model of Flow in the Unsaturated Zone, Yucca Mountain, Nevada," U. S. Geological Survey Water-Resources Investigations Report 84-4345 (1984).

Nitsche, H., and N. M. Edelstein, "Determination of the Solubilities and Complexation of Waste Radionuclides Pertinent to Geologic Disposal at the Nevada Tuff Site," Lawrence Berkeley Laboratory report LBL-18900 (January 1985).

Ogard, A. E., W. R. Daniels, and D. T. Vaniman, "Research and Development Related to the Nevada Nuclear Waste Storage Investigations, October 1 - December 31, 1982," Los Alamos National Laboratory report LA-9666-PR (May 1983).

Ogard, A. E. and J. F. Kerrisk, "Groundwater Chemistry Along Flow Paths Between a Proposed Repository Site and the Accessible Environment," Los Alamos National Laboratory report LA-10188-MS (November 1984).

Ogard A. E. and D. T. Vaniman, "Research and Development Related to the Nevada Nuclear Waste Storage Investigations, July 1 - September 30, 1984," Los Alamos National Laboratory report LA-10299-PR (May 1985).

Robison, J. H., "Ground-Water Level Data and Preliminary Potentiometric-Surface Maps, Yucca Mountain, Nye County, Nevada," U. S. Geological Survey Water-Resources Investigations Report 84-4197 (1984).

Rundberg, R. S., A. E. Ogard, and D. T. Vaniman, "Research and Development Related to the Nevada Nuclear Waste Storage Investigations, April 1 - June 30, 1984," Los Alamos National Laboratory report LA-10297-PR (May 1985).

Spaulding, W. G., "Vegetation and Climates of the Last 45,000 Years in the Vicinity of the Nevada Test Site, South-Central Nevada," U. S. Geological Survey Professional Paper 1329 (1985).

Stumm, W., and J. J. Morgan, Aquatic Chemistry (John Wiley and Sons, New York, 1981) p. 236.

Thorstenson, D. C., E. P. Weeks, H. Haas, and D. W. Fisher, "Distribution of Gaseous $^{12}\text{CO}_2$, $^{13}\text{CO}_2$, and $^{14}\text{CO}_2$ in the Sub-Soil Unsaturated zone of the Western US Great Plains," Radiocarbon, 25, 315-346 (1983).

Vaniman, D. T., "Progress Report on Research Conducted by the Earth and Space Science Division, Los Alamos National Laboratory, In Support of the Nevada Nuclear Waste Storage Investigations, January-June, 1985," Los Alamos National Laboratory report in preparation (1986).

Waddell, R. K., J. H. Robison, and R. K. Blankenagel, "Hydrology of Yucca Mountain and Vicinity, Nevada-California -- Investigative results through Mid-1983," U. S. Geological Survey Water-Resources Investigations Report 84-4267 (1984).

Warren, R. G. and D. E. Broxton, "Chemical Composition of Hydrated Glasses in Rock Units of the Nevada Test Site," Los Alamos National Laboratory report in preparation (1986).

White, A. F., "Geochemistry of Ground water Associated with Tuffaceous Rocks, Oasis Valley, Nevada," U. S. Geological Survey Professional Paper 712-E (1979).

White, A. F. and N. J. Chuma, "Isotope and Carbon Models of the Oasis Valley-Fortymile Canyon Ground Water Basin, Nevada," to be published in Water Resources Research (1986).

White, A. F., H. C. Claassen, and L. V. Benson, "The Effect of Dissolution of Volcanic Glass on the Water Chemistry in a Tuffaceous Aquifer, Rainier Mesa, Nevada," U. S. Geological Survey Water-Supply Paper 1535-Q (1980).

Wigley, T. M. L., L. N. Plummer, and F. J. Pearson, Jr., "Mass Transfer and Carbon Isotope Evolution in Natural Water Systems," Geochimica et Cosmochimica Acta, 42, 1117-1139 (1978).

Winograd, I. J. and W. Thordarson, "Hydrologic and Hydrochemical Framework, South-Central Great Basin, Nevada-California, with Special Reference to the Nevada Test Site," U. S. Geological Survey Professional Paper 712-C (1975).

Wolery, T. J., "EQ3NR, A Computer Program for Geochemical Aqueous Speciation-Solubility Calculations: Users Guide and Documentation," Lawrence Livermore National Laboratory report UCRL-53414 (April 1983).

Wolfsberg, K. and D. T. Vaniman, "Research and Development Related to the Nevada Nuclear Waste Storage Investigations, October 1 - December 31, 1983," Los Alamos National Laboratory report LA-10032-PR (August 1984).

Yang, In C., H. H. Haas, E. P. Weeks, D. C. Thorstenson, "Analysis of Gaseous-Phase Stable and Radioactive Isotopes in the Unsaturated Zone, Yucca Mountain, Nevada," Proc. National Water Well Association Conf. on Characterization and Monitoring the Vadose Zone, Denver, Colorado, November 19-21, 1985.

Yang, In C., U. S. Geological Survey, private communication (1986).

Young, R. A., "Water Supply for the Nuclear Rocket Development Station at the U. S. Atomic Energy Commission's Nevada Test Site," U. S. Geological Survey Water-Supply Paper 1938 (1972).

APPENDIX A
WATER CHEMISTRY DATA

This appendix contains a list of the water chemistry data reviewed for this report. The list is in the form of a SAS® output listing from the data base used to prepare the data for the plots contained in the report.

The various column headings in the data listing are described here. SITE refers to the well, spring, or sampling location where the water sample was obtained. REFERENCE refers to the source of the data; the references listed under this heading are in the reference section of the report. LOCATION refers to the location of the sampling site; the abbreviations used under this heading are listed in Table II of the report. DATE refers to the sampling date. INT SAMP (M) refers to the depth interval sampled in meters for wells; under this heading the entry INT refers to an integral well sample, in which the entire well bore was pumped, or to an integral sample taken from some other source such as a spring or seep. LITHOLOGY refers to the primary lithology of the well or the area sampled; under this heading T means tertiary rock, C means carbonate (Paleozoic) rock, QAL means quaternary alluvium, BULL means Bullfrog, and SOIL means soil zone. SURFACE ALTITUDE (M) refers to the ground surface elevation above sea level in meters at the well or sampling site. WELL DEPTH (M) refers to the depth of the well in meters. DEPTH TO WATER (M) refers to the depth to the static water level in the well in meters. TEMP (C) refers to the water temperature in °C. The headings CA, MG, NA, K, CL, SO4, SiO2, F, NO3, and HCO3, refer to the concentrations of calcium, magnesium, sodium, potassium, chloride, sulfate, aqueous silica, fluoride, nitrate, and total carbonate in mmoles/l. PH refers to water pH. O2 (MG/L) refers to dissolved oxygen in mg/l. DEL D refers to SD in %... DEL O18 refers to $\delta^{18}\text{O}$ in %... DEL C13 refers to $\delta^{13}\text{C}$ in %... PMC refers to percentage of modern carbon in %. LOG(CO2 PRESS) refers to \log_{10} (carbon dioxide pressure in atm). CALCITE LOG(Q/K), DOLOMITE LOG(Q/K), GYPSUM LOG(Q/K), MAGNESITE LOG(Q/K), and FLUORITE LOG(Q/K) refer to the \log_{10} (Q/K) for calcite, dolomite, gypsum, magnesite, and fluorite; the meaning of \log_{10} (Q/K) is discussed in Section VI(A) of this report and in Stumm and Morgan (1981). A blank or period (.) under any heading indicates that no data were available.

SITE	REFERENCE
WELL J-12	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL J-13	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL UE-25B#1	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL UE-25B#1	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL UE-25B#1	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL UE-25C#1	BENSON AND MCKINLEY (1985)
WELL UE-25C#2	BENSON AND MCKINLEY (1985)
WELL UE-25C#3	BENSON AND MCKINLEY (1985)
WELL UE-25P#1	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL UE-25P#1	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL UE-29A#2	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL UE-29A#2	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL USW G-4	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL USW H-1	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL USW H-1	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL USW H-3	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL USW H-4	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL USW H-5	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL USW H-5	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL USW H-6	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL USW H-6	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL USW H-6	BENSON AND MCKINLEY (1985), OGARD AND KERRISK (1984)
WELL USW VH-1	BENSON AND MCKINLEY (1985)
WELL USW VH-1	BENSON AND MCKINLEY (1985)
WELL USW VH-1	BENSON AND MCKINLEY (1985)
3. 15S/49E-22DC	CLAASSEN (1985)
4. 16S/49E-5ACC	CLAASSEN (1985)
5. 16S/49E-8ABB	CLAASSEN (1985)
6. 16S/49E-8ACC	CLAASSEN (1985)
7. 16S/49E-9CDA	CLAASSEN (1985)
8. 16S/49E-9DCC	CLAASSEN (1985)
9. 16S/49E-18DC	CLAASSEN (1985)
10. 16S/49E-16CCC	CLAASSEN (1985)
11. 16S/49E-19DAA	CLAASSEN (1985)
12. 16S/48E-24AAA	CLAASSEN (1985)
13. 16S/48E-25AA	CLAASSEN (1985)
14. 16S/48E-36AAA	CLAASSEN (1985)
15. 17S/48E-1AB	CLAASSEN (1985)
16. 17S/49E-7BB	CLAASSEN (1985)
17. 17S/49E-9AA	CLAASSEN (1985)
18. 17S/49E-8DD8	CLAASSEN (1985)
19. 17S/49E-15BBD	CLAASSEN (1985)
64. WELL 8, NTS	CLAASSEN (1985)
20. 17S/49E-350DC, ASH TR	CLAASSEN (1985)
21. 16S/49E-23ADD	CLAASSEN (1985)
23. 16S/48E-15AAA	CLAASSEN (1985)
25. 16S/48E-10CBA	CLAASSEN (1985)
27. 16S/50E-7BCD	CLAASSEN (1985)
29. 16S/49E-15AAA	CLAASSEN (1985)
30. 16S/49E-36AAA	CLAASSEN (1985)
45. 16S/48E-8BBA	CLAASSEN (1985)
46. 16S/48E-7BBA	CLAASSEN (1985)
47. 16S/48E-7CBC	CLAASSEN (1985)
48. 16S/48E-18BCC	CLAASSEN (1985)

SITE	REFERENCE
49, 16S/48E-17CCC	CLAASSEN (1985)
50, 16S/48E-18DAD	CLAASSEN (1985)
51, 16S/48E-8CDA	CLAASSEN (1985)
52, 16S/48E-17ABB	CLAASSEN (1985)
62, WHITEROCK SPRING	CLAASSEN (1985)
68, BUTTE SPRING	CLAASSEN (1985)
70, OAK SPRING	CLAASSEN (1985)
71, TOPOPAH SPRING	CLAASSEN (1965)
1, 10S/47E-14BAB	WHITE (1979)
2, 10S/47E-27CBA	WHITE (1979)
3, 10S/47E-31AAB	WHITE (1979)
4, 10S/47E-32DDA	WHITE (1979)
5, 10S/47E-33AAB	WHITE (1979)
6, 10S/47E-30DCC	WHITE (1979)
7, 11S/46E-26BBBB	WHITE (1979)
8, 10S/46E-26BCC	WHITE (1979)
9, 11S/47E-3CDB	WHITE (1979)
10, 11S/47E-4CAD	WHITE (1979)
11, 11S/47E-10CAA	WHITE (1979)
12, 11S/47E-10BCC	WHITE (1979)
13, 11S/47E-16DCD	WHITE (1979)
14, 11S/47E-16BDC	WHITE (1979)
15, 11S/47E-18ACD	WHITE (1979)
16, 11S/47E-21ACC	WHITE (1979)
17, 11S/47E-21DBB	WHITE (1979)
18, 11S/47E-21ABA	WHITE (1979)
19, 11S/47E-21ABA	WHITE (1979)
20, 11S/47E-27CBA	WHITE (1979)
21, 11S/47E-28AAC	WHITE (1979)
22, 11S/47E-28DAC	WHITE (1979)
23, 11S/47E-33BAC	WHITE (1979)
24, 11S/47E-10CCB	WHITE (1979)
25, 12S/47E-5CDA	WHITE (1979)
26, 12S/47E-6CDD	WHITE (1979)
27, 12S/47E-70BD	WHITE (1979)
28, 12S/47E-20BBB	WHITE (1979)
29, 12S/47E-19ADC	WHITE (1979)
18B, 12S/47E-20	WHITE AND CHUMA (1986)
19B, 12S/47E-20	WHITE AND CHUMA (1986)
20B, 12S/47E-20	WHITE AND CHUMA (1986)
21B, 12S/47E-20	WHITE AND CHUMA (1986)
22B, 12S/47E-20	WHITE AND CHUMA (1986)
WELL UE-19B-1	BLANKENNAGEL AND WEIR (1973)
WELL UE-19C	BLANKENNAGEL AND WEIR (1973)
WELL UE-19D	BLANKENNAGEL AND WEIR (1973)
WELL UE-19E	BLANKENNAGEL AND WEIR (1973)
WELL UE-19E	CLAASSEN (1973)
WELL UE-19GS	BLANKENNAGEL AND WEIR (1973)
WELL UE-19GS	CLAASSEN (1973)
WELL U-20A-2	BLANKENNAGEL AND WEIR (1973)
WELL U-20A-2	CLAASSEN (1973)
WELL UE-20D	BLANKENNAGEL AND WEIR (1973)
WELL UE-20E-1	BLANKENNAGEL AND WEIR (1973)
WELL UE-20H	BLANKENNAGEL AND WEIR (1973)

SITE	REFERENCE
WELL UE-20J	BLANKENAGEL AND WEIR (1973)
WELL ARMY-1	CLAASSEN (1973)
WELL 5A	CLAASSEN (1973)
WELL 5B	CLAASSEN (1973)
WELL 5C	CLAASSEN (1973)
WELL C	CLAASSEN (1973)
WELL C-1	CLAASSEN (1973)
WELL 3	CLAASSEN (1973)
WELL A	CLAASSEN (1973)
WELL 2	CLAASSEN (1973)
WELL UE-15D	CLAASSEN (1973)
UE12T#3-5, 134.4(M)	BENSON (1976)
UE12T#3-6, 169.6(M)	BENSON (1976)
UE12T#3-7, 199.2(M)	BENSON (1976)
UE12T#3-8, 202.4(M)	BENSON (1976)
UE12T#3-9, 257.6(M)	BENSON (1976)
UE12T#3-10, 260.0(M)	BENSON (1976)
UE12T#3-11, 441.4(M)	BENSON (1976)
UE12T#3-12, 442.4(M)	BENSON (1976)
UE12T#3-13, 291.4(M)	BENSON (1976)
UE12T#3-14, 320.3(M)	BENSON (1976)
UE12T#3-15, 320.6(M)	BENSON (1976)
UE12T#3-16, 321.3(M)	BENSON (1976)
UE12T#3-17, 350.8(M)	BENSON (1976)
UE12T#3-18, 411.2(M)	BENSON (1976)
UE12T#3-19, 470.6(M)	BENSON (1976)
UE12T#3-20, 472.4(M)	BENSON (1976)
UE12T#3-21, 501.7(M)	BENSON (1976)
UE12T#3-22, 503.2(M)	BENSON (1976)
UE12T#3-23, 532.8(M)	BENSON (1976)
RML 1A, SURFACE	BENSON (1976)
RML 1B, SURFACE	BENSON (1976)
1, U12N.05 BYPASS	WHITE, CLAASSEN AND BENSON (1980)
2, U12N.05	WHITE, CLAASSEN AND BENSON (1980)
3, U12N.05	WHITE, CLAASSEN AND BENSON (1980)
4, U12N MAIN	WHITE, CLAASSEN AND BENSON (1980)
5, U12N.07 BYPASS	WHITE, CLAASSEN AND BENSON (1980)
6, U12N.02	WHITE, CLAASSEN AND BENSON (1980)
7, U12T.02 BYPASS	WHITE, CLAASSEN AND BENSON (1980)
8, U12T.02	WHITE, CLAASSEN AND BENSON (1980)
9, U12T.02 BYPASS	WHITE, CLAASSEN AND BENSON (1980)
10, U12T MAIN	WHITE, CLAASSEN AND BENSON (1980)
11, U12T.03	WHITE, CLAASSEN AND BENSON (1980)
12, U12T.03	WHITE, CLAASSEN AND BENSON (1980)
13, U12T.04	WHITE, CLAASSEN AND BENSON (1980)
14, U12T.03	WHITE, CLAASSEN AND BENSON (1980)
15, U12T.03	WHITE, CLAASSEN AND BENSON (1980)
16, U12T.03	WHITE, CLAASSEN AND BENSON (1980)
17, U12E.07	WHITE, CLAASSEN AND BENSON (1980)
18, U12E.04	WHITE, CLAASSEN AND BENSON (1980)
19, U12E	WHITE, CLAASSEN AND BENSON (1980)
20, U12E	WHITE, CLAASSEN AND BENSON (1980)
21, U12E.04	WHITE, CLAASSEN AND BENSON (1980)
22, U12E.03	WHITE, CLAASSEN AND BENSON (1980)

SAS

14:49 THURSDAY, MAY 29, 1986 4

SITE	REFERENCE
23, U12E	WHITE, CLAASSEN AND BENSON (1980)
24, U12E.02	WHITE, CLAASSEN AND BENSON (1980)
25, U12E.05	WHITE, CLAASSEN AND BENSON (1980)
26, U12E.03	WHITE, CLAASSEN AND BENSON (1980)
27, U12E.03	WHITE, CLAASSEN AND BENSON (1980)
28, U12E.03	WHITE, CLAASSEN AND BENSON (1980)
29, U12E.07	WHITE, CLAASSEN AND BENSON (1980)
30, U12E	WHITE, CLAASSEN AND BENSON (1980)
31, U12E	WHITE, CLAASSEN AND BENSON (1980)
32, U12E	WHITE, CLAASSEN AND BENSON (1980)
33, U12E	WHITE, CLAASSEN AND BENSON (1980)
34, U12B	WHITE, CLAASSEN AND BENSON (1980)
35, U12B.03	WHITE, CLAASSEN AND BENSON (1980)
36, U12.04	WHITE, CLAASSEN AND BENSON (1980)
SEEP 1, U12T	HENNE (1982)
SEEP 2, U12N.03	HENNE (1982)
SEEP 3, U12N.05	HENNE (1982)
SEEP 4, U12N.03	HENNE (1982)
LYSIMETER 1, U12N.05	HENNE (1982)
LYSIMETER 2, U12N.05	HENNE (1982)
LYSIMETER 3, U12N.05	HENNE (1982)
LYSIMETER 4, U12N.05	HENNE (1982)
LYSIMETER 5, U12N.05	HENNE (1982)
LYSIMETER 6, SURF SOIL	HENNE (1982)
LYSIMETER 7, SURF SOIL	HENNE (1982)
LYSIMETER 9, SURF SOIL	HENNE (1982)
ANAL 2589, TUNNEL U12B, E	CLEBSCH AND BARKER (1960)
ANAL 2819, TUNNEL U12B, D	CLEBSCH AND BARKER (1960)
ANAL 2913, TUNNEL U12E, D	CLEBSCH AND BARKER (1960)
ANAL 3260, TUNNEL U12E, D	CLEBSCH AND BARKER (1960)
ANAL 3541, TUNNEL U12E, M	CLEBSCH AND BARKER (1960)

N= 193

SITE	LOCATION	DATE	INT SAMP (M)	LITHOLOGY
WELL J-12	FMW	03/26/71	INT	T
WELL J-13	FMW	03/26/71	INT	T
WELL UE-25B#1	YM	08/07/81	INT	T
WELL UE-25B#1	YM	09/01/81	INT	T
WELL UE-25B#1	YM	07/20/82	863-875	T
WELL UE-25C#1	YM	09/30/83	INT	T
WELL UE-25C#2	YM	03/13/84	INT	T
WELL UE-25C#3	YM	05/09/84	INT	T
WELL UE-25P#1	YM	02/09/83	381-1197	T/C
WELL UE-25P#1	YM	05/12/83	1297-1805	C
WELL UE-29A#2	YM	01/08/82	247-354	T
WELL UE-29A#2	YM	01/15/82	87-213	T
WELL USW G-4	YM	12/09/82	INT	T
WELL USW H-1	YM	10/20/80	572-687	T
WELL USW H-1	YM	12/08/80	687-1829	T
WELL USW H-3	YM	03/14/84	822-1220	T
WELL USW H-4	YM	05/17/82	INT	T
WELL USW H-5	YM	07/03/82	INT	T
WELL USW H-5	YM	07/26/82	INT	T
WELL USW H-6	YM	10/16/82	INT	T
WELL USW H-6	YM	06/20/84	753-835	T
WELL USW H-6	YM	07/06/84	608-646	T
WELL USW VH-1	CF	02/06/81	INT	T
WELL USW VH-1	CF	02/08/81	INT	T
WELL USW VH-1	CF	02/11/81	INT	T
3, 16S/49E-22DC	AD/FMW	11/20/72	INT	T/C?
4, 16S/49E-5ACC	AD/FMW	03/04/74	INT	T
5, 16S/49E-8ABB	AD/FMW	11/17/72	INT	T/C?
6, 16S/49E-8ACC	AD/FMW	04/01/71	INT	
7, 16S/49E-9CDA	AD/FMW	11/18/72	INT	
8, 16S/49E-9DCC	AD/FMW	03/01/74	INT	T/C?
9, 16S/49E-18DC	AD/FMW	03/01/74	INT	T
10, 16S/49E-16CCC	AD/FMW	06/26/79	INT	T/C?
11, 16S/49E-19DAA	AD/FMW	03/05/74	INT	T
12, 16S/48E-24AAA	AD/FMW	11/17/72	INT	
13, 16S/48E-25AA	AD/FMW	03/05/74	INT	T
14, 16S/48E-36AAA	AD/FMW	03/04/74	INT	T
15, 17S/48E-1AB	AD/FMW	03/05/74	INT	T
16, 17S/49E-7BB	AD/FMW	03/01/74	INT	T/C
17, 17S/49E-9AA	AD/FMW	03/01/74	INT	T/C?
18, 17S/49E-8DDB	AD/FMW	03/06/74	INT	T
19, 17S/49E-15BBD	AD/FMW	03/06/74	INT	T/C
64, WELL 8, NTS	RM	03/24/71	INT	T
20, 17S/49E-35000, ASH TR	AD/MISC	03/06/74	INT	T
21, 16S/49E-23ADD	AD/INT	06/25/79	INT	T
23, 16S/48E-15AAA	AD/INT	03/31/71	INT	T
25, 16S/48E-10CBA	AD/INT	03/31/71	INT	T
27, 16S/50E-7BCD	AD/MISC	04/01/71	INT	C/T?
29, 16S/49E-15AAA	AD/MISC	03/31/71	INT	C/T
30, 16S/49E-36AAA	AD/MISC	06/24/79	INT	C/T?
45, 16S/48E-8BBA	AD/OV	06/24/71	INT	
46, 16S/48E-7BBA	AD/OV	03/30/71	INT	

SITE	LOCATION	DATE	INT SAMP (M)	LITHOLOGY
47, 16S/48E-7CBC	AD/OV	03/31/71	INT	
48, 16S/48E-18BCC	AD/OV	06/24/79	INT	
49, 16S/48E-17CCC	AD/OV	06/25/79	INT	
50, 16S/48E-18DAD	AD/OV	06/25/79	INT	
51, 16S/48E-8CDA	AD/OV	03/31/71	INT	
52, 16S/48E-17AAB	AD/OV	08/18/62	INT	
62, WHITTEROCK SPRING	RM	04/10/72	INT	T
68, BUTTE SPRING	RM	11/10/60	INT	T
70, OAK SPRING	RM	04/28/58	INT	T
71, TOPPOPAH SPRING	CH	03/25/58	INT	T
1, 10S/47E-14BAB	OV		INT	QAL
2, 10S/47E-27CBA	OV		INT	QAL
3, 10S/47E-31AAB	OV		INT	QAL/BULL
4, 10S/47E-32DDA	OV		INT	QAL/BULL
5, 10S/47E-33AAB	OV		INT	QAL
6, 10S/47E-30DCC	OV		INT	QAL/BULL
7, 11S/46E-26BBB	OV		INT	T
8, 10S/46E-26BCC	OV		INT	T
9, 11S/47E-3CDB	OV		INT	T
10, 11S/47E-4CAD	OV		INT	QAL
11, 11S/47E-10CAA	OV		INT	T
12, 11S/47E-10BCC	OV		INT	T
13, 11S/47E-16DCD	OV		INT	T
14, 11S/47E-16BDC	OV		INT	T
15, 11S/47E-18ACD	OV		INT	T
16, 11S/47E-21ACC	OV		INT	QAL
17, 11S/47E-21DBB	OV		INT	QAL
18, 11S/47E-21ABA	OV		INT	QAL
19, 11S/47E-21ABA	OV		INT	T
20, 11S/47E-27CBA	OV		INT	T
21, 11S/47E-28AAC	OV		INT	QAL
22, 11S/47E-28DAC	OV		INT	QAL
23, 11S/47E-33BAC	OV		INT	T
24, 11S/47E-10CCB	OV		INT	T
25, 12S/47E-5CDA	OV		INT	T
26, 12S/47E-6CDD	OV		INT	QAL/BULL
27, 12S/47E-7DBD	OV		INT	QAL
28, 12S/47E-20BBB	OV		INT	QAL
29, 12S/47E-19ADC	OV		INT	QAL
18B, 12S/47E-20	OV		INT	QAL/SOIL
19B, 12S/47E-20	OV		INT	QAL/SOIL
20B, 12S/47E-20	OV		INT	QAL/SOIL
21B, 12S/47E-20	OV		INT	QAL/SOIL
22B, 12S/47E-20	OV		INT	QAL/SOIL
WELL UE-19B-1	PM	10/13/64	2190-4500	T
WELL UE-19C	PM	03/09/66	3040-3075	T
WELL UE-19D	PM	03/09/66	3300-3480	T
WELL UE-19E	PM	08/01/66	2475-6005	T
WELL UE-19E	PM	10/06/71	2475-6005	T
WELL UE-19GS	PM	08/02/66	2650-7500	T
WELL UE-19GS	PM	10/06/71	2650-7500	T
WELL U-20A-2	PM	03/10/66	2066-4500	T

SITE	LOCATION	DATE	INT SAMP (M)	LITHOLOGY
WELL U-20A-2	PM	10/06/71	2066-4500	T
WELL UE-20D	PM	07/27/66	2446-4500	T
WELL UE-20E-1	PM	03/08/66	2600	T
WELL UE-20H	PM	08/26/65	2506-7207	T
WELL UE-20J	PM	10/21/64	1740-5690	T
WELL ARMY-1	FF	03/18/71	INT	C
WELL 5A	FF	06/04/64	INT	QAL
WELL 5B	FF	03/25/71	INT	QAL
WELL 5C	FF	03/22/71	INT	QAL
WELL C	YF	04/11/69	INT	C
WELL C-1	YF	03/29/71	INT	C
WELL 3	YF	04/16/69	INT	QAL
WELL A	YF	03/23/71	INT	QAL
WELL 2	YF	03/21/71	INT	T/C
WELL UE-15D	YF	03/21/71	INT	T/C
UE12T#3-5, 134.4(M)	RM/PW		INT	T
UE12T#3-6, 169.6(M)	RM/PW		INT	T
UE12T#3-7, 199.2(M)	RM/PW		INT	T
UE12T#3-8, 202.4(M)	RM/PW		INT	T
UE12T#3-9, 257.5(M)	RM/PW		INT	T
UE12T#3-10, 260.0(M)	RM/PW		INT	T
UE12T#3-11, 441.4(M)	RM/PW		INT	T
UE12T#3-12, 442.4(M)	RM/PW		INT	T
UE12T#3-13, 291.4(M)	RM/PW		INT	T
UE12T#3-14, 320.3(M)	RM/PW		INT	T
UE12T#3-15, 320.6(M)	RM/PW		INT	T
UE12T#3-16, 321.3(M)	RM/PW		INT	T
UE12T#3-17, 350.8(M)	RM/PW		INT	T
UE12T#3-18, 411.2(M)	RM/PW		INT	T
UE12T#3-19, 470.6(M)	RM/PW		INT	T
UE12T#3-20, 472.4(M)	RM/PW		INT	T
UE12T#3-21, 501.7(M)	RM/PW		INT	T
UE12T#3-22, 503.2(M)	RM/PW		INT	T
UE12T#3-23, 532.8(M)	RM/PW		INT	T
RML 1A, SURFACE	RM/PW		INT	T
RML 1B, SURFACE	RM/PW		INT	T
1, U12N.05 BYPASS	RM/FW	06/02/71	INT	T
2, U12N.05	RM/FW	09/21/71	INT	T
3, U12N.05	RM/FW	09/21/71	INT	T
4, U12N MAIN	RM/FW	08/03/72	INT	T
5, U12N.07 BYPASS	RM/FW	08/03/72	INT	T
6, U12N.02	RM/FW	11/14/72	INT	T
7, U12T.02 BYPASS	RM/FW	09/22/71	INT	T
8, U12T.02	RM/FW	09/22/71	INT	T
9, U12T.02 BYPASS	RM/FW	09/22/71	INT	T
10, U12T MAIN	RM/FW	09/22/71	INT	T
11, U12T.03	RM/FW	02/15/73	INT	T
12, U12T.03	RM/FW	02/01/73	INT	T
13, U12T.04	RM/FW	08/22/73	INT	T
14, U12T.03	RM/FW	03/19/73	INT	T
15, U12T.03	RM/FW	11/16/72	INT	T
16, U12T.03	RM/FW	09/24/74	INT	T

SITE	LOCATION	DATE	INT SAMP (M)	LITHOLOGY
17, U12E.07	RM/FW	03/11/60	INT	T
18, U12E.04	RM/FW	06/02/59	INT	T
19, U12E	RM/FW	11/22/59	INT	T
20, U12E	RM/FW	11/29/59	INT	T
21, U12E.04	RM/FW	01/07/60	INT	T
22, U12E.03	RM/FW	12/14/59	INT	T
23, U12E	RM/FW	01/29/59	INT	T
24, U12E.02	RM/FW	10/11/58	INT	T
25, U12E.05	RM/FW	09/12/58	INT	T
26, U12E.03	RM/FW	05/27/59	INT	T
27, U12E.03	RM/FW	05/20/59	INT	T
28, U12E.03	RM/FW	12/03/59	INT	T
29, U12E.07	RM/FW	03/18/66	INT	T
30, U12E	RM/FW	01/22/59	INT	T
31, U12E	RM/FW	06/24/59	INT	T
32, U12E	RM/FW	07/18/59	INT	T
33, U12E	RM/FW	07/21/59	INT	T
34, U12B	RM/FW	06/06/58	INT	T
35, U12B.03	RM/FW	08/22/58	INT	T
36, U12.04	RM/FW	09/29/58	INT	T
SEEP 1, U12T	RM	AVERAGE	INT	T
SEEP 2, U12N.03	RM	AVERAGE	INT	T
SEEP 3, U12N.05	RM	AVERAGE	INT	T
SEEP 4, U12N.03	RM	AVERAGE	INT	T
LYSIMETER 1, U12N.05	RM	AVERAGE	INT	T
LYSIMETER 2, U12N.05	RM	AVERAGE	INT	T
LYSIMETER 3, U12N.05	RM	AVERAGE	INT	T
LYSIMETER 4, U12N.05	RM	AVERAGE	INT	T
LYSIMETER 5, U12N.05	RM	AVERAGE	INT	T
LYSIMETER 6, SURF SOIL	RM	AVERAGE	INT	T
LYSIMETER 7, SURF SOIL	RM	AVERAGE	INT	T
LYSIMETER 9, SURF SOIL	RM	AVERAGE	INT	T
ANAL 2589, TUNNEL U12B, E	RM	08/22/58	INT	T
ANAL 2819, TUNNEL U12B, D	RM	06/06/58	INT	T
ANAL 2913, TUNNEL U12E, D	RM	10/11/58	INT	T
ANAL 3260, TUNNEL U12E, D	RM	06/02/59	INT	T
ANAL 3541, TUNNEL U12E, M	RM	11/22/59	INT	T

SITE	SURFACE ALTITUDE (M)	WELL	DEPTH (M)	DEPTH TO WATER (M)	TEMP (C)
WELL J-12	953.50	347	225	27.0000	
WELL J-13	1011.30	1063	282	31.0000	
WELL UE-25B#1	1200.40	1220	470	36.0000	
WELL UE-25B#1	1200.40	1220	470	36.0000	
WELL UE-25B#1	1200.40	1220	470	37.2000	
WELL UE-25C#1	1131.00	914	400	41.5000	
WELL UE-25C#2	1132.00	913	401	40.5000	
WELL UE-25C#3	1132.00	913	402	40.8000	
WELL UE-25P#1	1114.00	1800	381	44.3000	
WELL UE-25P#1	1114.00	1800	381	56.0000	
WELL UE-29A#2	1215.10	422	29	25.1000	
WELL UE-29A#2	1215.10	422	29	22.7000	
WELL USW G-4	1270.00	915	541	35.6000	
WELL USW H-1	1302.20	1829	572	33.0000	
WELL USW H-1	1302.20	1829	572	34.7000	
WELL USW H-3	1483.00	1220	26.5000		
WELL USW H-4	1249.00	1220	519	34.8000	
WELL USW H-5	1477.80	1220	704	36.5000	
WELL USW H-5	1477.80	1220	704	35.3000	
WELL USW H-6	1302.00	1220	526	37.8000	
WELL USW H-6	1302.00	1220	526	41.6000	
WELL USW H-6	1302.00	1220	526	37.2000	
WELL USW VH-1	954.50	762	184	35.2000	
WELL USW VH-1	954.50	762	184	35.5000	
WELL USW VH-1	954.50	762	184	35.5000	
3. 155/49E-22DC	150	78			
4. 165/49E-5ACC	90	21			
5. 165/49E-8ABB	60	45	23.0000		
6. 165/49E-8ACC	90	45	25.8000		
7. 165/49E-9CDA	90	46	24.0000		
8. 165/49E-9DCC	60	49	23.3000		
9. 165/49E-18DC	110	33			
10. 165/49E-16CCC					
11. 165/49E-19DAA	90	30	26.4000		
12. 165/48E-24AAA	150	29	27.0000		
13. 165/48E-25AA	50	26	26.5000		
14. 165/48E-36AAA	50	21			
15. 175/48E-1AB	60	16			
16. 175/49E-7BB	150	12			
17. 175/49E-9AA					
18. 175/49E-8DDB	100	15	24.0000		
19. 175/49E-15BBBD	110	17	22.5000		
64. WELL B, NTS	1680	328	26.5000		
20. 175/49E-35DDD, ASH TR	0	51	30.6000		
21. 165/49E-23ADD	50	0	23.8000		
23. 165/48E-15AAA	29		25.5000		
25. 165/48E-10CBA			24.5000		
27. 165/50E-7BCD	60	43			
29. 165/49E-15AAA	120	51			
30. 165/49E-36AAA			25.0000		
45. 165/48E-8BBBA	80	34	24.7000		
46. 165/48E-7BBBA			24.2000		
47. 165/48E-7CBC					

48.	16S/48E-18BBC						
49.	16S/48E-17CC						
50.	16S/48E-18DAD						
51.	16S/48E-BCDA						
52.	16S/48E-17ABB						
62.	WHITE ROCK SPRING	1524.00					
68.	BJTE SPRING	1707.00					
70.	OAK SPRING	1768.00					
71.	TDPOPAH SPRING	1768.00					
1.	105/47E-14BAB						
2.	105/47E-27CBA						
3.	105/47E-31AAB	90.00	0.00	31.000	23.3000		
4.	105/47E-32DDA	0.00	0.00	0.000	24.0000		
5.	105/47E-33AAB	0.00	0.00	0.000	15.0000		
6.	105/47E-30DCC	0.00	0.00	0.000	12.8000		
7.	11S/46E-26BBBB	0.00	0.00	0.000	11.7000		
8.	10S/46E-26BBC	0.00	0.00	0.000	29.0000		
9.	11S/47E-3CDB	0.00	0.00	0.000	19.0000		
10.	11S/47E-4CAD	0.00	0.00	0.000	19.5000		
11.	11S/47E-10CAA	0.00	0.00	0.000	22.0000		
12.	11S/47E-10BBC	0.00	0.00	0.000	22.0000		
13.	11S/47E-16DCD	0.00	0.00	0.000	22.5000		
14.	11S/47E-16BDC	0.00	0.00	0.000	24.0000		
15.	11S/47E-18ACD	0.00	0.00	0.000	18.5000		
16.	11S/47E-21ACC	0.00	0.00	0.000	36.5000		
17.	11S/47E-21DBB	0.00	0.00	0.000	24.0000		
18.	11S/47E-21ABA	0.00	0.00	0.000	21.0000		
19.	11S/47E-21ABA	0.00	0.00	0.000	41.0000		
20.	11S/47E-27CBA	55.00	0.00	55.00	21.5000		
21.	11S/47E-28AAC	25.00	0.00	25.00	18.0000		
22.	11S/47E-28DAC	0.00	0.00	0.000	21.0000		
23.	11S/47E-33BAC	0.00	0.00	0.000	34.0000		
24.	11S/47E-10CCB	0.00	0.00	0.000	21.0000		
25.	12S/47E-5CDA	0.00	0.00	0.000	24.0000		
26.	12S/47E-6CDD	0.00	0.00	0.000	21.5000		
27.	12S/47E-7DBD	91.00	0.00	91.00	20.0000		
28.	12S/47E-20BBB	0.00	0.00	0.000	18.5000		
29.	12S/47E-19ADC	0.00	0.00	0.000	20.0000		
18B.	12S/47E-20						
19B.	12S/47E-20						
20B.	12S/47E-20						
21B.	12S/47E-20						
22B.	12S/47E-20						
WELL	UE-19B-1						
WELL	UE-19C						
WELL	UE-1SD						
WELL	UE-19E						
WELL	UE-19E						
WELL	UE-19GS						
WELL	UE-19GS						
WELL	U-20A-2						
WELL	UE-20D						

SITE	SURFACE ALTITUDE (M)	WELL DEPTH (M)	DEPTH TO WATER (M)	TEMP (C)
WELL UE-20E-1	.	1949.20	.	32.8000
WELL UE-20H	.	2196.69	.	32.2000
WELL UE-20J	.	1734.31	.	38.9000
WELL ARMY-1	961.30	593.00	240	31.0000
WELL 5A	943.00	277.00	212	23.0000
WELL 5B	943.00	274.00	209	25.0000
WELL 5C	939.00	366.00	210	24.5000
WELL C	1195.00	519.00	470	37.0000
WELL C-1	1195.00	503.00	470	38.0000
WELL 3	1209.00	548.00	486	21.5000
WELL A	1221.00	570.00	492	26.5000
WELL 2	1362.00	1043.00	626	34.5000
WELL UE-15D	1398.00	1810.00	203	34.5000
UE12T#3-5, 134.4(M)	.	134.40	.	.
UE12T#3-6, 169.6(M)	.	169.60	.	.
UE12T#3-7, 199.2(M)	.	199.20	.	.
UE12T#3-8, 202.4(M)	.	202.40	.	.
UE12T#3-9, 257.6(M)	.	257.60	.	.
UE12T#3-10, 260.0(M)	.	260.00	.	.
UE12T#3-11, 441.4(M)	.	441.40	.	.
UE12T#3-12, 442.4(M)	.	442.40	.	.
UE12T#3-13, 291.4(M)	.	291.40	.	.
UE12T#3-14, 320.3(M)	.	320.30	.	.
UE12T#3-15, 320.6(M)	.	320.60	.	.
UE12T#3-16, 321.3(M)	.	321.30	.	.
UE12T#3-17, 350.8(M)	.	350.80	.	.
UE12T#3-18, 411.2(M)	.	411.20	.	.
UE12T#3-19, 470.6(M)	.	470.60	.	.
UE12T#3-20, 472.4(M)	.	472.40	.	.
UE12T#3-21, 501.7(M)	.	501.70	.	.
UE12T#3-22, 503.2(M)	.	503.20	.	.
UE12T#3-23, 502.8(M)	.	502.80	.	.
RML 1A, SURFACE	.	0.00	.	.
RML 1B, SURFACE	.	0.00	.	.
1, U12N.05 BYPASS
2, U12N.05
3, U12N.05
4, U12N MAIN
5, U12N.07 BYPASS
6, U12N.02
7, U12T.02 BYPASS
8, U12T.02
9, U12T.02 BYPASS
10, U12T MAIN
11, U12T.03
12, U12T.03
13, U12T.04
14, U12T.03
15, U12T.03
16, U12T.03
17, U12E.07
18, U12E.04
19, U12E

SAS

14:49 THURSDAY, MAY 29, 1986 12

SITE	SURFACE ALTITUDE (M)	WELL DEPTH (M)	DEPTH TO WATER (M)	TEMP (C)
20, U12E
21, U12E.04
22, U12E.03
23, U12E
24, U12E.02
25, U12E.05
26, U12E.03
27, U12E.03
28, U12E.03
29, U12E.07
30, U12E
31, U12E
32, U12E
33, U12E
34, U12B
35, U12B.03
36, U12.04
SEEP 1, U12T	.	.	.	13.1000
SEEP 2, U12N.03	.	.	.	15.8000
SEEP 3, U12N.05	.	.	.	16.6000
SEEP 4, U12N.03	.	.	.	20.5000
LYSIMETER 1, U12N.05	.	.	.	14.1000
LYSIMETER 2, U12N.05	.	.	.	16.1000
LYSIMETER 3, U12N.05	.	.	.	15.5000
LYSIMETER 4, U12N.05	.	.	.	16.4000
LYSIMETER 5, U12N.05	.	.	.	18.7000
LYSIMETER 6, SURF SOIL	.	.	.	13.5000
LYSIMETER 7, SURF SOIL	.	.	.	9.3000
LYSIMETER 9, SURF SOIL	.	.	.	14.7000
ANAL 2529, TUNNEL U12B, E
ANAL 2819, TUNNEL U12B, D	.	.	.	18.0000
ANAL 2913, TUNNEL U12E, D
ANAL 3260, TUNNEL U12E, D
ANAL 3541, TUNNEL U12E, M	.	.	.	16.0000

N= 193

SITE	CA	MG	NA	K
WELL J-12	0.34930	0.08640	1.65291	0.130440
WELL J-13	0.29940	0.08640	1.82690	0.127883
WELL UE-25B#1	0.47405	0.03003	2.30537	0.094633
WELL UE-25B#1	0.42415	0.02427	2.00089	0.089518
WELL UE-25C#1	0.44910	0.02962	2.43586	0.071614
WELL UE-25C#2	0.27445	0.01399	2.34887	0.051153
WELL UE-25C#3	0.29940	0.01646	2.39237	0.053711
WELL UE-25P#1	0.27445	0.01646	4.00177	0.048595
WELL UE-25P#1	0.92315	0.41144	6.52463	0.306919
WELL UE-29A#2	0.24950	0.00823	1.91389	0.028134
WELL UE-29A#2	0.24950	0.01234	1.91389	0.033250
WELL USW G-4	0.32435	0.00823	2.47936	0.053711
WELL USW H-1	0.92315	0.00411	2.21838	0.061384
WELL USW H-1	0.15469	0.00411	2.21838	0.040922
WELL USW H-3	0.01996	0.00082	5.21971	0.028134
WELL USW H-4	0.42415	0.01193	3.17532	0.066499
WELL USW H-5	0.04741	0.00041	2.60985	0.053711
WELL USW H-5	0.04990	0.00411	3.74079	0.033250
WELL USW H-6	0.10230	0.00370	3.82778	0.033250
WELL USW H-6	0.03493	0.00082	3.82778	0.035807
WELL USW H-6	0.11727	0.00288	3.43631	0.048595
WELL USW VH-1	0.27445	0.06583	3.47980	0.048595
WELL USW VH-1	0.24950	0.06172	3.39281	0.046038
WELL USW VH-1	0.24701	0.06172	1.87000	0.120000
3. 15S/49E-22DC	0.67000	0.08000	1.52000	0.130000
4. 16S/49E-5ACC	0.72000	0.09000	1.61000	0.140000
5. 16S/49E-8ABB	0.75000	0.11000	1.61000	0.170000
6. 16S/49E-8ACC	0.57000	0.10000	2.22000	0.220000
7. 16S/49E-9CDA	0.76000	0.14000	2.44000	0.230000
8. 16S/49E-9DCC	0.57000	0.1000	1.83000	0.230000
9. 16S/49E-18DC	0.50000	0.11000	1.73000	0.110000
10. 16S/49E-16CCC	0.75000	0.08000	1.73000	0.110000
11. 16S/49E-19DAA	0.60000	0.05000	1.57000	0.210000
12. 16S/49E-24AAA	0.45000	0.03000	2.35000	0.180000
13. 16S/48E-25AA	0.47000	0.03000	1.87000	0.190000
14. 16S/48E-36AAA	0.42000	0.08000	1.74000	0.160000
15. 17S/48E-1AB	0.47000	0.06000	1.74000	0.180000
16. 17S/49E-7BB	0.60000	0.07000	2.09000	0.190000
17. 17S/49E-9AA	0.62000	0.15000	2.09000	0.250000
18. 17S/49E-80DB	0.52000	0.11000	1.57000	0.190000
19. 17S/49E-15BBD	0.52000	0.16000	1.36000	0.210000
64. WELL 8, NTS	0.21000	0.05000	1.35000	0.090000
20. 17S/49E-35DDD, ASH TR	0.38000	0.07000	2.09000	0.205000
21. 16S/49E-23ADD	0.40000	0.07000	0.165000	0.165000
23. 16S/48E-15AAA	0.24000	0.13000	2.52000	0.150000
25. 16S/48E-10CBA	0.23000	0.16000	2.65000	0.140000
27. 16S/50E-7BCC	1.19000	0.72000	4.85000	0.330000
29. 16S/49E-15AAA	0.02000	0.31000	3.48000	0.250000
30. 16S/49E-36AAA	1.30000	0.91000	3.22000	0.460000
45. 16S/48E-88BBA	1.46000	0.26000	7.85000	0.330000
46. 16S/48E-7BBA	1.32000	0.39000	6.09000	0.260000
47. 16S/48E-7BCC	1.17000	0.66000	5.66000	0.240000
48. 16S/48E-18BCC	1.37000	0.45000	6.53000	0.300000

SAS

14:49 THURSDAY, MAY 29, 1986 14

SITE	CA	MG	NA	K
49, 16S/48E-17CCC	1.65000	0.450000	7.3900	0.310000
50, 16S/48E-18DAD	1.32000	0.350000	6.5200	0.270000
51, 16S/48E-8CDA	1.20000	0.280000	6.9600	0.260000
52, 16S/48E-17ABB	1.50000	0.320000	6.8300	0.310000
62, WHITEROCK SPRING	0.10000	0.010000	1.9100	0.200000
68, BUTTE SPRING	0.52000	0.100000	1.4800	0.150000
70, OAK SPRING	0.45000	0.200000	0.9600	0.160000
71, TUPOPAH SPRING	0.18000	0.040000	0.6100	0.160000
1, 10S/47E-14BAB	0.18000	0.010000	6.2200	0.210000
2, 10S/47E-27CBA	0.55000	0.060000	7.4400	0.220000
3, 10S/47E-31AAB	0.58000	0.180000	4.3500	0.200000
4, 10S/47E-32DDA	0.75000	0.220000	5.9600	0.000000
5, 10S/47E-33AAB	0.75000	0.190000	7.3500	0.230000
6, 10S/47E-30DCC	0.60000	0.190000	4.3500	0.200000
7, 11S/46E-26BBBB	0.00000	0.010000	2.5700	0.040000
8, 10S/46E-26BCC	0.15000	0.040000	2.4800	0.040000
9, 11S/47E-3CDB	0.40000	0.040000	5.3100	0.120000
10, 11S/47E-4CAD	0.65000	0.180000	9.7000	0.220000
11, 11S/47E-10CAA	0.35000	0.030000	8.5300	0.060000
12, 11S/47E-10BCC	0.35000	0.020000	6.7900	0.180000
13, 11S/47E-16DCD	0.45000	0.020000	7.5300	0.200000
14, 11S/47E-16BDC	0.42000	0.020000	7.1300	0.190000
15, 11S/47E-18ACD	0.55000	0.150000	2.1800	0.090000
16, 11S/47E-21ACC	0.58000	0.120000	10.0900	0.220000
17, 11S/47E-21DBB	0.62000	0.130000	10.5700	0.210000
18, 11S/47E-21ABA	0.65000	0.130000	10.7000	0.210000
19, 11S/47E-21ABA	0.32000	0.020000	6.5300	0.200000
20, 11S/47E-27CBA	0.90000	0.210000	5.0000	0.280000
21, 11S/47E-28AAC	0.25000	0.190000	13.7000	0.230000
22, 11S/47E-28DAC	0.21000	0.000000	10.8300	0.060000
23, 11S/47E-33BAC	0.30000	0.030000	4.8700	0.120000
24, 11S/47E-10CCB	0.32000	0.110000	5.3900	0.150000
25, 12S/47E-5CDA	0.80000	0.180000	4.6100	0.190000
26, 12S/47E-6CDD	0.68000	0.130000	4.5700	0.260000
27, 12S/47E-7DBD	0.63000	0.150000	11.1400	0.260000
28, 12S/47E-20BBB	0.68000	0.150000	11.0500	0.260000
29, 12S/47E-19ADC	0.95000	0.230000	12.6200	0.260000
18B, 12S/47E-20
19B, 12S/47E-20
20B, 12S/47E-20
21B, 12S/47E-20
22B, 12S/47E-20
WELL UE-19B-1	0.59880	0.098745	1.8269	0.076730
WELL UE-19C	0.32435	0.004114	6.1332	0.005115
WELL UE-19D	1.42216	0.115203	6.6551	0.109979
WELL UE-19E	0.09232	0.002057	1.8704	0.020461
WELL UE-19E	0.00998	0.004114	1.6529	0.023019
WELL UE-19GS	0.06986	0.002057	3.6538	0.038365
WELL UE-19GS	0.87325	0.008229	2.9578	0.020461
WELL U-20A-2	0.15220	0.004114	2.3924	0.005115
WELL U-20A-2	0.14721	0.008229	2.3924	0.056268
WELL UE-20D	0.10729	0.004114	3.8278	0.043480
WELL UE-20E-1	0.00499	0.002057	3.6103	0.051153
WELL UE-20H	0.01497	0.002057	2.7838	0.046038

SITE	CA	MG	NA	K	SAS
WELL UE-20J	1.14770	0.04937	6.00266	0.163690	
WELL ARMY-1	1.09780	0.90516	1.60941	0.132998	
WELL 5A	0.06737	0.01646	7.09010	0.163690	
WELL 5B	0.17465	0.09052	3.91478	0.281342	
WELL 5C	0.02495	0.01646	5.65468	0.173921	
WELL C	1.87126	1.19317	5.43719	0.383648	
WELL C-1	1.79641	1.23431	5.21971	0.358072	
WELL 3	0.47405	0.53487	1.73990	0.217401	
WELL A	0.52395	0.30446	2.13138	0.225074	
WELL 2	0.77345	0.57601	1.17443	0.173633	
WELL UE-15D	1.39721	0.65830	3.47980	0.383648	
UE 12T #3-5	134.4(M)	0.27445	0.12343	1.06569	0.255766
UE 12T #3-6	169.6(M)	0.64870	0.28801	1.50937	0.358072
UE 12T #3-7	199.2(M)	0.57385	0.25921	1.89214	0.255766
UE 12T #3-8	202.4(M)	0.42415	0.17280	1.01784	0.104864
UE 12T #3-9	257.6(M)	0.39920	0.19749	1.30493	0.122767
UE 12T #3-10	260.0(M)	0.64870	0.29624	1.40932	0.143229
UE 12T #3-11	441.4(M)	0.21956	0.02880	2.00089	0.358072
UE 12T #3-12	442.4(M)	0.27445	0.04114	2.41846	0.485955
UE 12T #3-13	291.4(M)	0.19711	0.08640	1.10919	0.122767
UE 12T #3-14	320.3(M)	0.22954	0.06994	1.35147	0.186709
UE 12T #3-15	320.6(M)	0.44910	0.09052	1.37017	0.204612
UE 12T #3-16	321.3(M)	0.69860	0.18103	1.54416	0.209728
UE 12T #3-17	350.8(M)	0.32435	0.09052	1.01349	0.148344
UE 12T #3-18	411.2(M)	0.14222	0.03415	1.34407	0.140671
UE 12T #3-19	470.6(M)	0.02520	0.00082	1.73990	0.038365
UE 12T #3-20	472.4(M)	0.02250	0.00370	2.27057	0.030692
UE 12T #3-21	501.7(M)	0.05489	0.00617	2.62290	0.030692
UE 12T #3-22	503.2(M)	0.04491	0.00576	2.82134	0.023019
UE 12T #3-23	532.8(M)	0.02495	0.00453	3.09268	0.037191
RML 1A, SURFACE	0.24950	0.09463	0.66551	0.061384	
RML 1B, SURFACE	0.27445	0.13166	1.15703	0.081845	
1. U12N.05 BYPASS	0.18000	0.02000	1.91000	0.240000	
2. U12N.05	0.23000	0.02000	2.85000	0.280000	
3. U12N.05	0.40000	0.06000	3.04000	0.330000	
4. U12N MAIN	1.15000	0.01000	2.70000	0.190000	
5. U12N.07 BYPASS	0.18000	0.01000	0.96000	0.01000	
6. U12N.02	0.01000	0.00100	2.04000	0.140000	
7. U12T.02 BYPASS	0.01000	0.00100	0.96000	0.01000	
8. U12T.02	0.03000	0.01000	1.09000	0.01000	
9. U12T.02 BYPASS	0.07000	0.00100	1.13000	0.020000	
10. U12T MAIN	0.02000	0.01000	2.83000	0.090000	
11. U12T.03	0.50000	0.21000	1.44000	0.140000	
12. U12T.03	0.22000	0.06000	1.39000	0.140000	
13. U12T.04	0.24000	0.08000	1.44000	0.130000	
14. U12T.03	0.50000	0.17000	1.30000	0.170000	
15. U12T.03	0.23000	0.06000	1.65000	0.170000	
16. U12T.03	0.51000	0.19000	1.13000	0.140000	
17. U12E.07	0.43000	0.17000	1.44000	0.150000	
18. U12E.04	0.01000	0.00100	1.04000	0.020000	
19. U12E	0.06000	0.00100	1.13000	0.090000	
20. U12E	0.06000	0.00100	1.04000	0.200000	
21. U12E.04	0.04000	0.00100	1.26000	0.010000	
22. U12E.03	0.04000	0.00100	1.35000	0.010000	

SAS

14:49 THURSDAY, MAY 29, 1986 16

SITE	CA	MG	NA	K
23, U12E	0.200000	0.080000	0.87000	0.170000
24, U12E.02	0.001000	0.001000	1.39000	0.060000
25, U12E.05	0.060000	0.001000	1.44000	0.080000
26, U12E.03	0.060000	0.001000	1.74000	0.030000
27, U12E.03	0.020000	0.001000	1.91000	0.040000
28, U12E.03	0.040000	0.001000	1.74000	0.070000
29, U12E.07	0.430000	0.120000	1.13000	0.150000
30, U12E	0.330000	0.001000	1.48000	0.160000
31, U12E	0.100000	0.001000	2.04000	0.280000
32, U12E	0.400000	0.040000	1.61000	0.200000
33, U12E	0.080000	0.001000	3.04000	0.070000
34, U12B	0.320000	0.040000	0.78000	0.070000
35, U12B.03	0.240000	0.060000	0.65000	0.070000
36, U12.04	0.200000	0.340000	0.96000	0.080000
SEEP 1, U12T	0.037176		2.91869	0.092076
SEEP 2, U12N.03	0.069611	0.005349	2.03569	0.089518
SEEP 3, U12N.05	0.225549	0.006583	2.58810	0.181594
SEEP 4, U12N.03	0.018713	0.004937	2.20968	0.066499
LYSIMETER 1, U12N.05	0.983034	0.046081	8.09054	0.340168
LYSIMETER 2, U12N.05	0.104291	0.005760	2.58810	0.117652
LYSIMETER 3, U12N.05	0.146457	0.004937	2.84474	0.173921
LYSIMETER 4, U12N.05	0.103543	0.004937	2.48371	0.156017
LYSIMETER 5, U12N.05	0.004990	0.004937	0.92650	0.014834
LYSIMETER 6, SURF SDIL	0.441617	0.158404	1.01784	0.023530
LYSIMETER 7, SURF SOIL	0.244511	0.098745	0.66551	0.027111
LYSIMETER 9, SURF SOIL	0.763473	0.259617	1.85735	0.112537
ANAL 2589, TUNNEL U12B, E	0.324351	0.078173	0.78296	0.071614
ANAL 2819, TUNNEL U12B, D	0.239521	0.061716	0.65246	0.071614
ANAL 2913, TUNNEL U12E, D	0.059880	0.000041	1.39192	0.056268
ANAL 3260, TUNNEL U12E, D	0.004990	0.000041	1.04394	0.015346
ANAL 3541, TUNNEL U12E, M	0.059880	0.000041	1.13094	0.092076

N= 193

SITE	SAS	CL	SD4	S102	F	NO3
WELL J-12	0.20591	0.22903	0.89874	0.110536	0.112894	
WELL J-13	0.20027	0.17698	0.94867	0.126326	0.162890	
WELL UE-25B#1	0.36668	0.24985	0.88209	0.078954	0.009677	
WELL UE-25B#1	0.23975	0.22903	0.86545	0.084218	0.009677	
WELL UE-25B#1	0.21155	0.21862	0.84881	0.084218	0.053222	
WELL UE-25C#1	0.20873	0.23944	0.93202	0.110536		
WELL UE-25C#2	0.20027	0.22903	0.89874	0.110536		
WELL UE-25C#3	0.20309	0.22903	0.88209	0.105272		
WELL UE-25P#1	0.36668	0.39560	0.81552	0.178962		
WELL UE-25P#1	0.78978	1.66567	0.68237	0.247389	0.000806	
WELL UE-29A#2	0.31027	0.22903	0.73230	0.052636	0.301589	
WELL UE-29A#2	0.24822	0.21862	0.73230	0.047372	0.301589	
WELL USW G-4	0.16642	0.19780	0.74895	0.13590	0.088703	
WELL USW H-1	0.16078	0.18739	0.78223	0.063163		
WELL USW H-1	0.16360	0.19780	0.66573	0.052636		
WELL USW H-3	0.15513	0.32272	0.71566	0.289498	0.003226	
WELL USW H-4	0.19462	0.27067	0.76559	0.242126	0.075800	
WELL USW H-5	0.17206	0.16657	0.79888	0.073690	0.138699	
WELL USW H-5	0.17206	0.16657	0.79888	0.073690	0.138699	
WELL USW H-6	0.21437	0.30190	0.79888	0.247389	0.085477	
WELL USW H-6	0.20309	0.26026	0.78223	0.205280	0.085477	
WELL USW H-6	0.20873	0.33313	0.81552	0.247389	0.085477	
WELL USW VH-1	0.31027	0.45806	0.83216	0.142117		
WELL USW VH-1	0.28206	0.46847	0.83216	0.142117		
WELL USW VH-1	0.28206	0.45806	0.81552	0.142117		
3. 15S/49E-22DC	0.24000	0.34000	0.82000			
4. 16S/49E-5ACC	0.17000	0.27000	1.03000			
5. 16S/49E-8ABB	0.22000	0.31000	0.90000			
6. 16S/49E-8ACC	0.17000	0.30000	0.97000			
7. 16S/49E-9CDA	0.34000	0.67000	1.09000			
8. 16S/49E-9DCC	0.28000	0.70000	1.20000			
9. 16S/49E-18DC	0.21000	0.29000	0.98000			
10. 16S/49E-16CCC	0.23000	0.53000	1.28000			
11. 16S/49E-19DAA	0.19000	0.34000	1.25000			
12. 16S/49E-24AAA	0.22000	0.31000	1.31000			
13. 16S/48E-25AA	0.26000	0.29000	1.20000			
14. 16S/48E-36AAA	0.19000	0.26000	1.31000			
15. 17S/48E-1AB	0.18000	0.26000	1.31000			
16. 17S/49E-7BB	0.27000	0.32000	1.33000			
17. 17S/49E-9AA	0.28000	0.31000	1.17000			
18. 17S/49E-8DBB	0.18000	0.28000	1.35000			
19. 17S/49E-15BBD	0.28000	0.36000	1.21000			
64. WELL B. NTS	0.21000	0.15000	0.68000	0.040000		
20. 17S/49E-35DDD. ASH TR	0.19000	0.42000	1.34500			
21. 16S/49E-23ADD	0.25000	0.36000	1.27000			
23. 16S/48E-15AAA	0.21000	0.29000	1.13000			
25. 16S/48E-10CBA	0.23000	0.34000	1.07000			
27. 16S/50E-7BCD	0.82000	1.58000	0.48000			
29. 16S/49E-15AAA	0.65000	1.35000	0.77000			
30. 16S/49E-36AAA	0.76000	1.75000	0.63000			
45. 16S/48E-8BBBA	2.25000	2.11000	0.63000			
46. 16S/48E-7BBA	1.78000	1.87000	1.15000			
47. 16S/48E-7CBC	1.75000	1.87000	1.07000			
48. 16S/48E-18BCC	1.72000	1.98000	1.33000			

SITE	SAS	CL	SD4	S102	F	NO3
49.	16S/48E-17CCC	2.34000	2.45000	1.29000	1.29000	
50.	16S/48E-18DAD	1.78000	1.95000	1.28000	1.28000	
51.	16S/48E-8CDA	1.89000	1.87000	1.13000	1.13000	
52.	16S/48E-17ABB	1.95000	1.86000	1.25000	1.25000	
62.	WHITEROCK SPRING	0.31000	0.28000	0.77000	0.77000	
68.	BUTTE SPRING	0.34000	0.12000	0.50000	0.50000	
70.	OAK SPRING	0.25000	0.15000	0.95000	0.95000	
71.	TOPOPAH SPRING	0.08000	0.16000	0.87000	0.87000	
1.	10S/47E-14B4B	1.44000	0.86000	0.95000	0.95000	0.220000
2.	10S/47E-27CBA	1.83000	1.06000	1.03000	1.03000	0.990000
3.	10S/47E-31AAB	1.18000	0.55000	1.18000	1.18000	0.130000
4.	10S/47E-32DDA	1.04000	1.00000	1.03000	1.03000	0.120000
5.	10S/47E-33AAC	1.92000	1.07000	0.90000	0.90000	0.230000
6.	10S/47E-30DCC	1.13000	0.61000	1.20000	1.20000	0.090000
7.	11S/46E-26BBBB	0.39000	0.15000	0.73000	0.73000	0.020000
8.	10S/46E-26BCC	0.42000	0.18000	0.80000	0.80000	0.020000
9.	11S/47E-3CDB	0.98000	0.78000	0.15000	0.15000	
10.	11S/47E-4CAD	1.27000	1.27000	0.270000	0.270000	
11.	11S/47E-10CAA	2.26000	1.35000	1.03000	1.03000	0.320000
12.	11S/47E-10BCC	1.52000	1.13000	0.63000	0.63000	0.320000
13.	11S/47E-16DCD	1.33000	0.95000	0.85000	0.85000	0.240000
14.	11S/47E-16BDC	1.21000	1.25000	1.07000	1.07000	0.320000
15.	11S/47E-18ACD	0.59000	0.23000	0.75000	0.75000	0.030000
16.	11S/47E-21ACC	1.95000	1.65000	1.00000	1.00000	0.320000
17.	11S/47E-21DBB	2.03000	1.74000	0.93000	0.93000	0.320000
18.	11S/47E-21ABA	2.03000	1.74000	0.90000	0.90000	0.320000
19.	11S/47E-21ABA	0.99000	1.21000	0.90000	0.90000	0.320000
20.	11S/47E-27CBA	2.62000	2.27000	0.98000	0.98000	0.370000
21.	11S/47E-28AAC	1.92000	1.76000	0.75000	0.75000	0.340000
22.	11S/47E-28DAC	0.76000	0.73000	0.77000	0.77000	0.200000
23.	11S/47E-33BAC	1.27000	0.85000	0.83000	0.83000	0.140000
24.	11S/47E-10CCB	0.73000	0.73000	1.00000	1.00000	0.200000
25.	12S/47E-5CDA	1.07000	0.97000	0.90000	0.90000	0.020000
26.	12S/47E-6CDD	2.06000	1.86000	1.320000	1.320000	
27.	12S/47E-7DBD	2.06000	1.36000	1.10000	1.10000	0.320000
28.	12S/47E-20B8B	2.17000	1.91000	1.12000	1.12000	0.310000
29.	12S/47E-19ADC	2.82000	2.60000	1.12000	1.12000	0.330000
18B.	12S/47E-20
19B.	12S/47E-20
20B.	12S/47E-20
21B.	12S/47E-20
22B.	12S/47E-20
WELL	UE-19B-1	0.19180	0.21862	0.68237	0.68237	0.0064511
WELL	UE-19C	0.21719	0.00052	0.49930	0.226335	0.0032256
WELL	UE-19D	0.56413	0.59339	0.91538	0.257916	0.0096767
WELL	UE-19E	0.10436	0.16657	0.93202	0.278971	0.0274172
WELL	UE-19F	0.12975	0.08328	1.08181	0.036845	0.0451577
WELL	UE-19G	0.62054	0.44765	0.83216	0.157908	0.0096767
WELL	UE-19GS	0.25104	0.78078	0.99860	0.073690	0.0048383
WELL	U-20A-2	0.31027	0.28108	0.79888	0.142117	0.0225789
WELL	U-20A-2	0.28206	0.29149	0.73230	0.147381	0.0112894
WELL	UE-20D	0.64875	0.45806	0.78223	0.147381	0.0016128
WELL	UE-20E-1	0.56413	0.43724	0.59916	0.236862	0.0080639
WELL	UE-20H	0.42310	0.31231	0.81552	0.142117	0.0209661

SITE	SAS	CL	504	S102	F	NU3
WELL UE-20J	3. 24373	1. 40541	0. 73230	0. 115799	0. 014515	0. 014515
WELL ARMY-1	0. 42310	0. 53093	0. 31622	0. 052636	0. 014515	0. 014515
WELL 5A	0. 31027	0. 28108	0. 83216	0. 121063	0. 120958	0. 120958
WELL 5B	0. 59233	0. 54134	0. 68237	0. 036845	0. 0177405	0. 0177405
WELL 5C	0. 23411	0. 23944	0. 78223	0. 042109	0. 088703	0. 088703
WELL C	0. 93081	0. 68709	0. 48266	0. 057900	0. 000806	0. 000806
WELL C-1	0. 93081	0. 68709	0. 48266	0. 052636	0. 000806	0. 000806
WELL 3	0. 11283	0. 19780	1. 06517	0. 047372	0. 193533	0. 193533
WELL A	0. 13257	0. 17698	0. 14839	0. 031582	0. 111282	0. 111282
WELL 2	0. 16924	0. 21862	0. 73230	0. 021054	0. 082252	0. 082252
WELL UE-15D	0. 42310	0. 45806	0. 31622	0. 073690	0. 000806	0. 000806
UE 12T#3-5.	134.4 (M)	0. 76157	0. 21341	1. 03188	0. 73230	0. 73230
UE 12T#3-6.	169.6 (M)	0. 90260	0. 37790	0. 99860	0. 73230	0. 73230
UE 12T#3-7.	199.2 (M)	0. 76157	0. 42370	0. 96531	0. 73230	0. 73230
UE 12T#3-8.	202.4 (M)	1. 07184	0. 41954	1. 18167	0. 73230	0. 73230
UE 12T#3-9.	257.6 (M)	1. 46673	0. 40184	1. 01524	0. 73230	0. 73230
UE 12T#3-10.	260.0 (M)	1. 74879	0. 73602	1. 26489	0. 73230	0. 73230
UE 12T#3-11.	441.4 (M)	0. 56413	0. 57049	0. 83216	0. 73230	0. 73230
UE 12T#3-12.	442.4 (M)	0. 56413	0. 86823	0. 89874	0. 73230	0. 73230
UE 12T#3-13.	291.4 (M)	0. 59233	0. 22903	1. 26489	0. 73230	0. 73230
UE 12T#3-14.	320.3 (M)	0. 84619	0. 46847	0. 93202	0. 73230	0. 73230
UE 12T#3-15.	320.6 (M)	0. 98722	0. 52052	1. 29818	0. 73230	0. 73230
UE 12T#3-16.	321.3 (M)	0. 81793	0. 31752	0. 64909	0. 73230	0. 73230
UE 12T#3-17.	350.8 (M)	0. 84619	0. 22903	1. 26489	0. 73230	0. 73230
UE 12T#3-18.	411.2 (M)	0. 45130	0. 33834	0. 99860	0. 73230	0. 73230
UE 12T#3-19.	470.6 (M)	0. 26514	0. 21862	0. 76559	0. 73230	0. 73230
UE 12T#3-20.	472.4 (M)	0. 31027	0. 34354	0. 73230	0. 73230	0. 73230
UE 12T#3-21.	501.7 (M)	0. 33848	0. 31752	1. 29818	0. 73230	0. 73230
UE 12T#3-22.	503.2 (M)	0. 47951	0. 31231	0. 66573	0. 73230	0. 73230
UE 12T#3-23.	532.8 (M)	0. 47951	0. 37790	0. 69902	0. 73230	0. 73230
RML 1A . SURFACE	0. 08180	0. 04997	0. 32954	0. 32454	0. 05000	0. 01000
RML 1B . SURFACE	0. 04513	0. 04581	0. 32454	0. 05000	0. 01000	0. 01000
1. U12N.05 BYPASS	0. 16000	0. 06000	0. 85000	0. 01000	0. 01000	0. 01000
2. U12N.05	0. 37000	0. 15000	0. 72000	0. 01000	0. 01000	0. 01000
3. U12N.05	0. 37000	0. 28000	0. 70000	0. 01000	0. 01000	0. 01000
4. U12N MAIN	0. 90000	0. 66000	0. 68000	0. 03000	0. 02000	0. 02000
5. U12N.07 BYPASS	0. 06000	0. 05000	0. 63000	0. 01000	0. 01000	0. 01000
6. U12N.02	0. 20000	0. 09000	0. 88000	0. 01000	0. 02000	0. 02000
7. U12T.02 BYPASS	0. 02000	0. 01000	0. 65000	0. 02000	0. 01000	0. 01000
8. U12T.02 BYPASS	0. 05000	0. 01000	0. 63000	0. 01000	0. 01000	0. 01000
9. U12T.02 BYPASS	0. 28000	0. 14000	0. 65000	0. 02000	0. 01000	0. 01000
10. U12T MAIN	0. 27000	0. 18000	1. 35000	0. 01000	0. 01000	0. 01000
11. U12T.03	0. 26000	0. 17000	1. 00000	0. 92000	0. 01000	0. 01000
12. U12T.03	0. 31000	0. 19000	0. 90000	0. 88000	0. 01000	0. 01000
13. U12T.04	0. 28000	0. 16000	1. 00000	0. 02000	0. 01000	0. 01000
14. U12T.03	0. 31000	0. 16000	1. 00000	0. 90000	0. 02000	0. 02000
15. U12T.03	0. 28000	0. 16000	0. 83000	0. 01000	0. 01000	0. 01000
16. U12T.03	0. 26000	0. 17000	1. 00000	0. 92000	0. 01000	0. 01000
17. U12E.07	0. 06000	0. 15000	0. 90000	0. 01000	0. 01000	0. 01000
18. U12E.04	0. 10000	0. 13000	0. 60000	0. 01000	0. 01000	0. 01000
19. U12E	0. 17000	0. 12000	0. 90000	0. 02000	0. 02000	0. 02000
20. U12E	0. 11000	0. 10000	0. 87000	0. 01000	0. 01000	0. 01000
21. U12E.04	0. 17000	0. 09000	0. 60000	0. 01000	0. 01000	0. 01000
22. U12E.03	0. 06000	0. 09000	0. 67000	0. 02000	0. 02000	0. 02000

14:49 THURSDAY, MAY 29, 1986 20

SITE	SAS	SD4	CL	S102	F	NO3
23, U12E	0.230000	0.180000	0.92000	0.0100000	0.0100000	0.0100000
24, U12E.02	0.270000	0.120000	0.78000	0.0100000	0.0100000	0.0100000
25, U12E.05	0.280000	0.120000	1.23000	0.0100000	0.0100000	0.0100000
26, U12E.03	0.250000	0.120000	0.98000	0.0200000	0.0200000	0.0200000
27, U12E.03	0.280000	0.120000	0.63000	0.0100000	0.0100000	0.0100000
28, U12E.03	0.230000	0.120000	0.97000	0.0200000	0.0200000	0.0200000
29, U12E.07	0.110000	0.150000	0.90000	0.0100000	0.0100000	0.0100000
30, U12E	0.280000	0.150000	1.02000	0.0100000	0.0100000	0.0100000
31, U12E	0.340000	0.150000	0.67000	0.0100000	0.0100000	0.0100000
32, U12E	0.340000	0.150000	0.87000	0.0100000	0.0100000	0.0100000
33, U12E	0.450000	0.330000	2.10000	0.0200000	0.0200000	0.0200000
34, U12B	0.140000	0.120000	1.11000	0.0100000	0.0100000	0.0100000
35, U12B.03	0.210000	0.090000	1.10000	0.0100000	0.0100000	0.0100000
36, U12.04	0.340000	0.340000	0.156156	0.66573	0.66573	0.66573
SEEP 1, U12T	0.310270	0.183341	0.135335	0.73230	0.73230	0.73230
SEEP 2, U12N.03	0.183341	0.217189	0.218619	0.89874	0.89874	0.89874
SEEP 3, U12N.05	0.183341	0.256678	0.156156	0.78223	0.78223	0.78223
SEEP 4, U12N.03	0.183341	0.115646	0.012493	0.63244	0.63244	0.63244
LYSIMETER 1, U12N.05	0.115646	0.118467	0.017698	0.54923	0.54923	0.54923
LYSIMETER 2, U12N.05	0.118467	0.183341	0.045806	0.91538	0.91538	0.91538
LYSIMETER 3, U12N.05	0.183341	0.160776	0.067668	0.88209	0.88209	0.88209
LYSIMETER 4, U12N.05	0.160776	0.039489	0.023944	0.59916	0.59916	0.59916
LYSIMETER 5, U12N.05	0.039489	0.141032	0.016657	0.38280	0.38280	0.38280
LYSIMETER 6, SURF SOIL	0.141032	0.902603	0.093694	0.29958	0.29958	0.29958
LYSIMETER 7, SURF SOIL	0.902603	0.141032	0.124925	1.13174	0.0052636	0.033868
LYSIMETER 9, SURF SOIL	0.141032	0.211548	0.085365	1.11510	0.0105272	0.119345
ANAL 2589, TUNNEL U12B, E	0.085365	0.267960	0.114515	0.78223	0.035481	0.064545
ANAL 2819, TUNNEL U12B, D	0.267960	0.098722	0.098722	0.0105272	0.027417	0.053222
ANAL 2913, TUNNEL U12E, D	0.098722	0.114515	0.114515	0.0105272	0.027417	0.053222
ANAL 3260, TUNNEL U12E, M	0.114515	0.169238	0.89874	0.0157908	0.0157908	0.0157908

N= 193

SITE	HC03	PH	DEL C13	DEL PMC	SAS
WELL J-12	1.95027	7.10000	-7.900	32.2000	
WELL J-13	2.03222	7.20000	-7.300	29.2000	
WELL UE-25B#1	2.83527	7.10000	-10.700	16.7000	
WELL UE-25B#1	2.27805	7.50000	-10.400	18.9000	
WELL UE-25B#1	2.17972	7.10000	-8.600	15.0000	
WELL UE-25C#1	2.47472	7.60000	-7.100	16.6000	
WELL UE-25C#2	2.27805	7.70000	-7.000	15.7000	
WELL UE-25C#3	2.24527	7.70000	-7.500	3.5000	
WELL UE-25P#1	4.62166	6.80000	-4.200	2.3000	
WELL UE-25P#1	9.32525	6.60000	-2.300	62.3000	
WELL UE-29A#2	1.75361	7.20000	-13.000	60.0000	
WELL UE-29A#2	1.75361	7.00000	-13.100	22.0000	
WELL USW G-4	2.27805	7.70000	-9.100	19.9000	
WELL USW H-1	1.88472	7.70000	-11.400	23.9000	
WELL USW H-1	1.99944	7.50000	-11.400	10.5000	
WELL USW H-3	4.49054	9.20000	-4.900	11.8000	
WELL USW H-4	2.83527	7.40000	-7.400	18.2000	
WELL USW H-5	2.06499	7.80000	-10.300	21.4000	
WELL USW H-5	2.08138	7.90000	-10.300	16.3000	
WELL USW H-6	2.98277	8.10000	-7.500	10.0000	
WELL USW H-6	3.55638	8.30000	-7.300	12.4000	
WELL USW H-6	3.83499	8.30000	-7.100		
WELL USW VH-1	2.73694	7.90000			
WELL USW VH-1	2.70416	7.50000	-8.500	12.2000	
WELL USW VH-1	2.65499	7.50000	-8.500	15.6000	
3. 155/49E-22DC	2.44000	7.78000	-7.100	19.3000	
4. 165/49E-5ACC	2.21000	8.15000	-7.100	21.4000	
5. 165/49E-8ABB	2.49000	7.47000	-6.800		
6. 165/49E-8ACC	2.26000	7.90000			
7. 165/49E-9CDA	2.35000	7.61000			
8. 165/49E-9DCC	2.31000	8.16000	-7.300	21.9000	
9. 165/49E-18CC	2.46000	8.12000	-7.300	28.4000	
10. 165/49E-16CCC	2.17000	7.87000	-5.200	24.8000	
11. 165/49E-19DAA	2.20000	8.20000	-5.200	20.8000	
12. 165/48E-24AAA	2.41000	8.09000			
13. 165/48E-25AA	2.18000	8.06000		19.3000	
14. 165/48E-36AAA	2.18000	8.40000			
15. 175/48E-1AB	2.21000	8.15000		18.4000	
16. 175/49E-7BB	2.20000	8.30000		10.0000	
17. 175/49E-9AA	2.15000	8.02000		18.9000	
18. 175/49E-8DB	2.02000	8.35000		27.8000	
19. 175/49E-15BBB	1.97000	8.12000		40.3000	
64. WELL 8, NTS	1.31000	7.40000	-12.100	25.4000	
20. 175/49E-35DD, ASH TR	2.58000	7.96000	13.8000		
21. 165/49E-23ADD	2.08000	8.19000	-8.400	27.4000	
23. 165/48E-15AAA	2.51000	8.10000	-7.100	17.1000	
25. 165/48E-10CBA	2.72000	8.30000	-5.600	15.6000	
27. 165/50E-7BCD	4.78000	7.62000	-3.600	7.0000	
29. 165/49E-15AAA	3.20000	7.70000	-3.400		
30. 165/49E-36AAA	5.15000	7.76000	-4.400	10.3000	
45. 165/48E-8BBA	4.85000	7.90000			
46. 165/48E-7BBA	4.11000	7.40000			
47. 165/48E-7CBC	3.92000	7.70000	-6.200	31.4000	

14:49 THURSDAY, MAY 29, 1986 22

SITE	SAS	HC03	PH	DEL C13	PMC
48.	165/48E-18BBC	4.45000	7.98000	-	-
49.	165/48E-17CCC	3.92000	7.69000	-5.700	-
50.	165/48E-18DAD	3.87000	7.69000	-	-
51.	165/48E-8CDA	4.33000	7.60000	-	-
52.	165/48E-17ABB	4.95000	7.40000	-	-
62.	WHITEROCK SPRING	1.26000	7.30000	-	-
68.	BUTTE SPRING	2.28000	7.80000	-	-
70.	OAK SPRING	1.90000	7.50000	-	-
71.	TOPOPAH SPRING	0.79000	6.90000	-	-
1.	105/47E-14BAB	3.39000	8.10000	-2.520	-
2.	105/47E-27CBA	4.57000	7.70000	-5.330	38.9000
3.	105/47E-31AAB	3.75000	7.60000	-5.320	-
4.	105/47E-32DDA	5.08000	7.60000	-5.990	-
5.	105/47E-33AAB	4.85000	7.80000	-4.910	99.9900
6.	105/47E-30DCC	3.80000	7.80000	-	35.0000
7.	115/46E-26BBBB	1.90000	8.70000	-	-
8.	105/46E-26BBC	2.07000	7.90000	-	-
9.	115/47E-3CDB	3.00000	8.20000	-	-
10.	115/47E-4CAD	6.23000	7.70000	-5.020	-
11.	115/47E-10CAA	5.41000	8.10000	-	-
12.	115/47E-10BCC	4.79000	7.60000	-	-
13.	115/47E-16DCD	4.39000	7.80000	-	-
14.	115/47E-16BDC	4.15000	7.80000	-	-
15.	115/47E-18ACD	2.33000	7.70000	-	-
16.	115/47E-21ACC	6.10000	7.70000	-4.540	-
17.	115/47E-21DBB	6.44000	7.70000	-	-
18.	115/47E-21ABA	6.46000	7.90000	-	-
19.	115/47E-21ABA	3.65000	7.60000	-	-
20.	115/47E-27CBA	3.03000	8.00000	-6.260	-
21.	115/47E-28AAC	8.39000	9.10000	-7.070	-
22.	115/47E-28DAC	5.08000	8.20000	-6.920	-
23.	115/47E-33BAC	2.84000	8.30000	-	-
24.	115/47E-10CCB	3.03000	8.20000	-	-
25.	125/47E-5CDA	3.21000	7.90000	-6.290	-
26.	125/47E-6CDD	3.47000	7.90000	-6.580	-
27.	125/47E-7DBD	6.49000	7.70000	-6.950	64.7000
28.	125/47E-20BBB	6.39000	7.80000	-8.000	-
29.	125/47E-19ADC	7.20000	7.70000	-8.000	92.8000
18B.	125/47E-20	-	-	-13.050	-
19B.	125/47E-20	-	-	-11.140	-
20B.	125/47E-20	-	-	-11.350	-
21B.	125/47E-20	-	-	-9.590	-
22B.	125/47E-20	-	-	-11.390	-
WELL.	UE-19B-1	2.45833	7.40000	-	-
WELL.	UE-19C	6.55554	7.50000	-	-
WELL.	UE-19D	8.01415	7.90000	-	-
WELL.	UE-19E	1.31111	8.20000	-	-
WELL.	UE-19F	1.32750	7.70000	-	-
WELL.	UE-19G	2.01583	8.20000	-	-
WELL.	UE-19G	2.99916	8.20000	-	15.3000
WELL.	UE-19G	1.73722	7.70000	-	-13.470
WELL.	U-20A-2	1.80277	7.90000	-	-13.470
WELL.	U-20A-2	2.24527	7.80000	-	15.3000

WELL UE-20E-1
 WELL UE-20H
 WELL UE-20J
 WELL ARMY-1
 WELL 5A
 WELL 5B
 WELL 5C
 WELL C
 WELL C-1
 WELL 3
 WELL A
 WELL 2
 WELL UE-15D
 UE12T#3-5, 134.4 (M)
 UE12T#3-6, 169.6 (M)
 UE12T#3-7, 199.2 (M)
 UE12T#3-8, 202.4 (M)
 UE12T#3-9, 257.6 (M)
 UE12T#3-10, 260.0 (M)
 UE12T#3-11, 441.4 (M)
 UE12T#3-12, 442.4 (M)
 UE12T#3-13, 291.4 (M)
 UE12T#3-14, 320.3 (M)
 UE12T#3-15, 320.6 (M)
 UE12T#3-16, 321.3 (M)
 UE12T#3-17, 350.8 (M)
 UE12T#3-18, 411.2 (M)
 UE12T#3-19, 470.6 (M)
 UE12T#3-20, 472.4 (M)
 UE12T#3-21, 501.7 (M)
 UE12T#3-22, 503.2 (M)
 UE12T#3-23, 532.8 (M)
 RML 1A, SURFACE
 1. U12N.05 BYPASS
 2. U12N.05
 3. U12N.05
 4. U12N MAIN
 5. U12N.07 BYPASS
 6. U12N.02
 7. U12T.02 BYPASS
 8. U12T.02
 9. U12T.02 BYPASS
 10. U12T MAIN
 11. U12T.03
 12. U12T.03
 13. U12T.04
 14. U12T.03
 15. U12T.03
 16. U12T.03
 17. U12E.07
 18. U12E.04
 19. U12E

1. 95027 8. 50000
 1. 75361 8. 10000
 2. 45833 7. 00000
 4. 29388 8. 00000
 6. 29332 8. 20000
 2. 96638 7. 90000
 5. 17888 8. 10000
 9. 50553 7. 30000
 9. 65303 7. 60000
 3. 22860 7. 70000
 3. 52360 7. 80000
 3. 22860 7. 80000
 6. 52276 7. 90000
 0. 95055 7. 60000
 2. 24527 7. 70000
 2. 24527 7. 70000
 0. 44250 7. 60000
 0. 43103 7. 20000
 0. 37367 7. 50000
 1. 08986 7. 60000
 1. 06855 7. 60000
 0. 78830 7. 80000
 0. 83911 7. 70000
 0. 57197 7. 80000
 1. 78638 8. 20000
 0. 96530 7. 90000
 0. 60967 7. 80000
 0. 91450 7. 90000
 1. 40944 7. 90000
 1. 72083 8. 10000
 1. 83555 8. 10000
 1. 90111 8. 00000
 1. 22425 7. 80000
 1. 75361 7. 90000
 2. 29000 8. 00000
 2. 85000 7. 60000
 3. 39000 8. 30000
 3. 62000 8. 20000
 0. 87000 8. 04000
 2. 25000 7. 31000
 0. 92000 6. 90000
 1. 08000 7. 00000
 0. 82000 7. 50000
 2. 44000 7. 50000
 2. 23000
 1. 46000
 1. 61000
 2. 18000
 1. 85000
 2. 15000
 2. 00000
 2. 00000
 0. 79000
 0. 87000

14:49 THURSDAY, MAY 29, 1986 24

SITE	SAS	HC03	PH	DEL C13	PMC
20, U12E		0.9300	7.00000	.	.
21, U12E.04		0.8800	6.90000	.	.
22, U12E.03		1.1500	7.10000	.	.
23, U12E		0.9200	7.80000	.	.
24, U12E.02		0.9800	7.40000	.	.
25, U12E.05		1.0500	7.40000	.	.
26, U12E.03		1.3400	7.50000	.	.
27, U12E.03		1.3400	8.00000	.	.
28, U12E.03		1.3800	7.50000	.	.
29, U12E.07		2.0000	7.40000	.	.
30, U12E		1.6100	8.00000	.	.
31, U12E		1.7700	6.80000	.	.
32, U12E		2.0000	7.90000	.	.
33, U12E		2.1000	7.00000	.	.
34, U12B		1.2100	7.60000	.	.
35, U12B.03		0.7900	6.90000	.	.
36, U12.04		0.9800	7.50000	.	.
SEEP 1, U12T		2.4747	7.90000	.	.
SEEP 2, U12N.03		1.8192	6.80000	.	.
SEEP 3, U12N.05		2.6878	6.80000	.	.
SEEP 4, U12N.03		1.7536	7.20000	.	.
LYSIMETER 1, U12N.05		10.7347	6.60000	.	.
LYSIMETER 2, U12N.05		2.7205	6.80000	.	.
LYSIMETER 3, U12N.05		3.1794	6.80000	.	.
LYSIMETER 4, U12N.05		2.7205	6.80000	.	.
LYSIMETER 5, U12N.05		0.9047	7.10000	.	.
LYSIMETER 6, SURF SOIL	
LYSIMETER 7, SURF SOIL		1.7208	7.10000	.	.
LYSIMETER 9, SURF SOIL		3.0483	7.30000	.	.
ANAL 2589, TUNNEL U12B, E		1.2128	7.60000	.	.
ANAL 2819, TUNNEL U12B, D		0.7867	7.60000	.	.
ANAL 2913, TUNNEL U12E, D		0.9833	7.40000	.	.
ANAL 3260, TUNNEL U12E, D		0.7867	7.30000	.	.
ANAL 3541, TUNNEL U12E, M		0.8686	7.60000	.	.

N= 193

SITE	SAS	DEL D	DEL 018	DEL 02 (MG/L)
WELL J-12		-97.50	-12.800	5.70000
WELL J-13		-97.50	-13.000	1.80000
WELL UE-25B#1		-99.50	-13.400	
WELL UE-25B#1		-101.00	-13.400	
WELL UE-25B#1		-99.50	-13.500	1.60000
WELL UE-25C#1		-102.00	-13.500	
WELL UE-25C#2		-100.00	-13.400	
WELL UE-25C#3		-103.00	-13.500	
WELL UE-25P#1		-106.00	-13.500	
WELL UE-25P#1		-106.00	-13.800	1.30000
WELL UE-29A#2		-93.50	-12.800	5.40000
WELL UE-29A#2		-93.00	-12.800	5.60000
WELL USW G-4		-103.00	-13.800	6.40000
WELL USW H-1		-103.00	-13.400	
WELL USW H-1		-101.00	-13.500	
WELL USW H-3		-101.00	-13.900	-0.10000
WELL USW H-4		-104.00	-14.000	5.80000
WELL USW H-5		-102.00	-13.600	6.30000
WELL USW H-5		-102.00	-13.600	
WELL USW H-6		-106.00	-13.800	5.90000
WELL USW H-6		-105.00	-14.000	
WELL USW H-6		-107.00	-14.000	
WELL USW VH-1				
WELL USW VH-1		-108.00	-14.200	
WELL USW VH-1		-102.00	-12.800	
3. 165/49E-22DC		-103.00	-13.200	
4. 165/49E-5ACC		-99.50	-13.200	
5. 165/49E-8ABB				
6. 165/49E-8ACC				
7. 165/49E-9CDA				
8. 165/49E-9DCC		-103.00	-13.400	
9. 165/49E-18DC		-102.00	-12.600	
10. 165/49E-16CCC		-97.50	-13.200	
11. 165/49E-19DAA		-101.00	-13.100	
12. 165/49E-24AAA				
13. 165/49E-25AA		-102.00	-13.000	
14. 165/49E-36AAA		-98.50	-12.600	
15. 175/49E-1AB		-104.00	-13.000	
16. 175/49E-7BB		-104.00	-12.700	
17. 175/49E-9AA		-105.00	-12.800	
18. 175/49E-80DB		-102.00	-13.000	
19. 175/49E-15BBBD				
64. WELL 8, NTS		-104.00	-13.000	
20. 175/49E-35DDN, ASH TR		-102.00	-12.400	
21. 165/49E-23ADD		-99.00	-13.200	
23. 165/49E-15AAA		-103.00	-13.400	
25. 165/49E-10CBA		-102.00	-13.400	
27. 165/50E-7BCD		-105.00	-13.800	
29. 165/49E-15AAA		-105.00	-13.800	
30. 165/49E-36AAA		-104.00	-13.700	
45. 165/49E-8BBA				
46. 165/49E-7BBA				
47. 165/49E-7CBC				-13.100

SITE	SAS	DEL D	DEL 018	02 (MG/L)
48.	16S/48E-18BCC	.	.	.
49.	16S/48E-17CCC	- 104.00	- 13.600	
50.	16S/48E-18DAD	.	.	
51.	16S/48E-8CDA	.	.	
52.	16S/48E-17ABB	.	.	
62.	WHITEROCK SPRING	.	.	
68.	BUTTE SPRING	.	.	
70.	OAK SPRING	.	.	
71.	TOPOPAH SPRING	.	.	
1.	10S/47E-14BAB	- 112.50	- 14.520	
2.	10S/47E-27CBA	- 110.50	- 14.320	
3.	10S/47E-31AAB	- 102.00	- 13.390	
4.	10S/47E-32DDA	- 102.00	- 13.390	
5.	10S/47E-33AAB	- 108.00	- 14.020	
6.	10S/47E-30DCC	- 102.00	- 13.420	
7.	11S/46E-26BBB	.	.	
8.	10S/46E-26BCC	.	.	
9.	11S/47E-3CDB	.	.	
10.	11S/47E-4CAD	- 107.50	- 14.020	
11.	11S/47E-10CAA	.	.	
12.	11S/47E-10BCC	.	.	
13.	11S/47E-16DCD	- 110.00	.	
14.	11S/47E-16BDC	.	.	
15.	11S/47E-18ACD	.	.	
16.	11S/47E-21ACC	.	.	
17.	11S/47E-21DBB	.	.	
18.	11S/47E-21ABA	.	.	
19.	11S/47E-21ABA	.	.	
20.	11S/47E-27CBA	- 108.00	- 14.090	
21.	11S/47E-28AAC	- 109.00	- 14.140	
22.	11S/47E-28DAC	.	.	
23.	11S/47E-33BAC	.	.	
24.	11S/47E-10CCB	.	.	
25.	12S/47E-5CDA	- 108.00	- 14.090	
26.	12S/47E-6CDD	- 102.00	- 13.300	
27.	12S/47E-7DBD	- 107.50	- 13.880	
28.	12S/47E-20BBB	- 106.00	- 13.570	
29.	12S/47E-19ADC	- 104.00	- 13.300	
18B.	12S/47E-20	- 105.00	- 13.400	
19B.	12S/47E-20	- 101.00	- 13.800	
20B.	12S/47E-20	- 102.00	- 13.400	
21B.	12S/47E-20	- 102.00	- 13.400	
22B.	12S/47E-20	- 104.00	- 13.100	
WELL	UE-19B-1	.	.	
WELL	UE-19C	.	.	
WELL	UE-19D	.	.	
WELL	UE-19E	.	.	
WELL	UE-19E	.	.	
WELL	UE-19GS	- 113.50	- 14.500	
WELL	UE-19GS	- 114.00	- 14.750	
WELL	U-20A-2	- 114.00	- 14.750	
WELL	U-20A-2	.	.	
WELL	UE-20D	.	.	

SITE	DEL D	DEL 018	O2 (MG/L)
WELL UE-20E-1			
WELL UE-20H			
WELL UE-20J			
WELL ARMY-1			
WELL 5A			
WELL 5B			
WELL 5C			
WELL C			
WELL C-1			
WELL 3			
WELL A			
WELL 2			
WELL UE-15D			
UE12T#3-5, 134.4 (M)			
UE12T#3-6, 169.6 (M)			
UE12T#3-7, 199.2 (M)			
UE12T#3-8, 202.4 (M)			
UE12T#3-9, 257.6 (M)			
UE12T#3-10, 260.0 (M)			
UE12T#3-11, 441.4 (M)			
UE12T#3-12, 442.4 (M)			
UE12T#3-13, 291.4 (M)			
UE12T#3-14, 320.3 (M)			
UE12T#3-15, 320.6 (M)			
UE12T#3-16, 321.3 (M)			
UE12T#3-17, 350.8 (M)			
UE12T#3-18, 411.2 (M)			
UE12T#3-19, 470.6 (M)			
UE12T#3-20, 472.4 (M)			
UE12T#3-21, 501.7 (M)			
UE12T#3-22, 503.2 (M)			
UE12T#3-23, 532.8 (M)			
RML 1A, SURFACE			
RML 1B, SURFACE			
1. U12N.05 BYPASS			
2. U12N.05			
3. U12N.05			
4. U12N MAIN			
5. U12N.07 BYPASS			
6. U12N.02			
7. U12T.02 BYPASS			
8. U12T.02			
9. U12T.02 BYPASS			
10. U12T MAIN			
11. U12T.03			
12. U12T.03			
13. U12T.04			
14. U12T.03			
15. U12T.03			
16. U12T.03			
17. U12E.07			
18. U12E.04			
19. U12E			

SAS

14:50 THURSDAY, MAY 29, 1986 28

SITE	DEL D	DEL 018	O2 (MG/L)
20, U12E	.	.	.
21, U12E.04	.	.	.
22, U12E.03	.	.	.
23, U12E	.	.	.
24, U12E.02	.	.	.
25, U12E.05	.	.	.
26, U12E.03	.	.	.
27, U12E.03	.	.	.
28, U12E.03	.	.	.
29, U12E.07	.	.	.
30, U12E	.	.	.
31, U12E	.	.	.
32, U12E	.	.	.
33, U12E	.	.	.
34, U12B	.	.	.
35, U12B.03	.	.	.
36, U12.04	.	.	.
SEEP 1, U12T	.	.	.
SEEP 2, U12N.03	.	.	.
SEEP 3, U12N.05	.	.	.
SEEP 4, U12N.03	.	.	.
LYSIMETER 1, U12N.05	.	.	.
LYSIMETER 2, U12N.05	.	.	.
LYSIMETER 3, U12N.05	.	.	.
LYSIMETER 4, U12N.05	.	.	.
LYSIMETER 5, U12N.05	.	.	.
LYSIMETER 6, SURF SCIL	.	.	.
LYSIMETER 7, SURF SOIL	.	.	.
LYSIMETER 9, SURF SOIL	.	.	.
ANAL 2589, TUNNEL U12B, E	.	.	.
ANAL 2819, TUNNEL U12B, D	.	.	.
ANAL 2913, TUNNEL U12E, D	.	.	.
ANAL 3260, TUNNEL U12E, D	.	.	.
ANAL 3541, TUNNEL U12E, M	.	.	.

N= 193

SITE	SAS	LOG(CO2 PRESS)	CALCITE LOG(Q/K)	DOLOMITE LOG(Q/K)
WELL J-12	-2.0736	-0.9720	-1.6260	
WELL J-13	-2.1153	-0.8480	-1.2880	
WELL UE-25BN#1	-1.8502	-0.5670	-1.3520	
WELL UE-25BN#1	-2.3100	-0.2660	-0.7960	
WELL UE-25BN#1	-1.9533	-0.6710	-1.5390	
WELL UE-25C#1	-2.3302	-0.2400	-0.7640	
WELL UE-25C#2	-2.4698	-0.1470	-0.5500	
WELL UE-25C#3	-2.4737	-0.1860	-0.5880	
WELL UE-25P#1	-1.3322	-0.3450	-0.0130	
WELL UE-25P#1	-0.7946	0.1910	1.2930	
WELL UE-29A#2	-2.2186	-1.0740	-2.7150	
WELL UE-29A#2	-2.0633	-1.3350	-3.0750	
WELL USW G-4	-2.5041	-0.1790	-0.9780	
WELL USW H-1	-2.5995	-0.7400	-2.2570	
WELL USW H-1	-2.3714	-0.7620	-2.4300	
WELL USW H-3	-3.8009	0.0880	-0.2340	
WELL USW H-4	-2.1323	-0.3130	-1.2070	
WELL USW H-5	-2.6335	-0.9270	-2.9340	
WELL USW H-5	-2.7364	-0.8170	-3.0450	
WELL USW H-6	-2.7673	-0.1570	-0.7630	
WELL USW H-6	-2.8650	-0.3090	-1.2260	
WELL USW H-6	-2.8645	0.1790	-0.2560	
WELL USW VH-1	-2.6269	-0.0110	0.3370	
WELL USW VH-1	-2.2421	-0.4610	-0.5500	
WELL USW VH-1	-2.2498	-0.4710	-0.5660	
3. 15S/49E-22DC	-2.6300	0.0800	0.1500	
4. 16S/49E-5ACC	-3.0300	0.4500	0.9100	
5. 16S/49E-8ABB	-2.3300	-0.2100	-0.3600	
6. 16S/49E-8ACC	-	-	-	
7. 16S/49E-9CDA	-3.0400	0.3200	0.8300	
8. 16S/49E-9DCC	-2.9600	0.3000	0.8700	
9. 16S/49E-18DC	-2.7700	0.1600	0.2600	
10. 16S/49E-16CCC	-3.0800	0.4300	0.7100	
11. 16S/49E-19DAA	-2.9200	0.2500	0.2500	
12. 16S/48E-24AAA	-2.9400	0.1900	0.1200	
13. 16S/48E-25AA	-3.2900	0.4600	1.1200	
14. 16S/48E-36AAA	-3.0300	0.2700	0.5600	
15. 17S/48E-1AB	-3.1300	0.5600	1.1100	
16. 17S/49E-7BB	-	-	-	
17. 17S/49E-9AA	-3.2800	0.4600	1.1500	
18. 17S/49E-8DDB	-3.0700	0.1900	0.7800	
19. 17S/49E-15BBB	-	-	-	
64. WELL B. NTS	-	-	-	
20. 17S/49E-35DDD. ASH TR	-	-	-	
21. 16S/49E-23ADD	-	-	-	
23. 16S/48E-15AAA	-	-	-	
25. 16S/48E-10CBA	-	-	-	
27. 16S/50E-7BCD	-	-	-	
29. 16S/49E-15AAA	-	-	-	
30. 16S/49E-36AAA	-	-	-	1.5400
45. 16S/48E-BBBA	-2.4700	0.6800	0.5300	
46. 16S/48E-7BBA	-2.0600	0.0600	1.2800	
47. 16S/48E-7CBC	-2.3700	0.3000	-	

14:50 THURSDAY, MAY 29, 1986 30

 SITE
 SAS
 LOG(CO₂)
 PRESS)

 CALCITE LOG(Q/K)
 LOG(Q/K)

 DOLOMITE
 LOG(Q/K)

48.	165/48E-18BCC	-2.5900	0.7000	1.8600
49.	165/48E-17CCC	-2.3600	0.4200	1.2100
50.	165/48E-18DAD	-2.3600	0.3400	1.0400
51.	165/48E-8CDA	-2.2400	0.2300	0.7600
52.	165/48E-17AB3	-1.9900	0.1800	0.6200
62.	WHITE ROCK SPRING	.	.	.
68.	BUTTE SPRING	.	.	.
70.	OAK SPRING	.	.	.
71.	TOPOPAH SPRING	.	.	.
1.	105/47E-14BAB	-2.7800	-0.0420	-0.3880
2.	105/47E-27CBA	.	.	.
3.	105/47E-31AAB	-2.1700	0.1240	0.6260
4.	105/47E-32DDA	.	.	.
5.	105/47E-33AAB	.	.	.
6.	105/47E-30DCC	.	.	.
7.	115/46E-26BBBB	.	.	.
8.	105/46E-26BCC	-2.9700	0.2690	0.4540
9.	115/47E-3CDB	-2.1900	0.1970	0.7460
10.	115/47E-33AAB	.	.	.
11.	115/47E-10CAA	.	.	.
12.	115/47E-10BCC	.	.	.
13.	115/47E-16DCD	.	.	.
14.	115/47E-4CAD	.	.	.
15.	115/47E-18ACD	-2.1300	0.2720	0.9320
16.	115/47E-21ACC	.	.	.
17.	115/47E-210BB	.	.	.
18.	115/47E-21ABA	-2.1800	-0.1230	-0.4330
19.	115/47E-21ABA	.	.	.
20.	115/47E-27CBA	-3.5000	1.1230	3.0630
21.	115/47E-28AAC	.	.	.
22.	115/47E-28DAD	.	.	.
23.	115/47E-33BAC	.	.	.
24.	115/47E-10CBB	-2.6400	0.3090	0.8930
25.	125/47E-5CDA	.	.	.
26.	125/47E-6CDD	.	.	.
27.	125/47E-7DBD	.	.	.
28.	125/47E-20BBB	-2.2800	0.2690	0.7790
29.	125/47E-19ADC	.	.	.
18B.	125/47E-20	.	.	.
19B.	125/47E-20	.	.	.
20B.	125/47E-20	.	.	.
21B.	125/47E-20	.	.	.
22B.	125/47E-20	.	.	.
WELL	UE-19B-1	-2.2300	-0.2830	-0.4050
WELL	UE-19C	-2.2800	0.3520	-0.2310
WELL	UE-19D	-2.1900	1.0540	2.0050
WELL	UE-19E	-3.2300	-0.4390	-1.2620
WELL	UE-19E	-3.0100	-0.3400	-0.8970
WELL	UE-19GS	-2.6800	-0.7460	-2.1360
WELL	UE-19GS	-2.5800	-0.5320	-1.4760

14:50 THURSDAY, MAY 29, 1986 31

SITE	SAS LOG(CO2 PRESS)	CALCITE LOG(Q/K)	DOLOMITE LOG(Q/K)
WELL UE-20E-1	-3.3800	-1.3200	-1.7570
WELL UE-20H	-3.0300	-1.2700	-2.1430
WELL UE-20J	-1.8200	-0.4280	-1.2140
WELL ARMY-1	-2.5730	0.7790	2.4360
WELL 5A	-2.6480	-0.1780	-0.0490
WELL 5B	-2.6600	-0.3260	-0.0140
WELL 5C	-2.6190	-0.7460	-0.7470
WELL C	-1.5270	0.6410	2.1080
WELL C-1	-1.7980	0.9540	2.7580
WELL 3	-2.4510	-0.0890	0.7720
WELL A	-2.4780	0.1650	1.0230
WELL 2	-2.4680	0.4000	1.6480
WELL UE-15D	-2.2750	0.9840	2.6280
UE 12T#3-5, 134.4(M)	-2.8500	-0.8600	-1.1600
UE 12T#3-6, 169.6(M)	-2.5900	-0.0600	0.4400
UE 12T#3-7, 199.2(M)	-2.5900	-0.1200	0.3400
UE 12T#3-8, 202.4(M)	-3.1900	-1.0200	-1.5200
UE 12T#3-9, 257.6(M)	-2.8300	-1.5000	-2.3800
UE 12T#3-10, 260.0(M)	-3.1700	-1.0600	-1.5400
UE 12T#3-11, 441.4(M)	-2.8000	-0.9300	-1.8300
UE 12T#3-12, 442.4(M)	-2.8100	-0.8700	-1.6500
UE 12T#3-13, 291.4(M)	-3.1300	-0.8700	-1.1900
UE 12T#3-14, 320.3(M)	-3.0100	-0.9100	-1.4200
UE 12T#3-15, 320.6(M)	-3.2700	-0.6900	-1.1500
UE 12T#3-16, 321.3(M)	-3.1800	0.3700	1.0800
UE 12T#3-17, 350.8(M)	-3.1400	-0.5100	-0.6500
UE 12T#3-18, 411.2(M)	-3.2400	-1.1300	-1.9600
UE 12T#3-19, 470.6(M)	-3.1600	-2.2900	-4.0600
UE 12T#3-20, 472.4(M)	-2.9700	-2.1300	-3.4800
UE 12T#3-21, 501.7(M)	-3.0800	-0.8100	-1.6500
UE 12T#3-22, 503.2(M)	-3.0600	-0.8700	-1.7200
UE 12T#3-23, 532.8(M)	-2.9400	-1.2200	-2.6800
RML 1A, SURFACE	-2.9300	-0.5700	-0.6500
RML 1B, SURFACE	-2.8800	-0.3000	0.0030
1, U12N.05 BYPASS	-2.8600	-0.2600	-0.5500
2, U12N.05	-2.3800	-0.4900	-1.1200
3, U12N.05	-3.0000	0.5000	1.1100
4, U12N MAIN	.	.	.
5, U12N.07 BYPASS	.	.	.
6, U12N.02	-2.2100	-0.9800	-2.3000
7, U12T.02 BYPASS	-2.2400	-3.0300	.
8, U12T.02	-2.2500	-2.3700	-4.3200
9, U12T.02 BYPASS	.	.	.
10, U12T MAIN	-2.3500	-1.7100	-2.8000
11, U12T.03	.	.	.
12, U12T.03	.	.	.
13, U12T.04	.	.	.
14, U12T.03	.	.	.
15, U12T.03	-2.1800	-1.1000	-1.8800
16, U12T.03	.	.	.
17, U12E.07	-2.3500	-0.5600	-0.6200
18, U12E.04	.	.	.
19, U12E	-2.8900	-1.3500	.

14:50 THURSDAY, MAY 29, 1986 32

SITE	SAS LOG(CO ₂ PRESS)	CALCITE LOG(Q/K)	DOLOMITE LOG(Q/K)
20, U12E	.	.	.
21, U12E.04	.	.	.
22, U12E.03	-2.3100	-2.1200	.
23, U12E	.	.	.
24, U12E.02	.	.	.
25, U12E.05	-2.6200	-1.6600	.
26, U12E.03	.	.	.
27, U12E.03	.	.	.
28, U12E.03	-2.5900	-1.6200	.
29, U12E.07	-2.3400	-0.5600	-0.7700
30, U12E	.	.	.
31, U12E	-1.8800	-1.9200	.
32, U12E	.	.	.
33, U12E	.	.	.
34, U12B	-2.7500	-0.6700	-1.3300
35, U12B.03	-2.3100	-1.7400	-3.1800
36, U12.04	.	.	.
SEEP 1, U12T	-2.7900	-1.2860	.
SEEP 2, U12N.03	-1.9220	-2.2950	-4.4750
SEEP 3, U12N.05	-1.7510	-1.6310	-3.8290
SEEP 4, U12N.03	-2.2330	-2.3430	-4.2740
LYSIMETER 1, U12N.05	-1.0400	-0.7740	-1.9180
LYSIMETER 2, U12N.05	-1.7480	-1.9480	-4.1930
LYSIMETER 3, U12N.05	-1.6850	-1.7530	-4.0190
LYSIMETER 4, U12N.05	-1.7450	-1.9490	-4.2550
LYSIMETER 5, U12N.05	-2.4390	-3.2950	-5.6200
LYSIMETER 6, SURF SOIL	.	.	.
LYSIMETER 7, SURF SOIL	-2.2300	-1.4990	-2.4780
LYSIMETER 9, SURF SOIL	-2.1340	-0.5120	-0.5360
ANAL 2589, TUNNEL U12B, E	-2.7500	-0.6700	-1.0400
ANAL 2819, TUNNEL U12B, D	-2.3100	-1.7500	-3.1700
ANAL 2913, TUNNEL U12E, D	-2.6900	-1.7900	.
ANAL 3260, TUNNEL U12E, D	-2.6500	-2.9500	.
ANAL 3541, TUNNEL U12E, M	-2.9400	-1.6400	.

N= 193

SITE	SAS	GYPSUM LOG(Q/K)	MAGNESITE LOG(Q/K)	FLUORITE LOG(Q/K)
WELL J-12		-2. 5070	-2. 2710	-0. 6200
WELL J-13		-2. 6820	-2. 0330	-0. 6080
WELL UE-25B#1		-2. 3720	-2. 3500	-0. 8890
WELL UE-25B#1		-2. 4390	-2. 0950	-0. 8680
WELL UE-25B#1		-2. 4290	-2. 4260	-0. 8480
WELL UE-25C#1		-2. 6000	-2. 0590	-0. 8660
WELL UE-25C#2		-2. 5800	-1. 9430	-0. 8180
WELL UE-25C#3		-2. 6150	-1. 9410	-0. 8990
WELL UE-25P#1		-1. 9880	-1. 1870	-0. 0380
WELL UE-25P#1		-1. 1550	-0. 3580	0. 3940
WELL UE-29A#2		-2. 6350	-3. 2690	-1. 3780
WELL UE-29A#2		-2. 6500	-3. 3830	-1. 4390
WELL USW G-4		-2. 6130	-2. 3660	-0. 5910
WELL USW H-1		-3. 0600	-3. 0980	-1. 6440
WELL USW H-1		-2. 9020	-3. 2400	-1. 6810
WELL USW H-3		-3. 7920	-1. 9420	-1. 2000
WELL USW H-4		-2. 3960	-2. 4650	0. 0380
WELL USW H-5		-3. 4860	-3. 5690	-1. 9180
WELL USW H-5		-3. 4660	-3. 7960	-1. 8880
WELL USW H-6		-2. 9600	-2. 1600	-0. 5950
WELL USW H-6		-3. 4990	-2. 4510	-1. 2670
WELL USW H-6		-2. 8880	-1. 9930	-0. 5570
WELL USW VH-1		-2. 3720	-1. 2210	-0. 6380
WELL USW VH-1		-2. 3970	-1. 6560	-0. 6760
WELL USW VH-1		-2. 4080	-1. 6620	-0. 6780
3. 155/49E-22DC		-2. 1100	-1. 5600	.
4. 165/49E-5ACC		-2. 1700	-1. 1700	.
5. 165/49E-8ABB		-2. 1000	-1. 7800	.
6. 165/49E-8ACC		.	.	.
7. 165/49E-9CDA		.	.	.
8. 165/49E-9DCC		-1. 8900	-1. 1200	.
9. 165/49E-18DC		-2. 2900	-1. 0600	.
10. 165/49E-16CCC		-1. 8800	-1. 5200	.
11. 165/49E-19DAA		-2. 1500	-1. 3400	.
12. 165/48E-24AAA		-2. 3100	-1. 6200	.
13. 165/48E-25AA		-2. 3000	-1. 7000	.
14. 165/48E-36AAA		-2. 4000	-0. 9700	.
15. 175/48E-1AB		-2. 3500	-1. 3400	.
16. 175/49E-7BB		-2. 1900	-1. 0800	.
17. 175/49E-9AA		.	.	.
18. 175/49E-8DDB		-2. 2800	-0. 9400	.
19. 175/49E-15BBD		-2. 1800	-1. 0600	.
64. WELL B, NTS		.	.	.
20. 175/49E-35DDD, ASH TR		.	.	.
21. 165/49E-23ADD		.	.	.
23. 165/48E-15AAA		.	.	.
25. 165/48E-10CBA		.	.	.
27. 165/50E-7BCD		.	.	.
29. 165/49E-15AAA		.	.	.
30. 165/49E-36AAA		.	.	.
45. 165/48E-8BBA		-1. 2200	-1. 2800	-0. 7600
46. 165/48E-7BBA		-1. 3300	-1. 6500	-1. 1600
47. 165/48E-7CBC		.	.	.

14:50 THURSDAY, MAY 29, 1986 34

SITE	SAS	GYPSUM LOG(Q/K)	MAGNESITE LOG(Q/K)	FLUORITE LOG(Q/K)
48, 16S/48E-18BCC		-1.2600	0.4700	.
49, 16S/48E-17CCC		-1.1100	0.8400	.
50, 16S/48E-18DAD		-1.2700	-0.9300	.
51, 16S/48E-8CDA		-1.3200	-1.1100	.
52, 16S/48E-17ABB		-1.2400	-1.2000	.
62, WHITEROCK SPRING		.	.	.
68, BUTTE SPRING		.	.	.
70, OAK SPRING		.	.	.
71, TOPOPAH SPRING		.	.	.
1, 10S/47E-14BAB		-2.3580	-1.9510	-0.4430
2, 10S/47E-27CBA		.	.	.
3, 10S/47E-31AAB		.	.	.
4, 10S/47E-32DDA		-1.7210	-1.1450	-0.3020
5, 10S/47E-33AAB		.	.	.
6, 10S/47E-30DCC		.	.	.
7, 11S/46E-26BBB		.	.	.
8, 10S/46E-26BCC		.	.	.
9, 11S/47E-3CDB		-1.9560	-1.4560	-0.3630
10, 11S/47E-4CAD		-1.7120	-1.1050	0.3050
11, 11S/47E-10CAA		.	.	.
12, 11S/47E-10BCC		.	.	.
13, 11S/47E-16DCD		.	.	.
14, 11S/47E-16BDC		.	.	.
15, 11S/47E-18ACD		.	.	.
16, 11S/47E-21ACC		-1.6950	-1.0300	0.2720
17, 11S/47E-21DBB		.	.	.
18, 11S/47E-21ABA		.	.	.
19, 11S/47E-21ABA		-1.9880	-1.8480	0.0030
20, 11S/47E-27CBA		.	.	.
21, 11S/47E-28AAC		-2.1460	0.2670	-0.0030
22, 11S/47E-28DAC		.	.	.
23, 11S/47E-33BAC		.	.	.
24, 11S/47E-10CCB		.	.	.
25, 12S/47E-5CDA		-1.6800	-1.0510	-1.8310
26, 12S/47E-6CDD		.	.	.
27, 12S/47E-7DBD		.	.	.
28, 12S/47E-20 BBB		-1.5720	-1.1590	0.4500
29, 12S/47E-19ADC		.	.	.
18B, 12S/47E-20		.	.	.
19B, 12S/47E-20		.	.	.
20B, 12S/47E-20		.	.	.
21B, 12S/47E-20		.	.	.
22B, 12S/47E-20		.	.	.
WELL UE-19B-1		-2.3340	-1.7210	-0.0740
WELL UE-19C		-4.9970	-2.1760	-0.1580
WELL UE-19D		-1.7300	-0.6220	0.4780
WELL UE-19E		-3.1800	-2.3930	-0.4460
WELL UE-19E		.	.	.
WELL UE-19GS		-2.9480	-2.0900	-1.1790
WELL UE-19GS		.	.	.
WELL U-20A-2		-2.7700	-3.0090	-0.7560
WELL U-20A-2		.	.	.
WELL UE-20D		-2.7570	-2.4870	-1.0410

SITE	GYPSUM LOG(O/K)	MAGNESITE LOG(O/K)	FLUDRITE LOG(O/K)
WELL UE-20E-1	-4.1090	-2.0200	-1.9030
WELL UE-20H	-3.7350	-2.4590	-1.8280
WELL UE-20J	-1.4180	-2.3350	-0.3290
WELL ARMY-1	-1.8260	0.0640	-0.9490
WELL 5A	-3.2610	-1.5110	-1.3180
WELL 5B	-2.5090	-1.3170	-1.9180
WELL 5C	-3.7210	-1.6330	-2.6520
WELL C	-1.6130	-0.0930	-0.7900
WELL C-1	-1.6340	0.2510	-0.9050
WELL 3	-2.5080	-0.7890	-1.2170
WELL A	-2.5120	-0.7620	-1.5790
WELL 2	-2.2810	-0.3250	-1.8640
WELL UE-15D	-1.8280	0.0710	-0.6110
UE12T#3-5, 134.4(M)	-2.6200	-1.9200	.
UE12T#3-6, 169.6(M)	-2.0900	-1.1300	.
UE12T#3-7, 199.2(M)	-2.1000	-1.1300	.
UE12T#3-8, 202.4(M)	-2.1700	-2.1300	.
UE12T#3-9, 257.6(M)	-2.2200	-2.5200	.
UE12T#3-10, 260.0(M)	-1.8100	-2.1100	.
UE12T#3-11, 441.4(M)	-2.3300	-2.5300	.
UE12T#3-12, 442.4(M)	-2.0900	-2.4000	.
UE12T#3-13, 291.4(M)	-2.7200	-1.9500	.
UE12T#3-14, 320.3(M)	-2.3800	-2.1400	.
UE12T#3-15, 320.6(M)	-2.1100	-2.1000	.
UE12T#3-16, 321.3(M)	-1.9300	-0.9200	.
UE12T#3-17, 350.8(M)	-2.1600	-1.7700	.
UE12T#3-18, 411.2(M)	-2.6900	-2.4600	.
UE12T#3-19, 470.6(M)	-4.3100	-3.3900	.
UE12T#3-20, 472.4(M)	-4.1500	-2.9700	.
UE12T#3-21, 501.7(M)	-3.1600	-2.4700	.
UE12T#3-22, 503.2(M)	-3.2600	-2.4800	.
UE12T#3-23, 532.8(M)	-3.4400	-2.6700	.
RML 1A, SURFACE	-2.7600	-1.7100	.
RML 1B, SURFACE	-2.5200	-1.3300	.
1. U12N.05 BYPASS	-3.3500	-1.9200	.
2. U12N.05	-2.8900	-2.2600	.
3. U12N.05	-2.4300	-1.0300	.
4. U12N MAIN	.	.	.
5. U12N.07 BYPASS	.	.	.
6. U12N.02	-3.1700	-2.9500	.
7. U12T.02 BYPASS	.	.	.
8. U12T.02	.	-3.5700	.
9. U12T.02 BYPASS	.	.	.
10. U12T MAIN	-3.9400	-2.2700	.
11. U12T.03	.	.	.
12. U12T.03	.	.	.
13. U12T.04	.	.	.
14. U12T.03	-2.8200	-2.4000	.
15. U12T.03	-2.6000	-1.6800	.
16. U12T.03	-2.6000	.	.
17. U12E.07	-3.2900	.	.
18. U12E.04	.	.	.
19. U12E	.	.	.

14:50 THURSDAY, MAY 29, 1986 36

SITE	SAS GYPSUM LOG(Q/K)	MAGNESITE LOG(Q/K)	FLUORITE LOG(Q/K)
20, U12E	.	.	.
21, U12E.04	.	.	.
22, U12E.03	-3.7600	.	.
23, U12E	.	.	.
24, U12E.02	.	.	.
25, U12E.05	-3.4700	.	.
26, U12E.03	.	.	.
27, U12E.03	.	.	.
28, U12E.03	-3.6600	.	.
29, U12E.07	-2.6000	-1.8300	.
30, U12E	.	.	.
31, U12E	-3.1900	.	.
32, U12E	.	.	.
33, U12E	.	.	.
34, U12B	-2.7700	-2.2900	.
35, U12B.03	-3.0000	-3.0600	.
36, U12.04	.	.	.
SEEP 1, U12T	-3.6170	.	.
SEEP 2, U12N.03	-3.3700	-4.1360	.
SEEP 3, U12N.05	-2.7030	-3.8790	.
SEEP 4, U12N.03	-3.8900	-3.5870	.
LYSIMETER 1, U12N.05	-3.4720	-2.8410	.
LYSIMETER 2, U12N.05	-4.0940	-3.9300	.
LYSIMETER 3, U12N.05	-3.5530	-3.9540	.
LYSIMETER 4, U12N.05	-3.5200	-3.9890	.
LYSIMETER 5, U12N.05	-5.1980	-3.9920	.
LYSIMETER 6, SURF SOIL	.	.	.
LYSIMETER 7, SURF SOIL	-3.7120	-2.7080	.
LYSIMETER 9, SURF SOIL	-2.5950	-1.7180	.
ANAL 2589, TUNNEL U12B, E	-2.7600	-2.0000	-3.2400
ANAL 2819, TUNNEL U12B, O	-3.0200	-3.0600	-2.7500
ANAL 2913, TUNNEL U12E, D	-3.4900	.	.
ANAL 3260, TUNNEL U12E, D	-4.7800	.	.
ANAL 3541, TUNNEL U12E, M	-3.4800	.	.

N= 193

APPENDIX B

IMPORTANCE OF RADIONUCLIDE TRANSPORT BY PARTICULATES ENTRAINED IN FLOWING GROUNDWATERS

Allen Ogard

INTRODUCTION

The Nevada Nuclear Waste Storage Investigations Project of the US Department of Energy is studying the suitability of Yucca Mountain (Nye County, Southern Nevada) as a potential repository for high-level nuclear waste. The possibility that flowing groundwaters in this area might transport potentially hazardous radionuclides or other harmful elements (both hereafter referred to as waste elements) from a waste repository to the accessible environment requires careful evaluation, and a considerable amount of study is currently being devoted to understanding this potential problem. Leached waste elements could potentially migrate as dissolved species with the groundwater, but it is also conceivable that particulates (perhaps small mineral fragments from tuffs) or natural colloids (that is, iron hydroxide) moving with the groundwaters could strongly sorb various waste species and transport these elements through fractures or open matrix porosity. The intent of this appendix is to qualitatively assess the potential of any particulates that may be entrained in flowing groundwaters to transport important quantities of sorbed waste elements as these groundwaters migrate through the Yucca Mountain environment.

EXPERIMENT AND RESULTS

Well J-13 is the closest well to Yucca Mountain from which groundwater is being routinely pumped, and is located approximately 4 miles to the southeast of the crest of Yucca Mountain, on the east side of Fortymile Wash. As J-13 may intersect groundwater flow paths from the candidate repository site to the accessible environment, water from this well was chosen for particulate content studies. Water was diverted from the well into a mobile laboratory containing filtration equipment at a rate of approximately 1 l/min. A prefilter which removes material larger than 10 μm from the water was positioned upstream from a large stainless steel One-Sevener Nuclepore Membrane Filter Assembly which was normally loaded with seven 0.4 μm membrane filters, mounted in parallel.

Approximately one half of the water discharged from this assembly subsequently passed through an Amicon Hollow Fiber Filter system which removes particulates with diameters greater than ~5 nm.

A filtration run was conducted for 14 days, during which time 9300 l of water were passed through the 0.4 μm membrane filters, and 5300 l through the 5 nm hollow fiber system. The material collected on the membrane filters was removed by ultrasonic treatment in a small quantity of Nanopure water, and the resulting suspension was centrifuged at 10,000 rpm for 20 minutes. The liquid was then decanted and discarded, and the sediment was washed with ethyl alcohol and allowed to dry in air. A total of 0.25 g of solid material was thus obtained which, when divided by the quantity of water that was filtered, corresponds to a sediment concentration of $\sim 2.7 \times 10^{-5}$ g/l. This particulate fraction was dissolved in a mixture of HNO_3 , HCl and HF acids, and the solution was then diluted and analyzed. The particulates collected by the hollow fiber system were removed by backflushing with the minimum amount of Nanopure water and were subsequently analyzed in solution. It was calculated from the concentrations of species in the two solutions that the amount of material in the smaller-size particulate fraction was only about 1% that in the larger-size fraction.

Both solutions were analyzed for cation composition by means of emission spectroscopy. The detectable cations in the $>0.4 \mu\text{m}$ fraction were (in wt%) Si(60), Fe(20), Ca(11) and Al(4), while analysis of the smaller-sized fraction gave somewhat different results: Na(44), Si(42), Ca(8), and Fe(4), with no detectable Al. Because the amount of Fe in the Yucca Mountain tuffs and groundwaters is very low, it is possible that the iron-rich particulates (and perhaps others) could have been contamination from the steel piping and pumping systems. However, as we wish to make a conservative assessment of the importance of particulates in waste element transport, we will assume that all recovered particulates are natural and were originally entrained in the groundwater pumped from J-13.

DISCUSSION AND CONCLUSIONS

To assess the importance of particulates in the transport of waste elements, we need to determine the amount of a given species that is sorbed on particulates, and compare this quantity to the amount which is dissolved in the groundwater. Let

$$R_d = \text{sorption ratio} = \frac{\text{moles of particulate-sorbed species/g of particulate}}{\text{moles dissolved species/ml of groundwater}}$$

[units of ml/g]

$$x = \text{concentration of species-sorbing particulates in groundwater}$$

[units of g/l]

$$c = \text{concentration of species dissolved in groundwater}$$

[units of moles/l]

For the purposes of this simple analysis we will assume the system attains steady state, and that particulate surface area and ground water chemistry effects, etc. can be neglected. From the above then,

$$y = \text{moles of particulate-sorbed species/g of particulate} = \frac{R_d \cdot c}{1000}$$

and

$$z = \text{moles of particulate-sorbed species/liter groundwater} = y \cdot x.$$

Therefore, to determine the relative distribution of waste elements between sorbed and dissolved species, we need only examine the ratio z/c , which from above is also equal to $R_d x/1000$.

Figure B-1 is a plot of the sorption ratio, R_d , versus the groundwater particulate concentration, x , and shows a trajectory for the value of $z/c = 0.1$. This value has been arbitrarily chosen, and implies that the quantity of species sorbed on particulates is only 10% of that dissolved in the groundwater. The value of c in groundwater can range anywhere from essentially zero to the solubility limit of the species, and as concentrations are frequently not known to better than an order of magnitude, an additional contribution of 10% to the total waste element concentration because of the presence of particulate-sorbed species, should constitute a negligible source of error. Examination of Fig. B-1, therefore, indicates that for any combination of R_d and x lying below the $z/c = 0.1$ line (that is, particulates sorb less than 10% of the total species), transport of waste elements as sorbed species on particulates entrained in flowing groundwater should be of little consequence as the bulk of the waste element will be present as dissolved species. This is actually a very conservative analysis in that we are assuming that the particulate velocity is essentially equal to the groundwater velocity. In reality, particulate transport is a strong function

of size (among other things) and is probably very significantly retarded relative to the groundwater flux for the bulk of the entrained particulates.

As determined above, the approximate concentration of $>0.4 \mu\text{m}$ particulates in J-13 water was $2.7 \times 10^{-5} \text{ g/l}$. Examination of Fig. B-1 indicates that these particulates would have to exhibit a sorption ratio greater than $\sim 4 \times 10^6 \text{ mol/g}$ for the species of interest in order for particulates to contribute more than 10% to the total waste element flux. The smaller-sized particulate fraction would have to exhibit sorption ratios greater than $\sim 4 \times 10^8 \text{ mol/g}$ to have a similar effect. These sorption ratios are extremely high and have seldom even been approached in sorption experiments using Yucca Mountain tuffs. However, no sorption ratios have ever been directly determined for the particulates, and it may be possible that they would exhibit sorption ratios of this magnitude or even higher; this is considered highly unlikely as the particulates are probably directly derived from Yucca Mountain tuffs and would be expected to yield similar sorption ratios. To assess this possibility, additional quantities of particulates have been collected from J-13, and we will attempt to experimentally determine sorption ratios for these materials in the future.

Considerably more work would be needed to quantitatively establish the particulate-sorbed contribution to total radionuclide transport. Particulate concentrations, sorption ratios, relative transport velocities, etc., would need to be determined at a number of locations along representative groundwater flow paths to arrive at a more realistic assessment of sorbed-radionuclide fluxes at Yucca Mountain. However, based on the above analysis, and considering the conservative assumptions employed throughout, it can probably be safely assumed that the transport of particulate-sorbed radionuclides by groundwater flow will constitute a negligible component of the total waste element flux at Yucca Mountain. It is highly probable that dissolved species or natural colloids will comprise a much more important component of the overall flux.




Fig. B-1. Relative importance of particulate-sorbed vs dissolved waste element transport as a function of sorption ratio and particulate concentration. $z/c = 0.1$ implies that 10% of the total amount of waste element present is sorbed, and 90% is dissolved.