

716
PATENTS-US--A7146622

146,622
January 21, 1988

PATENTS-US--A7146622

DE89 010888

DE-AC05-840R21400

Gerald W. Strandberg
Susan N. Lewis

MICROBIAL SOLUBILIZATION OF COAL

146,622

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MICROBIAL SOLUBILIZATION OF COAL

Inventors: Gerald W. Strandberg
11534 S. Monticello Drive
Farragut, Tennessee 37922
U.S. Citizen

Susan N. Lewis
2939 W. Gallaher Ferry Drive
Knoxville, Tennessee 37932
U.S. Citizen

ab
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

TITLE OF THE INVENTION

MICROBIAL SOLUBILIZATION OF COAL

BACKGROUND OF THE INVENTION

Field of the Invention

5 The present invention relates to a cell-free preparation and process for the microbial solubilization of coal into solubilized coal products. More specifically, the present invention relates to bacterial solubilization of coal into solubilized coal products and a cell-free bacterial byproduct useful for solubilizing coal.

Description of Related Art

15 Thermal and chemical processes for the conversion of coal to liquid and gaseous products generally require somewhat extreme temperature, pressure and chemical conditions. The severity of the operating conditions may commonly include pressures in excess of 3,000 psi and temperatures in excess of 800 degrees Fahrenheit. These processes usually also require a significant capital investment. Because of the relatively mild operating conditions associated with many biological processes, there has been a recurring

20

interest in the potential use of microorganisms for coal processing.

There have been earlier suggestions that microorganisms may be able to solubilize native coal.

5 M.H. Rogoff et al, "The Microbiology of Coal in Bacterial Oxidation of Phenanthrene," J. Bacteriol., Vol. 73, pp. 264-68 (1957); M.H. Rogoff et al, "Microbiology of Coal," U.S. Bureau of Mines, Information Circular 8075 (1962); J.A. Korbinger,

10 "Microbiology of Coal: Growth of Bacteria in Plain and Oxidized Coal Slurries," Proc. W. Va. Acad. Sci., Vol. 36, pp. 26-30 (1964). In recent experiments, certain strains of fungi have been shown to produce a liquid product when cultured on the surface of lignite coal in the presence of humid air. M.S. Cohen et al,

15 "Degradation of Coal by the Fungi Polyporous versicolor and Poria monticola," Appl. Environ. Microbiol., Vol. 24, pp. 23-27 (1982); C.D. Scott, G.W. Strandberg, and S.N. Lewis, "Microbial

20 Solubilization of Coal," Biotechnol. Prog., Vol. 2, p. 131 (1986). The coal solubilization product is highly polar, water-soluble material of moderately high molecular weight having a high degree of aromaticity. Coal solubilization by these fungi appears, at this

25 time, to be limited to conditions in which the coal must be in contact with aerial mycelia.

Spectroscopic evidence has also been found for the use of coal as a substrate for a bacterial strain of Pseudomonas isolated from a mixed enrichment

30 culture growing in the presence of finely ground bituminous coal. Fakoussa et al, "Koehle als Microbielles Substrat unter Aeroben Bedingungen," Kolloquium in der Bergbau-Forschung Gmbh, Essen, pp.

41-49 (1983). Fakoussa et al suggested that both an enzyme and a surface active agent produced by the organism were involved in attacking the coal. Fakoussa et al does not appear to demonstrate that the organism causes both a substantial decrease in the weight of coal and an increase in the carbon content of the culture broth. Fakoussa et al further appears to fail to demonstrate that coal solubilizing activity is produced in the absence of coal.

It is therefore desirable to effect microbial solubilization by using bacteria. It is further desirable to accomplish bacterial solubilization of coal with a submerged bacterial culture that does not require fungal aerial mycelia. It is further desirable to obtain a cell-free substance capable of solubilizing coal since such a substance would significantly enhance potential microbial solubilization techniques.

SUMMARY OF THE INVENTION

Therefore, it is an object of the present invention to provide for the solubilization of coal by the use of bacteria.

It is a further object of the present invention to provide a means to accomplish coal solubilization in a submerged process.

Another object of the present invention is to provide a cell-free substance capable of solubilizing coal.

Still, a further object of the present invention is to provide a bacterial byproduct capable of solubilizing coal.

The foregoing objects and others are accomplished in accordance with the present invention, generally

speaking, by providing a process for the bacterial solubilization of coal which may be carried out in submerged conditions and which includes introducing coal to a medium containing a bacterial culture so as to cause microbial solubilization and so as to produce solubilized coal products which may be removed from the medium.

The present invention is also generally described as encompassing a non-cellular bacterial byproduct capable of solubilizing coal so that solid portions of coal are converted into liquid form.

Further scope of the applicability of the present invention will become apparent from the detailed description given hereinafter. However, it should be understood that the detailed description and specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the present invention will become apparent to those skilled in the art from this detailed description.

DETAILED DESCRIPTION OF THE INVENTION

According to a preferred embodiment of the present invention, microbial solubilization of coal is accomplished by employing the bacterial species Streptomyces setonii 75 Vi2 and Streptomyces viridosporous T7A. These bacteria are known lignin-degrading bacteria. Both organisms were obtained from Prof. Don L. Crawford, Dept. of Bacteriology and Biochemistry, Univ. of Idaho, Moscow, Idaho. S. setonii 75 Vi2 was deposited into the American Type Culture Collection (ATCC) and was designated as ATCC strain 39116. This strain was

tested for solubilizing activity and the test results indicated that its solubilizing characteristics do not differ from those of the organisms obtained directly from D. Crawford.

The microbial solubilization process of the present invention is capable of solubilizing various kinds of coal. Coals that are preferred for use in the present invention include North Dakota lignite, Mississippi lignite, Texas lignite, and Wyoming-Dakota (Wyodak) Subbituminous coal. The origins and properties of these coals are described, for example, in Scott et al, "Microbial Solubilization of Coal," Biotechnol. Prog., Vol. 2, p. 131 (1986). Some of these coals are pretreated or preoxidized with an oxidizing agent so as to render the coal uniformly and more readily susceptible to microbial solubilization. Suitable oxidizing agents include, for example, nitric acid, hydrogen peroxide, and ozone. During the oxidization pretreatment process the coal is exposed to the oxidizing agent by, for example, immersing the coal in an oxidizing agent solution or exposing the coal to an oxidizing agent gas. The coal may require sterilization after pretreatment and before exposure to microbial solubilization. Other suitable oxidation pretreatment conditions may be employed. Preferably the coal is in the form of coal particles in the particle size range of from 10 to 100 mesh and, most preferably, from 10 to 20 mesh.

The medium employed in the process of the present invention must be capable of supporting microbial growth, such as bacterial growth, and maintaining the coal-solubilizing activity of the microorganisms employed in the process. An essential feature of the

medium appears to be the provision of an appropriate organic nitrogen source which the microorganism can modify or metabolize in order to exhibit coal-solubilizing activity. Suitable medium ingredients include, for example, agar, Neopeptone (manufactured by Difco Laboratories, Detroit, MI), Neopeptone plus maltose, Yeast extract (Difco), Amberex-Crude (manufactured by Amber Laboratories, Juneau, WI), cheese whey, and Casamino acids (Difco). Suitable nitrogen sources also include complex nitrogen compounds, such as Peptone, Proteose-peptone, Tryptone, and Tryptose (complex nitrogen nutrients obtained from Difco, Detroit, MI). Further nitrogen sources include amino acids, such as Glutamic acid, Valine, and Asparagine. The medium does not require a carbohydrate. The coal-solubilizing activity of the microorganisms in the medium is usually accompanied by an increase in alkalinity of the medium.

Other conditions associated with the medium of the process of the present invention include temperature, humidity, and the time period during which the microorganisms are cultured. These conditions can be determined by one of ordinary skill in the art with reference to the examples indicated below.

The present invention also encompasses an extracellular bacterial preparation that is capable of solubilizing coal. In a preferred embodiment, this extracellular component is isolated from a bacterial culture of Streptomyces setonii 75 Vi2 which produces this component. The presence of this soluble, cell-free component capable of solubilizing coal was conclusively demonstrated by using a sterilizing

filter (0.22 μ or 0.45 μ , Millipore filter) to remove all cells from a seven day old S. setonii 75 Vi2 culture broth and contacting coal with the sterilized broth which resulted in a solubilizing activity on the coal. The active cell-free component in the broth is likely not a protein, enzyme, or some combination since solubilizing activity cannot be eliminated by treating the sterilized broth with any of several proteases. The cell-free component has an unusually high heat stability since autoclaving of the sterilized broth for up to one hour at 121 degrees Celcius did not completely destroy solubilizing activity.

The cell-free component capable of solubilizing coal may be isolated by various methods. For example, a sterilizing filter, such as a 0.22 μ or 0.45 μ Millipore filter, may be used in a filtering method in order to isolate the component. Gel permeation chromatography may possibly also be used to isolate the cell-free component. With this method, sodium dodecyl sulfate (SDS) may be used to enhance separation and recovery of the cell-free component.

EXAMPLES

In order to further define the specifics of the present invention, the following examples are provided and are intended to illustrate the concepts of the coal-solubilizing processes and preparations associated with the present invention and not limit the particulars of the present invention.

Example 1

Coal is microbially solubilized on the surface of a bacterial culture. The bacterial species Streptomyces setonii 75 Vi2 and S. viridosporous T7A

are cultured on the surface of Sabouraud Maltose agar (Neopeptone by Difco, 10 g/L; Maltose, 40 g/L; agar, 15 g/L) at 30° C and greater than 80% relative humidity for 7 to 10 days. Coal solubilization of low-ranked coals such as North Dakota lignite and preoxidized Wyodak subbituminous coal is evidenced by the production of a black liquid commencing within a few hours after adding the coals to the surface of the cultures. Up to 55% solubilization of the Wyodak coal was observed after 14 days contact with S. setonii 75 Vi2.

Example 2

In order to test the solubilization of coal by a submerged bacterial culture, coal was added to shake flask cultures of S. setonii 75 Vi2. The culture was grown in Sabouraud Maltose broth at 30° C and the flask was shaken at 100 RPM with a two inch stroke. The coals used were presized (1-2 mm) particles of either a North Dakota lignite (American Colloid Co., Skokie, IL) or a subbituminous coal (Wyodak, Amax Coal Co., Indianapolis, IN). The coals were previously sterilized by autoclaving at 121° C for 40 minutes. The Wyodak subbituminous coal had been previously treated by soaking in 8 N HNO₃ for 48 hours in order to render the coal more susceptible to solubilization. After two days incubation with the cultures, the coal and microorganisms were removed from the culture broth by centrifugation. The residual coal was separated from the microorganisms by repeated, gentle agitation in deionized, distilled water followed by decanting the liquid with the lighter microorganism cells suspended in it. The dry weight of the remaining coal residue was determined after oven drying at 95 to

100° C for 16-18 hours. The results of these experiments are presented in Table 1. It is noted that blackening of the broth began within 3-4 hours after adding the coals to the seven day shake cultures. No apparent solubilization occurred in shake flasks of S. viridosporous T7A.

Table 1

The solubilization of coal by a submerged culture of Streptomyces setonii 75 Vi2

Conditions	Coal ^a	Dry weight of coal (mg)		Weight change(%)
		initial	final ^b	
Sterile medium	NDV	542	517	-4.6
	Wyodak-PT	509	518	+2.0
Submerged cultures ^c	NDV	508	289	-43.0
		511	371	-27.5
	Wyodak-PT	513	98	-81.0

^aNDV is a North Dakota lignite obtained from American Colloid Co., Skokie, IL. Wyodak-Pt is a subbituminous coal obtained from Amax Coal Co., Indianapolis, IN. It was pretreated by soaking for 48 h in 8 M HNO₃.

^bFinal dry weights of residual coal after 48 h incubation with cultures.

^cCoal was added to 7 day shake cultures (Sabouraud Maltose medium, 30° C, 100 rpm, 2 inch stroke).

Example 3

In order to demonstrate the coal-solubilizing activity of a cell-free component of the present invention, coal solubilization was carried out after isolating the cell-free extract from the bacterium Streptomyces setonii 75 Vi2. The results of coal solubilization experiments are indicated in Table 2.

Table 2
Results of Treatment of Coal with an Extracellular
Product from Streptomyces setonii 75 Vi2

	Coal		<u>Dry Weight of Coal (mg)</u>		% Weight loss
			<u>Initial</u>	<u>Final</u>	
5	North Dakota Lignite	Extract ² + coal	550	423	23.0
		Autoclaved filtered extract + coal	512	444	12.3
10		Control - fresh sterile medium	534	497	6.9
15	Wyodak Subbit- uminous ³	Extract + coal	513	162	68.5
		Autoclaved filtered extract + coal	506	256	49.3
20		Control - fresh sterile medium	509	518	(2.0) gain

25 ¹Dry weight of residual coal following two days incubation with filtered broths (shaken at 30° C, 100 RPM, 2 inch stroke). Residual coal was washed several times with distilled water prior to oven drying (95-100° C, 16-18 h). The coals were sterilized by autoclaving (40 min, 121° C) prior to addition to broths.

30 ²Extracts from 7 day cultures of S. setonii 75 Vi2 grown on Sabouraud Maltose broth, 30° C, 100 RPM.

35 ³Wyodak subbituminous coal was preoxidized by soaking in 8 N HNO₃, 48 h followed by extensive washing in deionized, distilled H₂O.

Example 4

Various media were tested for their ability to support growth and the production of coal-solubilizing activity by a bacterial culture. In this case, Streptomyces setonii 75 Vi2 was used. The results of the various media tested are indicated in Table 3. It is noted from the results that a carbohydrate is not required in the medium and can be detrimental. For example, Neopeptone plus maltose or Neopeptone alone supported the production of coal-solubilizing activity, whereas no activity is found when glucose was substituted for maltose.

5

10

Table 3
Results of Employing Various Media with a Bacterial
Culture of Streptomyces setonii 75 Vi2

Medium ¹	Medium pH		Coal solubilization- % loss
	Initial	Final (7 d)	
Neopeptone ³ (1%) plus maltose (4%)	5.8	8.9	43,50; 69,64; 65,62
Neopeptone ³ (1%)	7.4	8.9 8.5	39.29 38,40
Neopeptone ³ (1%) plus glucose (1%)	7.4	7.9	0
Yeast extract ³ (1%)	7.0	8.8	62,60
Amberex- EHC ⁴ (1%)	5.8	6.0	0
Amberex- crude ⁴ (2%)	5.1	8.8	43,47
Whey ⁵ (2%)	5.9	8.2	32,36
Casamino acids ³ (3%)	5.9	9.0	68,72

¹Medium components were dissolved at the concentrations indicated in demineralized, distilled water and sterilized at 121° C, 20 min.

²Coal solubilization was measured using cell-free broths of S. setonii 75 Vi2 grown on the medium for 7 days (filter sterilized with 0.45 μ filter). A known quantity of nitric acid-treated Wyodak subbituminous coal was shaken in the cell-free broths for 2 days (30° C). The residual non-solubilized coal was recovered by filtration and the dry weight determined.

³Difco Laboratories, Detroit MI.

⁴Amber Laboratories, Juneau, WI.

⁵Whole powdered cheese whey (partially demineralized), Flav-O-Rich, Inc. Knoxville, TN.

Example 5

In order to further test media which support bacterial culture growth and coal-solubilizing activity, various complex nitrogen sources and amino acids were introduced to a bacterial culture of Streptomyces setonii 75 Vi2. The results of these further media tests are summarized in Table 4.

Table 4
The Suitability of Various Nitrogen Sources for Growth
and Coal Solubilization by Streptomyces setonii 75 Vi2

5	Nitrogen Source ¹	(mg dry wt) ² Coal		Solubilized Coal ³ (%)
		Initial	Final	
Complex N Compounds				
10	Peptone ⁴	341	173	49
		306	168	45
	Peptose-peptone ⁴	312	129	59
		312	136	56
	Tryptone ⁴	312	150	52
		320	164	49
	Tryptose ⁴	315	139	56
		313	125	60
	Amino Acids			
	Glutamic acid ⁵	329	71	78
(after cell growth)	346	80	77	
15	Glutamic acid	181	177	2
	(sterile control)	177	174	2
	Valine ⁵	333	306	8
	(after cell growth)	356	322	10
20	Valine	179	179	0
	(sterile control)	204	202	1
	Asparagine ⁵	314	158	50
	(after cell growth)	321	149	54
	Asparagine	193	190	2
	(sterile control)	193	189	2

¹Streptomyces setonii 75 Vi2 was cultured on the nitrogen sources for 7 days at 30° C. After 7 day growth the cells were removed by filtration (0.45 μ Millipore filter), and the sterile culture broths tested for coal solubilizing activity.

²The coal used was a nitric acid treated (2 days, 8N HNO₃).

³Percent coal solubilized after 2 days at 30° C in 50 ml broth.

⁴The complex nitrogen sources were the only nutrients supplied (3% w/v in 100 ml water). All were obtained from Difco, Detroit MI.

⁵The amino acids (0.5% w/v) were supplemented with 0.05% KCl, 0.05% MgSO₄·7H₂O, 0.01% FeSO₄·7 H₂O, and K₂HPO₄ as inorganic nutrients. The volume of culture broth used was 100 ml.

Example 6

The effect of SDS on coal-solubilizing activity of the cell-free culture broth was determined. It was found that SDS enhanced coal solubilization by approximately 20%. The results are shown in Table 5.

Table 5

Results of Gel Permeation Chromatography
Using SDS

<u>Addition</u>	<u>Coal Solubilization % loss</u>
Cell-free culture broth	50, 47
Cell-free culture broth plus 0.1% SDS	59, 63
SDS only	6

The invention being thus described, it will be obvious that the same be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention and all such modifications as would be obvious to one skilled in the art are intended to be included within the scope of the following claims.