

ORGANOMETALLIC CHEMISTRY OF BIMETALLIC COMPOUNDS

DOE/ER/13406--5

DE90 017601

Department of Energy
Division of Basic Energy Sciences

Grant Number DE-FG-02-85ER13406

Progress Report

August 1990

Progress Report Number DOE/ER/13406-5

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Charles P. Casey, Principal Investigator
University of Wisconsin - Madison
Madison, Wisconsin 53706

Received by OSTI
SEP 21 1990

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

cb


DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.


DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.


A. Heterobimetallic Dihydrides. A full paper describing hydrogen transfer from $\text{Cp}(\text{CO})_2\text{ReH}_2$ to *trans*- $\text{Ir}(\text{CO})(\text{Cl})(\text{PPh}_3)_2$ via a heterobimetallic Re-Ir intermediate was published.¹




A communication describing the reaction of the rhenium-platinum dihydride complex $\text{Cp}(\text{CO})_2\text{Re}(\mu\text{-H})\text{PtH}(\text{PPh}_3)_2$ with alkynes was published.² The mechanism of this rare example of heterobimetallic catalysis is under active investigation.




In the course of probing whether rhenium-alkyne complexes are intermediates in this reaction, we discovered the interesting RePt alkyne complex shown below.³




The reaction of Cp(CO)₂ReH⁻ with cationic vinylplatinum(II) complexes was investigated as a possible route to a proposed RePt vinyl intermediate in the alkyne hydrogenation reaction. In agreement with the proposed mechanism, this reaction produced rhenium alkene complexes.³



In an attempt to isolate a compound with a Re-Pt bond, the reaction of $\text{Cp}(\text{CO})_2\text{ReCH}_3^-$ with vinylplatinum(II) compounds was studied. An interesting $\mu\text{-CH}_2$ RePt compound was produced; the reaction is proposed to proceed as shown below with key steps involving Re-Pt bond formation and insertion of Pt into a C-H bond of the ReCH_3 group.³




Kinetic investigations of the reaction of $\text{Cp}(\text{CO})_2\text{Re}(\mu\text{-H})\text{Pt}(\text{H})(\text{PPh}_3)_2$ with propyne have been initiated. We have found that increasing alkyne concentrations results in slower rates of reaction. To explain this, we propose that an equilibrium between RePtH_2 and $\text{Cp}(\text{CO})_2\text{ReH}_2$ and $\text{Pt}(\text{PPh}_3)_2(\text{alkyne})$ is set up and that the slow step in the reaction is a unimolecular rearrangement of RePtH_2 to a reactive intermediate of unknown structure.³





B. Early-Late Transition Metal Heterobimetallic Compounds.

We published a communication describing the syntheses of the Zr-Re μ -alkylidene complexes shown below by reaction of rhenium carbene complexes with Cp_2ZrH_2 .⁴



We also published the synthesis of TaRh and ReRh hydride complexes.⁵




An article describing our work with ZrRu heterobimetallic compounds will appear in Volume 400 of the Journal of Organometallic Chemistry.⁶


C. Diphosphine Chelates with Natural Bite Angles near 120°.

We have defined the "natural bite angle" of a chelating diphosphine as the preferred chelation angle determined only by ligand backbone constraints and not by metal valence angles. We have used molecular mechanics to calculate the natural bite angle of chelates and to estimate their flexibility.⁷ We are particularly interested in ligands with wide natural bite angles near 120°. Such ligands should preferentially occupy equatorial sites in trigonal bipyramids.


Three of the ligands we have studied in detail are BISBI, T-BDCP, and the norbornyl diphosphine shown below which we recently designed and synthesized.⁸




BISBI



T-BDCP



We have compared our calculated natural bite angles with observed bite angles for $(\text{CO})_3\text{Fe}(\text{diphosphine})$ complexes.⁹ The observed bite angles are systematically larger than the calculated natural bite angles but the calculations indicate that the ligands are relatively flexible.



We have employed a series of chelating diphosphines with wide bite angles as ligands in rhodium catalyzed hydroformylation.¹⁰ a rough correlation between the regioselectivity of linear aldehyde formation and the size of the natural bite angle of chelating diphosphines was found.

References

- (1) Synthesis of Rhenium-Alkene Complexes from the Reaction of the Heterobimetallic Dihydride $\text{C}_5\text{H}_5(\text{CO})_2\text{HRePtH}(\text{PPh}_3)_2$ with Alkynes, Casey, C. P.; Rutter, Jr., E. W. *J. Am. Chem. Soc.* **1989**, *111*, 8917-8919.

(2) Hydrogen Transfer from $\text{CpRe}(\text{CO})_2\text{H}_2$ to *trans*- $\text{Ir}(\text{CO})(\text{Cl})(\text{PPh}_3)_2$ via a Heterobimetallic Re-Ir Intermediate, Casey, C. P.; Rutter, Jr., E. W. *Inorg. Chem.* **1990**, *29*, 2333-2335.

(3) New Reactions of Organorhenium Compounds, Casey, C.P. in "Organic Synthesis Via Organometallics" Ed. Hoffmann, R. W.; Dötz, K. H., Vieweg & Verlag, in press.

(4) A New Route to Early-Late Heterobimetallic μ -Alkylidene Complexes, Casey, C. P.; Askham, F. R.; Petrovich, L. M. *J. Organomet.Chem.* **1990**, *387*, C31-C35.

(5) Synthesis of Rhodium Containing Heterobimetallic Hydride Complexes, Casey, C. P.; Whiteker, G. T. *Inorg. Chem.* **1990**, *29*, 876-879.

(6) Synthesis and Reactions of Directly Bonded Zirconium-Ruthenium Heterbimetallic Complexes, Casey, C. P. *J. Organometal. Chem.*, in press.

(7) The Natural Bite Angle of Chelating Diphosphines, Casey, C. P.; Whiteker, G. T. *Israel J. Chem.*, in press.

(8) Synthesis of endo, endo-2,5-Bis(diphenylphosphinomethyl) bicyclo[2.2.1]heptane, a Chelating Diphosphine with a Natural Bite Angle of 120° , Casey, C. P.; Whiteker, G. T. *J. Org. Chem.* **1990**, *55*, 1394-1396.

(9) Pentacoordinate $\text{Fe}(\text{CO})_3$ Complexes of Diphosphine Ligands with Bite Angles Greater than 120° , Casey, C. P.; Whiteker, G. T.; Campana, C. F.; Powell, D. R. *Inorg. Chem.* in press.

(10) Diphosphines with Natural Bite Angles Greater than 120° Increase Selectivity for α -Aldehyde Formation in Rhodium-Catalyzed Hydroformylation, Casey, C. P.; Whiteker, G. T.; Powell, D. R., *J. Am. Chem. Soc.*, submitted.