



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

EARTH SCIENCES DIVISION

DOCUMENTATION FOR COMPUTER CODE NACL

O. Weres, J.C. Peiper, K.S. Pitzer,
and R. Pabalan

February 1987



Prepared for the U.S. Department of Energy under Contract DE-AC03-76SF00098

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

LBL--21859

DE87 007906

Documentation for Computer Code NACL

O. Weres

Earth Sciences Division, Lawrence Berkeley Laboratory, 50E
University of California, Berkeley, California 94720

and

J.C. Peiper, K.S. Pitzer, and R. Pabalan

Department of Chemistry and Lawrence Berkeley Laboratory
University of California, Berkeley, California 94720

February 1987

MASTER

Introduction

The computer program NACL incorporates the empirical model of the thermodynamic properties of the system NaCl - H₂O recently published by Pitzer *et al.* (1). NACL is derived from the research codes used by Pitzer *et al.* to analyze the experimental data and fix the parameters in their model. NACL calculates values for all thermodynamic properties which are identical to values tabulated in Ref. 1.

NACL is written in VAX/VMS FORTRAN, and was developed on a VAX 8600 computer. Machine specific features have been avoided, and NACL should require few changes to compile and run with other compilers and computers.

A sample output and full code listing of NACL are appended to this document. For one year following the publication of this document, the code will be made available to interested users on 5.25" floppy diskette in MS-DOS 2.11 format. Please send a formatted diskette and a stamped, self-addressed mailer to Oleh Weres, Lawrence Berkeley Laboratory, 50E, Berkeley, CA 94720. Please put your name and address on the diskette.

Input and output

NACL accepts interactive input, and produces an output file (NACL.OUT) in print format. After invoking NACL, the user is asked to provide values of temperature, pressure, and NaCl molality. If the pressure specified is less than the vapor pressure of pure water at the temperature specified, the pressure is reset to equal the saturation pressure. NACL will then calculate many thermodynamic properties of the solution.

Two tabular formats are available for the output, and the user is interactively asked to select one or both of these. The first format ("Table 1") presents a listing of all quantities that have been calculated, properly identified and logically grouped. The symbols used in Table 1 are the variable names in the code. The second format ("Table 2"), presents just the quantities that are tabulated in Ref. 1. The values in Table 2 are explicitly related to the tables in Ref. 1. Table 2 may be extended by removing statement 105 in the subroutine TABLE2. With this statement removed, Table 2 will be extended to include "Class 2" and "Class 3" quantities, which are not tabulated in Ref. 1, and cannot be validated by comparison with Ref. 1. "Class 2" quantities have been used in calculating quantities tabulated in Ref. 1, and have thereby been indirectly validated. "Class 3" quantities have not been validated by either means.

The calculation may be repeated, with output for additional values of t , P , c appended to the same output file.

Program structure

NACL is clearly structured and well commented. The code is provided as three files. The first file (NACL.FOR) contains the main program NACL which manages interactive input, and subroutines TABLE1 and TABLE2 which generate the corresponding output. The subroutines which actually calculate the thermodynamic properties are called from the main program. If so desired, the first file may be discarded, and the remainder of the program supplied with another main program or integrated into a larger program. The second file (NACLB.FOR) contains the heart of the model. The third file (NACLC.FOR) calculates the required properties of pure water based on the empirical formulas of Haar *et al.* (2). The dielectric constant of pure water and its derivatives are calculate in subroutine DIELEC using the empirical formula of Bradley and Pitzer (3).

Acknowledgements

This work was supported by the Office of Basic Energy Sciences of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.

Final validation and correction of NACL against the Tables in Ref. 1 were done by Mr. C.H. Jun.

References

1. K.S. Pitzer, J.C. Peiper, and R.H. Busey, Thermodynamic properties of aqueous sodium chloride solutions. J. Phys. Chem. Ref. Data, **13**, 1-102 (1984).
2. L. Haar, J.S. Gallagher, and G.S. Kell, Proc. 9th international conference on the properties of steam, J. Straub and K. Scheffler, eds., p. 69, Pergamon Press (Oxford, 1980).
3. D.J. Bradley and K.S. Pitzer, J. Phys. Chem. **83**, 1599 (1979).

List of Symbols

The following is a list of constants and variables used in the NaCl program. With few exceptions, constants and variables assigned to a common block in the program are grouped together in this list. The common block names are listed alphabetically.

The Gibbs energy, enthalpy, entropy, and heat capacity are dimensionless (divided by R or RT), whereas the volume and its derivatives with respect to temperature or pressure have dimensions of cc/mol, cc/mol/K, cc/mol/bar, except where noted otherwise.

Common Block/ cval2/

tr	reference temperature, 298.15 K
pr	reference pressure, 177 bars
mr	reference molality, 5.550825 moles NaCl/kg. water
mw	molar mass of water, 18.01534 g.
n1	moles of solvent

Common Block / debye/

ap,ah,as,	Debye-Hückel limiting law parameters for the
aj,av,aa,	osmotic coefficient, enthalpy, entropy, heat
ab	capacity, volume, expansivity, and compressibility, respectively.

Common Block/ delta/

v	specific volume of water, cc/g
vt	= $(\partial v / \partial T)_P$
vp	= $(\partial v / \partial P)_T$
vtt	= $(\partial^2 v / \partial T^2)_P$
vpp	= $(\partial^2 v / \partial P^2)_T$
vtp	= $(\partial^2 v / \partial T \partial P)$

Common Block/ dielec/

d	dielectric constant of water
dt	= $(\partial D / \partial T)_P$
dt	= $(\partial^2 D / \partial T^2)_P$
dp	= $(\partial D / \partial P)_T$
dpp	= $(\partial^2 D / \partial P^2)_T$
dtp	= $(\partial^2 D / \partial T \partial P)$

Common Block / excess/

gex	excess Gibbs energy of the solution
g1ex, g2ex	partial molal excess Gibbs energy of solvent and solute
hex	excess enthalpy of the solution
h1ex, h2ex	partial molal excess enthalpy of solvent and solute
sex	excess entropy of the solution
s1ex, s2ex	partial molal excess entropy of solvent and solute
cpex	excess heat capacity of the solution
cp1ex, cp2ex	partial molal excess heat capacity of solvent and solute
vex	excess volume of the solution
v1ex, v2ex	partial molal excess volume of solvent and solute
vtex, vt1ex, vt2ex	temperature derivatives of vex, v1ex and v2ex, respectively
vpex, vp1ex, vp2ex	pressure derivatives of vex, v1ex and v2ex, respectively

**Common Block/ haar/ – standard state properties
of pure liquid water at any T and P**

f10	Helmholtz free energy
g10	Gibbs free energy
u10	internal energy
h10	enthalpy
s10	entropy
cv10	heat capacity at constant volume
cp10	heat capacity at constant pressure
cs10	heat capacity along saturation curve
v10	specific volume, cc/g
a10	expansivity, per Kelvin
b10	compressibility, per (10^5 pascals)

**Common Block/ ideal/ – thermodynamic properties of water
as an ideal gas at T and P and referenced to 0 K**

f0	Helmholtz free energy
g0	Gibbs free energy
u0	internal energy
h0	enthalpy
s0	entropy
cv0	heat capacity at constant volume
cp0	heat capacity at constant pressure

Common Block/ partial/ – partial molal properties of water and NaCl(aq)

g1,g2	partial molal Gibbs energy of solvent and solute
s1,s2	partial molal entropy of solvent and solute
h1,h2	partial molal enthalpy of solvent and solute
cp1,cp2	partial molal heat capacity of solvent and solute
v1,v2	partial molal volume of solvent and solute
vt1,vt2	temperature derivatives of v1 and v2, respectively
vp1,vp2	pressure derivatives of v1 and v2, respectively

Common block/ pg/

phi	osmotic coefficient of NaCl(aq)
gam	activity coefficient of NaCl(aq)

Common Block / ref/ – thermodynamic properties of NaCl(aq) in its standard state of the hypothetical one molal ideal solution at any T and P.

g20	standard state Gibbs energy of NaCl(aq), $(\bar{G}_{2(T,P)}^{\circ} - \bar{H}_{2(298,1 \text{ atm})})/RT$
s20	standard state entropy of NaCl(aq)
h20	standard state enthalpy of NaCl(aq)
cp20	standard state heat capacity of NaCl(aq)
v20	standard state molal volume of NaCl(aq)
vt20,vp20	temperature and pressure derivatives of v20, respectively

Common Block/ solid/ – standard state properties of pure solid NaCl

g2s	Gibbs energy of solid NaCl, $(G_{T,P}^{\circ} - H_{298,1 \text{ atm}})/RT$
h2s	enthalpy of solid NaCl
s2s	entropy of solid NaCl
cp2s	heat capacity of solid NaCl
v2s	molal volume of solid NaCl
vt2s,vp2s	temperature and pressure derivatives of v2s, respectively.

Common Block/ soln/

dgs0	standard Gibbs energy of solution, ΔG_s°
dss0	standard entropy of solution, ΔS_s°
dhs0	standard enthalpy of solution, ΔH_s°
dcps0	standard heat capacity of solution, $\Delta C_{p,s}^\circ$
dvs0	standard volume of solution, ΔV_s°
dvts0,dvps0	temperature and pressure derivatives of dvs0, respectively

Common Block / virial/ – virial coefficients for the Pitzer equations

b0,b1,cp	b0,b1 are pair-wise ion-interaction parameters and cp is the triplet ion-interaction parameter for Pitzer's equations for Gibbs energy. Symbols with letters L,s,j,v,a and b refer to parameters for enthalpy, entropy, heat capacity, volume, expansivity, and compressibility, respectively.
b0L,b1L,cpL	
b0s,b1s,cps	
b0j,b1j,cpj	
b0v,b1v,cpv	
b0a,b1a,cpa	
b0b,b1b,cpb	

Others

i	ionic strength
m	molality
p	pressure, bars
ps	saturation pressure at temperature T
r	gas constant = 8.31440 J/mol/K = 83.1440 cc-bar/mol/K = 4.61518 cc-bar/g/K
si	square root of ionic strength
t	temperature, Kelvin

PROGRAM LISTING

program nacl

This program calculates the properties of NaCl solutions using the accurate empirical model described by Pitzer, Peiper, and Busey, J. Phys. Chem. Ref. Data, V.13, 1-102 (1984) and in Lawrence Berkeley Laboratory Report LBL-15512.

The tables in the cited references were generated using an earlier version of this program, which was written by J.C. Peiper.

The present version of NACL was developed by O. Weres with assistance from C.H. Jun.

Version 2.00.

- O. Weres, March 1986

The main program accepts interactive input, calls the subroutines that perform the actual calculations, and calls the subroutines that produce printed output.

```
implicit real*8(a-h,o-z)
character*1 ans
common /beta/ ps,dpdt,er,b,bt,btt,bb,bbt,bbtt,y,yr,yt,ytt
open(unit=6,file='nacl.out',status='unknown')
```

Read tc,p,c interactively

```
write(*,101)
101 format('Welcome to program NACL, v. 2.00. '/
1 ' Interactive input. Output is written to file NACL.OUT'/)
27 continue
write(*,114)
114 format(' Enter T(C).')
accept 102,tc
t = tc + 273.15
if(tc.lt.0d0.or.tc.gt.300d0) write (*,112)
112 format(' Caution. T(C) < 0. or T(C) > 300. ')
call psat(t)
102 format(f10.1)
write(*,103)
103 format(' Enter P(bar)')
accept 102, p
if(p.gt.1000d0) write (*,108)
108 format(' Caution. P > 1000 bar')
if(p.lt.ps) then
write(*,109) ps
109 format(' Caution. P < Psat(T). P is set = Psat(T) =',f10.3)
p = ps
endif
write(*,104)
104 format(' Enter C(molal)')
accept 102,c
if(c.lt.1d-20) then
write(*,110)
110 format(' Caution. C must be positive. C is reset to 1e-20')
```

```

c = 1d-20
endif
if(c.gt.6d0) write (*,111)
111 format(' Caution.  C > 6 molal')
c
write(6,30)
call haar(t,p)
call nacl1(t,p)
call nacl2(t,p,c)
call nacl3(t,p,c)
write (*,107)
107 format(' Do you want Table 1 (y or n)?')
accept 106, ans
if(ans.eq.'y'.or.ans.eq.'Y') call table1(t,p,c)
write (*,113)
113 format(' Do you want Table 2 (y or n)?')
accept 106, ans
if(ans.eq.'y'.or.ans.eq.'Y') call table2(t,p,c)
write(*,105)
105 format(' Do you want to go again (y or n) ?')
accept 106,ans
106 format(a1)
if(ans.eq.'y' .or. ans.eq.'Y') go to 27
30 format('1NACL calculates the properties of NaCl solutions using'
1/' the accurate empirical model published by Pitzer, Peiper, and'
2/' Busey, J. Phys. Chem. Ref. Data v. 13, pp. 1-102 (1984).'
3/' The valid range is 0-300 C, 0-6 molal, Psat(T)-1000 bar.')
stop
end
c
c
c
c
subroutine table1(t,p,c)
c
implicit real*8(a-h,o-z)
real*8 i,m,mr,mw,n1
common /beta/ ps,dpdt,er,b,bt,btt,bb,bbt,bbtt,y,yr,yt,ytt
common /debye/ ap,ah,as,aj,av,aa,ab
common/delta/v,vt,vp,vtt,vpp,vtp
common /diel/ d,dt,dt,dp,dpp,dt
common /excess/ gex,glex,g2ex,hex,hlex,h2ex,sex,slex,
1 s2ex,cpex,cplex,cp2ex,vex,vlex,v2ex,vtex,vtlex,vt2ex,
2 vpex,vplex,vp2ex
common /haar/ f10,g10,u10,h10,s10,cv10,cp10,cs10,v10,a10,b10
common /ideal/ f0,g0,u0,h0,s0,cv0,cp0
common /partial/ g1,g2,s1,s2,h1,h2,cp1,cp2,v1,v2,vt1,vt2,
1 vp1,vp2
common /pg/ phi,gam
common /ref/ g20,s20,h20,cp20,v20,vt20,vp20
common /solid/ g2s,h2s,s2s,cp2s,v2s,vt2s,vp2s
common /soln/ dgs0,dss0,dhs0,dcps0,dvs0,dvts0,dvps0
common /vir/ b0,b0l,b0s,b0j,b0v,b0a,b0b,b1,b1l,b1s,b1j,
1 b1v,b1a,b1b,cp,cpl,cps,cpj,cpv,cpa,cpb
tc = t - 273.15
write(6,2) t,tc,p,c
write(6,28)
write(6,29)
write(6,21) ps,dpdt
write(6,7) f10,g10,u10,h10,s10,cv10,cp10,cs10,v10,a10,b10
write(6,8) f0,g0,u0,h0,s0,cv0,cp0

```

```
write(6,5) v,vt,vp,vtt,vpp,vtp
write(6,6) d,dt,dt,dp,dpp,ntp
write(6,4) ap,ah,as,aj,av,aa,ab
write(6,31)
write(6,9) b0,b0l,b0s,b0j,b0v,b0a,b0b
write(6,10) b1,b1l,b1s,b1j,b1v,b1a,b1b
write(6,11) cp,cpl,cps,cpj,cpv,cpa,cpb
write(6,12) g20,s20,h20,cp20,v20,vt20,vp20
write(6,13) gex,gl,ex,g2ex,hex,hlex,h2ex
write(6,14) sex,slex,s2ex,cpex,cplex,cp2ex
write(6,15) vex,vlex,v2ex,vtex,vtlex,vt2ex
write(6,26) vpex,vplex,vp2ex
write(6,16) g1,g2,s1,s2,h1,h2,cp1,cp2
write(6,17) v1,v2,vt1,vt2,vp1,vp2
write(6,18) phi,gam
write(6,19) g2s,h2s,s2s,cp2s,v2s,vt2s,vp2s
write(6,20) dgs0,dss0,dhs0,dcps0,dvs0,dvts0,dvps0
write(6,32)
32 format('1')
return
2 format('0Temperature, pressure and molality are'//
1 ' t,tc = ',f10.2,' p = ',f10.3,' m = ',f7.3/)
28 format('0Table 1 contains all calculated quantities.'// See comme
ints in HAAR, RH0, NACL1 and NACL3 for further explanation.')
29 format('0Properties of pure water are calculated first.')
21 format('0Saturation pressure of pure water and temperature derivat
live from PSAT'// ps,dpdt = ',1p2e13.4/)
7 format('0Reference state properties for liquid water from HAAR'//
1 ' f10,g10 = ',1p2e13.4,' u10,h10,s10 = ',1p3e13.4//
2 ' cv10,cp10,cs10 = ',1p3e13.4,' v10,a10,b10 = ',1p3e13.4/)
8 format('0Properties of water as ideal gas from HAAR'//
1 ' f0,g0 = ',1p2e13.4,' u0,h0,s0 = ',1p3e13.4,
2 ' cv0,cp0 = ',1p2e13.4/)
5 format('0Specific volume of pure water and its derivatives over
1 T and P from RH0'//
2 ' v,vt,vp = ',1p3e13.4,' vtt,vpp,vtp = ',1p3e13.4/)
6 format('0Dielectric constant and its derivatives from DIELEC'//
1 ' d,dt,dt = ',1p3e13.4,' dp,dpp,ntp = ',1p3e13.4/)
4 format('0Debye-Huckel coefficients from DEBYE'//
1 ' ap,ah,as = ',1p3e13.3,' aj,av = ',1p2e13.4,
2 ' aa,ab = ',1p2e13.4/)
31 format('0Virial coefficients from NACL1')
9 format('0b0,b0l = ',1p2e13.4,' b0s,b0j = ',1p2e13.4,
1 ' b0v,b0a,b0b = ',1p3e13.4)
10 format('0b1,b1l = ',1p2e13.4,' b1s,b1j = ',1p2e13.4,
1 ' b1v,b1a,b1b = ',1p3e13.4)
11 format('0cp,cpl = ',1p2e13.4,' cps,cpj = ',1p2e13.4,
1 ' cpv,cpa,cpb = ',1p3e13.4/)
12 format('0Reference state properties for NaCl from NACL1'//
1 ' g20,s20 = ',1p2e13.4,' h20,cp20 = ',1p2e13.4,
2 ' v20,vt20,vp20 = ',1p3e13.4/)
13 format('0Excess properties from EXCESS'//
1 ' gex,gl,ex,g2ex = ',1p3e13.4,' hex,hlex,h2ex = ',1p3e13.4)
14 format('0sex,slex,s2ex = ',1p3e14.4,' cpex,cplex,cp2ex = ',
1 1p3e13.4)
15 format('0vex,vlex,v2ex = ',1p3e15.4,' vtex,vtlex,vt2ex = ',
1 1p3e13.4)
26 format('0vpex,vplex,vp2ex = ',1p3e13.4/)
16 format('0Partial molal quantities from NACL2'//
1 ' g1,g2 = ',1p2e13.4,' s1,s2 = ',1p2e13.4,
```

```

2 ' '//' h1,h2 =',1p2e13.4,' cp1,cp2 =',1p2e13.4)
17 format('0v1,v2 =',1p2e13.4,' vt1,vt2 =',1p2e13.4,
1 ' vp1,vp2 =',1p2e13.4/)
18 format('00smotic coefficient and mean ion activity coefficient fro
1m NaCL2.'//' phi,gam =',1p2e13.4/)
19 format('0Properties of solid NaCl from NaCL3'//
1 ' g2s,h2s =',1p2e13.4,' s2s,cp2s =',1p2e13.4,//
2 ' v2s,vt2s,vp2s =',1p3e13.4/)
20 format('0Change of free energy and other thermodynamic functions a
1ccompanying dissolution of NaCl in water from NaCL3'//
2 ' dgs0,dss0 =',1p2e13.4,' dhs0,dcps0 =',1p2e13.4,//
3 ' dvs0,dvts0,dvps0 =',1p3e13.4/)
end

```

```

subroutine table2(t,p,c)

```

```

This subroutine generates output which may be compared to
the tables in report LBL-15512.

```

created by c.h.jun

```

implicit real*8(a-h,o-z)
real*8 i,m,mr,mw,n1,m2
common /debye/ ap,ah,as,aj,av,aa,ab
common /excess/ gex,glex,g2ex,hex,hlex,h2ex,sex,slex,
1 s2ex,cpex,cplex,cp2ex,vex,vlex,v2ex,vtex,vtlex,
2 vt2ex,vpex,vplex,vp2ex
common /haar/ f10,g10,u10,h10,s10,cv10,cp10,cs10,v10,a10,b10
common /partial/ g1,g2,s1,s2,h1,h2,cp1,cp2,v1,v2,vt1,vt2,
1 vp1,vp2
common /pg/ phi,gam
common /ref/ g20,s20,h20,cp20,v20,vt20,vp20
common /soln/ dgs0,dss0,dhs0,dcps0,dvs0,dvts0,dvps0
common /vir/ b0,b0l,b0s,b0j,b0v,b0a,b0b,b1,b1l,b1s,b1j,
1 b1v,b1a,b1b,cp,cpl,cps,cpj,cpv,cpa,cpb
common /cval2/ tr,pr,mr,mw,n1
common /beta/ ps,dpdt,er,b,bt,btt,bb,bbt,bbtt,y,yr,yt,ytt
common /delta/ v,vt,vp,vtt,vpp,vtp
common /diel/ d,dt,dttd,dp,dpp,dtp
common /ideal/ f0,g0,u0,h0,s0,cv0,cp0
common /solid/ g2s,h2s,s2s,cp2s,v2s,vt2s,vp2s

```

```

in this routine r = 8.3144 j/k-mol

```

```

data r/8.3144/
data m2/58.44/

```

```

This subroutine generates a table whose entries correspond to the
tables in Report LBL-15512. The original purpose of this
subroutine was to check values generated by this version of the
program against tables in the report. Class 1 values, which are
normally printed out are those presented in the report,
and have been directly verified. Note: the values for specific
entropy and specific enthalpy (tables 16 and 17) were incorrect
in LBL-15512, but have been corrected in the program and in the
journal article.

```

```

tc=t-273.15

```

```

f15= ( 1.e3*v10 + c*v20 + vex )/( 1.e3 + c*m2 )

```

```
f16 = ( n1*s10 + c*s20 + sex + 2.e0*c*(1.e0-dlog(c)) )/
1      ( 1.e3 + c*m2 )
f17 = ( n1*h1 + c*h2 )/( 1.e3 + c*m2 )
```

c

```
write(6,100) t,tc,p,c
write(6,101) g10,g20,ap,b0,b1,1.e3*cp/2.
write(6,102) s10,s20,as,b0s,b1s,1.e3*cps/2.
write(6,103) h10,h20,ah,1.e3*b0l,1.e3*b1l,1.e3*cpl/2.
write(6,104) cp10,cp20,aj,1.e6*b0j,1.e6*b1j,1.e6*cpj/2.
write(6,105) mw*v10,v20,av,1.e6*b0v,1.e6*cpv/2.
write(6,106) 1.e3*mw*v10*a10,vt20,aa,1.e6*b0a,1.e9*cpa/2.
write(6,107) -1.e3*mw*v10*b10,1.e3*vp20,1.e3*ab,1.e9*b0b
write(6,108) gam
write(6,109) phi
write(6,110) dss0
write(6,111) dhs0
write(6,112) sex/c
write(6,113) hex/c
write(6,114) cpex/c
write(6,115) 1./f15
write(6,116) r*f16
write(6,117) r*t*f17
100 format('0Table 2. The following variables (class 1) are tabul
lated in report LBL-15512'' and are presented here in the same ord
2er.'// ' t,tc =',2f9.2, ' p =',f10.3, ' m =',f8.3/)
101 format('//1x,'table 1 g10,g20,ap,b0,b1,1e3*cp/2',
1 25x,5f10.4,f10.2)
102 format('//1x,'table 2 s10,s20,as,b0s,b1s,1e3*cps/2',
1 22x,5f10.4,f10.2)
103 format('//1x,'table 3 h10,h20,ah,1e3*b0l,1e3*b1l,1e3*cpl/2',
1 14x,f10.4,5f10.3)
104 format('//1x,'table 4 cp10,cp20,aj,1e6*b0j,1e6*b1j,1e6*cpj/2',
1 12x,6f10.3)
105 format('//1x,'table 5 mw*v10,v20,av,1e6*b0v,1e6*cpv/2',
1 19x,5f10.3)
106 format('//1x,'table 6 1e3*mw*v10*a10,vt20,aa,1e6*b0a,1e9*cpa/2',
1 10x,5f10.3)
107 format('//1x,'table 7 -1e3*mw*v10*b10,1e3*vp20,1e3*ab,1e9*b0b',
1 11x,4f10.3)
108 format('//1x,'table 8 gam',47x,f10.3)
109 format('/1x,'table 9 phi',47x,f10.3)
110 format('/1x,'table 10 dss0',46x,f10.3)
111 format('/1x,'table 11 dhs0',46x,f10.3)
112 format('/1x,'table 12 sex/m',45x,f10.3)
113 format('/1x,'table 13 hex/m',45x,f10.3)
114 format('/1x,'table 14 cpex/m',44x,f10.3)
115 format('/1x,'table 15 density',43x,f10.5)
116 format('/1x,'table 16 specific entropy(j/g-k)',27x,f10.3)
117 format('/1x,'table 17 specific enthalpy(j/g)',28x,f10.1)
```

c

c The remaining code is normally skipped over. Class 2 and 3
c variables were not verified directly. Class 2 variables
c were used in the calculation of class 1 variables and have
c thereby been verified indirectly.

c

c To execute the remaining code, remove the following statement.

c

```
if(tc.lt.1d10) return
write(6,200)
write(6,201)v,vt,vp,vtt,vpp,vtp
```

```
write(6,202)d,dt,dt,dp,dpp,dt  
write(6,203)f10,u10,cv10  
write(6,204)f0,u0,cv0  
write(6,205)b1v  
write(6,206)gex,h1ex,h2ex,vex  
write(6,207)h1,h2  
write(6,208)h2s,s2s
```

c

```
write(6,300)  
write(6,301)cs10  
write(6,302)g0,h0,s0,cp0  
write(6,303)b1a,b1b,cpb  
write(6,304)glex,g2ex  
write(6,305)slex,s2ex,cp1ex,cp2ex  
write(6,306)v1ex,v2ex,vtex,vt1ex,vt2ex  
write(6,307)g1,g2,s1,s2,cp1,cp2  
write(6,308)v1,v2,vt1,vt2,vp1,vp2  
write(6,309)g2s,cp2s,v2s,vt2s,vp2s  
write(6,310)dgs0,dcps0,dvs0,dvts0,dvps0  
write(6,311)ps,dpdt
```

c

```
200 format(1x/////1  
1' Class 2; variables used in calculation of class 1 variables.'  
2 ///)  
201 format(/1x,'v,vt,vp,vtt,vpp,vtp',6x,1p6e13.4)  
202 format(/1x,'d,dt,dt,dp,dpp,dt',6x,1p6e13.4)  
203 format(/1x,'f10,u10,cv10',12x,1p3e13.4)  
204 format(/1x,'f0,u0,cv0',16x,1p3e13.4)  
205 format(/1x,'b1v',22x,1p1e13.4)  
206 format(/1x,'gex,h1ex,h2ex,vex',7x,1p4e13.4)  
207 format(/1x,'h1,h2',20x,1p2e13.4)  
208 format(/1x,'h2s,s2s',18x,1p2e13.4)
```

c

```
300 format(/////1x,'Class 3 ; values not verified'///)  
301 format(/1x,'cs10',21x,1p1e13.4)  
302 format(/1x,'g0,h0,s0,cp0',13x,1p4e13.4)  
303 format(/1x,'b1a,b1b,cpb',14x,1p3e13.4)  
304 format(/1x,'glex,g2ex',16x,1p2e13.4)  
305 format(/1x,'slex,s2ex,cp1ex,cp2ex',3x,1p4e13.4)  
306 format(/1x,'v1ex,v2ex,vtex,vt1ex,vt2ex',12x,1p5e13.4)  
307 format(/1x,'g1,g2,s1,s2,cp1,cp2',6x,1p6e13.4)  
308 format(/1x,'v1,v2,vt1,vt2,vp1,vp2',4x,1p6e13.4)  
309 format(/1x,'g2s,cp2s,v2s,vt2s,vp2s',3x,1p5e13.4)  
310 format(/1x,'dgs0,dcps0,dvs0,dvts0,dvps0',11x,1p5e13.4)  
311 format(/1x,'psat,dpdt',16x,1p2e13.4)  
return  
end
```

```

subroutine nacl1(t,p)

implicit real*8(a-h,o-z)
real*8 i,m,mr,mw,n1
common /cval2/ tr,pr,mr,mw,n1
common /excess/ gex,glex,g2ex,hex,h1ex,h2ex,sex,slex,
1 s2ex,cpex,cplex,cp2ex,vex,vlex,v2ex,vtex,vtlex,vt2ex,
2 vpex,vplex,vp2ex
common /haar/ f10,g10,u10,h10,s10,cv10,cp10,cs10,v10,a10,b10
common /ref/ g20,s20,h20,cp20,v20,vt20,vp20
common /vir/ b0,b0l,b0s,b0j,b0v,b0a,b0b,b1,b1l,b1s,b1j,
1 b1v,b1a,b1b,cp,cpl,cps,cpj,cpv,cpa,cpb
common /zz/ z(53),zl(53),zh(53)

This program calculates the properties of NaCl solutions
using the accurate empirical model published by Pitzer,
Peiper, and Busey, J. Phys. Chem. Ref. Data, v.13, 1-102 (1984)
and in Lawrence Berkeley Laboratory Report LBL-15512.

The tables in the cited references were generated using
an earlier version of this program, which was written by
J.C. Peiper.

The present version of NACL was developed by O. Weres with
assistance from C.H. Jun.

- O. Weres, March 1986

data zl /-71659.53087, 2.348333613, -8.366848370e-5,
1 2.401816892e-9, 624.8820839, -5.369711934e-4,
2 3.512696524e-7, 0.0, -110.74702, 0.03890080081,
3 2.697345512e-6, -6.274687539e-10, -1.5267612e-5, 0.0,
4 516.99706, -5.9960301e6, -656.81518, 24.87918316,
5 -2.155273113e-5, 5.016685500e-8, 0.0, -4.4640952,
6 0.01108709891, -6.447976135e-8, -2.323403150e-10, 0.0,
7 -5.219487066e-6, 2.444520990e-10, 2.852706550e-13,
8 -1.569623077, 2.233786380e-3, -6.393389110e-7,
9 4.527057333e-11, 5.4151933, 0.0, 0.0, 0.0, 119.31966,
x -0.48309327, 1.4068095e-3, -4.2345814, -6.1084589,
1 0.4074380280, -6.8152430e-6, -0.075354649,
2 1.260901375e-4, 6.2480692e-8, 1.899437268e-8,
3 -1.0731284e-10, 0.3213657227e0, -2.5382945e-4, 0.0, 0.0/
data zh/ -71637.20299, 2.220901258, -7.799139523e-5,
5 -4.809927222e-9, 624.6812520, 6.015978715e-4,
6 3.406907417e-7, 2.196204308e-11, -110.74702,
7 0.03949447337, -6.531347462e-7, -6.478189444e-10,
8 -1.584201250e-5, 3.245200554e-9, 516.99706,
9 -5.9960301e6, -656.81518, 24.86912950, 5.381275267e-5,
x -5.588746990e-8, 6.589326333e-12, -4.4640952,
1 0.01110991383, -2.657339906e-7, 1.746006963e-10,
2 1.046261900e-14, -5.307012889e-6, 8.634023325e-10,
3 -4.178596200e-13, -1.579365943, 2.202282079e-3,
4 -1.310550324e-7, -6.381368333e-11, 9.706578079,
5 -0.02686039622, 1.534474401e-5, -3.215398267e-9,
6 119.31966, -0.48309327, 1.4068095e-3, -4.2345814,
7 -6.1084589, 0.4021779279, 2.2902837e-5, -0.075354649,
8 1.531767295e-4, -9.0550901e-8, -1.538600820e-8,

```



```

9      8.6926600e-11, 0.3531041360, -4.3314252e-4,
x      -0.09187145529, 5.1904777e-4/

```

```

c      In this subroutine r = 83.144 bar-cu.cm/k-mol
c

```

```

      data mw,n1 /18.01533, 55.50825/
      data tr,pr,mr,r /298.15, 1.01325, 5.550825, 83.144/

```

```

c      Set array of z values.
c

```

```

      if(t.ge.338.15) then
      do 1 k=1,53
1  z(k)=zh(k)
      else
      do 2 k=1,53
2  z(k)=zl(k)
      end if

```

```

c      Calculate virial coefficients.
c

```

```

      b0 = z(17)/t + z(18) + z(19)*p + z(20)*p**2
1      + z(21)*p**3 + z(22)*dlog(t)
2      + ( z(23) + z(24)*p + z(25)*p**2
3      + z(26)*p**3 )*t
4      + ( z(27) + z(28)*p + z(29)*p**2 )*t**2
5      + ( z(30) + z(31)*p + z(32)*p**2
6      + z(33)*p**3 )/(t-227e0)
7      + ( z(34) + z(35)*p + z(36)*p**2
8      + z(37)*p**3 )/(680e0-t)
      b0l = -z(17)/t**2 + z(22)/t + z(23) + z(24)*p
1      + z(25)*p**2 + z(26)*p**3
2      + 2e0*( z(27) + z(28)*p + z(29)*p**2 )*t
3      - ( z(30) + z(31)*p + z(32)*p**2
4      + z(33)*p**3 )/(t-227e0)**2
5      + ( z(34) + z(35)*p + z(36)*p**2
6      + z(37)*p**3 )/(680e0-t)**2
      b0s = z(18) + z(19)*p + z(20)*p**2 + z(21)*p**3
1      + z(22)*(1e0+dlog(t))
2      + 2e0*( z(23) + z(24)*p + z(25)*p**2
3      + z(26)*p**3 )*t
4      + 3e0*( z(27) + z(28)*p + z(29)*p**2 )*t**2
5      + ( z(30) + z(31)*p + z(32)*p**2
6      + z(33)*p**3 )*(1e0-t/(t-227e0))/(t-227e0)
7      + ( z(34) + z(35)*p + z(36)*p**2
8      + z(37)*p**3 )*(1e0+t/(680e0-t))/(680e0-t)
      b0j = z(22)/t**2 + 2e0*( z(23) + z(24)*p + z(25)*p**2
1      + z(26)*p**3 )/t
2      + 6e0*( z(27) + z(28)*p + z(29)*p**2 )
3      + 454e0*( z(30) + z(31)*p + z(32)*p**2
4      + z(33)*p**3 )/(t*(t-227e0)**3)
5      + 1360e0*( z(34) + z(35)*p + z(36)*p**2
6      + z(37)*p**3 )/(t*(680e0-t)**3)
      b0v = z(19) + 2e0*z(20)*p + 3e0*z(21)*p**2
1      + ( z(24) + 2e0*z(25)*p + 3e0*z(26)*p**2 )*t
2      + ( z(28) + 2e0*z(29)*p )*t**2
3      + ( z(31) + 2e0*z(32)*p + 3e0*z(33)*p**2 )/(t-227e0)
4      + ( z(35) + 2e0*z(36)*p + 3e0*z(37)*p**2 )/(680e0-t)
      b0a = ( z(19) + 2e0*z(20)*p + 3e0*z(21)*p**2 )/t
1      + 2e0*z(24) + 4e0*z(25)*p + 6e0*z(26)*p**2
2      + 3e0*( z(28) + 2e0*z(29)*p )*t

```

```

4      - 227e0*( z(31) + 2e0*z(32)*p + 3e0*z(33)*p**2 )
5          / (t*(t-227e0)**2)
6      + 680e0*( z(35) + 2e0*z(36)*p + 3e0*z(37)*p**2 )
7          / (t*(680e0-t)**2)
      b0b = 2e0*z(20) + 6e0*z(21)*p
1          + ( 2e0*z(25) + 6e0*z(26)*p )*t
2          + 2e0*z(29)*t**2
3          + ( 2e0*z(32) + 6e0*z(33)*p )/(t-227e0)
4          + ( 2e0*z(36) + 6e0*z(37)*p )/(680e0-t)
      b1 = z(38)/t + z(39) + z(40)*t + z(41)/(t-227e0)
      b1l = -z(38)/t**2 + z(40) - z(41)/(t-227e0)**2
      b1s = z(39) + 2e0*z(40)*t + z(41)*(1e0-t/(t-227e0))/(t-227e0)
      b1j = 2e0*z(40)/t + 454e0*z(41)/(t*(t-227e0)**3)
      b1v=0.
      b1a=0.
      b1b=0.
      cp = z(42)/t + z(43) + z(44)*p + z(45)*dlog(t)
1          + ( z(46) + z(47)*p )*t
2          + ( z(48) + z(49)*p )*t**2
3          + ( z(50) + z(51)*p )/(t-227e0)
4          + ( z(52) + z(53)*p )/(680e0-t)
      cpl = -z(42)/t**2 + z(45)/t + z(46) + z(47)*p
1          + 2e0*( z(48) + z(49)*p )*t
2          - ( z(50) + z(51)*p )/(t-227e0)**2
3          + ( z(52) + z(53)*p )/(680e0-t)**2
      cps = z(43) + z(44)*p + z(45)*(1e0+dlog(t))
1          + 2e0*( z(46) + z(47)*p )*t
2          + 3e0*( z(48) + z(49)*p )*t**2
3          + ( z(50) + z(51)*p )*(1e0-t/(t-227e0))/(t-227e0)
4          + ( z(52) + z(53)*p )*(1e0+t/(680e0-t))/(680e0-t)
      cpj = z(45)/t**2 + 2e0*( z(46) + z(47)*p )/t
1          + 6e0*( z(48) + z(49)*p )
2          + 454e0*( z(50) + z(51)*p )/(t*(t-227e0)**3)
3          + 1360e0*( z(52) + z(53)*p )/(t*(680e0-t)**3)
      cpv = z(44) + z(47)*t + z(49)*t**2 + z(51)/(t-227e0)
1          + z(53)/(680e0-t)
      cpa = z(44)/t + 2e0*z(47) + 3e0*z(49)*t
1          - 227e0*z(51)/(t*(t-227e0)**2)
2          + 680e0*z(53)/(t*(680e0-t)**2)
      cpb=0.

```

c
c
c
c
c
c
c
c
c
c
c

Calculate reference state properties.

This routine calculates the properties of nacl(aq) in its reference state (hypothetical 1 mol/kg solution at t and p). The gibbs energy, enthalpy, entropy and heat capacity are dimensionless, whereas the volume and its derivatives have dimensions of cc/mol, cc/mol-k and cc/mol-bar (e.g., a20 and b20 are the derivatives (d v20 /dt) and (d v20 /dp), not the usual quantities (d ln v20 /dt) and (d ln v20 /dp).

```

      call excess(t,p,mr)
      g20 = -10e0*g10 - gex/mr
1          + ( z(1) + z(2)*p + z(3)*p**2 + z(4)*p**3 )/t
2          + z(5) + z(6)*p + z(7)*p**2 + z(8)*p**3
3          + z(9)*dlog(t)
4          + ( z(10) + z(11)*p + z(12)*p**2 )*t
5          + ( z(13) + z(14)*p )*t**2
6          + z(15)/(t*(t-227e0)) + z(16)/(t*(680e0-t)**3)
      s20 = -10e0*s10 - sex/mr

```

```

1      - z(5) - z(6)*p - z(7)*p**2 - z(8)*p**3
2      - z(9)*(1e0 + dlog(t))
3      - 2e0*( z(10) + z(11)*p + z(12)*p**2 )*t
4      - 3e0*( z(13) + z(14)*p )*t**2
5      + z(15)/(t-227e0)**2 - 3e0*z(16)/(680e0-t)**4
h20 = -10e0*h10 - hex/mr
1      + ( z(1) + z(2)*p + z(3)*p**2 + z(4)*p**3 )/t
2      - z(9) - ( z(10) + z(11)*p + z(12)*p**2 )*t
3      - 2e0*( z(13) + z(14)*p )*t**2
4      + z(15)*(2e0*t-227e0)/(t*(t-227e0)**2)
5      - z(16)*(4e0*t-680e0)/(t*(680e0-t)**4)
cp20 = -10e0*cp10 - cpex/mr
1      - z(9) - 2e0*( z(10) + z(11)*p + z(12)*p**2 )*t
2      - 6e0*( z(13) + z(14)*p )*t**2
3      - 2e0*z(15)*t/(t-227e0)**3
4      - 12e0*z(16)*t/(680e0-t)**5
v20 = -10e0*mw*v10 - vex/mr
1      + r*( z(2) + 2e0*z(3)*p + 3e0*z(4)*p**2 )
2      + r*t*( z(6) + 2e0*z(7)*p + 3e0*z(8)*p**2 )
3      + r*t**2*( z(11) + 2e0*z(12)*p ) + r*t**3*z(14)
vt20 = -10e0*mw*v10*a10 - vtex/mr
1      + r*( z(6) + 2e0*z(7)*p + 3e0*z(8)*p**2 )
2      + 2e0*r*t*( z(11) + 2e0*z(12)*p ) + 3e0*r*t**2*z(14)
vp20 = 10e0*mw*v10*b10 - vpex/mr
1      + r*( 2e0*z(3) + 6e0*z(4)*p )
2      + r*t*( 2e0*z(7) + 6e0*z(8)*p ) + 2e0*r*t**2*z(12)
return
end

```

```

subroutine nacl2(t,p,m)

```

```

implicit real*8(a-h,o-z)
real*8 i,m,mr,mw,n1
common /cval2/ tr,pr,mr,mw,n1
common /debye/ ap,ah,as,aj,av,aa,ab
common /excess/ gex,glex,g2ex,hex,hlex,h2ex,sex,slex,
1 s2ex,cpex,cplex,cp2ex,vex,vlex,v2ex,vtex,vtlex,vt2ex,
2 vpex,vplex,vp2ex
common /haar/ f10,g10,u10,h10,s10,cv10,cp10,cs10,v10,a10,b10
common /partial/ g1,g2,s1,s2,h1,h2,cp1,cp2,v1,v2,vt1,vt2,
1 vp1,vp2
common /pg/ phi,gam
common /ref/ g20,s20,h20,cp20,v20,vt20,vp20
common /vir/ b0,b0l,b0s,b0j,b0v,b0a,b0b,b1,b1l,b1s,b1j,
1 b1v,b1a,b1b,cp,cpl,cps,cpj,cpv,cpa,cpb

```

```

In this routine r = 83.144 bar-cu.cm/k-mol

```

```

data r/83.144/
i = m
si = dsqrt(i)

```

```

Calculate partial molal quantities.

```

```

call excess(t,p,i)
g1 = g10 - 2e0*m/n1 + glex
g2 = g20 - 2e0*dlog(m) + g2ex
s1 = s10 + 2e0*m/n1 + sllex

```

```

s2 = s20 - 2e0*dlog(m) + s2ex
h1 = h10 + h1ex
h2 = h20 + h2ex
cp1 = cp10 + cp1ex
cp2 = cp20 + cp2ex
v1 = mw*v10 + v1ex
v2 = v20 + v2ex
vt1 = mw*v10*a10 + vt1ex
vt2 = vt20 + vt2ex
vp1 = -mw*v10*b10 + vp1ex
vp2 = vp20 + vp2ex

```

Calculate phi and gam

```

b = 2e0*b0 + b1 *(1e0-(1e0+2e0*si-2e0*i)*dexp(-2e0*si))/(2e0*i)
alng = -ap*( si/(1e0+1.2*si) + dlog(1e0+1.2*si)/0.6 )
1      + b*m + 1.5*cp *m**2
gam = dexp(alng)
phi = 1e0 - ap*si/(1e0+1.2*si)
1      + m*( b0 + b1 *dexp(-2e0*si) )
2      + cp *m**2
return
end

```

subroutine excess(t,p,i)

```

implicit real*8(a-h,o-z)
real*8 i,m,mr,mw,n1
common /cval2/ tr,pr,mr,mw,n1
common /debye/ ap,ah,as,aj,av,aa,ab
common /excess/ gex,glex,g2ex,hex,h1ex,h2ex,sex,s1ex,
1 s2ex,cpex,cplex,cp2ex,vex,v1ex,v2ex,vtex,vt1ex,vt2ex,
2 vpex,vplex,vp2ex
common /vir/ b0,b0l,b0s,b0j,b0v,b0a,b0b,b1,b1l,b1s,b1j,
1 b1v,b1a,b1b,cp,cpl,cps,cpj,cpv,cpa,cpb

```

In this routine r=83.144 bar-cu.cm/k-mol

data r/83.144/

All quantities are dimensionless, except
for the volume and its derivatives.

```

m=i
si=dsqrt(i)
f0 = dexp(-2e0*si)
f1 = dlog(1e0+1.2*si)
f2 = (1e0-(1e0+2e0*si)*f0)/(2e0*i)
f3 = (1e0-(1e0+2e0*si-2e0*i)*f0)/(4e0*i)
f4 = i**1.5
gex = -4e0*ap*i*f1/1.2
1      + 2e0*m**2*( b0
2      + b1 *f2
3      + m*cp /2e0 )
glex = 2e0*ap*f4/(1e0+1.2*si)
1      - 2e0*m**2*( b0 + b1 *f0 + m*cp )
g2ex = -2e0*ap*( si/(1e0+1.2*si) + f1/0.6 )
1      + 4e0*m*( b0

```

```

2          + b1 *f3 )
3      + 3e0*m**2*cp
hex = ah*i*f1/1.2
1      - 2e0*t*m**2*( b0l
2          + b1l *f2
3          + m*cpl /2e0 )
hlex = (-ah*f4/(2e0*(1e0+1.2*si))
1      + 2e0*t*m**2*( b0l + b1l *f0
2          + m*cpl ))/n1
h2ex = ah*( si/(1e0+1.2*si) + f1/0.6 )/2e0
1      - 4e0*t*m*( b0l
2          + b1l *f3 )
3      - 3e0*t*m**2*cpl
sex=hex-gex
slex=hlex-glex
s2ex=h2ex-g2ex
cpex = aj*i*f1/1.2
1      - 2e0*(t*m)**2*( b0j
2          + b1j *f2
3          + cpj *m/2e0 )
cplex = -aj*f4/(2e0*(1e0+1.2*si))
1      + 2e0*(t*m)**2*( b0j + b1j *f0
2          + cpj *m )
cp2ex = aj*( si/(1e0+1.2*si) + f1/0.6 )/2e0
1      - 4e0*t**2*m*( b0j
2          + b1j *f3 )
3      - 3e0*(t*m)**2*cpj
vex = av*i*f1/1.2
1      + 2e0*r*t*m**2*( b0v
2          + b1v *f2
3          + cpv *m/2e0 )
vlex = av*f4/(2e0*(1e0+1.2*si))
1      - 2e0*r*t*m**2*( b0v + b1v *f0
2          + cpv *m )
v2ex = -av*( si/(1e0+1.2*si) + f1/0.6 )/2e0
1      + 4e0*r*t*m*( b0v
2          + b1v *f3 )
3      + 3e0*r*t*m**2*cpv
vtex = aa*i*f1/1.2
1      + 2e0*r*t*m**2*( b0a
2          + b1a *f2
3          + cpa *m/2e0 )
vtlex = aa*f4/(2e0*(1e0+1.2*si))
1      - 2e0*r*t*m**2*( b0a + b1a *f0
2          + cpa *m )
vt2ex = -aa*( si/(1e0+1.2*si) + f1/0.6 )/2e0
1      + 4e0*r*t*m*( b0a
2          + b1a *f3 )
3      + 3e0*r*t*m**2*cpa
vpex = ab*i*f1/1.2
1      + 2e0*r*t*m**2*( b0b
2          + b1b *f2
3          + cpb *m/2e0 )
vp1ex = ab*f4/(2e0*(1e0+1.2*si))
1      - 2e0*r*t*m**2*( b0b + b1b *f0
2          + cpb *m )
vp2ex = -ab*( si/(1e0+1.2*si) + f1/0.6 )/2e0
1      + 4e0*r*t*m*( b0b
2          + b1b *f3 )
3      + 3e0*r*t*m**2*cpb

```

```

return
end

```

```

subroutine nacl3(t,p,m)

```

```

implicit real*8(a-h,o-z)
real*8 i,m,mr,mw,n1
common /cval2/ tr,pr,mr,mw,n1
common /ref/ g20,s20,h20,cp20,v20,vt20,vp20
common /solid/ g2s,h2s,s2s,cp2s,v2s,vt2s,vp2s
common /soln/ dgs0,dss0,dhs0,dcps0,dvs0,dvts0,dvps0
data s2sr,a,b,aa,bb /8.676, -1734.4, 37.589, -5.5254, -9.81e-4/
data v2r /26.993/
data cc,dd,ee /4.5e-8, 9.30e-5, -0.031728/
data dhs0r /466.99/

```

```

In this routine r = 83.144 bar-cu.cm/k-mol

```

```

data r/83.144/
save s2sr,a,b,aa,bb,v2r,cc,dd,ee,dhs0r

```

This routine calculates the thermal properties of NaCl(c) based on information from K.K. Kelley (U.S. Bur. Mines Report #584, 1960) and from the American Institute of Physics Handbook (3rd ed., McGraw-Hill, New York, 1972). The gibbs energy, enthalpy, entropy and heat capacity are dimensionless. The volume and its derivatives have units of cc/mol, cc/mol-k and cc/mol-bar, e.g. a2s and b2s are the derivatives (dv/dt) and (dv/dp), not the quantities (d ln v/dt) and (d ln v/dp).

```

g2s = -s2sr + a/t + b + aa*dlog(t) + bb*t
1      + v2r*(p-pr)*dexp(cc*t**2 + dd*t + ee)/(r*t)
h2s = a/t - aa - bb*t + v2r*(p-pr)*(2e0*cc*t**2 + dd*t + 1)
1      *dexp(cc*t**2 + dd*t + ee)/(r*t)
s2s = s2sr - b - aa*(1e0+dlog(t)) - 2e0*bb*t
1      + v2r*(p-pr)*(2e0*cc*t + dd)*dexp(cc*t**2 + dd*t + ee)/r
cp2s = -aa-2e0*bb*t+v2r*(p-pr)*(4e0*cc**2e0*t**3+4e0*cc*dd*t**2
1      + (dd**2 + 6e0*cc)*t + 2e0*dd)*dexp(cc*t**2 + dd*t + ee)/r
v2s = v2r*dexp(cc*t**2 + dd*t + ee)
vt2s = v2r*(2e0*cc*t + dd)*dexp(cc*t**2 + dd*t + ee)
vp2s=0

```

```

Calculate properties of solution.

```

```

dgs0 = g20 - g2s + dhs0r/t
dss0 = s20 - s2s
dhs0 = h20 - h2s + dhs0r/t
dcps0 = cp20 - cp2s
dvs0 = v20 - v2s
dvts0 = vt20 - vt2s
dvps0 = vp20 - vp2s
return
end

```

```
implicit real*8 (a-h,o-z)
real*8 q(18),Int
common /alpha/ a(9,7),al,bet,gam,b0,c,d,e,bb0,b1,b2,b4
common /beta/ ps,dpdt,er,b,bt,btt,bb,bbt,bbtt,y,yr,yt,ytt
common/delta/v,vt,vp,vtt,vpp,vtp
common /haar/ f,g,u,h,s,cv,cp,cs,v10,alfa,beta
common /ideal/ f0,g0,u0,h0,s0,cv0,cp0
data q /20.9662681977,1973.0271018,-.483429455355,6.05743189245,
1  -68.3010170092,-106.4285576,-4.3135538513,.458155781,
2  -.047754901883,.0041238460633,-.00027929052852,1.4481695261e-5,
3  -5.6473658748e-7,1.6200446e-8,-3.303822796e-10,
4  4.51916067368e-12,-3.70734122708e-14,1.37546068238e-16/
```

```
data r/4.61518/
save q,r
```

The quantities returned by HAAR are --

f	helmholtz free energy
g	gibbs free energy
u	internal energy
h	enthalpy
s	entropy
cv	heat capacity at constant volume
cp	heat capacity at constant pressure
cs	heat capacity along the saturation curve (cs is set equal to cp for pressures greater than saturation pressure)
v	volume, in cubic centimeters per gram
alfa	expansivity, per kelvin
beta	compressibility, per (10**5 pa)

J. Christopher Peiper
29 March, 1982

```

C *****
C
C
C      call rho(t,p)
C      call debye(t,p)
C
C      v10=v
C      rh=1.0/v
C      tau=t/647.073
C      tt=t/100e0
C      lnt=dlog(t)
C      gz=-(q(1)+q(2)/t)*lnt
C      hz=q(1)+q(2)*(1e0-lnt)/t
C      cpz=q(1)-q(2)/t
C
C      do 7 i=3,18
C      gz=gz-q(i)*tt**(i-6)
C      hz=hz+q(i)*(i-6)*tt**(i-6)
17 cpz=cpz+q(i)*(i-6)*(i-5)*tt**(i-6)
C      f0=gz-1e0
C      g0=gz
C      u0=hz-1e0
C      h0=hz
C      s0=h0-g0
C      cv0=cpz-1e0
C      cp0=cpz
C
C      fz = f0 -
1 dlog(1e0-y) - (bet-1e0)/(1e0-y) + (1e0+a1+bet)/(2e0*(1e0-y)**2)
2   + rh*(bb-gam*b) - (a1-bet+3)/2e0 + dlog(r*rh*t/1.01325)
C      uz = u0 + t*
1 ((1e0/(y-1e0)+(bet-1e0)/(1e0-y)**2-(1e0+a1+bet)/(1e0-y)**3)
2   *yt
3   - rh*(bbt-gam*bt))
C      cvz=cv0 + 2e0*t*((1e0/(y-1e0) + (bet-1e0)/(1e0-y)**2
1   - (1e0+a1+bet)/(1e0-y)**3)*yt - rh*(bbt-gam*bt))
2   +t**2*((2e0*(bet-1e0)/(1e0-y)**3 - 1e0/(1e0-y)**2
3   - 3e0*(1e0+a1+bet)/(1e0-y)**4)*yt**2
4   + (1e0/(y-1e0)+(bet-1e0)/(1e0-y)**2-(1e0+a1+bet)/(1e0-y)**3)*ytt
5   -rh*(bbtt-gam*bt))
C
C      df= 0e0
C      du= 0e0
C      dcv= 0e0
C      do 20 n=1,9
C      s1=a(n,1)
C      s2=a(n,1)
C      s3= 0e0
C
C      do 17 j=2,7
C      s1=s1+a(n,j)/tau**(j-1)
C      s2=s2+j*a(n,j)/tau**(j-1)
17 s3=s3+j*(1e0-j)*a(n,j)/tau**(j-1)
C
C      df=df+s1*(1e0-er)**n/n
C      du=du+s2*(1e0-er)**n/n
20 dcv=dcv+s3*(1e0-er)**n/n
C
C      f=fz+df/(r*t)
C      u=uz+du/(r*t)

```



```

cv=cvz+dcv/(r*t)
g=f+p*v/(r*t)
h=u+p*v/(r*t)
s=h-g
cp=cv-t*vt**2/(vp*r)
cs=cp
if (p.eq.ps) cs=cp-t*vt*dpdt/r
alfa=vt/v
beta=-vp/v

```

```

return
end

```

```

subroutine debye(t,p)

```

```

implicit real*8(a-h,o-z)
real*8 n0,k
common /debye/ ap,ah,as,aj,av,aa,ab
common/delta/v,vt,vp,vtt,vpp,vtp
common /diel/ d,dt,dt,dp,dpp,ntp
data pi,n0,e,k/3.141592658,6.022045e23,4.803242e-10,1.380662e-16/

```

```

In this routine r = 83.144 bar-cu.cm/k-mol

```

```

data r /83.144/
save pi,n0,e,k,r

```

```

This subroutine calculates the Debye-Huckel coefficients.

```

```

rh=1e0/v
call dielec(t,p)
ap=dsqrt(2.*pi*n0*rh/1000.)*e**3/(d*k*t)**1.5/3.
ah=-6.*t*ap*(dt/d +1./t + vt/(3.*v))
aj=ah*t*(2e0/t-1.5*(dt/d+1e0/t+vt/(3e0*v))) - 6e0*t**2*ap*(dtt/d
1 -(dt/d)**2-1e0/t**2+vtt/(3e0*v)-(vt/v)**2/3e0)
av=2.*r*t*ap*(3.*dp/d + vp/v)
aa=av*(1.+ah/(4.*ap))/t
1 +2.*r*t*ap*(3.*(ntp/d -dt*dp/d**2)
2 +vtp/v -vt*vp/v**2)
ab=-av**2/(4.*r*t*ap)
1 +2.*r*t*ap*(3.*(dpp/d -(dp/d)**2) +vpp/v -(vp/v)**2)
as=ah+4.*ap
return
end

```

```

subroutine dielec(t,p)

```

```

implicit real*8(a-h,o-z)
common /diel/ d,dt,dt,dp,dpp,ntp
dimension u(9)
data u/342.79,-.0050866,9.4690e-7,-2.0525,3115.9,-182.89,
1-8032.5,4.2142e6,2.1417/
save u

```

```

This subroutine calculates the dielectric constant of water
and its derivatives using the empirical formula presented by

```

c Bradley and Pitzer, J. Phys. Chem. v. 83, 1599 (1979).

c

```

b=u(7) + u(8)/t +u(9)*t
bt=u(9)-u(8)/t**2
btt=2.*u(8)/t**3
c=u(4)+u(5)/(u(6)+t)
ct=-u(5)/(u(6)+t)**2
ctt=2.*u(5)/(u(6)+t)**3
d1k=u(1)*dexp(u(2)*t+u(3)*t**2)
d1kt=(u(2) + 2.*u(3)*t)*d1k
d1ktt=(2.*u(3)+(u(2)+2.*u(3)*t)**2)*d1k
d=d1k+c*dlog((b+p)/(b+1000.))
dt=d1kt+c*(1./(b+p)-1./(b+1000.))*bt + ct*dlog((b+p)/(b+1000.))
dtt=d1ktt-c*(1./(b+p)**2-1./(b+1000.))**2*bt**2+(1./(b+p)-1./
1(b+1000.))*(2.*bt*ct + c*btt)+ctt*dlog((b+p)/(b+1000.))
dp=c/(b+p)
dpp=-c/(b+p)**2
dtp=-c*bt/(b+p)**2 + ct/(b+p)
return
end

```

c

c

c

subroutine rho(t,p)

c

```

implicit real*8 (a-h,o-z)
common /alpha/ a(9,7),al,bet,gam,b0,c,d,e,bb0,b1,b2,b4
common /beta/ ps,dpdt,er,b,bt,btt,bb,bbt,bbtt,y,yr,yt,ytt
common/delta/v,vb,vp,vtt,vpp,vtp
data al,bet,gam /11.0,44.3333333,3.5/
data b0,c,d,e / .747862916,-.354078223,.007159876,-.003528426/
data bb0,b1,b2,b4 /1.127833441,-.594400123,-5.010995915,
1 .636842555/

```

c

c

In this routine r = 4.61518 bar-cu.cm/k-mol

c

```

data r /4.61518/
data a /0.0,0.0,68833.257944332,0.0,0.0,0.0,0.0,0.0,0.0,
1 -5306.2968529023,178638.32875422,-2563743.6613260,
1 11797433.655832,-31377774.947767,46561826.115608,
1 -31555231.392127,0.0,6962522.0862664,
2 22744.901424408,-395147.31563338,4821257.5981415,
2 -21734810.110373,52911910.757704,-72752773.275387,
2 47929666.384584,0.0,-10834900.096447,
3 -26627.944829770,0.0,217572.45522644,0.0,
3 -707304.18082074,0.0,0.0,0.0,0.0,
4 7877.9333020687,338038.84280753,-3418301.6969660,
4 10829952.168620,-13802577.177877,4177424.6148294,
4 4091266.4781209,0.0,-2272282.7401688,
5 0.0,0.0,0.0,0.0,0.0,0.0,0.0,0.0,0.0,
6 -698.30527374994,-138550.50202703,1222315.6417448,
6 -2544199.8064049,-2510991.4369001,14016358.244614,
6 -13626369.388386,0.0,3836548.6000660/
save r

```

c

c

```

c *****
c *
c * RHO calculates the density of water at the specified temperature *
c * (t, in kelvins) and pressure (p, in 10**5 pa), using the *

```

```
c * equations of Haar, Gallagher and Kell (Ninth International
c * Conference on the Properties of Steam, Munich, 1979). These
c * equations are valid for 273.15 K .le. t .le. 623.15 K and
c * saturation pressure .le. p .le. 100 mpa. The volume (which is
c * simply the reciprocal of the density) and its temperature and
c * pressure derivatives are also computed. The density has units
c * of grams per cubic centimeter.
c *
c *
c * J. Christopher Peiper
c * 29 march, 1982
c *
c *****
c
c
c tau=t/647.073
c call psat(t)
c if (p.le.0) p=ps
c b=b0+c*dlog(tau)+d/tau**3+e/tau**5
c bt=(c-3e0*d/tau**3-5e0*e/tau**5)/t
c btt=-(c-12e0*d/tau**3-30e0*e/tau**5)/t**2
c bb=bb0+b1/tau+b2/tau**2+b4/tau**4
c bbt=-(b1/tau+2e0*b2/tau**2+4e0*b4/tau**4)/t
c bbtt=(2e0*b1/tau+6e0*b2/tau**2+20e0*b4/tau**4)/t**2
c rh=1.0
c if (p.lt.ps) rh=.001
c
c do 30 i=1,100
c y=rh*b/4e0
c er=dexp(-rh)
c pr = 0e0
c dpr = 0e0
c
c do 15 n=1,9
c s=a(n,1)
c
c do 10 j=2,7
10 s=s+a(n,j)/tau**(j-1)
c
c pr=pr+s*(1e0-er)**(n-1)
15 dpr=dpr+s*(1e0-er)**(n-1)*(2e0+rh*(n*er-1e0)/(1e0-er))
c
c pr=rh**2*er*pr+rh*r*t*((1e0+al*y+bet*y**2)/(1e0-y)**3+4e0*y*
1 (bb/b-gam))
c dpr=rh*er*dpr + r*t*((1e0+2e0*al*y+3e0*bet*y**2)/(1e0-y)**3
1 +3e0*y*(1e0+al*y+bet*y**2)/(1e0-y)**4 + 8e0*y*(bb/b-gam))
c rhn=rh+(p-pr)/dpr
c if (abs(rhn-rh).lt.abs(rhn/1.e6).and.abs(p-pr).lt.abs(p/1.e5))
1 go to 35
30 rh=rhn
c
c print 100, t,p,rhn
c
c 35 rh=rhn
c y=rh*b/4e0
c yt=rh*bt/4e0
c ytt=rh*btt/4e0
c yr=b/4e0
c er=dexp(-rh)
c pr = 0e0
c dpr = 0e0
```

```
dpt = 0e0
dpr = 0e0
dptt = 0e0
dprt = 0e0

c
do 50 n=1,9
  an = n
  s=a(n,1)
  s1 = 0e0
  s2 = 0e0

c
do 40 j=2,7
  aj = j
  s=s+a(n,j)/tau**(j-1)
  s1=s1+(1e0-aj)*a(n,j)/tau**(j-1)
40 s2=s2+aj*(aj-1e0)*a(n,j)/tau**(j-1)

c
pr=pr+s*(1e0-er)**(n-1)
dpr=dpr+s*(2e0+rh*(an*er-1e0)/(1e0-er))*(1e0-er)**(n-1)
dpt=dpt+s1*(1e0-er)**(n-1)
dpr=dpr+s*((1e0+rh*(an*er-1e0)/(1e0-er))*
1 (2e0+rh*(an*er-1e0)/(1e0-er))
2 *(1e0-er)**(n-1)+rh*(1e0-er)**(n-2)
3 *(an*er*(1e0-rh)-1e0-rh*er*(an*er-1e0)/(1e0-er)))
dptt=dptt+s2*(1e0-er)**(n-1)
50 dprt=dprt+s1*(2e0+rh*(an*er-1e0)/(1e0-er))*(1e0-er)**(n-1)

c
pr=rh**2*er*pr+rh*r*t*((1e0+al*y+bet*y**2)/(1e0-y)**3
1 +rh*(bb-gam*b))
dpr=rh*er*dpr+r*t*((1e0+2e0*al*y+3e0*bet*y**2)/(1e0-y)**3
1 +3e0*y*(1e0+al*y+bet*y**2)/(1e0-y)**4
2 +2e0*rh*(bb-gam*b))
dpt=rh**2*er*dpt/t+rh*r*t*((1e0+al*y+bet*y**2)/(1e0-y)**3
1 +rh*(bb-gam*b))
2 +rh*r*t*((al+2e0*bet*y)/(1e0-y)**3
3 +3e0*(1e0+al*y+bet*y**2)/(1e0-y)**4)*yt
4 +rh**2*r*t*(bbt-gam*bt)
dptt=rh**2*er*dptt/t**2
1 +2e0*rh*r*t*((al+2e0*bet*y)/(1e0-y)**3
2 +3e0*(1e0+al*y+bet*y**2)/(1e0-y)**4)*yt
3 +2e0*rh**2*r*(bbt-gam*bt)
4 +rh*r*t*(2e0*bet/(1e0-y)**3+6e0*(al+2e0*bet*y)/(1e0-y)**4
5 +12e0*(1e0+al*y+bet*y**2)/(1e0-y)**5)*yt**2
6 +rh*r*t*((al+2e0*bet*y)/(1e0-y)**3
7 +3e0*(1e0+al*y+bet*y**2)/(1e0-y)**4)*ytt
8 +rh**2*r*t*(bbtt-gam*btt)
dpr=er*dpr + 2e0*r*t*((al+3e0*bet*y)/(1e0-y)**3
1 +3e0*(1e0+2e0*al*y+3e0*bet*y**2)/(1e0-y)**4
2 +6e0*y*(1e0+al*y+bet*y**2)/(1e0-y)**5)*yr
3 +2e0*r*t*(bb-gam*b)
dprt=rh*er*dprt/t + 2e0*rh*r*t*(bbt-gam*bt)
1 +r*((1e0+2e0*al*y+3e0*bet*y**2)/(1e0-y)**3
2 +3e0*y*(1e0+al*y+bet*y**2)/(1e0-y)**4 + 2e0*rh*(bb-gam*b))
3 +2e0*r*t*((al+3e0*bet*y)/(1e0-y)**3
4 +3e0*(1e0+2e0*al*y+3e0*bet*y**2)/(1e0-y)**4
5 +6e0*y*(1e0+al*y+bet*y**2)/(1e0-y)**5)*yt

c
v=1e0/rh
vt=dpt/(rh**2*dpr)
vp=-1e0/(rh**2*dpr)
```

```

    vtt=2e0*(dpt/dpr)**2/rh**3 + dptt/(dpr*rh**2)
1    - 2.*dprt*dpt/(rh*dpr)**2
3    +dpr*dpt**2/(rh**2*dpr**3)
    vpp=(2e0*dpr+rh*dpr)/(rh*dpr)**3
    vtp=-2e0*dpt/(rh**3*dpr**2) - dpr*dpt/(rh**2*dpr**3)
1    + dprt/(rh*dpr)**2
    return
C
100 format ('0',5x,'at',f7.2,'k and',f10.4,'bars convergence of ',
1    'the density was not achieved --',/10x,'the density is',f8.6)
    end
C
C
C
    subroutine psat(t)
C
    implicit real*8(a-h,o-z)
    common /beta/ ps,dpdt,er,b,bt,btt,bb,bbt,bbtt,y,yr,yt,ytt
    dimension a(8)
    data a/-7.8889166e0,2.5514255e0,-6.716169e0
1    ,33.239495e0,-105.38479e0,174.35319e0,-148.39348e0
2    ,48.631602e0/
    save a
C
C    This subroutine calculates the saturation pressure
C    of pure water and its temperature derivative.
C
    if(t.gt.314.e0) go to 2
    pl=6.3573118e0-8858.843e0/t+607.56335/t**.6
    ps=dexp(pl)
    dpdt=dexp(pl)*(8858.843e0/t**2 - 0.6*607.56335/t**1.6)
    return
2    v=t/647.25
    w=1.-v
    c=0.
    r=0.
    do 4 i=1,8
    z=i
    y=a(i)*w**((z+1.)/2.)
    r=r+ (.5*z+.5)*a(i)*w**((z-1.)/2.)
4    c=c+y
    q=c/v
    dpdt=220.93*dexp(q)*(-r/v/647.25-c/v/v/647.25)
    ps=220.93*dexp(q)
    return
    end

```

NACL calculates the properties of NaCl solutions using the accurate empirical model published by Pitzer, Peiper, and Busey, J. Phys. Chem. Ref. Data v. 13, pp. 1-102 (1984). The valid range is 0-300 C, 0-6 molal, Psat(T)-1000 bar.

Temperature, pressure and molality are

t,tc = 298.15 25.00 p = 1.000 m = 1.000

Table 1 contains all calculated quantities.
See comments in HAAR, RHO, NACL1 and NACL3 for further explanation.

Properties of pure water are calculated first.

Saturation pressure of pure water and temperature derivaive from PSAT

ps,dpdt = 3.1687E-02 1.8887E-03

Reference state properties for liquid water from HAAR

f10,g10 = -2.2170E+01 -2.2169E+01 u10,h10,s10 = -1.3757E+01 -1.3756E+01 8.4130E+00

cv10,cp10,cs10 = 8.9681E+00 9.0645E+00 9.0645E+00 v10,a10,b10 = 1.0029E+00 2.5942E-04 4.5218E-05

Properties of water as ideal gas from HAAR

f0,g0 = -1.9703E+01 -1.8703E+01 u0,h0,s0 = 2.9953E+00 3.9953E+00 2.2698E+01 cv0,cp0 = 3.0396E+00 4.0396E+00

Specific volume of pure water and its derivatives over T and P from RHO

v,vt,vp = 1.0029E+00 2.6019E-04 -4.5352E-05 vtt,vpp,vtp = 9.6806E-06 1.4528E-08 9.5806E-08

Dielectric constant and its derivatives from DIELEC

d,dt,dttd = 7.8384E+01 -3.5930E-01 1.5526E-03 dp,dpp,dtp = 3.7056E-03 -5.4966E-07 -9.9103E-06

Debye-Huckel coefficients from DEBYE

ap,ah,as = 3.915E-01 8.006E-01 2.366E+00 aj,av = 3.9449E+00 1.8749E+00 aa,ab = 1.6841E-02 -3.8752E-04

Virial coefficients from NACL1

b0,b0l = 7.5374E-02 7.3770E-04 b0s,b0j = 2.9532E-01 -1.3344E-05 b0v,b0a,b0b = 1.2343E-05 -3.1844E-07 -5.4612E-09

b1,b1l = 2.7703E-01 9.0102E-04 b1s,b1j = 5.4567E-01 -8.4653E-06 b1v,b1a,b1b = 0.0000E+00 0.0000E+00 0.0000E+00

cp,cp1 = 1.4074E-03 -1.1004E-04 cps,cpj = -3.1401E-02 1.4694E-06 cpv,cpa,cpb = -1.2938E-06 4.4292E-08 0.0000E+00

Reference state properties for NaCl from NACL1

g20,s20 = -1.3886E+01 1.3886E+01 h20,cp20 = 3.4132E-06 -1.0077E+01 v20,vt20,vp20 = 1.6679E+01 7.7179E-02 4.3351E-03

Excess properties from EXCESS

gex,glex,g2ex = -7.1210E-01 1.2732E-01 -8.3942E-01 hex,hlex,h2ex = -4.0604E-02 4.7746E-03 -3.0563E-01
sex,slex,s2ex = 6.7149E-01 -1.2254E-01 5.3379E-01 cpex,cplex,cp2ex = 5.2808E+00 -3.2115E+00 8.4923E+00
vex,vlex,v2ex = 1.8117E+00 -1.2170E-01 -5.3030E-01 vtex,vtlex,vt2ex = -3.6250E-03 1.7420E-02 -4.3175E-02
vpex,vplex,vp2ex = -5.2538E-04 1.8269E-04 -1.9883E-04

Partial molal quantities from NACL2

g1,g2 = -2.2078E+01 -1.4725E+01 s1,s2 = 8.3265E+00 1.4420E+01
h1,h2 = -1.3751E+01 -3.0563E-01 cp1,cp2 = 5.8531E+00 -1.5849E+00
v1,v2 = 1.7947E+01 1.6149E+01 vt1,vt2 = 2.2107E-02 3.4004E-02 vp1,vp2 = -6.3434E-04 4.1362E-03

Osmotic coefficient and mean ion activity coefficient from NACL2.

phi,gam = 9.3634E-01 6.5724E-01

Properties of solid NaCl from NACL3

g2s,h2s = -8.6782E+00 6.6410E-04 s2s,cp2s = 8.6789E+00 6.1104E+00
v2s,vt2s,vp2s = 2.6993E+01 3.2347E-03 0.0000E+00

Change of free energy and other thermodynamic functions accompanying dissolution of NaCl in water from NACL3

dgs0,dss0 = -3.6415E+00 5.2071E+00 dhs0,dcps0 = 1.5656E+00 -1.6188E+01
dvs0,dvts0,dvps0 = -1.0314E+01 7.3944E-02 4.3351E-03

Table 2. The following quantities (class 1) are tabulated in report LBL-15512 and are presented here in the same order.

t,tc = 298.15 25.00 p = 1.000 m = 1.000

table 1	g10,g20,ap,b0,b1,1e3*cp/2	-22.1689	-13.8860	0.3915	0.0754	0.2770	0.70
table 2	s10,s20,as,b0s,b1s,1e3*cps/2	8.4130	13.8860	2.3664	0.2953	0.5457	-15.70
table 3	h10,h20,ah,1e3*b0l,1e3*b1l,1e3*cp/2	-13.7560	0.000	0.801	0.738	0.901	-0.055
table 4	cp10,cp20,aj,1e6*b0j,1e6*b1j,1e6*cpj/2	9.065	-10.077	3.945	-13.344	-8.465	0.735
table 5	mw*v10,v20,av,1e6*b0v,1e6*cpv/2	18.068	18.679	1.875	12.343	-0.647	
table 6	1e3*mw*v10*a10,vt20,aa,1e6*b0a,1e9*cpa/2	4.687	0.077	0.017	-0.318	22.146	
table 7	-1e3*mw*v10*b10,1e3*vp20,1e3*ab,1e9*b0b	-0.817	4.335	-0.388	-5.461		
table 8	gam	0.657					
table 9	phi	0.936					
table 10	dss0	5.207					
table 11	dhs0	1.566					
table 12	sex/m	0.671					
table 13	hex/m	-0.041					
table 14	cpex/m	5.281					
table 15	density	1.03623					
table 16	specific entropy(j/g-k)	3.798					
table 17	specific enthalpy(j/g)	-1788.4					

NACL calculates the properties of NaCl solutions using the accurate empirical model published by Pitzer, Peiper, and Busey, J. Phys. Chem. Ref. Data v. 13, pp. 1-102 (1984). The valid range is 0-300 C, 0-6 molal, Psat(T)-1000 bar.

Table 2. The following quantities (class 1) are tabulated in report LBL-15512 and are presented here in the same order.

t,tc = 573.15 300.00 p = 85.832 m = 6.000

table 1	g10,g20,ap,b0,b1,1e3*cp/2	-17.1382	-11.3597	0.9595	0.0768	0.5192	-1.95
table 2	s10,s20,as,b0s,b1s,1e3*cps/2	14.6673	-5.1276	17.8896	0.0577	1.1376	2.83
table 3	h10,h20,ah,1e3*b0l,1e3*b1l,1e3*cp/2	-2.4710	-16.487	14.052	-0.033	1.079	0.008
table 4	cp10,cp20,aj,1e6*b0j,1e6*b1j,1e6*cpj/2	12.449	-197.332	183.418	8.273	4.828	0.072
table 5	mw*v10,v20,av,1e6*b0v,1e6*cpv/2	25.288	-95.681	98.730	-51.668	1.583	
table 6	1e3*mw*v10*a10,vt20,aa,1e6*b0a,1e9*cpa/2	82.760	-2.882	2.715	-1.565	31.847	
table 7	-1e3*mw*v10*b10,1e3*vp20,1e3*ab,1e9*b0b	-8.120	261.187	-217.969	91.188		
table 8	gam	0.164					
table 9	phi	0.747					
table 10	dss0	-17.961					
table 11	dhs0	-18.788					
table 12	sex/m	18.455					
table 13	hex/m	15.348					
table 14	cpex/m	173.723					
table 15	density	0.99187					
table 16	specific entropy (j/g-k)	5.446					
table 17	specific enthalpy (j/g)	-508.0					