

ANL/NDM--96

DE87 002518

ANL/NDM-96

RATIO OF THE PROMPT-FISSION-NEUTRON SPECTRUM OF PLUTONIUM 239
TO THAT OF URANIUM 235*

by

Masayoshi Sugimoto[†], Alan B. Smith and Peter T. Guenther

September 1986

Keywords:

Measured ratio of ^{239}Pu prompt-fission-neutron spectrum to that of ^{235}U , 1-10 MeV.

[†]Permanent address: Japan Atomic Energy Research Institute,
Tokai-Ibaraki, Japan.

* This work supported by the U. S. Department of Energy, Nuclear Energy
Programs under contract W-31-109-Eng-38.

Applied Physics Division
Argonne National Laboratory
9700 S. Cass Avenue
Argonne, Illinois 60439
U.S.A.

MASTER

mtR
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Argonne National Laboratory, with facilities in the states of Illinois and Idaho, is owned by the United States government, and operated by The University of Chicago under the provisions of a contract with the Department of Energy.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NUCLEAR DATA AND MEASUREMENTS SERIES

The Nuclear Data and Measurements Series presents results of studies in the field of microscopic nuclear data. The primary objective is the dissemination of information in the comprehensive form required for nuclear technology applications. This Series is devoted to: a) measured microscopic nuclear parameters, b) experimental techniques and facilities employed in measurements, c) the analysis, correlation and interpretation of nuclear data, and d) the evaluation of nuclear data. Contributions to this Series are reviewed to assure technical competence and, unless otherwise stated, the contents can be formally referenced. This Series does not supplant formal journal publication but it does provide the more extensive information required for technological applications (e.g., tabulated numerical data) in a timely manner.

Copies on microfiche can be obtained by contacting:

National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22161
U.S.A.

INFORMATION ABOUT OTHER ISSUES IN THE ANL/NDM SERIES

A list of titles and authors for reports ANL/NDM-1 through ANL/NDM-50 can be obtained by referring to any report of this Series numbered ANL/NDM-51 through ANL/NDM-76. Requests for a complete list of titles or for copies of previous reports should be directed to:

Section Secretary
Applied Nuclear Physics Section
Applied Physics Division
Building 316
Argonne National Laboratory
9700 South Cass Avenue
Argonne, Illinois 60439
U.S.A.

ANL/NDM-51 A. Smith, P. Guenther, D. Smith and J. Whalen, *Measured and Evaluated Cross Sections of Elemental Bismuth*, April 1980.

ANL/NDM-52 P. Guenther, A. Smith and J. Whalen, *Neutron Total and Scattering Cross Sections of ^6Li in the Few-MeV Region*, February 1980.

ANL/NDM-53 James W. Meadows and Donald L. Smith, *Neutron-source Investigations in Support of the Cross-section Program at the Argonne Fast-neutron Generator*, May 1980.

ANL/NDM-54 A.B. Smith, P.T. Guenther and J.F. Whalen, *The Nonelastic-scattering Cross Sections of Elemental Nickel*, June 1980.

ANL/NDM-55 M.M. Bretscher and D.L. Smith, *Thermal-neutron Calibration of a Tritium Extraction Facility using the $^6\text{Li}(n,t)^4\text{He}/\text{Au}(n,\gamma)^{198}\text{Au}$ Cross-section Ratio for Standardization*, August 1980.

ANL/NDM-56 P.T. Guenther, A.B. Smith and J.F. Whalen, *Fast-neutron Interaction with ^{182}W , ^{184}W and ^{186}W* , December 1980.

ANL/NDM-57 Peter T. Guenther, Alan B. Smith and James F. Whalen, *The Total, Elastic- and Inelastic-scattering Fast-neutron Cross Sections of Natural Chromium*, January 1981.

ANL/NDM-58 W.P. Poenitz, *Review of Measurement Techniques for the Neutron-capture Process*, August 1981.

ANL/NDM-59 Wolfgang P. Poenitz, *Review of the Importance of the Neutron-capture Process in Fission Reactors*, July 1981.

ANL/NDM-60 James W. Meadows and Donald L. Smith, *Gamma-ray Detector Calibration Methods Utilized in the Argonne FNG Group Activation Cross-section Measurement Program*, April 1984.

ANL/NDM-61 Carl Budtz-Joergensen, Peter T. Guenther, Alan B. Smith and James F. Whalen, *Fast-neutron Total and Scattering Cross Sections of ^{58}Ni* , September 1981.

ANL/NDM-62 Donald L. Smith, *Covariance Matrices and Applications to the Field of Nuclear Data*, November 1981.

ANL/NDM-63 Alan B. Smith and Peter T. Guenther, *On Neutron Inelastic-scattering Cross Sections of ^{232}Th , ^{233}U , ^{235}U , ^{238}U , ^{239}U and ^{239}Pu and ^{240}Pu* , January 1982.

ANL/NDM-64 James W. Meadows and Carl Budtz-Joergensen. *The Fission-fragment Angular Distributions and Total Kinetic Energies for $^{235}U(n,f)$ from 0.18 to 8.83 MeV.* January 1982.

ANL/NDM-65 Alan B. Smith and Peter T. Guenther. *Note on the Elastic Scattering of Several-MeV Neutrons from Elemental Calcium.* March 1982.

ANL/NDM-66 Alan B. Smith and Peter T. Guenther. *Fast-neutron Scattering Cross Sections of Elemental Silver.* May 1982.

ANL/NDM-67 Donald L. Smith. *Non-evaluation Applications for Covariance Matrices.* July 1982.

ANL/NDM-68 Alan B. Smith, Peter T. Guenther and James F. Whalen. *Fast-neutron Total and Scattering Cross Sections of ^{103}Rh .* July 1982.

ANL/NDM-69 Alan B. Smith and Peter T. Guenther. *Fast-neutron Scattering Cross Sections of Elemental Zirconium.* December 1982.

ANL/NDM-70 Alan B. Smith, Peter T. Guenther and James F. Whalen. *Fast-neutron Total and Scattering Cross Sections of Niobium.* July 1982.

ANL/NDM-71 Alan B. Smith, Peter T. Guenther and James F. Whalen. *Fast-neutron Total and Scattering Cross Sections of Elemental Palladium.* June 1982.

ANL/NDM-72 Alan B. Smith and Peter T. Guenther. *Fast-neutron Scattering from Elemental Cadmium.* July 1982.

ANL/NDM-73 C. Budtz-Joergensen, Peter T. Guenther and Alan B. Smith, *Fast-neutron Elastic-scattering Cross Sections of Elemental Tin*, July 1982.

ANL/NDM-74 Wolfgang Poenitz, Alan B. Smith and Robert Howerton, *Evaluation of the ^{238}U Neutron Total Cross Section*, December 1982.

ANL/NDM-75 A.B. Smith, P.T. Guenther and J.F. Whalen, *Neutron Total and Scattering Cross Sections of Elemental Antimony*, September 1982.

ANL/NDM-76 Alan B. Smith and Peter T. Guenther, *Scattering of Fast Neutrons from Elemental Molybdenum*, November 1982.

ANL/NDM-77 Donald L. Smith, *A Least-squares Method for Deriving Reaction Differential Cross Section Information from Measurements Performed in Diverse Neutron Fields*, November 1982.

ANL/NDM-78 A.B. Smith, P.T. Guenther and J.F. Whalen, *Fast-neutron Total and Elastic-scattering Cross Sections of Elemental Indium*, November 1982.

ANL/NDM-79 C. Budtz-Joergensen, P. Guenther, A. Smith and J. Whalen, *Few-MeV Neutrons Incident on Yttrium*, June 1983.

ANL/NDM-80 W.P. Poenitz and J.F. Whalen, *Neutron Total Cross Section Measurements in the Energy Region from 47 keV to 20 MeV*, July 1983.

ANL/NDM-81 D.L. Smith and P.T. Guenther, *Covariances for Neutron Cross Sections Calculated Using a Regional Model Based on Elemental-model Fits to Experimental Data*, November 1983.

ANL/NDM-82 D.L. Smith, *Reaction Differential Cross Sections from the Least-squares Unfolding of Ratio Data Measured in Diverse Neutrons Fields*, January 1984.

ANL/NDM-83 J.W. Meadows, *The Fission Cross Sections of Some Thorium, Uranium, Neptunium and Plutonium Isotopes Relative to ^{235}U* , October 1983.

ANL/NDM-84 W.P. Poenitz and J.W. Meadows, *^{235}U and ^{239}Pu Sample-mass Determinations and Intercomparisons*, November 1983.

ANL/NDM-85 D.L. Smith, J.W. Meadows and I. Kanno, *Measurement of the $^{51}V(n,p)$ ^{51}Ti Reaction Cross Section from Threshold to 9.3 MeV*, June 1984.

ANL/NDM-86 I. Kanno, J.W. Meadows and D.L. Smith, *Energy-differential Cross-section Measurement for the $^{51}V(n,\alpha)$ ^{48}Sc Reaction*, July 1984.

ANL/NDM-87 D.L. Smith, J.W. Meadows, M.M. Bretscher and S.A. Cox, *Cross-section Measurement for the $^7Li(n,n't)$ 4He Reaction at 14.74 MeV*, September 1984.

ANL/NDM-88 A.B. Smith, D.L. Smith and R.J. Howerton, *An Evaluated Nuclear Data File for Niobium*, March 1985.

ANL/NDM-89 Bernard P. Evain, Donald L. Smith and Paul Lucchese, *Compilation and Evaluation of 14-MeV Neutron-activation Cross Sections for Nuclear Technology Applications: Set I*, April 1985.

ANL/NDM-90 D.L. Smith, J.W. Meadows and P.T. Guenther, *Fast-neutron-spectrum Measurements for the Thick-target $^9Be(d,n)$ ^{10}B Reaction at $E_d = 7$ MeV*, April 1985.

ANL/NDM-91 A.B. Smith, P.T. Guenther and R.D. Lawson, *On the Energy Dependence of the Optical Model of Neutron Scattering from Niobium*, May 1985.

ANL/NDM-92 Donald L. Smith, *Nuclear Data Uncertainties (Vol.-I): Basic Concepts of Probability*, April 1986.

ANL/NDM-93 D.L. Smith, J.W. Meadows and M.M. Bretscher, *Integral Cross-section Measurements for $^7\text{Li}(n,n't)^4\text{He}$, $^{27}\text{Al}(n,p)^{27}\text{Mg}$, $^{27}\text{Al}(n,\alpha)^{24}\text{Na}$, $^{58}\text{Ni}(n,p)^{58}\text{Co}$ and $^{60}\text{Ni}(n,p)^{60}\text{Co}$ Relative to ^{238}U Neutron Fission in the Thick-target $^9\text{Be}(d,n)^{10}\text{B}$ Spectrum at $E_d = 7$ MeV*, October 1985.

ANL/NDM-94 A.B. Smith, D.L. Smith, P. Rousset, R.D. Lawson and R.J. Howerton, *Evaluated Neutronic Data File for Yttrium*, January 1986.

ANL/NDM-95 Donald L. Smith and James W. Meadows, *A Facility for High-intensity Neutron Irradiations Using Thick-target Sources at the Argonne Fast-neutron Generator*, May 1986.

TABLE OF CONTENTS

ABSTRACT	1
I. INTRODUCTION	2
II. EXPERIMENTAL METHOD.	3
III. ANALYSIS	4
IV. RESULTS.	5
V. CONCLUSIONS	6
APPENDIX	7
VI. REFERENCES.	9
TABLES (2).	10
FIGURES (2)	12

RATIO OF THE PROMPT-FISSION-NEUTRON SPECTRUM OF
PLUTONIUM 239 TO THAT OF URANIUM 235*

by

Masayoshi Sugimoto⁺, Alan B. Smith and Peter T. Guenther

Applied Physics Division
Argonne National Laboratory

ABSTRACT

The prompt-fission-neutron spectrum resulting from ²³⁹Pu fission induced by 0.55 MeV incident neutrons is measured from 1.0 to 10.0 MeV relative to that of ²³⁵U fission induced by the same incident-energy neutrons. The measurements employ the time-of-flight technique. Energy-dependent ratios of the two spectra are deduced from the measured values over the energy range 1.0 to 10.0 MeV. The experimentally-derived ratio results are compared with those calculated from ENDF/B-V, revision-2, and with results of recent microscopic measurements. Using the ENDF/B-V ²³⁵U Watt parameters for the ²³⁵U spectrum, the experimental measurements imply a ratio of average fission-spectrum energies of ²³⁹Pu/²³⁵U = 1.045 ± 0.003, compared to the value 1.046 calculated from ENDF/B-V, revision 2.

⁺Permanent address: Japan Atomic Energy Research Institute,
Tokai-Ibaraki, Japan.

*Work supported by the U. S. Department of Energy, Nuclear Energy Programs under contract W-31-109-Eng-38.

I. INTRODUCTION

Precise knowledge of the prompt-fission-neutron spectrum of ^{239}Pu is important in fast-reactor neutronic studies as it is an essential property of the neutron-source term. Therefore, the process has been extensively investigated using a variety of experimental methods (1). A convenient index for the comparison of ^{239}Pu fission spectra is the average fission-neutron energy, \bar{E} ; or the ratio, R , of \bar{E} to that of ^{235}U assumed as a standard, where

$$R = \bar{E} (^{239}\text{Pu}) / \bar{E} (^{235}\text{U}). \quad (1)$$

The ratios R obtained from the older microscopic measurements scatter about the value 1.08 compared to 1.03 - 1.04 suggested from integral measurements (2). More recent microscopic measurements, using fission-inducing neutron energies of either thermal or ≈ 0.55 MeV, generally tend toward R values close to those deduced from integral studies (3,4,5). Sophisticated methods have been developed for the calculation of prompt-fission-neutron spectra (6) and these have been used to calculate the spectrum given in the ^{239}Pu ENDF/B-V, revision-2, evaluation (7).

Recent integral measurements at Lawrence Livermore National Laboratory using ^{239}Pu and ^{235}U spheres having radii of 3.22 cm and 3.145 cm, respectively, provide a new and very detailed integral basis for the comparative assessment of evaluated ^{239}Pu and ^{235}U prompt-fission neutron spectra, as well as of integral calculational methods (8). The measurements involved incident, fission-inducing, neutron energies of 0.3 - 2.7 MeV and fission-neutron-emission energies of 3.0 - 10.0 MeV. The interpretation of these integral measurements has raised some questions and their quality permits new precision in integral interpretation. Therefore the present microscopic measurements were initiated in an effort to test the results of the integral measurements and to provide an improved basis for their interpretation. This study is a direct measurement of the ratio, R , of the ^{239}Pu to ^{235}U fission-neutron spectra resulting from fission induced by ≈ 0.55 MeV incident-energy neutrons, over a spectrum-energy range of 1.0 - 10.0 MeV. The experimental results are compared with those calculated from ENDF/B-V (revision-2) and with the results of other microscopic measurements. The ratio determination does not require any knowledge of absolute detector efficiency, and the experimental results require no assumption as to the shape of the ^{239}Pu fission-spectrum.

II. EXPERIMENTAL METHODS

The prompt-fission-neutron spectra were measured using a 10-angle time-of-flight system (9) long employed in fast-neutron spectral measurements at Argonne. A pulsed-proton beam with a burst duration of approximately 1 nsec and a repetition rate of 2 MHz was incident on a metallic lithium target. The proton energy and the lithium target thickness were adjusted so that neutrons produced by the $^7\text{Li}(\text{p},\text{n})^7\text{Be}$ reaction (10) had an energy of 0.55 ± 0.05 MeV at the zero-degree reaction angle. The uranium and plutonium samples were both solid metal cylinders of identical size, 2 cm in diameter and 2 cm long. The uranium sample was enriched to 94.6% in ^{235}U , with the remainder distributed among the other uranium isotopes. The plutonium sample was 95.5% ^{239}Pu , and the remainder ^{240}Pu . Chemical impurities in the samples were negligible. In these experiments we ignore the approximately 5% abundance of the minor isotopes in either sample. This is a valid assumption as the minor isotopes have negligible fission cross sections at the incident-neutron energies of the present experiment, and their abundance is small and essentially equal in each sample. The uranium sample was not clad and the plutonium sample was enclosed in a 0.1 mm thick welded-steel can. Throughout the measurements an identical empty steel can was used for background determinations. The samples were placed at a zero-degree reaction angle, 15 cm from the neutron source, at the focus of the ten flight paths. Neutrons emitted from the samples, and proceeding down the flight paths, were detected by NE213 liquid scintillators 12.7 cm in diameter and approximately 2 cm thick. The low-energy-sensitivity cutoff of these detectors was set at approximately 800 keV so that the maximum energy range of detection was obtained while at the same time certainly avoiding neutrons elastically scattered from the samples. Pulse-shape-sensitive circuitry discriminated against unwanted gamma-ray backgrounds. The flight paths varied in length from 4.976 - 5.077 m and were measured to $\pm .5$ cm, including the mean detector thickness. The detectors were approximately uniformly distributed over the angular range 25 - 150 degrees. An eleventh and independent time-of-flight channel and detector, arranged to directly observe the source, served as a monitor and also controlled dead-time effects. The monitor response was reasonably consistent with the integrated proton-beam current, given the inevitable variations in the lithium target.

The eleven measured flight times between the neutron burst and the neutron detection were multiplexed into a computer and sorted into spectra of 500 channels, each with a channel width of approximately 1 nsec. The time scale was determined using a set of calibrated delay lines, and the time zero established by the observation of the prompt-fission gamma-ray with the detector gamma-suppression circuitry turned off. Conversion of time-to-energy scale throughout the measurements was relativistically. The experiments were carried out in cyclic sets,

rotating uranium, no-sample, plutonium and empty-can measurements for approximately an hour each through many cycles. The results obtained for the separate cycles were consistent. A set was continued until there were at least a 3×10^3 events per channel at the spectrum maximum, and 2×10^2 at a channel corresponding to approximately 10 MeV neutrons. This procedure properly subtracted the ambient room background and the background due to the neutron source. However, there is a small additional background due to the presence of a fissile sample. Tests showed that this was essentially time uncorrelated. Therefore it was subtracted by linearly extrapolating from low energies, well below the detector bias point, to regions corresponding to neutron energies of 15 MeV or above (e.g., extrapolations between regions where there are no significant fission-spectrum contributions). This extrapolation also would tend to correct for non-linear dead-time effects (11) but these were negligible in the present experiments. The background treatment is further described in the Appendix.

III. ANALYSIS

First the measured time spectra were integrated from 1 to 10 MeV and then normalized to make that integral unity. The normalized spectrum of ^{239}Pu was divided by that of ^{235}U channel-by-channel to obtain

$$R_j = \frac{F_{1j} - B_{1j}}{F_{2j} - B_{2j}} \times \frac{I_2}{I_1} \quad (2)$$

where j is the channel range of the spectrum extending from 1 to 10 MeV. F_{kj} and B_{kj} are respectively foreground and background in channel j , and

$$I_k = \sum_{j=\min}^{\max} (F_{kj} - B_{kj}) \quad (3)$$

If correlations between uncertainties in F_{kj} , B_{kj} and I_k are negligible, the uncertainty in R_j is

$$\delta R_j = R_j \sqrt{\left(\frac{\delta F_{1j}^2 + \delta B_{1j}^2}{(F_{1j} - B_{1j})^2} + \frac{\delta F_{2j}^2 + \delta B_{2j}^2}{(F_{2j} - B_{2j})^2} + \frac{\delta I_1^2}{I_1^2} + \frac{\delta I_2^2}{I_2^2} \right)}^{1/2} \quad (4)$$

where δX_i represent the uncertainties in quantities X_i . The resulting R_j values have relatively large uncertainties near the 1 and 10 MeV energy extremes due to low counting statistics. The energy dependence of R was described by a power-series expansion of the form

$$R(E) = \sum_{n=0}^m a_n E^n. \quad (5)$$

Least-square fits were carried out with m increasing from 1 to 3 (i.e., linear to cubic fits). The cubic fit ($m = 3$) was abandoned as nonsignificant since the uncertainty in the a_3 coefficient exceeded its value for the results obtained with each of the ten detectors. An example of both linear and quadratic fits to $R(E)$ obtained for one of the detectors is shown in Fig. 1. The quadratic fit ($m = 3$) gave a better description of the measured values below approximately 5 MeV, an important region for the calculation of the average fission-spectrum energy. However, it systematically deviated from the measured values above about 6 MeV. Typical chi-square values for both orders of fit were in the range 1.1 - 1.3. The final expression for $R(E)$ was the weighted average of the parameters a_0 , a_1 and a_2 from the $R(E)$ least-square fits to results from the ten independent detectors. The resulting values, their standard deviations and their correlations are shown in Table 1. With these parameters, assuming the Watt ^{235}U fission-neutron spectrum (ENDF/B-V, revision-2), the average energy of the ^{239}Pu fission neutron spectrum over the 1-10 MeV energy range can be calculated as discussed in the following section.

Multiple-scattering of the fission neutrons in the fissile samples softens the observed fission spectra. However, in the present measurements the effect is small due to the small size of the samples, and the very similar inelastic-scattering cross sections, leading to appreciable energy transfer, of ^{235}U and ^{239}Pu (12). These similar and small effects are essentially completely cancelled in the ratio interpretation used in this study. This conclusion is supported by Monte-Carlo calculations carried out in a similar context (12).

IV. RESULTS

In Fig. 2 the ratio of the ^{239}Pu fission-neutron spectrum relative to that of ^{235}U is compared with the values obtained from ENDF/B-V (revision-2). Generally, the agreement is good, though the experimentally derived values imply a harder ^{239}Pu spectrum. The average energy between E_{min} and E_{max} can be calculated from

$$\bar{E} = \frac{\int_{E_{\min}}^{E_{\max}} N(E) E dE}{\int_{E_{\min}}^{E_{\max}} N(E) dE}, \quad (6)$$

where $N(E)$ is the fission-neutron spectrum, with an uncertainty in the quadratic case of

$$\delta\bar{E} = G^+ M G \quad (7)$$

where $G^+ = \left(\frac{d\bar{E}}{da_0}, \frac{d\bar{E}}{da_1}, \frac{d\bar{E}}{da_2} \right)^+$, and M = the covariance matrixes of a_0 , a_1 and a_2 , and where

$$\begin{aligned} M_{ii} &= (\delta a_i)^2, & i = 0, 1, 2 \\ M_{ij} &= \delta a_i \delta a_j \quad \text{Corr}(a_i, a_j) \end{aligned} \quad (8)$$

with $i = 1, 3$
 $j = 1, 3$
and $i \neq j$.

The respective values are given in Table 1. When $N(E)$ for ^{235}U is assumed (e.g. from ENDF/B-V which has the Watt form), then $N(E)$ for ^{239}Pu follows from the experimentally-derived parameters and the $N(E)$ of ^{235}U . The resulting \bar{E} values for the experimental energy range of 1 - 10 MeV are given in Table 2, together with the comparable quantities derived from ENDF/B-V (revision-2). The table also gives the same values derived by extrapolating over the energy range 0 to 20 MeV, as well as some values deduced from recent experimental results. The agreement with the results of the present measurements and the evaluation is very good.

V. CONCLUSIONS

The energy dependence of the prompt-fission-neutron spectrum of ^{239}Pu

(induced by ≈ 550 keV incident neutrons), relative to that of ^{235}U , was determined over the emitted-neutron energy range of 1 to 10 MeV with uncertainties of less than 4%. The energy dependence of this ratio is reasonably described by either linear or quadratic energy dependences. With either description, assuming the ENDF/B-V ^{235}U fission spectrum, the average fission-neutron energy of ^{239}Pu is defined to 0.15% accuracy. Our result is in excellent agreement with that predicted from ENDF/B-V (revision-2) and with values deduced from recent measurements. The agreement with ENDF/B-V is marginally better when the experimentally deduced results are extrapolated over the full 0 - 20 MeV energy range. There are small differences between results calculated with ENDF/B revision-1 and revision-2. The present experimental results tend to support the latter. The recent integral measurements (8) span a somewhat different and wider incident-neutron energy range (0.3 - 2.7 MeV) than the present work and imply a slightly harder fission-neutron spectrum of ^{239}Pu (relative to that of ^{235}U) than indicated by the present work, or the ENDF/B-V evaluation, over approximately the same emission-energy range. The difference is beyond the respective uncertainties. We also examined the fission spectra of ^{239}Pu and ^{235}U given by ENDF/B-V (revision-2) for a number of values of the incident energy between 10^{-5} eV and 2.0 MeV and found that ^{239}Pu is always significantly harder than that of ^{235}U by similar amounts. Thus the variation in the incident energy in the integral experiments is probably not an important factor in their interpretation.

The ^{239}Pu average fission-neutron energy derived from the present work depends upon the assumed ^{235}U spectrum but the ratio values can be very useful for precise testing of relative fission-neutron energies. The present test suggests that the values given in ENDF/B (revision-2) are correct to remarkably high accuracy and suitable for the analysis of integral benchmark measurements.

Appendix

The linear backgrounds of the time-of-flight spectra were determined by least-square fitting the spectral regions before and after the fission-neutron spectra (i.e., below 0.6 MeV and above 15 MeV). The fitting function was

$$B(j) = c + d(j - J_0) \quad (\text{A-1})$$

$$\text{where } J_0 = \frac{1}{M} \left(\frac{M_2}{M_1} + \frac{M_4}{M_3} \right), \quad M = M_2 - M_1 + M_4 - M_3 + 2, \quad \text{and } (M_1, M_2)$$

and (M_3, M_4) are the respective low- and high-energy regions selected for the background determination. The uncertainty in the background determination is given by

$$\delta B(j) = ((\delta c)^2 + (\delta d)^2 (j - J_0)^2)^{1/2} \quad (A-2)$$

and the correlation between δc and δd is proportional to

$$\sum_j w_j (j - J_0) \approx \bar{w} \sum_j (j - J_0) = 0 \quad (A-3)$$

when all weights, w_j , are not very different from their average, \bar{w} . The correlations calculated from the fitted results were always close to zero. In this case the uncertainties in the fitted parameters are simply expressed as

$$\delta c \approx (\sum_j w_j (j - J_0)^2)^{-1/2} \quad (A-4)$$

and

$$\delta d \approx (\sum_j w_j)^{-1/2} .$$

References

1. See for example, Proc. Consultant's Meeting on Prompt Fission Neutron Spectra, IAEA Press, Vienna (1971).
2. A. Smith, P. Guenther, G. Winkler and R. McKnight, Nucl. Sci. and Eng. 76 357 (1980).
3. P. Johansson, B. Holmqvist, T. Wielding and L. Jeki, Proc. Conf. Nucl. Cross Sections and Tech., NBS-425, p-572 (1975).
4. J. M. Adams, Proc. Specialist's Meeting on Inelastic Scattering and Fission-Neutron Spectra, Harwell Report AERE-R-8636 (1977).
5. L. Andreichuck, B. Basova, V. Kovostylov, V. Nefedov and D. Ryazarov, Sov. Atom. Energy 92 23 (1977).
6. D. Madland and J. Nix, Nucl. Sci. and Eng. 81 213 (1982).
7. E. Arthur, P. Young, D. Madland and R. MacFarlane, Nucl. Sci. and Eng. 88 56 (1986).
8. R. Haight, J. Anderson, J. Davis, S. Grimes, R. Howerton, T. Komoto, C. Logan, J. McClure, B. Pohl and C. Wong, Lawrence Livermore National Laboratory Report UCRL-93591 (1985).
9. A. Smith, P. Guenther, R. Larsen, C. Nelson, P. Walker and J. Whalen Nucl. Instr. and Methods, 50 277 (1967).
10. See for example, Proc. Advisory Group Meeting on Neutron Sources, Leningrad, June 1986, IAEA Press, Vienna; also Neutron Sources for Basic Physics and Applications, S. Cierjacks ed., Pergamon Press Oxford (1983).
11. W. Poenitz and T. Tamura, Proc. Nucl. Data for Sci. and Tech., Antwerp, p-465, Keidel Pub. Co. (1982).
12. A. Smith, P. Guenther and R. McKnight, Proc. Conf. Nucl. Data for Sci. and Tech., Antwerp, p-39, Reidel Pub. Co. (1982).

Table 1. Linear and Quadratic Parameters Derived from Least-Square Fits to the Energy Dependence of the Ratio of $^{239}\text{Pu}/^{235}\text{U}$ Fission-Neutron Spectra.

Parameter	Linear ($a_0 + a_1 E$)	Quadratic ($a_0 + a_1 E + a_2 E^2$)
a_0	0.902 ± 0.006	0.868 ± 0.001
a_1	0.035 ± 0.002	0.059 ± 0.008
a_2	-----	-0.004 ± 0.001
Corr(a_0, a_1)	-0.913	-0.960
Corr(a_0, a_2)	-----	0.875
Corr(a_1, a_2)	-----	-0.965

Table 2. Average Energies of the Prompt-Fission-neutron Spectra of ^{235}U and ^{239}Pu .

Source	$\bar{E}(^{235}\text{U})$ ^a	$\bar{E}^{239}\text{Pu}$	Ratio
1 - 10 MeV Energy Range			
Present (lin. fit)	(2.652) ^b	2.727 \pm .004	1.028
Present (quad. fit)	(2.652) ^b	2.715 \pm .004	1.024
ENDF/B-V (rev. 2)	2.652	2.711	1.022
0 - 20 MeV Energy Range			
Present (lin. fit)	(2.031) ^b	2.123 \pm .005	1.045
Present (quad. fit)	(2.031) ^b	2.119 \pm .006	1.043
ENDF/B-V (rev. 2)	2.031	2.125	1.046
ENDF/B-V (rev. 1)	2.031	2.112	1.039
Smith et al. (2) ^d	1.973 \pm .024	2.054 \pm .029	1.041
Johansson et al. (3)	2.028 \pm .030 ^c	2.118	1.044
Integral Benchmark			
Haight et al. (8) approx. energy range = 3 - 10 MeV	-----	-----	1.040 \pm .005

a. Average energies in MeV.
 b. ENDF/B-V values assumed.
 c. As corrected by Adams (4) and fitted with a Watt form over the energy range 0.975 - 10.439 MeV.
 d. Relative to ^{252}Cf .




Figure 1. An example of linear and quadratic fits to measured spectral ratios. Measured values have been binned for clarity.




Figure 2. Comparison of measured and evaluated ratios of the $^{239}\text{Pu}/^{235}\text{U}$ prompt-fission-neutron spectra.