

LA-UR- 00-2972

Approved for public release;  
distribution is unlimited.

*Title:* Nuclear Data for Accelerator-Transmutation of Waste

*Author(s):* M. B. Chadwick

*Submitted to:* Summary of invited talk to be given at the  
ANS/ENS 2000 Winter Meeting,  
November 12-16, 2000. Summary to be  
published in "ANS Transactions".

## Los Alamos

NATIONAL LABORATORY

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-36. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

## **DISCLAIMER**

**This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, make any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.**

## **DISCLAIMER**

**Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.**

RECEIVED

OCT 26 2000

OSTI

Paper summary for ANS/ENS 2000 Meeting, Nov. 12-16 2000

## Nuclear Data for Accelerator-Transmutation of Waste

M.B. Chadwick

University of California, Los Alamos National Laboratory,  
Theoretical Division, Los Alamos, New Mexico 87545, USA

### 1 Introduction

The development of an accelerator-transmutation of waste (ATW) capability requires a number of nuclear technology developments for the design of the spallation target and the subcritical transmuter. ATW design is guided by radiation transport code simulations, which are used to model particle transport, criticality, activation, and radiation heating and damage. This paper focuses on nuclear data needed to insure the accuracy of such simulations.

### 2 Results

New ENDF/B-VI data files have been produced for incident neutrons and protons for energies that extend up to 150 MeV (the LA150 library) [1]. By using

advanced nuclear models that account for details of nuclear structure and the quantum nature of the nuclear scattering, significant gains in simulation accuracy can be achieved below 150 MeV, where intranuclear cascade calculations become less accurate. To date, evaluations have been completed for isotopes of the following target/blanket and shielding materials: H, Li, C, N, O, Al, Si, P, Ca, Cr, Fe, Ni, Cu, Nb, W, Hg, Pb, Bi. The evaluations are based upon measured data as well as predictions from the GNASH nuclear model code, which calculates cross sections using Hauser-Feshbach, exciton and Feshbach-Kerman-Koonin preequilibrium models. Elastic scattering distributions and direct reactions are calculated from the optical model.

I describe recent developments to the MCNPX radiation transport code [2], which merges MCNP and LAHET in one code and uses the LA150 evaluated data. A number of benchmark comparisons against integral experiments are described, for thick-target neutron production, neutron transmission through macroscopic slabs, and neutron kerma coefficients. Good agreement between calculation and experiment is found. These benchmarks help validate the transport code and the evaluated data for use in accelerator-driven-system simulations of neutron production in a spallation target (n/p), radiation shielding, heating, and damage. As an illustrative example, Fig. 1 shows the calculated (solid line) thick-target neutron production from 68 MeV protons stopping in aluminum, compared with measured data. It is evident that MCNPX simulations that use our new evaluated LA150 data better agree with measurements compared to those that use intranuclear cascade LAHET-type physics (dashed-line).

For an ATW transmuter, the fundamental nuclear data for minor actinides are in many cases poorly understood. This affects our ability to predict the criticality,  $k_{eff}$ . Additionally, data uncertainties affect our ability to predict the rates of transmutation of the various radioactive species. Since most designs have, at present, focussed on transmuters that operate at fast neutron energies, the most important energy range is below 20 MeV, and primarily energies in the hundreds-of-keV to few-MeV region for the minor actinides, and lower energies for the fission products. More accurate data are desirable for the minor actinides, for the fission products, and for the coolants such as lead and bismuth [3]. Data at higher energies are needed in certain cases, such as for estimations of radiation damage, and shielding requirements.

For an ATW spallation target, accurate nuclear data are needed in order to predict the number of spallation neutrons produced per incident proton,  $n/p$ , which affects the overall economy. Data are also needed to better understand the radionuclides produced through spallation and fission nuclear reactions, for studies of activation, and materials properties. I provide an overview of the status of nuclear theory and modeling for calculations of spallation and sub-actinide fission.

This research is supported by the Department of Energy under contract W-7405-ENG-36.

## References

- [1] M. B. Chadwick, P. G. Young, S. Chiba, S. Frankle, G. M. Hale, H. G. Hughes, A. J. Koning, R. C. Little, R. E. MacFarlane, R. E. Prael, and

- L. S. Waters, "Cross section evaluations to 150 MeV for accelerator-driven systems and implementation in MCNPX," *Nucl. Sci. Eng.* **131**, 293 (1999).
- [2] H. G. Hughes, K. J. Adams, M. B. Chadwick, J. C. Comly, H. W. Egdorf, S. C. Frankle, J. S. Hendricks, R. C. Little, R. E. MacFarlane, S. G. Mashnki, R. E. Prael, A. J. Sierk, L. S. Waters, M. C. White, P. G. Young, F. X. Gallmeier, and E. C. Snow, "MCNPX for neutron-proton transport," in *Proc. of the Mathematics and Computation, Reactor Physics and Environmental Analysis in Nuclear Applications*, Madrid, Spain, September 27-30, 1999, edited by J. M. Aragonés (Senda Editorial, S. A., Madrid, Spain, 1999), p. 939.
- [3] G. Palmiotti, P. J. Finck, I. Gomes, B. Micklich, and M. Salvatores, "Uncertainty assessment for accelerator-driven systems," in *Proc. of the ANS GLOBAL '99 International Conference on future nuclear systems*, Jackson Hole, WY, August 29-September 3 (American Nuclear Society, La Grange Park, IL, 1999).
- [4] S. Meigo, H. Takada, H. Nakashima, T. Sasa, K. Tanaka, S. Shin, and S. Ono, "Measurements of neutron spectra from stopping-length targets irradiated by 68-MeV protons, 100 MeV alpha and 220 MeV carbon particles," in *Proc. of the International Conference on Nuclear Data for Science and Technology*, Trieste, Italy, May 18-24, 1997, edited by G. Reffo (Societa Italiana di Fisica, Bologna, Italy, 1997), pp. 413-415.




Figure 1: Thick target neutron production spectra for 68 MeV protons incident on aluminum [4] compared with calculations.