

LA-UR-02-5527

Approved for public release;
distribution is unlimited.

Title:

Acceleration of O+ ions in the dynamic magnetotail

Author(s):

Joachim Birn and Michelle F. Thomsen

Submitted to:

<http://lib-www.lanl.gov/cgi-bin/getfile?00852054.pdf>

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-36. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

FORM 836 (10/96)

Acceleration of O^+ ions in the dynamic magnetotail

Joachim Birn, Michelle F. Thomsen

Los Alamos National Laboratory, Los Alamos, New Mexico

Michael Hesse

NASA Goddard Space Flight Center, Greenbelt, Maryland

Short title: ACCELERATION OF O^+

Abstract. The acceleration and flux increases of energetic oxygen ions are studied on the basis of test particle orbits in the fields obtained from a MHD simulation of plasmoid formation and ejection and the collapse (dipolarization) of the inner tail. The simulated fluxes show large anisotropies and nongyrotropic effects, phase bunching, and spatially and temporally localized beams. The energy distribution of O^+ in the region of an earthward beam in the near tail becomes significantly harder, more pronounced than for protons, in qualitative agreement with observations. The simulation also shows tailward beams of energetic O^+ ions closely associated with the passage of a plasmoid, both inside the plasma sheet boundary and inside the central plasma sheet, consistent with observations in the far tail. The acceleration at the near-Earth x-type neutral line produces a narrow duskward beam of energetic O^+ in the duskward extension of the x-line, which was not found in proton test particle simulations.

1. Introduction

Large changes in energetic ion fluxes from a few tens of keV to hundreds of keV are often observed in the tail, commonly in association with substorms. Most of these observations of flux changes in the near and far tail concerned protons or could not distinguish protons from heavier ions. *Frits and Wilken* [1976] concluded from ATS-6 observations (with supporting solar wind data from IMP-7 and IMP-8) that heavy ions are energized inside the magnetosphere during substorm type injection events. *Ipatiich et al.* [1984] demonstrated substantial flux increases of 112-157 keV/e O⁺ ions in ISEE-1 observations of the tail plasma sheet, apparently correlated with activity. *Möbius et al.* [1987] used AMPTE/IRM data to investigate heavy energetic ion fluxes before and after substorm onset. They found large earthward directed plasma flows concurrent with energetic proton fluxes. The fluxes of ions of ionospheric origin, such as oxygen, were also enhanced and the spectra became considerably harder, more pronounced for oxygen than for protons.

Characteristic features of energetic heavy ions are also found in the more distant tail. ISEE-3 and, more recently, Geotail observations have led to a picture of tailward propagating plasmoids that are enveloped by a sheath of tailward streaming energetic ions, presumably mostly protons [e.g., *Scholer*, 1984]. The characteristics of heavier ions have been studied more recently. Investigating plasmoid or flux rope encounters $\approx 100 R_E$ downtail, *Lai et al.* [1994] reported high variability of the ion composition but did not find a consistent trend in the composition changes. In contrast, several

authors found tailward energetic oxygen streams (and other heavy ions) in association with substorms or pseudo breakups and the passage of plasmoids or flux ropes. The oxygen beams tend to be highly collimated in the tailward direction [Wilken *et al.*, 1995; Song *et al.*, 1998]. They can be detected in both the central plasma sheet and the plasma sheet boundary layer [Jacques *et al.*, 1994], and they are associated with tailward plasma flow and southward magnetic field components [Chrissis *et al.*, 1994; Jacques *et al.*, 1994; Wilken *et al.*, 1995; Song *et al.*, 1998].

Test particle simulations in dynamically changing fields, modeling substorm effects in the inner tail, have been successfully used to study the acceleration of energetic charged particles in the energy range from tens to hundreds of keV [Delcourt and Sazeard, 1994; Birn and Hesse, 1994; Birn *et al.*, 1997b, 1998; Li *et al.*, 1998; Saharia *et al.*, 2000]. These simulations have demonstrated that substorm related injections of energetic particles in the inner tail can be understood on the basis of the localised induced electric field associated with the collapse (dipolarisation) of the magnetic field in the inner tail during the expansive phase of substorms. Delcourt and Sazeard [1994] also studied the acceleration of O^+ ions and found dramatic spectral changes with energy increases exceeding those of protons. A combination of test particle orbit integration with a two-dimensional MHD simulation of plasmoid formation and ejection has also successfully reproduced energetic proton signatures in the more distant tail [Scholes and Jamilsky, 1987, 1989; Sachsenweber *et al.*, 1989].

Although enhanced oxygen outflow from the ionosphere presumably plays a role in the observed flux increases, additional acceleration appears necessary to achieve the

observed high energies of tens or hundreds of keV. In this paper we will extend previous test particle simulations to investigate heavier ions, specifically, O^+ . We will focus on the acceleration mechanisms and the corresponding flux increases, taking the presence of some seed population of oxygen ions in the plasma sheet as granted.

1. Background fields and numerical procedure

As in earlier test particle simulations, we use the magnetic and electric fields obtained by an MHD simulation of magnetotail dynamics [Birn and Hesse, 1996] as the basis for the orbit integration. This simulation covers the region from $z = -5 R_E$ out to $z = -65 R_E$. The dynamic breakup phase is preceded by a simulated growth phase during which an external electric field is applied at the high-latitude boundary of the simulation box. This leads not only to the energy buildup in the tail lobes but also to the formation of a thin current sheet in the near tail, characteristic for the late growth phase and presumably crucial for the onset of an instability that initiates the breakup phase.

The breakup of this current sheet is initiated by imposing finite resistivity (defining the time $t = 0$). A neutral line forms as a closed curve at $t \approx 2$ min. It subsequently expands across and down the tail, corresponding to the tailward motion of a plasmoid. The near-Earth part of the neutral line, representing the x line, remains located near $z \approx -13$ (units are in R_E). Rapid increases of temperature, flow, and electric field start at $t \approx 5$ min, leading eventually to peak values of E_y of 10-20 mV/m. Plate 1 shows the color-coded cross-tail electric field E_y from the MHD simulation for two different times.

The two figures on the left show E_y , together with magnetic field lines in the x, z plane ($y = 0$), the figures on the right show E_y in the equatorial plane ($z = 0$) together with the magnetic neutral line. The peak of the electric field is only initially located near the x line but subsequently moves earthward, while the magnitude increases. The earthward propagation stops near $x \approx -13$, where the flow from the x line is slowed down by the increasing magnetic field, leading to a pile-up of magnetic flux from approximately $x \approx -10$ tailward. It is noteworthy that the region of enhanced electric field is localised in all three space dimensions.

Using the magnetic and electric fields from the MHD simulation, we integrated the full O^+ ion orbits backward in time until they reached the boundary of the simulation box or the initial state. At the near-Earth boundary the particles were reflected back into the box, simulating mirroring closer to the Earth, but neglecting the loss cone and the flight time from the boundary to the actual mirror point and back. This boundary condition also disregards the potential new injection of ionospheric particles. The instantaneous local electric and magnetic fields were interpolated from the results of the MHD run, which were stored for time steps of 0.5 Alfvén times (corresponding to approximately 3 seconds).

Fluxes of particles at particular locations and times can be estimated by applying Liouville's theorem and imposing distribution functions at the boundaries of the simulation box and for the initial state. The conservation of the phase space density f along the particle trajectory then gives the value of the distribution function at the final

location. The imposed initial distributions are kappa distributions [Vasyliunas, 1968]

$$f = \hat{f} \left[1 + \frac{W_{i*}}{(s-1.5)W_s} \right]^{-s-1} \quad (1)$$

where W_{i*} is the initial kinetic energy and W_s is the average energy. We found that the typical source locations of accelerated O^+ ions were inside the dawn flank plasma sheet. Since observations in the tail indicate similar average energies for plasma sheet protons and oxygen ions [e.g., Frank et al., 1977; Petersen et al., 1981], we used the same temperature for both species and chose parameters typical for protons in the moderately active plasma sheet [Christer et al., 1988, 1991], consistent with the initial and boundary temperature of the MHD run

$$s \approx 5.5 \quad W_s \approx 5 \text{ keV} \quad (2)$$

A spatial variation is now contained only in the factor \hat{f} .

Through phase space mapping, the final values of the distribution function can be derived from the initial distribution. Using (1) we find

$$f = \hat{f} \left[1 + \frac{W_{f*} - \Delta W}{(s-1.5)W_s} \right]^{-s-1} \quad (3)$$

where W_{f*} is the final energy of a particle and ΔW the energy gain (or loss, for $\Delta W < 0$). If we neglect the weak variation of \hat{f} , we find that the major variation of fluxes at a given energy W_{f*} stems from the energy gain ΔW . Particles that start out with low energy are more abundant than higher-energy particles. If they gain a lot of energy, their high initial f values map to the final high-energy phase point. Therefore, even without imposing specific initial distribution functions, the energy gain can be

taken as a measure of flux enhancement when we compare fluxes at a given energy $W_{f,i}$ for different times and phase space locations. In fact, if we had chosen a Maxwellian instead of a kappa distribution, the logarithm of the phase space density, and hence of the flux, would be a linear function of the energy gain.

Ideally, one phase space trajectory can provide one f value of the final distribution function at a chosen location in space and time. This procedure works very well when the phase space mapping between the initial and final locations is a smooth function of the phase coordinates [cf., *Carron and Goeris*, 1989]. However, in a highly nonlinear system, such as the spatially and temporally variable magnetotail, chaotic effects may dominate in some regions of the phase space. That means that a slight variation in the final location or velocity can lead to vastly different initial locations and velocities and vice versa. In this case the mapped value of the distribution function can fluctuate significantly in a small neighborhood of the chosen phase space location. The actual value to be compared with observations then has to be an average of f over such a neighborhood, using a sufficient number of phase space trajectories. (This closely resembles the collection of phase space distributions from observed counts of particles.)

3. Overview

Plate 2 provides an overview of the characteristics of O^+ fluxes in the equatorial plane at $t \approx 9$ min at a typical energy of 180 keV. The four panels represent four different flux directions. Here no phase space averaging has been used. The color coding in this and subsequent figures hence can be interpreted in two ways. The color

directly represents the energy gain of an ion traced backwards from the given location. As discussed in Section 2, this energy gain is also a measure of the flux increase at the given location, since the initial phase space density is higher for smaller initial energy. The white contours represent the magnitude of the cross-tail electric field, and the black line shows the magnetic neutral line at $t = 9$ min. The length unit here and in all other simulation figures is $1 R_E$. The colored individual lines show projections of selected orbits of accelerated O^+ ions ending at the point of the arrow at $t = 9$ min. The colors for the trajectories were chosen for maximum contrast only and have no specific meaning.

Plate 2 demonstrates that regions of enhanced energetic O^+ fluxes are generated in the near tail for all four flux directions, primarily in the area earthward and duskward from the region of enhanced cross-tail electric field. However, the spatial extent of these regions differs for the different flux directions. Specifically, the third panel of Plate 1 shows a strong extended earthward beam near $z = -10$ that does not show up in the other flux directions. We will look at more of the characteristics of that beam in Section 4.

The signatures in the more distant tail are even more different for the different flux directions. The top panel of Plate 2 shows a tailward beam extending slightly duskward and tailward from the region of enhanced electric field (white contour) near midnight. The tailward end of this beam is given by the neutral line (black contour), which represents the α -line in the center of the tailward moving plasmoid. Such beams have indeed been observed in the distant tail, associated with the passage of a plasmoid

[Christon et al., 1994; Jaegley et al., 1994; Wilken et al., 1995; Song et al., 1998]. The particle trajectory ending in this region shows the typical source region, located on the down side of the near tail. Although this particle comes from the vicinity of the location of the near-Earth x-line, its acceleration takes place much farther tailward, between about 40 and 50 R_E near midnight, when it crosses the region of enhanced E , associated with fast tailward plasma flow.

Plate 2 (second panel) also shows a well focused duskward beam extending duskward from the near-Earth x line. Such a beam was not seen in proton test particle simulations [Birn et al., 1997b]. Enhanced duskward fluxes are also seen over most of the extent of the region of enhanced E , tailward of the x-line, whereas there are no increases of earthward and downward fluxes tailward of about $z = -15R_E$.

While the regions of enhanced fluxes differ considerably for the four panels of Plate 2, the trajectories of the O^+ ions that comprise these flux regions in the inner tail are quite similar, so that one might consider the different regions as belonging to different phases of a bunch of particles, accelerated in the duskward direction through the localised electric field in the inner tail, then turning subsequently earthward (third panel), downward (bottom panel), and tailward (top panel) in distinct spatial regions, as defined by the finite Larmor radius.

Plate 3 demonstrates the characteristics of the duskward fluxes in the x, z plane at $y = 3$, duskward of the acceleration region, using again the energy gain as a proxy for the fluxes, for both 180 keV protons and O^+ ions at $t = 9\text{min}$. The top panel shows proton fluxes and the two bottom panels represent the same O^+ fluxes with

different color scales. The color scale for the middle panel is chosen to be the same as for the protons, while the one for the bottom panel extends to the maximum energy gain, to emphasize the peak values. The red and orange lines represent the separatrices (boundaries between closed and open field lines) at $y = 0$ (wider contour) and $y = \delta$, respectively.

A comparison of the top two panels of Plate 3 demonstrates that in the inner tail the cross-tail electric field associated with the earthward flow provides similar acceleration for protons and oxygen ions. In the more distant tail, the electric field associated with fast tailward flows tailward of the x -line is much more effective in accelerating oxygen ions than protons. It should hence lead to a relatively more significant flux enhancement for oxygen ions. (For the absolute fluxes, however, we have to take into account that oxygen is much less abundant in the source region.) The bottom panel of Plate 3 shows more clearly the peak value of accelerated O^+ . This peak flux is highly localized, extending duskward from the near-midnight portion of the near-Earth neutral line. (The neutral line bends tailward away from midnight and does not extend to $y = \delta$.)

4. Inner tail

Now we look more closely at the properties of the earthward beam on the dusk side in the inner tail shown in Plate 2. Plate 4 shows the characteristics of 180 keV O^+ fluxes in the earthward direction in a tail cross-section at $s = -10$ for $t = 9$ min. The black lines represent the separatrices (boundaries between closed and open field lines) at this location. The region of enhanced earthward flux is centered around the equatorial plane

and occupies a significant portion of the duskside plasma sheet. Closer to midnight it extends into two layers just inside the plasma sheet boundary. The boundary here forms a bulge due to the advance of reconnection into the lobes and the corresponding expansion of the closed field line region. There are also some more chaotic occurrences of accelerated O^+ ions near the equatorial plane in this region.

Plate 5 shows the temporal variation of earthward O^+ fluxes in this region, for various energies, at $s = -10, z = 0$ and three locations in y . The fluxes are now calculated using initial kappa distributions (1) with the parameters given in (2). The absolute magnitude, however, is not calibrated. Plate 5 demonstrates that enhanced earthward fluxes occur at all energies above approximately 20 keV up to more than 300 keV. These fluxes are, however, highly variable in time.

Figure 1 provides a comparison of the energy distributions of proton and oxygen ion fluxes in this region, showing the fluxes as a function of energy at $s = -10, y = 3, z = 0$. The particle fluxes are again calculated using the initial kappa distribution (1) with the parameters given in (2). The dotted and solid lines represent omnidirectional fluxes averaged over 14 different directions before and after the injection, respectively, while the dash-dotted line gives the distribution of the earthward O^+ fluxes only. Figure 1 demonstrates that the oxygen flux enhancements are more significant than those of the energetic protons in this region, leading to a flatter (harder) distribution up to more than 100 keV. This result is also in qualitative agreement with observations [Möbius et al., 1985].

5. Outer tail

We now look at the properties of the tailward oxygen ion beam shown in the top panel of Plate 1. Plate 6 shows the spatial characteristics of 180 keV proton and O⁺ fluxes in the tailward direction at $z = -55$ and $t = 9$ min. The color-coding again represents the energy gain of ions traced backwards in time from the given location as a proxy of the flux increase. The black lines represent the separatrices (boundaries between closed and open field lines) at this location. The energetic protons (top panel) form two layers just inside the separatrix on the dusk side. The O⁺ ions also show similar layers. They are located farther inside the separatrix, because the oxygen ions, for a given energy, have lower speed along the magnetic field, which makes the inward displacement from $\mathbf{E} \times \mathbf{B}$ drift more effective. The oxygen ions, however, also show a stronger intensification around the duskward, equatorward edge of the boundary layers and enhanced fluxes near the equatorial plane closer to midnight, which are not seen for the protons.

The close temporal association between the tailward energetic O⁺ beam and the passage of the o-line is demonstrated in Figure 2, which shows v_x and B_z from the MHD simulation, together with the tailward 180 keV O⁺ flux at $z = -55$, $y = 1$, $z = 0$ as functions of time. Modest tailward plasma flow is present in the tailward part of the plasmoid where $B_z > 0$; however, the main increase occurs close to the passage of the o-line, when B_z turns southward. The increase of the O⁺ flux occurs basically simultaneously.

Plate 7 demonstrates the velocity space distributions of these beams, showing 130 keV O^+ fluxes as a function of flux direction, at $x = -55$, $y = 5$ and three locations in z for $t = 9$ min. The angle θ measures the elevation angle of the flux direction against the equatorial plane (with $\theta = 90^\circ$ for northward flux) and ϕ is the azimuth in the equatorial plane (with $\phi = 0$ corresponding to tailward flux and $\phi = 90^\circ$ corresponding to downward flux). The color-coding again represents the energy gain as a measure of the flux increase. The black crosses indicate the local magnetic field direction at each location. Plate 7 demonstrates that, at each location in z , the beams are narrowly focused in velocity space with a tailward and slightly duskward direction, independent of the magnetic field direction. At present it is not clear whether the structure (ridges) shown by the distribution functions in Plate 7 is real or an artifact of the finite simulation box together with the single energy considered. They resemble the ridges predicted by *Martin and Speiser* [1988]. However, these ridges were predicted for the vicinity of the (x-type) neutral line, whereas our structures are found far away.

Figure 3 demonstrates the source regions of the tailward beam, showing the origins of the O^+ ions contributing to the fluxes in Plate 6. The full circles represent only those orbits with an energy gain above 50 keV (that is, an initial energy below 130 keV), indicating the sources of accelerated ions contributing to the regions of enhanced flux. The source locations are widely distributed, although most come from the vicinity of the location of the near-Earth x-line ($x \approx -23$). However, as illustrated by the orbit in Plate 2, the acceleration takes place mostly not near the x-line but in the more extended region of enhanced E_y tailward from it.

6. Summary and Conclusions

We have used the electric and magnetic fields from a MHD simulation of magnetotail dynamics, modeling reconnection, plasmoid ejection, and the collapse (dipolarisation) of the near tail, to investigate the effects on heavy ions, specifically, O^+ . This is an extension of previous work, studying energetic protons and electrons as test particles in the same fields [Birn *et al.*, 1997b, 1998]. These studies have demonstrated that the electric field associated with the collapse of the inner tail is able to cause the acceleration and the typical injection characteristics of energetic protons and electrons in the inner tail [e.g., Birn *et al.*, 1997a].

Oxygen ions in the considered energy range of tens to hundreds of keV are highly nonadiabatic. Therefore, as might be expected, the simulated fluxes show large anisotropies and nongyrotropic effects, phase bunching, and spatially and temporally localised beams. Similar effects were found by Delcourt and Saze [1994]. The energy distribution of O^+ in the region of an earthward beam in the near tail becomes significantly harder, more pronounced than for protons, in qualitative agreement with observations [Möbius *et al.*, 1985].

The duskward electric field at the near-Earth neutral line has a pronounced, albeit localised, effect on O^+ , causing a narrow beam of duskward flux in the duskward extension of the near-Earth x-line. This effect was not found in the earlier investigations of protons, so that oxygen ions might provide a better indicator for the presence of the near-Earth neutral line. Enhanced duskward fluxes of energetic O^+ (at a somewhat lower

level) were also found for most of the region duskward of the region of tailward plasma flow. This tailward flow is present primarily only in the earthward part of the departing plasmoid, where $B_r < 0$, so that a corresponding effect of downward acceleration was not found in the part of the plasmoid that is tailward of the o-line. (This might change, however, at larger distances than considered here, when the plasmoid moves tailward with more uniform speed.) The energetic O⁺ flux increase on the dusk side is relatively stronger than for protons.

The test particle simulations also show tailward beams of O⁺ in the more distant tail. These beams are found inside the duskside plasma sheet boundary with a somewhat larger offset away from the boundary than for the protons. This offset can be easily understood because oxygen ions, for a given energy, have lower speed along the magnetic field, which makes the inward displacement from $E \times B$ drift more effective. We also found a tailward beam closer to midnight and closer to the equatorial plane, which was not seen in the energetic protons at the same location and time. This beam was found just earthward of the o-type magnetic neutral line in the center of the passing plasmoid, associated with negative B_r . These features are also consistent with observed energetic ion beams in the distant tail [Christon *et al.*, 1994; Jacquey *et al.*, 1994; Wilken *et al.*, 1995; Song *et al.*, 1998].

The primary source region for all investigated accelerated O⁺ ions is the dawn flank plasma sheet. However, our approach did not account for an ionospheric source, which was excluded by the mirroring condition at the near-Earth boundary of the simulation box. We found three major acceleration regions. As mentioned before,

acceleration at the near-Earth x-line itself is the cause of a narrow duskward beam in the duskward extension of the x-line. The oxygen ions in the inner tail were accelerated in the enhanced cross-tail electric fields earthward of the x-line, associated with the earthward plasma flow from the x-line and the collapse of the inner tail. This is similar to the results for protons and electrons [Birn et al., 1997b, 1998]. The energetic O⁺ ions contributing to the tailward beam in the distant tail were accelerated in the cross-tail electric fields associated with the extended tailward plasma flow tailward of the x-line. It is remarkable that the tailward flow, and hence the acceleration, in the simulation was restricted largely to the region of southward B_z , and hence duskward E_y ; we did not find a similar effect in the part of the plasmoid that is tailward of the x-line. However, this might change for larger distances than those considered in the simulation, which extended only to about 60-70 R_E down the tail.

Acknowledgments. This work was supported by the U.S. Department of Energy's Office of Basic Energy Sciences through its Geosciences Research Program, by NASA's Sun-Earth Connection Theory Program, and by the National Science Foundation through grant ATM-9901010.

References

Baker, D. N., E. W. Hones, Jr., D. T. Young, J. Birn, The possible role of ionospheric oxygen in the initiation and development of plasma sheet instabilities, *Geophys. Lett.*, **9**, 1337, 1982.

Baker, D. N., T. A. Frits, W. Lennartsson, B. Wilken, M. W. Krochl, and J. Birn, The role of heavy ionospheric ions in the localisation of substorm disturbances on March 22, 1979: CDAW-6, *J. Geophys. Res.*, **90**, 1273-1281, 1985.

Birn, J., and M. Hesse, Particle acceleration in the dynamic magnetotail: Orbits in self-consistent three-dimensional MHD fields, *J. Geophys. Res.*, **99**, 109, 1994.

Birn, J., and M. Hesse, Details of current disruption and diversion in simulations of magnetotail dynamics, *J. Geophys. Res.*, **101**, 15,345, 1996.

Birn, J., M. F. Thomson, J. E. Borovsky, G. D. Reeves, D. J. McComas, and R. D. Behan, Characteristic plasma properties of dispersionless substorm injections at geosynchronous orbit, *J. Geophys. Res.*, **101**, 2309, 1996a.

Birn, J., M. F. Thomson, J. E. Borovsky, G. D. Reeves, D. J. McComas, R. D. Behan, and M. Hesse, Substorm ion injections: Geosynchronous observations and test particle orbits in three-dimensional dynamic MHD fields, *J. Geophys. Res.*, **101**, 2325, 1996b.

Birn, J., M. F. Thomson, J. E. Borovsky, G. D. Reeves, D. J. McComas, R. D. Behan, and M. Hesse, Substorm electron injections: Geosynchronous observations and test particle simulations, *J. Geophys. Res.*, **104**, 9235, 1998.

Christon, S. P., D. G. Mitchell, D. J. Williams, C. Y. Huang, L. A. Frank, and T. E. Eastman, Energy spectra of plasma sheet ions and electrons from ~ 50 eV/c to ~ 1 MeV during plasma sheet temperature transitions, *J. Geophys. Res.*, **94**, 2862, 1988.

Christon, S. P., D. J. Williams, and D. G. Mitchell, Spectral characteristics of plasma sheet ion and electron populations during disturbed geomagnetic conditions, *J. Geophys. Lett.*, , 96, 1, 1991.

Christon, S. P., G. Gloeckler, D. J. Williams, T. Mukai, R. W. McEntire, C. Jacquay, V. Angelopoulos, A. T. Y. Lui, S. Kokubun, D. H. Fairfield, M. Hoshino, and T. Yamamoto, Energetic atomic and molecular ions of ionospheric origin observed in distant magnetotail flow reversal events, *Geophys. Res. Lett.*, , 17, 3023, 1990.

Curran, D. B., and C. K. Goertz, Particle distributions in a two-dimensional reconnection field geometry, *J. Geophys. Lett.*, , 94, 272, 1989.

Dalcourt, D. C., and J. A. Sauvage, Plasma sheet ion energisation during dipolarisation events, *J. Geophys. Lett.*, , 99, 97, 1994.

Frank, L. A., K. L. Ackerson, and D. M. Yeager, Observations of atomic oxygen (O^+) in the Earth's magnetotail, *J. Geophys. Lett.*, , 41, 129, 1977.

Fuji, T. A., and B. Wilken, Substorm generated fluxes of heavy ions at the geostationary orbit, in *Magnetospheric Particles and Fields*, edited by B. M. McCormac, p. 171, D. Reidel, Dordrecht-Holland, 1976.

Ivanich, F. M., A. B. Galvin, G. Gloeckler, D. Horvestadt, B. Klecker, and M. Scholer, Energetic (> 100 keV) O^+ ions in the plasma sheet, *Geophys. Res. Lett.*, , 11, 501, 1984.

Jacquay, C., D. J. Williams, R. W. McEntire, A. T. Y. Lui, V. Angelopoulos, S. P. Christon, S. Kokubun, T. Yamamoto, G. D. Reeves, and R. D. Behan, Tailward energetic ion streams observed at $\approx 100 R_E$ by GEOTAIL-EPIC associated with geomagnetic activity intensification, *Geophys. Res. Lett.*, , 21, 3015, 1994.

Lennartsson, W., R. D. Sharp, and R. D. Zwicky, Substorm effects on the plasma sheet ion

composition on March 22, 1979 (CDAW 6), *J. Geophys. Res.*, **90**, 1243, 1985.

Li, X., D. N. Baker, M. Temerin, and G. D. Reeves, Simulation of dispersionless injections and subsequent drift echoes of energetic electrons associated with substorms, *Geophys. Res. Lett.*, **25**, 3763, 1998.

Lui, A. T. Y., D. J. Williams, S. P. Christon, R. W. McEntire, V. Angelopoulos, C. Jacquey, T. Yamamoto, and S. Kokubun, A preliminary assessment of energetic ion species in flux ropes/plasmoids in the distant tail, *Geophys. Res. Lett.*, **21**, 3019, 1994.

Martin, R. F., Jr., and T. W. Spence, A predicted energetic ion signature of a neutral line in the geomagnetic tail, *J. Geophys. Res.*, **94**, 11,821, 1988.

Möbius, E., M. Scholer, B. Klecker, D. Horwitz, G. Gloeckler, and F. M. Ipavich, Acceleration of ions of ionospheric origin in the plasma sheet during substorm activity, in *Magnetotail Physics*, edited by A. T. Y. Lui, p. 231, The Johns Hopkins University Press, Baltimore, MD, 1987.

Peterson, W. K., R. D. Sharp, E. G. Shelley, and R. G. Johnson, Energetic ion composition of the plasma sheet, *J. Geophys. Res.*, **46**, 761, 1981.

Sachsenweiser, D., M. Scholer, and E. Möbius, Test particle acceleration in a magnetotail reconnection configuration, *Geophys. Res. Lett.*, **16**, 1027, 1989.

Sato, T., H. Matsumoto, and K. Nagai, Particle acceleration in time-developing magnetic reconnection process, *J. Geophys. Res.*, **47**, 6089, 1982.

Scholer, M., Energetic ions and electrons and their acceleration processes in the magnetotail, in *Magnetic Reconnection in Space and Laboratory Plasmas*, *Geophys. Monogr. Ser.*, vol. 30, edited by E. W. Hones, Jr., p. 216, AGU, Washington, D. C., 1984.

Scholer, M., Earthward plasma flow during near-Earth reconnection: Numerical simulations,

J. Geophys. Lett., **91**, 12,425, 1987.

Scholer, M., and F. Janitsky, Particle orbits during the development of plasmoids, *J. Geophys. Lett.*, **91**, 12,181, 1987.

Scholer, M., and F. Janitsky, Plasmoid-associated energetic ion bursts in the deep magnetotail: Numerical modeling of the boundary layer, *J. Geophys. Lett.*, **94**, 2459, 1989.

Vasyliunas, V. M., A survey of low-energy electrons in the evening sector of the magnetosphere with OGO 1 and OGO 3, *J. Geophys. Lett.*, **74**, 2839, 1968.

Wilken, B., Q. G. Zong, I. A. Daglis, T. Dole, S. Livi, K. Maesawa, L. Y. Pu, S. Ullaland, and T. Yamamoto, Tailward flowing energetic oxygen ion bursts associated with multiple flux ropes in the distant magnetotail: GEOTAIL observations, *Geophys. Lett.*, **21**, 3267, 1995.

Zehnert, S., C. L. Cheng, and J. R. Johnson, Particle transport and energization associated with substorms, *J. Geophys. Lett.*, **103**, 18,741, 2000.

Zong, Q. G., B. Wilken, I. Woch, T. Mukai, T. Yamamoto, G. D. Reeves, T. Dole, K. Maesawa, D. J. Williams, S. Kokubun, and S. Ullaland, Energetic oxygen bursts in the distant magnetotail as a product of intense substorms: Three case studies, *J. Geophys. Lett.*, **104**, 20,339, 1998.

Joachim Birn, Los Alamos National Laboratory, Los Alamos, New Mexico (e-mail: jbirn@lanl.gov)

Received _____

Figure 1. O^+ fluxes as function of energy at $x = -10, y = 3, z = 0$. The fluxes are calculated using initial kappa distributions (see text) and averaged over 14 different directions. The dash-dotted line represents earthward flux only.

Figure 2. Temporal evolution of characteristic parameters at $x = -55, y = 2, z = 0$; (top) plasma velocity component v_x , (center) magnetic field component B_x , and (bottom) 180 keV tailward O^+ fluxes. The fluxes are calculated using initial kappa distributions (see text).

Figure 3. Origins of the O^+ ions contributing to the tailward fluxes in Plate 6. The full circles represent only those orbits with an energy gain above 50 keV, indicating the sources of accelerated ions contributing to the regions of enhanced flux.

Plate 1. Color-coded cross-tail electric field E_x obtained from the MHD simulation [Birn and Hesse, 1996] at two different times, as indicated. The left-hand side figures show E_x , together with magnetic field lines, the right-hand side figures show E_x , together with the magnetic neutral line. Note that the color scale is different for the different times. The electric field unit corresponds to ~ 100 mV/m.

Plate 2. Characteristics of 180 keV O⁺ fluxes in the equatorial plane at $t = 9$ min. The color-coding can be interpreted in two ways. The color directly represents the energy gain of an ion traced backwards from the given location. Since the initial phase space density is higher for smaller initial energy, this energy gain is also a measure of the flux increase at the given location. The four panels correspond to four different flux directions. The white contours represent the magnitude of the cross-tail electric field, and the black line shows the magnetic neutral line at $t = 9$ min. The length unit here and in all other simulation figures is $1 R_{\oplus}$. The colored individual lines show projections of orbits ending at the point of the arrow at $t = 9$ min. The colors for the trajectories are chosen for maximum contrast only and have no specific meaning.

Plate 3. Characteristics of 180 keV protons and O⁺ fluxes in the duskward direction at $y = \delta$ and $t = 9$ min. The color-coding again represents the energy gain of an ion traced backwards from the given location, taken as a measure of the flux increase at the given location. The two bottom panels represent the same O⁺ fluxes shown with different color scales. The one for the middle panel is chosen to be the same as for the protons, while the one for the bottom panel extends to the maximum energy gain, to emphasize the peak values. The red and orange lines represent the separatrices (boundaries between closed and open field lines) at $y = 0$ and $y = \delta$, respectively.

Plate 4. Characteristics of 180 keV O⁺ fluxes in the earthward direction at $s = -10$ and $t = 9$ min. The color-coding again represents the energy gain of an ion traced backwards from the given location, taken as a measure of the flux increase at the given location. The black lines represent the separatrices (boundaries between closed and open field lines) at this location.

Plate 5. O⁺ fluxes in the earthward direction as function of time for various energies at $s = -10$, $z = 0$ and three locations in y . The fluxes are calculated using initial kappa distributions (see text).

Plate 6. Characteristics of 180 keV proton and O⁺ fluxes in the tailward direction at $s = -55$ and $t = 9$ min. The color-coding again represents the energy gain of ions traced backwards in time from the given location, taken as a measure of the flux increase at the given location. The black lines represent the separatrices (boundaries between closed and open field lines) at this location.

Plate 7. Phase space distribution of 180 keV O⁺ fluxes in the tailward direction at $x = -55$, $y = 5$ and three locations in z for $t = 9$ min. The color-coding again represents the energy gain of ions traced backwards in time from the given phase space location, taken as a measure of the flux. The black crosses represent the local magnetic field direction at each location.




Fig. 1




Fig. 2









Fig. 3





180 keV O⁺ flux, z = 0, t = 9 min




Duskward flux of 180 keV ions, $y = 8$, $t = 9$ min







Tailward moving 180 keV ions, $t = 9$ min, $x = -55$



180 keV O⁺, x = -55, y = 5, t = 9 min

