DECLASSIFIED

## TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

RADIATION LABORATORY



## UNIVERSITY OF CALIFORNIA Radiation Laboratory

Contract No. W-7405-eng-48

## DECLASSIFIED

Californium Isotopes from Bombardment of Uranium with Carbon Ions


A. Ghiorso, S. G. Thompson, K. Street, Jr., and G. T. Seaborg

September 6, 1950

This document contains information affecting the National Defense of the United States.

Its transmission or the disclosure of his contents in any manner to an unauthorized person is probleted and may result in severe criminal condities under applicable Federal Laws.

Berkeley, California



Standard Distribution		Copy Nos.
Argonne National Laboratory Atomic Energy Commission, Washington Brookhaven National Laboratory Carbide and Carbon Chemicals Division (K-25 Plant) Carbide and Carbon Chemicals Division (Y-12 Plant) General Electric Company, Richland Hanford Operations Office Iowa State College Kellex Corporation Knolls Atomic Power Laboratory Los Alamos Mound Laboratory Naval Radiological Defense Laboratory NEPA Project New York Operations Office Oak Ridge National Laboratory Patent Branch, Washington Technical Information Division, ORE UCLA Medical Research Laboratory (Warren) University of California Radiation Laboratory University of Rochester	•	1-10 11-12 13-16 17-18 19 20-25 26 27 28 29-32 33-35 36-38 39 40 41-42 43-48 49 50-64 65 66-68 69-70
	Total	70


Information Division Radiation Laboratory University of California Berkeley, California



## CALIFORNIUM ISOTOPES FROM BOMBARDMENT OF URANIUM WITH CARBON IONS

A. Ghiorso, S. G. Thompson, K. Street, Jr., and G. T. Seaborg Radiation Laboratory and Department of Chemistry University of California, Berkeley, California

September 6, 1950



The recent production and identification of isotopes of elements with atomic numbers up to six higher than the target element through bombardment with hexapositive 120-Mev carbon ions made it seem worthwhile to apply this technique to the transuranium region.

Accordingly, small pieces of natural uranium metal (about 0.5 mil thick and 2.5 cm by 0.6 cm area) were irradiated in the internal carbon ion beam in the Berkeley 60-inch cyclotron. Following the irradiations, the uranium was dissolved in dilute hydrochloric acid containing hydrogen peroxide and a transplutonium fraction was isolated through the use of lanthanum fluoride, and lanthanum hydroxide precipitation steps followed by the ion exchange adsorption column procedure in which concentrated hydrochloric acid is used to separate the tripositive actinide elements from the rare earth elements.<sup>2</sup>

1

The transplutonium fractions in hydrochloric acid were evaporated as weightless films on platinum plates which were placed in the ionization chamber of the 48 channel pulse analyzer apparatus in order to measure the yield and energies of any alpha-particles which might be present. In the best experiment at about one hour after the end of the 90-minute bombardment, some 50 disintegrations per minute of the distinctive 7.1-Mev alpha-particles  $^3$  of  $^{244}$  were observed to be present and to decay with the 45-minute half-life. The  $^{c}$   $^{$ 

After the decay of the alpha-particles due to Cf<sup>244</sup>, about five disintegrations per minute of alpha-particles with 6.8-Mev energy was observed and

this alpha-radioactivity decayed with a half-life of about 35 hours. A consideration of the systematics of alpha-radioactivity leads us to the view that this activity is due to the new isotope Cf<sup>246</sup> formed by the reaction U<sup>238</sup>(C<sup>12</sup>,4n). The measured half-life agrees with the expected alpha half-life for the observed energy for an even-even isotope of the element with atomic number 98.

It is not possible at this time to obtain a good estimate of the intensity of the carbon ion beam. Therefore, the cross sections for the reactions given above cannot be calculated. However, it is interesting to note that the indicated yields for the (C,6n) and (C,4n) reactions are comparable.

If the isotopic assignment of the new 35-hour activity to  $Cf^{246}$  is correct, these new data give a better idea as to the slope of the alpha-energy  $v_5$  mass number line for californium, which in turn makes it possible to make better predictions of the radicactive properties of the nuclides in this region.

We wish to express our appreciation to Professor J. G. Hamilton, J. F. Miller, G. B. Rossi, T. M. Putnam, Jr., M. T. Webb, and the operating crew of the 60-inch cyclotron in the Crocker Laboratory for their help in the carbon ion bombardments. We would also like to thank Mr. E. K. Hulet for his help in the chemical separations. This work was performed under the auspices of the U. S. Atomic Energy Commission.

<sup>&</sup>lt;sup>1</sup>Miller, Hamilton, Putnam, Haymond, and Rossi, Phys. Rev. (in press).

<sup>&</sup>lt;sup>2</sup>K. Street, Jr. and G. T. Seaborg, J. Am. Chem. Soc. <u>72</u>, 2790 (1950).

<sup>&</sup>lt;sup>3</sup>Thompson, Street, Jr., Ghiorso, and Seaborg, Phys. Rev.  $\frac{78}{2}$ , 298 (1950).

<sup>4</sup>Perlman, Ghiorso, and Seaborg, Phys. Rev. 77, 26 (1950).