

51/95js②

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76-CHO-3073

PPPL-3119
UC-426

PPPL-3119

ALFVÉNIC BEHAVIOR OF ALPHA PARTICLE DRIVEN ION CYCLOTRON EMISSION IN TFTR


BY

S. CAUFFMAN, R. MAJESKI, K.G. MCCLEMENTS AND R.O. DENDY

JULY 1995

PPPL

PRINCETON
PLASMA PHYSICS
LABORATORY



PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial produce, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE

This report has been reproduced from the best available copy.
Available in paper copy and microfiche.

Number of pages in this report: 23

DOE and DOE contractors can obtain copies of this report from:

Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831;
(615) 576-8401.

This report is publicly available from the:

National Technical Information Service
Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22161
(703) 487-4650

DISCLAIMER

**Portions of this document may be illegible
in electronic image products. Images are
produced from the best available original
document.**

Alfvénic Behavior of Alpha Particle Driven Ion Cyclotron Emission in TFTR

S. Cauffman and R. Majeski

Princeton Plasma Physics Laboratory, P.O. Box 451, Princeton, New Jersey 08543

K. G. McClements and R. O. Dendy

*UKAEA Government Division, Fusion, Culham, Abingdon, Oxfordshire, OX14 3DB UK
(Euratom/UKAEA Fusion Association)*

ABSTRACT

Ion cyclotron emission (ICE) has been observed during D-T discharges in the Tokamak Fusion Test Reactor (TFTR), using rf probes located near the top and bottom of the vacuum vessel. Harmonics of the alpha cyclotron frequency (Ω_α) evaluated at the outer midplane plasma edge are observed at the onset of the beam injection phase of TFTR supershots, and persist for approximately 100-250 ms. These results are in contrast with observations of ICE in JET, in which harmonics of Ω_α evolve with the alpha population in the plasma edge. Such differences are believed to be due to the fact that newly-born fusion alpha particles are super-Alfvénic near the edge of JET plasmas, while they are sub-Alfvénic near the edge of TFTR supershot plasmas. In TFTR discharges with edge densities such that newly-born alpha particles are super-Alfvénic, alpha cyclotron harmonics are observed to persist. These results are in qualitative agreement with numerical calculations of growth rates due to the magnetoacoustic cyclotron instability.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

INTRODUCTION

Cyclotronic emission has been observed during deuterium-tritium operation in both JET¹ and TFTR.^{2,3} In both cases, multiple harmonics of Ω_α (the alpha cyclotron frequency evaluated near the outer midplane edge of the plasma) are observed. On JET, the evolution of these harmonics follows that of the edge alpha particle population, and shows clear correlations with MHD events. On TFTR, emission at harmonics of Ω_α is also observed, but its temporal evolution is very different. In a typical TFTR supershot, Ω_α harmonics are observed within 50 ms of beam injection, but then disappear within 100-250 ms. These modes are then replaced by a new, distinct set of harmonics of Ω_D and Ω_T , the cyclotron frequencies of the injected species. Like the Ω_α emission, these frequencies correspond to a locus of emission near the outer midplane plasma edge, but at a slightly smaller major radius. While the magnetic field inferred from Ω_α (neglecting the Doppler shift) corresponds to a location slightly *beyond* the plasma edge (in the vacuum region), the field inferred from Ω_D and Ω_T corresponds to a location just *inside* the plasma edge. Typical early (Ω_α) and late (Ω_D and Ω_T) spectra, as well as the locations of cyclotron resonances within the plasma, are presented in figure 1. For comparison, ICE observations during DD plasmas in TFTR are described elsewhere.^{2,3,4}

Before discussing measurements of alpha particle driven ICE in TFTR in further detail, we present a brief description of the magnetoacoustic (or Alfvén) cyclotron instability (MCI), and its implications for expected emission in TFTR and in JET.

THE MAGNETOACOUSTIC CYCLOTRON INSTABILITY

The MCI has been identified as the probable mechanism behind fusion product driven ICE in both JET and TFTR.^{5,6,7} While ICE can be modelled with varying degrees of sophistication, much of the relevant physics is captured by simply solving the local dispersion relation for a uniform plasma with conditions comparable to those in the plasma edge. Such a treatment verifies that ICE can, under the right conditions, be excited by fusion products^{5,6} or by beam ions.⁸ We limit the present discussion to the issue of alpha particle driven ICE. At the onset of neutral beam injection, the alpha particle population in the plasma edge is initially dominated by barely trapped particles born near the core on very wide banana orbits. Figure 2 depicts such an orbit in TFTR, while figure 3 presents the pitch angle distribution of alpha particles at the TFTR plasma edge, as calculated by the ORBIT particle-following code,⁹ which integrates fusion source profiles (calculated by TRANSP¹⁰) over particle orbits. A simple but reasonable model for the distribution of birth alpha particles at the plasma edge is thus $f_\alpha \propto \exp\{-(v_{\parallel} - v_d)^2/v_{\parallel}^2 - (v_{\perp} - u)^2/v_{\perp}^2\}$, where $v_d^2 + u^2 = v_{\alpha 0}^2$ ($v_{\alpha 0}$ is the alpha particle birth velocity) and $u/v_d = \tan(\theta_0)$ (θ_0 is a pitch angle lying close to the trapped/passing boundary). Local dispersion relations for such distributions can be solved numerically, and in certain limits, analytically.^{5,6} In particular, as was shown in references 5 and 6, if one lets $f_\alpha \propto \exp\{-(v_{\parallel} - v_d)^2/v_{\parallel}^2\} \delta(v_{\perp} - u)$, takes the electrostatic limit, ignores Landau damping (appropriate if $k_{\parallel} \ll k_{\perp}$), and assumes $n_{\alpha} \ll n_i$, one recovers, to lowest order, the fast Alfvén wave dispersion relation:

$$\omega^2 = \frac{1}{2} c_A^2 \left[k^2 + k_{\parallel}^2 + k^2 k_{\parallel}^2 \frac{c_A^2}{\Omega_i^2} + \sqrt{\left(k^2 + k_{\parallel}^2 + k^2 k_{\parallel}^2 \frac{c_A^2}{\Omega_i^2} \right)^2 - 4k^2 k_{\parallel}^2} \right] \quad (1)$$

with the first order correction yielding the growth rate:

$$\gamma = \frac{\omega_{p\alpha}^2}{\omega_{pi}^2} \frac{\Omega_i^4}{(\Omega_i + (\omega - \Omega_i) N_{||}^2)(\Omega_i - (\omega + \Omega_i) N_{||}^2)} \left[\frac{n\Omega_\alpha}{k_{||}v_r} M_n - \frac{2u^2}{v_r^2} \eta_n N_n \right] \frac{\sqrt{\pi}}{2\omega} e^{-\eta_n^2} \quad (2)$$

where M_n and N_n are given by:

$$M_n = 2n \frac{\omega}{\Omega_i} \left[J_n'^2 + \frac{1}{z_a^2} (n^2 - z_a^2) J_n^2 \right] - 2 \frac{\omega^2 - \Omega_i^2}{\Omega_i^2} \frac{J_n J_n'}{z_a} \left[n^2 N_{||}^2 - (z_a^2 - 2n^2) N_{||}^2 \right] + \frac{2J_n J_n'}{z_a} (z_a^2 - 2n^2) \quad (3)$$

$$N_n = -2n \frac{\omega}{\Omega_i} \frac{J_n J_n'}{z_a} + \frac{\omega^2 - \Omega_i^2}{\Omega_i^2} \left[N_{||}^2 \left(\frac{n^2 J_n^2}{z_a^2} + J_n'^2 \right) + N_{\perp}^2 \frac{n^2 J_n^2}{z_a^2} \right] + \frac{n^2 J_n^2}{z_a^2} + J_n'^2 \quad (4)$$

and where $N_{||} = k_{||}c_A/\omega$, $N_{\perp} = k_{\perp}c_A/\omega$, $\eta_n = (\omega - k_{||}v_d - n\Omega_\alpha)/k_{||}v_r$, J_n is the Bessel function of order n with argument $z_a = k_{\perp}u/\Omega_\alpha$, and n is the integer for which $n\Omega_\alpha$ lies closest to ω . These equations can be expressed in terms of the following dimensionless parameters: the ratio of the alpha particle density to the background ion density ($\xi \equiv n_\alpha/n_i$), the mean alpha particle pitch angle (θ_0), the propagation angle ($\tan(\theta_k) \equiv k_{\perp}/k_{||}$), the ratio of the alpha birth velocity to the Alfvén speed ($\zeta \equiv v_{\alpha 0}/v_A$), and the narrowness of f_α ($\sigma \equiv v_r/u$). Numerical calculations indicate that while the growth rate is proportional to ξ , the value of σ required for instability is strongly dependent on ζ (see figures 4 and 5). If the alpha particles are super-Alfvénic (e.g. $\zeta = 1.5$), the second alpha particle harmonic, for example, can be destabilized for values of σ below 0.14; if the alpha particles are sub-Alfvénic (e.g. $\zeta = 0.5$), the same harmonic is stabilized for values of σ above 0.06.⁶ One would expect, therefore, that while the initial edge alpha particle population is narrow enough, ICE would be observed as ξ increases, but that as the distribution widens, the ICE would eventually be stabilized. Furthermore, this stabilization would be more easily achieved for smaller values of ζ .

IMPLICATIONS FOR JET AND TFTR

Due to the flatness of JET density profiles, 3.5 MeV alpha particles are super-Alfvénic even near the plasma edge.¹¹ TFTR supershots, on the other hand, are characterized by highly peaked density profiles, such as the one in figure 6. So while alpha particles are born super-Alfvénic in the core, they are sub-Alfvénic near the plasma edge. MCI theory would thus suggest that alpha particle driven ICE harmonics would be more easily excited in JET than in TFTR (which, indeed, is the case). In addition, the excitation of the MCI in TFTR depends on two competing processes: the rapid growth of the alpha particle population once beam heating commences, and the gradual widening of the alpha particle distribution function due to collisions. When alpha particle production begins, the distribution function at the plasma edge is very narrow, because the alpha particles have not yet slowed down and because only those with a narrow range of pitch angles (i.e. those that are barely trapped) will follow orbits carrying them through the plasma edge. As the alpha particle population continues to grow, however, the distribution function widens, eventually reaching the point at which the instability can no longer be excited. A simple analytic model of the evolution of the alpha particle distribution function (see reference 6 for a detailed description) predicts that for parameters appropriate for the edge plasma in TFTR supershots, alpha particle driven ICE should be observed during approximately the first 100 ms of beam injection, with the growth rate peaking near 40 ms. This prediction is qualitatively consistent with experimental observations.

ICE IN TFTR L-MODES

Recently, ICE data were obtained during several TFTR discharges with edge densities high enough to ensure that 3.5 MeV alpha particles were super-Alfvénic even near the plasma edge. MCI theory suggests that alpha particle driven ICE should be more easily excited under such conditions, and, in fact, this turns out to be the case. We present in figures 7, 8, and 9 a comparison of data from three different discharges. The first is a typical supershot, in which edge alpha particles are sub-Alfvénic throughout the discharge. In this case, alpha particle driven ICE is observed only at the onset of beams. The second is a supershot in which an L-mode transition is induced by means of a large He puff. In this case, alpha particle driven ICE is observed briefly at the onset of beams, but also *reappears* after the puff, when the density is high enough that $\zeta > 1$. The final case is an L-mode discharge for which the alpha particle birth speed is greater than the edge Alfvén speed throughout the beam phase. In this case, alpha particle driven ICE is observed immediately after the onset of beams, but persists until the beams are turned off. In all three cases, alpha particle driven ICE is observed only when the edge alpha particles have a very narrow distribution function, or when they are super-Alfvénic.

DISCUSSION

The data presented above provide valuable support for the MCI theory of ion cyclotron emission. While the local treatment described above is simplistic by construction, it nevertheless is qualitatively consistent with empirical observation. More elaborate models (such as the global eigenmode calculation presented in reference 7) should lead to further improvements in our understanding. At present, several aspects of ICE in TFTR remain poorly understood. First, spatial localization to one region of the plasma seems

insufficient, by itself, to account for the extreme narrowness of the observed modes. The alpha particle distribution function is described by a narrow drifting ring over a larger range of radii than the spectral line widths would indicate. We should also note, however, that a small fraction of alpha particle orbits extend beyond the outer midplane plasma edge. It may be that the least damped eigenmode is that with the lowest frequency accessible to the energetic species. (Recall that the locus of emission, neglecting any Doppler shift, is consistent with magnetic fields in the vacuum region beyond the plasma edge.) Second, the way in which the amplitude of alpha particle driven ICE in TFTR depends on global plasma parameters (e.g. fusion rate, beam power, plasma current) remains poorly understood. The ICE signal in JET, for example, has been demonstrated to be linearly proportional to the fusion rate over six orders of magnitude.¹ On TFTR, while a comparison of the peak Ω_α power with the peak D-T neutron rate also yields a regression coefficient close to unity, a large amount of scatter is also apparent.³ Given that the excitation of ICE is much more sensitive to the shape of the alpha particle distribution function in TFTR than it is in JET, its dependence on plasma conditions might be more complex as well. Finally, alpha particle driven ICE in TFTR exhibits an extremely rapid turn off at the end of the beam phase during the L-mode discharge (figure 6) presented earlier. Additional measurements of ICE during L-mode plasmas are planned.

SUMMARY

Emission at multiples of Ω_α is observed during D-T discharges in TFTR. While this emission is highly transient when $\nu_{\alpha 0} < \nu_A$, it reappears (or persists) when $\nu_{\alpha 0} > \nu_A$. These observations provide evidence for the

Alfvénic nature of ICE, and are consistent with predictions based on local growth rate calculations for the Magnetoacoustic Cyclotron Instability.

ACKNOWLEDGEMENTS

This work was supported by US DoE contract number DE-AC02-76-CH03073, the United Kingdom Department of Trade and Industry, and Euratom.

REFERENCES

- [1] COTTRELL, G.A., et al., Nucl. Fusion **33** (1993) 1365.
- [2] CAUFFMAN, S., MAJESKI, R., Rev. Sci. Instrum. **66** (1995) 817.
- [3] CAUFFMAN, S., MAJESKI, R., Bull. Am. Phys. Soc. **39** (1994) 1675.
- [4] GREENE, G., et al., Proceedings of the Seventeenth EPS, Amsterdam, **14B-IV** (1990) 1540.
- [5] DENDY, R.O., et al., Phys. Plasmas **1** (1994) 1918.
- [6] MCCLEMENTS, K.G., et al., An interpretation of ion cyclotron emission from sub-Alfvénic fusion products in the Tokamak Fusion Test Reactor, submitted to Phys. Plasmas.
- [7] GORELENKOV, N., CHENG, C.Z., to be published in Phys. Plasmas (June 1995).
- [8] DENDY, R.O., et al., Phys. Plasmas **1** (1994) 3407.
- [9] FELT, J., et al., Rev. Sci. Instrum. **61** (1990) 3262.
- [10] BUDNY, R., et al., Nucl. Fusion **34** (1994) 1247.
- [11] REBUT, P., et al., Nucl. Fusion **32** (1992) 187.

FIGURE CAPTIONS

FIG. 1. *Early and late ICE spectra from a DT supershot.* The apparent locus of emission (a) and the spectral structure at 66 ms (b) is distinct from that at 243 ms (c) after the onset of neutral beam injection.

FIG. 2. *Barely trapped alpha orbit passing through the plasma edge.* Orbit near the plasma edge will sample a significant fraction of the plasma core if they are barely trapped.

FIG. 3. *Alpha pitch angle distribution at the plasma edge.* Because most alpha particles are born near the core, their population near the edge will be dominated by those that are barely trapped.

FIG. 4. *MCI growth rates for super-Alfvénic alpha particles.* Shown here are growth rates for $n=2$, $\xi=10^{-4}$, $\theta_k=80^\circ$, $\theta_0=135^\circ$, $\zeta=1.5$, and three values of σ : 0.12, 0.13, and 0.14.


FIG. 5. *MCI growth rates for sub-Alfvénic alpha particles.* Shown here are growth rates for the same parameters as in figure 4, except $\zeta=0.5$, and $\sigma=0.050, 0.055$, and 0.060.


FIG. 6. *Supershot density profile.* The electron density, as well as the critical value above which 3.5 MeV alpha particles will be super-Alfvénic, is plotted as a function of major radius. Near the edge, alpha particles are sub-Alfvénic.

FIG. 7. *Alpha particle driven ICE during a typical supershot.* The neutron source rate (a), edge electron density (b), and ICE power in the Ω_α fundamental (c) are plotted as functions of time. Birth alpha particles are sub-Alfvénic in the plasma edge, and the Ω_α fundamental is only excited briefly at the onset of neutral beam injection.

FIG. 8. *Alpha particle driven ICE during a supershot-to-L-mode transition.* The same quantities are shown as in figure 7. The Ω_α fundamental is excited briefly at the onset of neutral beam injection, but reappears after a He puff raises the edge density so that $v_{\alpha 0} > v_A$.

FIG. 9. *Alpha particle driven ICE during an L-mode shot.* The same quantities are shown as in figures 7 and 8. In this case, $v_{\alpha 0} > v_A$ throughout the beam injection phase. The Ω_α fundamental is excited at the onset of neutral beam injection, and persists until the beams are turned off.








Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

- Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA
Prof. R.C. Cross, Univ. of Sydney, AUSTRALIA
Plasma Research Lab., Australian Nat. Univ., AUSTRALIA
Prof. I.R. Jones, Flinders Univ, AUSTRALIA
Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA
Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA
Prof. M. Goossens, Astronomisch Instituut, BELGIUM
Ecole Royale Militaire, Lab. de Phys. Plasmas, BELGIUM
Commission-European, DG. XII-Fusion Prog., BELGIUM
Prof. R. Boucqué, Rijksuniversiteit Gent, BELGIUM
Dr. P.H. Sakanaka, Instituto Fisica, BRAZIL
Prof. Dr. I.C. Nascimento, Instituto Fisica, Sao Paulo, BRAZIL
Instituto Nacional De Pesquisas Espaciais-INPE, BRAZIL
Documents Office, Atomic Energy of Canada Ltd., CANADA
Ms. M. Morin, CCFM/Tokamak de Varennes, CANADA
Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA
Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA
Prof. J. Teichmann, Univ. of Montreal, CANADA
Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA
Prof. R. Marchand, INRS-Energie et Matériaux, CANADA
Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA
Dr. C.R. James, Univ. of Alberta, CANADA
Dr. P. Lukáč, Komenského Universzita, CZECHO-SLOVAKIA
The Librarian, Culham Laboratory, ENGLAND
Library, R61, Rutherford Appleton Laboratory, ENGLAND
Mrs. S.A. Hutchinson, JET Library, ENGLAND
Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS
P. Mähönen, Univ. of Helsinki, FINLAND
Prof. M.N. Bussac, Ecole Polytechnique, FRANCE
C. Mouttet, Lab. de Physique des Milieux Ionisés, FRANCE
J. Radet, CEN/CADARACHE - Bat 506, FRANCE
Prof. E. Economou, Univ. of Crete, GREECE
Ms. C. Rinni, Univ. of Ioannina, GREECE
Preprint Library, Hungarian Academy of Sci., HUNGARY
Dr. B. DasGupta, Saha Inst. of Nuclear Physics, INDIA
Dr. P. Kaw, Inst. for Plasma Research, INDIA
Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL
Librarian, International Center for Theo Physics, ITALY
Miss C. De Palo, Associazione EURATOM-ENEA, ITALY
Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY
Prof. G. Rostangni, Istituto Gas Ionizzati Del Cnr, ITALY
Dr. H. Yamato, Toshiba Res & Devel Center, JAPAN
Prof. I. Kawakami, Hiroshima Univ., JAPAN
Prof. K. Nishikawa, Hiroshima Univ., JAPAN
Librarian, Naka Fusion Research Establishment, JAERI, JAPAN
Director, Japan Atomic Energy Research Inst., JAPAN
Prof. S. Itoh, Kyushu Univ., JAPAN
Research Info. Ctr., National Instit. for Fusion Science, JAPAN
Prof. S. Tanaka, Kyoto Univ., JAPAN
Library, Kyoto Univ., JAPAN
Prof. N. Inoue, Univ. of Tokyo, JAPAN
Secretary, Plasma Section, Electrotechnical Lab., JAPAN
Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN
Dr. G.S. Lee, Korea Basic Sci. Ctr., KOREA
J. Hyon-Sook, Korea Atomic Energy Research Inst., KOREA
D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
Leandro Melendez Lugo, Inst. Nac1. de Inves. Nucl, MEXICO
Prof. B.S. Liley, Univ. of Waikato, NEW ZEALAND
Inst of Physics, Chinese Acad Sci PEOPLE'S REP. OF CHINA
Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA
Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA
Z. Li, S.W. Inst Physics, PEOPLE'S REPUBLIC OF CHINA
Prof. J.A.C. Cabral, Instituto Superior Técnico, PORTUGAL
Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA
Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN
Dr. L. Stenflo, Univ. of UMEA, SWEDEN
Library, Royal Inst. of Technology, SWEDEN
Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY
Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys.I Apparatus, USSR
Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR
Librarian, The Ukr.SSR Academy of Sciences, USSR
Dr. L.M. Kovrzhnykh, Inst. of General Physics, USSR
Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
Librarian, Max-Planck-Institut, W. GERMANY