

10/18/95 JSC 1

LBL-36852  
UC-403



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

## EARTH SCIENCES DIVISION

**Features and Dimensions of the Hayward Fault  
Zone in the Strawberry and Blackberry Creek  
Area, Berkeley, California**

P.L. Williams

March 1995



#### DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof, or The Regents of the University of California.

Available to DOE and DOE Contractors  
from the Office of Scientific and Technical Information  
P.O. Box 62, Oak Ridge, TN 37831  
Prices available from (615) 576-8401

Available to the public from the  
National Technical Information Service  
U.S. Department of Commerce  
5285 Port Royal Road, Springfield, VA 22161

Lawrence Berkeley Laboratory is an equal opportunity employer.

## **DISCLAIMER**

**Portions of this document may be illegible  
in electronic image products. Images are  
produced from the best available original  
document.**

LBL-36852  
UC-403

**Features and Dimensions of the Hayward Fault Zone in the  
Strawberry and Blackberry Creek Area, Berkeley, California**

Patrick L. Williams

Earth Sciences Division  
Lawrence Berkeley Laboratory  
University of California  
Berkeley, California 94720

March 1995

This work was supported by the Director, Office of Energy Research, Division of Basic Energy Sciences, Geosciences Program, of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

*me*

MASTER

# Features and Dimensions of the Hayward fault zone in the Strawberry and Blackberry Creek area, Berkeley California

## *Introduction*

This report presents an examination of the geometry of the Hayward fault adjacent to the Lawrence Berkeley Laboratory and University of California campuses in central Berkeley. The fault crosses inside the eastern border of the UC campus. Most subtle geomorphic (landform) expressions of the fault have been removed by development, and by the natural processes of landsliding and erosion. Some clear expressions of the fault remain however, and these are key to mapping the main trace through the campus area. In addition, original geomorphic evidence of the fault's location was recovered from large scale mapping of the site dating from 1873 to 1897. Before construction obscured and removed natural landforms, the fault was expressed by a linear, northwest-trending zone of fault-related geomorphic features. There existed well-defined and subtle stream offsets and beheaded channels, fault scarps, and a prominent "shutter ridge". To improve our confidence in fault locations interpreted from landforms, we referred to clear fault exposures revealed in trenching, revealed during the construction of the Foothill Housing Complex, and revealed along the length of the Lawson Adit mining tunnel. Also utilized were the locations of offset cultural features. At several locations across the study area, distress features in buildings and streets have been used to precisely locate the fault. Recent published mapping of the fault (Lienkaemper, 1992) was principally used for reference to evidence of the fault's location to the northwest and southeast of Lawrence Berkeley Laboratory.

## *Fault-Related Landforms*

*Offset and beheaded streams.* One of the most direct means of mapping strike-slip (lateral slip) faults is by the identification and alignment of deflected drainages along faultlines. Many drainages are offset by the Hayward fault, and these provide primary evidence of the fault's location. Before construction of the California Memorial Stadium in 1923, the largest intact channel offset along the fault's length was that of Strawberry Creek (Figure 1, location A-A'). This large, competent stream is offset 1100 feet by right-lateral motion. The channel of Blackberry Creek is offset 190 feet by faulting. These two channel offsets are the most prominent evidence of the fault's location adjacent to Lawrence Berkeley Lab. A number of more subtle stream offsets and deflections have been interpreted from the 1897 UC Berkeley Building and Grounds basemap upon which more recent cultural features and the Hayward fault have been superimposed (Figure 1.).

Two former channels of Strawberry Creek cross the Berkeley Campus. These offset channels are termed "beheaded" or "abandoned", since they have been cut off from the Strawberry Creek watershed. The beheaded channels (locations B, C; Figure 1) were described by Buwalda (1929). One is located along a line between the east entrance to UC Berkeley and the Mining Circle. This "Mining Circle Channel" is offset 1900 feet from Strawberry Canyon (Location B, Figure 1). A second (and older) beheaded channel of Strawberry Creek is located beneath Hearst Avenue (Figure 1, location C). The "Hearst

"Avenue Channel" is offset 2400 feet from Strawberry Canyon. These landforms, and their deposits, provide clear evidence of the fault's location. Analysis of the morphology and ages of the beheaded and active Strawberry Creek channels also enables the quantification of both vertical and lateral deformation rates across the Hayward fault in Berkeley.

Several additional abandoned and offset stream channels are interpreted on Figure 1. The confidence of these interpretations varies, but where the offsetting fault is also located by additional evidence, the abandoned and offset streams increase confidence in the analysis.

*Shutter ridge.* Buwalda (1929) noted that the course of Strawberry Creek is deflected by a fault-bordering ridge of slope-base deposits, a *shutter ridge* (Located at sr'-sr", Figure 1). The 1000-foot-long shutter ridge ranged in height from 16 to 28 feet. Buwalda documented both the fault's location and lithology of the shutter ridge during construction of the California Memorial Stadium. He found that the ridge contained serpentine fragments. No source of serpentine exists in the Strawberry Creek drainage, but serpentine is present in Hamilton Gulch, 1400 feet southeast. This required that "at a date so recent that the topography of the region was essentially that of today" the ridge deposits had moved at least 1,400 feet from the gulch.

*Fault scarps.* Fault scarps near the foot of the Berkeley Hills appear as abrupt increases in slope angle as shown by tight contour spacing (Figure 1). The crisp definition of several of these scarps indicates the recency of fault movement. The linear trend of these scarps also aids in locating the fault trace. Particularly clear is the east-facing scarp (at sr, Figure 1) on the uphill side of the Strawberry Creek shutter ridge

#### *Offsets of Built Features.*

Along the Hayward fault, built features such as roads, utilities, fences, and buildings are offset by the slow (0.2"/yr) "creep" motion of the Hayward fault. The phenomena of creep and attendant distress of built features provides an additional method for locating fault lines. A number of buildings in the UC Berkeley area have been offset. Most notably, a total of 14 inches of fault motion has accumulated across the UC Berkeley Memorial Stadium since construction in 1923. Fault motion is expressed as distress of interior and exterior walls and joints, and is particularly apparent in the offset of an expansion joint along the stadium's south rim. This deformation, together with pre-construction landform data, are combined to precisely locate the fault just southwest of the Lawrence Berkeley Laboratory (Figure 2).

Near the northwestern corner of the Lawrence Berkeley Laboratory site, the Highlander Apartment complex is offset by fault motion. The Highlander offset, in conjunction with the discovery of offset sidewalks, streets and foundations north of Blackberry Creek, and landform data, provide a high confidence fault location to the north and west of Lawrence Berkeley Laboratory (Figure 2).

### *Fault Exposures*

Geotechnical investigations prior to construction of the Foothill Housing Project (Harding Lawson Associates, 1988) and study of exposures during construction provide additional information for the location of Hayward fault traces adjacent to Lawrence Berkeley Laboratory. Summary maps compiled from those observations (Williams, 1990; Williams and Hosokawa, 1992) were incorporated in this report. Evidence from direct exposure of the Hayward fault's primary traces are compiled in Figure 2. Additional information of the fault's location from pre-modification landforms and springs, and new evidence of fault creep, have confirmed many of the prior interpretations that were based primarily on exposures prior to and during construction in the fault zone.

*Acknowledgments.* This research was supported by the Director, Office of Energy Research, Division of Basic Energy Sciences, Geosciences Program, through U.S. Department of Energy contract DE-AC03-76SF00098. Support was also provided by the Lawrence Berkeley Laboratory Department of Facilities. Data Processing was carried out at the Center for Computational Seismology, Lawrence Berkeley Laboratory.

### *References cited*

Buwalda, J.P., 1929. Nature of the Late Movements on the Haywards Rift, Central California. *Bulletin of the Seismological Society of America*, 19, 187-199.

Harding Lawson Associates, 1988. Geologic and Fault Hazard Investigation, Phase II, Foothill Student Housing, University of California, Berkeley, California [on file at California Division of Mines and Geology, San Francisco, California; file no. AP-2529].

Lienkaemper, J.J., 1992. Map of recently active traces of the Hayward fault, Alameda and Contra Costa Counties, California. U.S. Geological Survey map, mf-2196, 1 sheet.

Williams, P.L., 1990, Hayward fault investigations, 1989, un-numbered report to the Director, Lawrence Berkeley Laboratory, 25 pp.

Williams, P.L. and A.M. Hosokawa, 1992, Geomorphic features related to the Hayward fault at the University of California at Berkeley, in Taylor, C.L., Hall N.T. and Melody, M. eds., *Field Trip Guidebook, Second Conference on Earthquake Hazards of the Eastern San Francisco Bay Area*, 65-71.

WGCEP 1990, (Working Group on California Earthquake Probabilities) J.H. Dietrich chair, C.R. Allen, L.S. Cluff, C.A. Cornell, W.L. Ellsworth, L.R. Johnson, A.G. Lindh, S.P. Nishenko, C.H. Scholz, D.P. Schwartz, W. Thatcher, and P.L. Williams, *Probabilities of large earthquakes in the San Francisco Bay Region, California*. USGS Circ. v. 1053, p. 51.

## *Figures*

Figure 1. Landforms and culture in the area of the Hayward fault zone, University of California, Berkeley (from Williams and Hosokawa, 1992). Locations of the fault traces have been improved subsequent to the publication of this map. University of California structures as of AD 1897 are solid. Selected later University of California structures outlined for reference. Major fault-related landforms include: A-A': Strawberry Creek channel offset; B: beheaded Mining Circle Channel; C: beheaded Hearst Avenue Channel; D: capture(?) of unnamed stream, E: offset of Blackberry Creek.

Figure 2. High resolution map of the Hayward fault trace in the Vicinity of Lawrence Berkeley Laboratory. Note two prominent traces of the fault, expressed by landforms, fault exposures, and distress to built structures. Where the fault trace is drawn as a solid line, the location has been determined from multiple lines of evidence, and the fault's location is deemed reliable. Where the fault's trace is dashed, the location is inferred from nearby evidence, and the location and continuity of the fault less certain in these locations.

### **Key to Numbering, Figure 2:**


1. Offset and distress to California Memorial Stadium (Harding-Lawson and Associates, 1988; Williams and Hosokawa, 1992).
2. Distress within Strawberry Creek bypass culvert (Williams and Hosokawa, 1992).
3. Offset of tributary channel of Strawberry Creek (Williams and Hosokawa, 1992).
4. Direct exposure of faulting of young sediments exposed in trenches or during excavation of Foothill Housing site (Harding-Lawson and Associates, 1988; Williams, 1990; Williams and Hosokawa, 1992).
5. Fault scarp interpreted from UC Berkeley basemap dated 1897 (Williams and Hosokawa, 1992).
6. Spring, from UC Berkeley basemap dated 1897 (Williams and Hosokawa, 1992).
7. Small offset within active channel of Strawberry Creek from excavation plan for California Memorial Stadium (UC Berkeley Facilities Drawing No. 713, dated 3/15/1922; interpreted in this study).
8. Exposure of active fault in Lawson Adit (Professor. George. D. Louderback, written communication to Luther A. Nichols, Comptroller, University of California, Berkeley, dated 12/13/1939).
9. Beheaded channel of Strawberry Ck. (Williams and Hosokawa, 1992).
10. Distress and en echelon cracks in foundation of Highlander Apartments (this study).
11. 190' offset of Blackberry Creek (Williams and Hosokawa, 1992).
12. Offset of sidewalk, pavement, curb, foundation or retaining wall (this study).

Landforms in the Area of the Hayward Fault Zone,  
UC Berkeley, Modified from UC Berkeley  
Grounds and Buildings map, 1897



0 200 400 Feet

Contours below 400 feet: 4 ft. intervals  
Contours above 400 feet: 8 ft. intervals






Figure 1.




Figure 2.