

Title: Developing Microbe-Plant Interactions for Applications in Plant-Growth Promotion and Disease Control, Production of Useful Compounds, Remediation, and Carbon Sequestration

Authors

Cindy H. Wu ^{1*}

Stéphanie M. Bernard ¹

Gary L. Andersen ¹

Wilfred Chen ²

Affiliations

1. Lawrence Berkeley National Laboratory. Earth Sciences Division. One Cyclotron Rd. Berkeley, CA 94720, USA
2. Department of Chemical and Environmental Engineering, University of California, Riverside, CA 92521, USA

* Corresponding Author

E-mail: CHWu@lbl.gov

Phone: 510.486.4594

Fax: 510.486.7152

Summary

Interactions between plants and microbes are an integral part of our terrestrial ecosystem. Microbe-plant interactions are being applied in many areas. In this review, we present recent reports of applications in the areas of plant-growth promotion, biocontrol, bioactive compound and biomaterial production, remediation and carbon sequestration. Challenges, limitations and future outlook for each field are discussed.

Introduction

Interactions between plants and microbes are an integral part of our terrestrial ecosystem. There are several types of plant-microbe interactions: competition, commensalism, mutualism, and parasitism. The more common interactions are commensalism or mutualism, where either one or both species benefit from the relationship, respectively (Campbell, 1993). There are several excellent reviews reporting current research on lifestyles and molecular interactions of plant-associated bacteria (Sorensen and Sessitsch, 2007), rhizosphere interactions (Singh et al., 2004a), plant responses to bacterial quorum sensing (QS) signals (Bauer and Mathesius, 2004), endophyte applications (Ryan et al., 2008), and rhizosphere bacteria responses to transgenic plants (Fillion, 2008).

Examination of these interactions helps us to understand natural phenomena that affect our daily lives and could lead to applications resulting in sustainable resources, less impact on the environment, cleanup of pollution and influence on atmospheric gases on a global-scale. Advantages of using these interactions for biotechnological applications are many-fold. The use of naturally existing plant-microbe symbiosis for plant growth and

biocontrol reduces synthetic fertilizer and pesticide treatments leading to cost-effectiveness and less impact by nutrients (Boddey et al., 2003) and pesticides (Whipps et al., 2004; Elmer and Reglinski, 2006) on surrounding fauna and flora. The production of useful compounds with pharmaceutical and industrial relevance using plant-bacteria symbiosis is energy efficient (Wu et al., 2007; Del Giudice et al., 2008) and diminishes the need to add expensive precursors and catalysts. Remediation through conventional method, such as excavate and treat, is expensive and labor intensive. Conversely, plant-microbial remediation strategies can be less intrusive and much more economical (Anderson et al., 1993). Carbon sequestration through plant-rhizosphere processes is a potentially sustainable method to lowering atmospheric carbon (Kumar et al., 2006). This review focuses on recent progress in the fields of plant-growth promotion, plant disease control, production of bioactive compounds and biomaterials, remediation of contaminated sites, and carbon sequestration. The potential of applying these new developments are discussed. Figure 1 summarizes applications resulting from microbe-shoot and microbe-root interactions and techniques used. Table 1 is a glossary of the techniques mentioned in this review.

Plant-growth promotion

Plant-microbe interactions have been utilized to improve plant growth for the production of food, fiber, biofuels and key metabolites. The mutualistic interaction can be beneficial in directly providing nutrients to the plant (biofertilizer) or increasing the availability of compounds such as iron or phosphate. Free-living plant-growth promoting bacteria also produce compounds that directly affect plant metabolism or modulate phytohormone

production or degradation. The phytohormones: auxins, cytokinins, gibberellic acid (GA_3), abscisic acid (ABA) and ethylene are signaling molecules essential for growth which mediate a range of developmental processes in plants. Recent studies on each of these areas are presented in the following section.

Biofertilizers

As chemical fertilizers are costly both to the agricultural businesses and the environment, development of biofertilizers is an important and exciting area. Symbiotic relationships established between plants and bacteria such as rhizobium and some actinomycetes (e.g. *Frankia*) provide most of the nitrogen (N) available to legumes and actinorhizal plant species. A number of studies attempted to develop new association between non-legume plants and N-fixing bacteria, however, these attempts were mostly unsuccessful (Preininger and Cyurjan, 2001). The discovery of endophytic diazotrophs that provided reduced N to the non-legume plant species represented an alternative way of exploiting plant-microbe interaction for N nutrition. Brazilian sugarcane plants harboring N-fixing endophytes were grown for many years with low fertilizer inputs, and showed no symptoms of N deficiencies (Boddey et al., 2003). Other non-leguminosae plant species were also shown to benefit from association with diazotrophic endophytes, such as wheat (*Triticum aestivum*; Iniguez et al., 2004) and rice (*Oryza sativa*; James et al., 2002). Iniguez et al. (2004) generated a *nifH* mutant from *Klebsiella pneumoniae* and compared this to the wild type strain in order to demonstrate that the wild type *K. pneumoniae* strain provided N to the wheat plant thus alleviating N deficiency. The wheat plants were supplemented with stable N-isotope labeled nutrients. Mass spectrometry was used to

show that less ^{15}N was present in wheat plants inoculated with the wild type *K. pneumoniae* strain. Since these plants obtained a higher percentage of their N from non-isotopically labeled atmospheric N_2 , the majority of the plant nitrogen was derived from the wild type strain. In a different study, the transcript abundance of the *nifH* gene was measured by RT-PCR in endophytic *Herbaspirillum* sp. during colonization of internal regions of wild rice plants (You et al., 2005). Interestingly, they showed higher *nifH* transcript abundance during the light period compared to the dark period. Given the tight relationship between N-fixing activity and *nifH* gene transcription, the high-level of *nifH* mRNA during the light period indicated that the endophytes produced higher level of reduced N at a time when they could benefit most from photosynthate production. This study described a promising avenue for non-legume plants to obtain N from the reduction of atmospheric N_2 . It remains to be seen that similar results are achievable with soil grown plants or in the field where plants may be less limited for growth than in sand-filled pots. In addition, competition from other bacteria may prevent colonization by endophytic diazotroph.

Plant genetic contribution to this symbiosis is an important area of research leading to selection of agriculturally important plant genotypes with increased capacity to establish symbiosis for nutrient acquisition. Several studies have begun to pave the way in this area. The identification of genetic loci underlying complex traits in *Lotus* spp and *Medicago* led to the sequencing of specific genes (Stougaard, 2001; Salvi and Tuberrosa, 2005) essential for the establishment of root symbiosis. Kistner et al. (2005) characterized seven mutants of *Lotus japonicus* that were impaired in nodulation and were also defective in arbuscular mycorrhiza formation. Kanamori et al. (2006) used a F2 mapping

population established by crossing the *Lotus japonicus* mutant and wild type for positional cloning of a gene essential for symbiosis that encoded a plant nucleoporin gene required for Ca^{2+} spiking occurring after contact between *Nod*-factor molecules and root hair cells. The combination of genetic and genomic analysis resulted in a deeper understanding of the plant genetic contribution to the establishment of symbiosis with rhizosphere bacteria. One drawback of this approach is that it may take many years, from the identification of the genetic basis for complex traits to the development of new cultivars with enhanced capacity for establishing and maintaining symbiosis.

Phytohormone modulation

Rhizosphere bacteria and phyllosphere colonizing epiphytes have been shown to produce a range of plant growth stimulating phytohormones. A recent study by Boiero et al. (2007) evaluated the phytohormone synthesis of three commercially available strains of *Bradyrhizobium japonicum* grown in pure culture. They showed that the three strains have differential capability to produce the five major phytohormones: auxins, cytokinins, gibberellic acid (GA_3), abscisic acid (ABA) and ethylene. This is important to consider as each individual and combinations of phytohormones may have different impact on plant growth.

The auxin compound indole-3-acetic acid (IAA) can be generated in bacteria through different biosynthetic pathways (Spaepen et al., 2007). Saravanan et al. (2008) reviewed the roles in plant growth promotion of two N- fixing organisms, *Gluconacetobacter diazotrophicus* and an *Acetobacteraceae* strain. In particular they discussed additional roles for *G. diazotrophicus*, including the production of plant

hormones such as IAA, and gibberellins. Idris et al. (2007) showed for the first time that the gram-positive bacterium *Bacillus amyloliquefaciens* produced and secreted significant amounts of IAA. They also showed the positive effect of IAA on the growth of *Lemna minor*. Patten and Glick (2002) demonstrated the direct positive effect of IAA produced by *Pseudomonas putida* through the indolepyruvic acid pathway, on root development. Roots from Canola seeds treated with the wild-type *P. putida* were longer in length than that of seeds treated with an IAA-deficient mutant. However, the bacterial production of IAA may not always be beneficial for plant as it is involved in pathogenesis and that high concentration of IAA can also inhibit root cell growth (Spaepen et al. 2007 for review).

Ethylene has many physiological effects on plant growth, development and modulation of responses to biotic and abiotic stresses. Bacteria such as *Pseudomonas* spp., *Burkholderia caryophylli*, *Achromobacter piechaudii* were shown to lower the endogenous ethylene level in planta by producing a degradative enzyme 1-aminocyclopropane-1-carboxylic acid (ACC)-deaminase (Shahroona et al., 2007; Mayak et al., 2004a, b). The effects of ACC-deaminase producing rhizobacteria on plants included increased root growth, and improved tolerance of salt and water stress (Mayak et al., 2004a, b). These effects were noted in axenic conditions and more recently in field conditions. Inoculation of wheat (*Triticum aestivum* L.) plants with *Pseudomonas* spp. and *B. caryophylli* improved grain and straw yield by as much as 43% and 44%, respectively (Shahroona et al., 2007). However no measurements of ethylene concentration were carried out on plants in pots or in the field to show direct correlation of ACC-deaminase production and ethylene degradation. It is therefore possible that other bacteria-associated effects might have improved wheat plant growth.

Cytokinins constitute a group of plant hormones that promote cell division in conjunction with auxin and are known to induce stomata-opening. Rhizosphere bacteria and fungi associated with plants were shown to produce cytokinins. Arkhipova et al. (2005) studied the production of different cytokinins by *Bacillus subtilis* and showed improved growth for lettuce plants (*Lactuca sativa* L.) after inoculation with cytokinin-producing bacteria. This was also the case when plants were grown under water-stress (Arkhipova et al., 2007). The authors mainly considered the production of cytokinins by *B. subtilis* as mechanism of plant production. It would be interesting to investigate possible alternative mechanism by which this species is able to promote plant growth. Lettuce plants were grown in sand and the study did not address whether similar results could be obtained in the field where competition from other rhizosphere bacteria might prevent *B. subtilis* growth and where bacterial production of other phytohormones might interfere with the effect of cytokinins.

The modulation of phytohormone by bacteria that lead to disease resistance is addressed in the Biocontrol section.

Future outlook

Plant growth promoting bacteria have been shown to positively impact plant performance through different mechanisms. For future development of commercial inoculant, it is therefore important to consider all potential metabolic activity of the phytohormone-producing bacteria. In addition, components of the inoculant growth media could be transformed into metabolites (phytohormones) that could impact plant growth in the early stages of plant development (Boiero et al., 2007). For successful application of plant-

growth promotion using bacterial inoculant, many aspects of the plant environment have to be considered in the field. In particular, it is important that the inoculant be beneficial to the crop, and do not improve the growth of the weedy species and do not render the plants more susceptible to biotic and abiotic stresses. Bacterial colonization efficiency is also critical for successful inoculation. Further development of plant growth promoting bacteria could benefit from selection, through breeding programs of plant genotypes that respond better to the plant growth promoting bacteria.

Plant disease control

The efficiency of biocontrol agents for plant disease has been demonstrated and some are commercially available (Whipps, 2004; Elmer and Reglinski, 2006; Whipps and Gerhardson, 2007). General mechanisms of action for plant pathogen control include competition for nutrients and space at the infection site, antibiosis, parasitism, production of cell wall degrading enzymes, induced resistance in the plant, and manipulation of bacterial signaling molecules. It is likely that several mechanisms of action are at work in many biocontrol agents. Compant et al. (2005) reviews the mechanisms of plant disease biocontrol by plant growth-promoting bacteria. Here, we discuss two mechanisms of disease control: induced resistance and manipulation of signaling molecules.

Induced Resistance

Interactions with bacteria can induce two types of plant defense responses that help protect against further infection. Systemic acquired resistance (SAR) is a specific response that triggers both a local increase in phytohormone accumulation and the

formation of phloem mobile signal. Non-pathogenic free-living rhizosphere bacteria and endophytes can trigger the second type of plant defense called induced systemic resistance (ISR). A major distinction between SAR and ISR is the involvement of salicylic acid (SA), with ISR being activated via a SA-independent pathway.

One example of plant disease control is the use of *Pseudomonas* strains with biocontrol activity to induce resistance in apple (*Malus domestica*) against the pathogenic fungus *Venturia inaequalis*, which causes apple scab. The movement of *Pseudomonas* on apple leaves was studied in order to understand its antagonistic interactions against *V. inaequalis* (Gau et al., 2002). Using confocal laser scanning microscopy (CLSM), it was demonstrated that *P. fluorescens* Bk3 localized near stomatal openings. The *P. fluorescens* Bk3 traversed the cuticle through secretion of cutinases, and acquired nutrient from fluid isolated from the apoplastic space. Also, isolated leaf cuticles stimulated bacterial extracellular proteins (Singh et al., 2004b). In order to understand the interaction between the *V. inaequalis* and apple tree, cDNA libraries using suppression subtractive hybridization were constructed for the resistant and susceptible cultivars (Degenhardt et al., 2005). Many plant defense related transcripts such as those encoding for β -1,3-glucanase, cystein protease inhibitor and metallothioneins were at higher levels in the resistant cultivar. On the other hand, more RuBisCo transcripts were found in the susceptible cultivar than in the resistant one. Similar proteins were expressed in apple trees treated with the biocontrol strain *P. fluorescens* Bk3 and *V. inaequalis* (Kurkcuoglu et al., 2007). The presence of *P. fluorescens* Bk3 was demonstrated to elevate the defense mechanisms in the apple trees and could serve as effective biocontrol strategy. The

authors implied that low levels of RuBisCo and high levels of metallothioneins resembled that of “old leaves” and were “unattractive” to *V. inaequalis* (Degenhardt et al., 2005). The authors did not investigate whether *P. fluorescens* Bk3 induced SAR or ISR type of plant resistance.

Manipulation of bacterial signaling molecules

Some bacteria rely on signaling molecules for the development of pathogenesis. Other microbes living in the same environment may degrade these signaling molecules. Recombinant biocontrol strains producing *carAB*, genes required for degradation of a fatty acid signaling molecule, were shown to reduce virulence caused by *Xanthomonas* sp. and *Xylella fastidiosa* (Newman et al., 2008). Secondary metabolites produced by plants were shown to mimic or inhibit quorum sensing (QS) molecules. *M. truncatula* was shown to produce more than a dozen compounds that stimulated or inhibited QS (Gao et al., 2003). Plants have been manipulated to produce molecules that mimic or block QS signal, and enzymes that degrade QS molecules or quorum-quenching strategy. Scott et al. (2006) engineered tobacco plants to synthesize acyl-homoserine-lactones (AHL) in the chloroplast and demonstrated that AHL was transported and secreted on the phyllosphere and in the rhizosphere. Pre-treatment of potato slices with *Bacillus thuriensis* resulted in reduced maceration from *Erwinia carotovora* virulence (Dong et al., 2004). The authors showed that the decreased pathogenesis involved the disruption of *E. carotovora* quorum sensing by deterioration of QS molecules with AHL-lactonase. Pre-treatment of potato slices with *B. thuriensis* lacking the ability to produce AHL-lactonase did not reduce maceration. Manipulating plants to express AHL-lactonase has

shown positive effect in protecting plants against pathogens (Zhang, 2003, for review). Recently, a gene called *qsda* (for quorum-sensing signal degradation) from *Rhodococcus erythropolis*, encoding a new type of AHL-lactonase has been characterized. The gene was able to confer quorum-quenching capacity to *P. fluorescens* that led to enhanced protection of potato tuber against the soft rot pathogen *P. carotovorum* (Uroz et al., 2008).

Future outlook

Elucidating the mechanism by which bacteria elicit ISR is important for the development of commercial biocontrol agent. Flagellin, siderophores, lipopolysaccharides and more recently volatile organic compounds have been proposed as determinants that trigger ISR (Comptant et al. 2005 for review). Recent studies have also suggested that exopolysaccharides (EPS) produced by *Bukholderia gladioli* are able to elicit ISR in cucumber plants (Park et al., 2008). A better understanding of bacterial elicitor of ISR may eventually lead to the development of disease resistance strategies for crop protection.

Despite their potential use as biocontrol agent, bacterial inoculant are not widely used in agricultural setting. While their success has been demonstrated in some field trials, they are not consistently efficient in diverse field conditions (Mark et al., 2006). One characteristic that is paramount to the success of bacterial inoculant is a high efficiency in colonizing the rhizosphere. The disruption of quorum sensing either by plants or bacteria have potential use in plant disease protection. However, it has relied on

the genetic manipulation of plants or microbes thus far. The release of these genetically modified organisms might lead to potential risk to the crop supplies. Furthermore, it has been shown that plants can detect AHL and take this cue to enhance their defense against pathogens (Mathesius et al., 2003). Therefore, the artificial quenching of AHL molecules could also interfere with natural adaptation of the plant's defense system. Overall, it seems that while quorum quenching may delay the onset of pathogenesis it.

Production of bioactive compounds and biomaterials

Secondary metabolites produced by plants constitute a major source of bioactive compounds that can be used as therapeutic agents or for biomaterial production. Recent studies have shown that plant-microbial interaction can be exploited to enhance the production of important secondary metabolites. The infection of plants by *Agrobacterium rhizogenes* produced hairy roots that showed high growth rate and branching, and were used for production of useful compounds (Guillon et al., 2006, for review). Plant-bacteria interaction can be used to improve production of secondary metabolites by hairy roots. For example, Wu et al. (2007) established co-cultures of *Salvia miltiorrhiza* bunge hairy roots with *Bacillus* sp. and demonstrated enhanced production of tashinone. Tashinone is a bioactive compound in *S. miltiorrhiza* bunge root, used in Chinese medicine, for the treatment of menstrual disorders and cardiovascular diseases and prevention of inflammation. The hairy root culture is an exciting example of plant-microbe interaction that is now moving towards large-scale industrial application. Scaling up hairy roots culture for production of secondary metabolites may not be straightforward due to the

complex features of hairy roots. Please refer to other reviews that discussed aspects of this process (Srivastava and Srivastava, 2007; Georgiev et al., 2007).

Rhizobacteria was shown to induce the accumulation of sesquiterperne synthase transcripts (Del Giudice et al., 2008). Sesquiterperne, used in cosmetics and perfumery, is one component of vetiver root essential oil synthesis. The authors showed that the root-associated bacteria metabolized the vetiver oil and produced additional compounds, which suggested that each distinct rhizobacteria community contributed to a signature composition of commercial vetiver oil.

For two decades, it was thought that the plant pathogenic fungus *Rhizopus microsporus* produced the antimitotic polyketide macrolide rhizoxin. Partida-Martinez and Hertweck (2005) uncovered that it was actually an endosymbiont, *Burkholderia* sp., of the fungus that produced rhizoxin. The authors demonstrated that in the absence of *Burkholderia*, no rhizoxin was produced in the fungal culture. Transmission electron microscopy demonstrated that *Burkholderia* sp. was localized in the fungal cytosol (Partida-Martinez et al., 2007). Rhizoxin inhibits mitosis and leads to cell cycle arrest, and has potential as an antitumor drug. Further investigation elucidated rhizoxin derivative structures and obtained stable analogues by inhibition of a putative P-450 monooxygenase (Scherlach et al., 2006).

Ryan et al. (2008) summarized applications for bacterial endophytes, including production of biomaterials such as poly-3-hydroxybutyrate (PHB) and poly-3-

hydroxyalkanoate (PHA). Catalán et al. (2007) showed that diazotrophic endophyte *Herbaspirillum seropedicae* could accumulate 36% of its biomass as PHB and could constitute a cost-effective mean for producing biomaterial.

Future outlook

The above example of three-way interaction between plant-bacterium-fungus provides a reason to step beyond pair-wise interactions and currently existing model systems (Douglas 2008; Heath, 2008). Marra et al. (2006) examined the three-way interactions among bean plant (*Phaseolus vulgaris*), two pathogenic fungi and an antagonistic *Trichoderma* spp. and found that induced systemic resistance was achieved. Focusing on interactions from a community perspective may reveal further useful applicable mechanisms.

Remediation

Researchers have explored the use of plant-microbe symbiosis for remediation of pollutants since the early 90's (Anderson et al., 1993; Brazil et al., 1995). Investigations in this area have progressed from bench-scale experiments towards addressing more *in planta* field remediation efficiencies in recent years. Current phyto- or rhizo-remediation systems targeting contaminants such as recalcitrant chlorinated compounds, volatile organic carbons and heavy metals are highlighted here.

Recalcitrant chlorinated compounds

Organic contaminants naturally biodegrade, except for chlorinated compounds such as polychlorinated biphenyl (PCB) and 4-chloronitrobenzene (4CNB), which tend to be persistent and recalcitrant in the environment. Recent reports on chlorinated compound degrading rhizosphere bacteria are presented. Natural PCB-degrader bacterial populations were cultured from several plant species growing in a contaminated site. High numbers of culturable PCB-degrader colonies were isolated from roots of Austrian pine (*Pinus nigra*) and goat willow (*Salix caprea*) (Leigh et al., 2006). Most of the PCB-metabolizing bacteria are *Rhodococcus* sp. However, the isolates were first selected based on cultivability and morphology thus possibly leading to a biased representation of PCB-degrading bacterial communities. In addition, since the PCB degradation capability of these strains was tested in liquid medium, the benefits of plant association was not addressed.

Narasimhan et al. (2003) investigated the role of plant secondary compounds in stimulating rhizobacteria growth and PCB removal efficiency. The authors identified that the *Arabidopsis* root exudates consisted mostly of phenylpropanoids, such as flavonoids, lignins and indole compounds. Wild type and mutant *Arabidopsis* lines over producing flavonoids sustained higher counts of flavonoid-utilizing *P. putida* strain than bacteria grown on *Arabidopsis* mutant not producing flavonoids. More interestingly, close to 90% of PCB was degraded in soil adhering to the roots indicating that direct contact with roots and the exudates resulted in bacteria growth and biodegradation enhancement. Genetically engineered microbes were also applied towards remediation. The PCB degradation efficiency of a recombinant strain of *P. fluorescens* expressing a *bph* operon

under the control of nodulation (*nod*) genes from *Sinorhizobium meliloti*, was examined (Villacieros et al., 2005). Resting cell PCB-degradation experiments indicated that the recombinant strain metabolized different cogeners of PCB more efficiently than the *Burkholderia* sp. strain LB400. However, no plant-bacteria potted experiments were performed to measure PCB degradation.

Böltner et al. (2008) enriched rhizosphere microbes and demonstrated that four *Sphingomonas* strains were capable of rhizoremediating hexachlorocyclohexane (Lindane). Potted experiments showed that 30% of lindane was removed with corn seedlings inoculated with the *Sphingomonas* strains. Whereas, in unplanted soil, sterile planted soil, and uninoculated-unplanted soil, less than 3% of lindane was removed.

Liu et al. (2007) successfully inoculated a *gfp*-tagged strain of *Comamonas* sp. onto the roots of alfalfa as demonstrated by quantitative competitive (QC)-PCR and confocal laser scanning microscopy (CLSM). The rhizosphere community shifted due to addition of 4CNB and the *Comamonas* strain was characterized using denaturing gradient gel electrophoresis (DGGE). In outdoor potted experiments, the symbiosis enhanced phytotoxicity resistance, and removed 4CNB faster than the plant control without the strain in a 24-hour period. However, the control without the strain removed 60% of the 4CNB compared to 99% in experiment with strain inoculation. It seemed likely that if the experiment was extended beyond the 24-hour measurements, the 4CNB removal might be similar between the control and the treatment.

Volatile organic compounds

Phytoremediation has been used as a treatment method to remove contaminants from groundwater. However, certain volatile organic compounds such as trichloroethylene (TCE) and BTEX (benzene, toluene, ethylbenzene, xylene) are released into the atmosphere through the plant's vascular system. The endophyte-plant interaction was used to degrade these volatile organic pollutants and minimize evapotranspiration (Barac et al., 2004; Taghavi et al., 2005; Moore et al., 2006). Barac et al. (2004) demonstrated that the genetically modified endophytic strain of *Burkholderia cepacia* together with yellow lupine (*Lupinus luteus L.*) reduced evapotranspiration of toluene by 50-70%, and reduced phytotoxicity. The same group inoculated poplar cuttings with two strains of *B. cepacia*, an endophyte and a soil isolate, expressing the toluene monooxygenase (*tom*), and showed significantly less toluene being transpired in the poplar inoculated with the endophytic *B. cepacia* (Taghavi et al., 2005). In addition, the authors indicated evidence for horizontal gene transfer of the recombinant plasmid encoding the toluene degradation pathway from the inoculant *B. cepacia* to the endogenous microbial community, and suggested that it would be possible to eliminate selection of an appropriate endophyte since the biodegradation gene will be transferred to the endogenous microorganisms. However, the authors touched on the issue of low horizontal transfer efficiency but did not elaborate on solutions for the lack of control over the time scale of the transfer and recipients of the genes. The authors also did not comment on the risks associated with the spread of exogenous genes into a new environment. However, the risk caused by a gene originally isolated from the environment, such as the *tom* gene should be low compared to risks from synthetic genes.

Moore et al. (2006) isolated endophytic bacteria from two poplar varieties that were grown at a field trial site phytoremediating BTEX in the groundwater. There were differences in spatial compartmentalization of strain localization in the root, shoot and leaves suggesting there were species-specific and non-specific associations between bacteria and plants. However, the study did not examine the reasons for the strain localization and association with the plants and whether remediation efficiency was affected by bacterial community.

Metals

Heavy metal contamination is a persistent problem since the metals, unlike organic compounds, do not biodegrade. Several recent reviews address role of plant growth promoting rhizobacteria in metals remediation (Khan et al., 2009), and various plant-bacteria systems that have been applied towards remediation of metals contamination (Kamaludeen and Ramasamy, 2008). Removal of metals from soil or groundwater using rhizosphere bacteria and plants has been demonstrated recently. Willow (*Salix caprea*) seedlings inoculated with different strains of rhizosphere *Streptomyces* and *Agromyces* and the strains ability to produce indole acetic acid (IAA) and siderophore was measured. No correlation was observed between IAA and siderophore production and metal accumulation. Only plants inoculated with *Agromyces terreus* exhibited reduced phytotoxicity due to cadmium and zinc. Metal accumulation in leaf biomass was measured (Kuffner et al., 2008). However, the data did not demonstrate a marked difference between the inoculated plants and the non-inoculated control. Perhaps, most of

the phytoaccumulated metals were immobilized in the roots where most of the rhizosphere strains would be located. No root metal concentrations were measured.

Wu et al. (2006) utilized an engineered symbiosis between recombinant *P. putida* and sunflower plants (*Helianthus annuus*) for adsorption of cadmium. The recombinant *P. putida* expressed a synthetic phytochelatin, EC20, and exhibited inherent cadmium resistance. The inoculation of the *P. putida* strain enhanced plant growth and resulted in 40 % more cadmium accumulation from hydroponic solutions than the non-inoculated control. No field study was conducted. Ryan et al. (2007) demonstrated that a recombinant strain of *P. fluorescens* F113rifPCB with arsenic resistance genes and PCB degradation capabilities protected *M. sativa* when grown in soil supplemented with sodium arsenate. No arsenic removal nor PCB degradation efficiency were assessed with this plant-bacteria system. Since most of the metals tend to accumulate in the top 20 cm layer of the soil (Li et al., 2009), grass species with high fine-root biomass providing large surface area for bacteria colonization and metal accumulation, in the top soil layer, would be ideal for remediation of metal contaminated sites. The plants serve as concentrators of metals, and further treatment of the plant biomass accumulated with metals would be required. Overall, the use of the plant-bacteria system would be more cost-effective than excavation of the soil contaminated with low levels of heavy metals.

Future outlook

The successful demonstrations of laboratory-scale phyto- or rhizo- remediation do not always translate into adequate removal of contaminants in field-scale. The reasons

include heterogeneity of the field site, unpredictability of environmental conditions, inability to sustain bacterial population and the remediating microbes being outcompeted by endogenous organisms. There are several reasons that plant-bacteria interactions are advantageous when applied to remediation. The plant-associated bacteria population would be more competitive than the native soil microorganisms since plant exudates provide nutrients. Plants would act as natural pumps that draw contaminated soil pore water towards the plant-associated remediating bacteria. Rhizoremediation is an ideal strategy for cleanup of mixed-contaminants. Recombinant rhizosphere bacteria with specific genes targeting pollutants present at the site can be inoculated into plant roots. For example, Lee et al. (2006) engineered two strains of rhizobacteria with TCE degradation capability, and surface expression of synthetic phytochelatins for improved heavy metal resistance, thus enhancing the rate of TCE degradation.

An excellent review by Gerhardt et al. (2009) addressed strategies to overcoming challenges of field application of phyto- or rhizo-remediation such as stressors to the microbes, complexity of the field conditions, regulatory acceptability and the use of genetically modified organisms (GMOs). Research that look at the impact of GMOs on native bacteria (de Cácer et al., 2007) and bacterial community changes post-introduction of willow trees (*S. viminalis*) for rhizoremediation of PCB contamination (de Cácer et al., 2007) are starting to emerge and will elucidate questions regarding the impact on the native microfauna. Studies that compare remediation efficiencies of natural attenuation, bioaugmentation, phytoremediation and rhizoremediation, under field conditions (van Dillewijn et al., 2007), assess bacterial colonization (Molina et al., 2000;

Watt et al., 2006) and activity (Wu et al., 2008) on plant roots, and measure allometric relationships of tree trunk size and various root parameters (Olson et al., 2003) provide invaluable information for determining field application options and evaluating contaminant removal. More extensive, long-term field comparisons of control plots with treatment plots for remediation efficiency are necessary to validate laboratory observations and to gain public and regulatory agency confidence for this promising application.

Carbon sequestration

Another application of plant-microbe interaction drawing increasing attention is carbon (C) sequestration, where atmospheric C is deposited as plant root material, incorporated into the soil microorganisms and soil organic matter. It is hypothesized that increase in CO₂ leads to an increase in rhizodeposition, and wider C/N ratio thus retarding decomposition (Norby and Cotrufo, 1998). Several reviews are dedicated to presenting methods for measuring CO₂ fluxes in different soil compartments (Kuzyakov, 2006), mechanism of root carbon stabilization (Rasse et al., 2005), effects of elevated CO₂ on belowground carbon storage (Pendall et al., 2004), and carbon sequestration by roots (Kumar et al., 2006). This nascent field is still at the exploratory stage where most of the research is focusing on understanding the effects of elevated CO₂ levels on microbial community, below-ground plant material production, and biomass decomposition, as well as land management practices and plant species on the long term potential of C rhizodeposition.

Affects of land management practices and fertilizer application

Several researchers explored the possibility of different land management practices to enhance C sequestration. Bailey et al. (2002) compared *ex situ* incubations of soil samples from five different ecosystems (desert, restored tallgrass prairie, Douglas fir forest, loblolly pine forest, and agricultural land). The restored prairie samples had the highest total soil carbon and also the highest fungal-to-bacterial activities (F:B). The authors asserted that increased F:B ratios correlated with higher amount of carbon stored in the soil. Also, that invasive land management decreased fungal biomass and thus the carbon stored in the soil. Soil samples were collected and CO₂ respiration experiments were conducted in a laboratory setting over a 6-hour period, thus the measurements might not be representative of field conditions.

Heinemeyer et al. (2007) described contradicting results in their *in situ* study of the ability of Lodgepole pine associated mycorrhiza to store soil C over a period of 1-year, where the fungus was thought to return plant surplus C directly back to the atmosphere. Verburg et al. (2004) compared the net ecosystem carbon exchange between the atmosphere and two experimental non-native cheatgrass (*Bromus tectorum*) varieties in a 2-year study in the Desert Research Institute's (Nevada, USA) Ecologically Controlled Enclosed Lysimeter Laboratories. They showed that fertilization increased C uptake initially, however, C loss through soil respiration was also enhanced.

Bazot et al. (2006) found similar results, at a grassland ecosystem of Free Air CO₂ Enrichment (FACE) in Eschikon, Switzerland, where increased N supply to the plants

enhanced allocation of fixed C to the shoots and reduced belowground carbon allocation and rhizodeposition. The plots were enriched with CO₂ for 9-years in this study. At the same Swiss FACE facility, changes in microorganism structural diversity and function were also examined after 9-years of CO₂ enrichment in monocultures and mixed cultures of *Trifolium repens* L. cv. Milkanowa and *Lolium perenne* L. cv. Bastion with and without N treatments (Drissner et al., 2007). The authors concluded that increased atmospheric CO₂ stimulated microbial enzymatic activities and changed structural diversity. Subsequently, the increase in microbial activity led to higher mineralization rate of soil organic matter and thus would reduce the C sequestered in the soil. However, the soil C concentrations were not quantified in this study.

Long-term field studies

Long-term effects are essential for assessing the applicability of plant-rhizosphere C sequestration potential, and are investigated in the following field studies. Ingram et al. (2008) investigated microbial community changes to cattle grazing practices over a 10-year period at the High Plains Grasslands Research Station (Wyoming, USA). The authors found that even moderate grazing impacted microbial community structures and vegetation composition that could lead to loss of soil organic C to the atmosphere. In the heavily grazed area, there was a 30% loss of soil organic C and the measured N-mineralization rate was the lowest among the different grazing regimes.

A 5-year study of loblolly pine-associated fungi at a FACE site (North Carolina, USA) resulted in an increase in fungal biomass and a shift in distribution to deeper soils

(Pritchard et al., 2008). Leake et al. (2006) presented a comprehensive study on C fluxes from plants to soil microbiota using $^{13}\text{CO}_2$ pulse-labeling on a Scottish upland grassland over a period of 4-years. It was found that there were two distinct pools of soil carbon: one with fast turnover and one with slow turnover time. Mycorrhizal mycelial system contributed to the pool with the fast turnover, by consuming approximately 9% of the fixed C, and returning much of it to the atmosphere within 16 hours. Liming was found to increase shoot and root productivity. However, liming enhanced soil respiration and microbial biomass production, thus decreasing the amounts of C retained in the soil.

Bremer et al. (2008) compared soil organic C 6- and 12- years after a crop rotation study commenced and found that the C sequestration of the different soil treatments did not increase after the first 6-years for the carbon-conserving practices. However, fertilized grass continued to gain soil organic C over the 12-year period. Niklaus and Falloon (2006) studied the capacity of calcareous grassland exposed to high levels of CO_2 for 6-years and found that the C sequestration potential was limited due to processes that were unaccounted for such as increased soil moisture due to reduced leaf conductance, soil disaggregation due to increased moisture and accelerated soil organic matter decomposition. A 30-year soil C projection modeling using field measurements was used to assessing affects of ambient and elevated CO_2 level (Sindhøj et al., 2006). The results did not support the hypothesis that decreased litter quality due to increased CO_2 level would lead to lower decomposition rate (Norby and Cotrufo, 1998). The authors suggested that it would be more effective to reduce the rate of decomposition, rather than increase storage of plant litter C.

Future outlook

The studies presented in this review do not encompass all possible plant-microbe C sequestration research available. However, most of the studies discussed here indicate that optimal conditions for belowground C storage have not been found. There are many possible direct and indirect effects on the belowground C pools and processes resulting from elevated CO₂ and increased temperature (Pendall et al., 2004). Microbial activity is affected by increased temperature leading to increased N availability and net primary production. There might be differences in C sequestration capacity for different plant species, and whether they are perennial or annual. The water content of the soil and N-limitation would affect microbial mineralization of organic soil C. The challenges may lie in improving measurement methods of C stored below ground and de-convoluting the myriads of confounding factors that could affect C and N cycling. In addition, the scale of the application required to have an impact on the atmospheric C level is unclear at this point. More long-term and standardized studies, under different environmental conditions, of belowground carbon fluxes, integrating models and measurements are needed. C sequestration through plant-microbe interaction is still in its exploratory phase. As more worldwide attention is drawn towards mitigating elevated atmospheric C level, hopefully more global collaborative interdisciplinary research efforts will be directed towards assessing the conditions required for successful application of plant-microbe C sequestration.

Conclusion

The biotechnological applications presented in this review are evident of a human-plant-microbe symbiosis where all three benefit from one another. Applications with plant-microbe interactions provide more economical and environmentally sound alternatives to conventional processes. Successful practice of these applications will depend on enhancements in the following areas. The sometimes confounding results in field applications might result from varying environmental conditions or species specificity and, thus, warrant comprehensive investigations of basic mechanisms and evaluation of responses. Knowledge of available techniques and continual improvement are important first steps to addressing the unknowns in this complex network. Research in microbe-plant interactions will benefit greatly from development of new techniques. A global approach in quantifying all possible changes is necessary, and is now feasible with the development of high throughput techniques in genomics, metagenomics, transcriptomics, proteomics and metabolomics. These techniques produce a sizeable dataset and require sophisticated statistical analyses and modeling (Fiehn, 2002). The results from the global analyses would elucidate key pathways or genes. Genetic manipulation in plants and microbes altering metabolic pathways and genes is needed for validation. Most importantly, long-term field treatment efficacy and impacts associated with use of genetically modified organisms need to be assessed thoroughly (Fillion et al., 2008). Expansion of plant transgenic and microbial recombinant protocols would facilitate the validation process, as well as the development of new applications.

Acknowledgment

In memoriam of Bo Bodvarsson, LBL Earth Sciences Division Director, 2001 to 2006.

We are grateful to Janet Jansson and two anonymous reviewers for critically reviewing this manuscript. We thank Walter Denn for drawing the plant figure.

We would like to acknowledge funding support from the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. SMB acknowledges support from the U.S. Department of Energy (DOE), Office of Science, Biological and Environmental Research, Program for Ecosystem Research.

References

Anderson, T.A., Guthrie, E.A., and Walton, B.T. (1993) Bioremediation in the rhizosphere. *Environ Sci Technol* **27**: 2630-2636.

Arkhipova, T. N., Prinsen, E., Veselov, S. U., Martinenko, E. V., Melentiev, A. I. and Kudoyarova, G. R. (2007) Cytokinin producing bacteria enhance plant growth in drying soil. *Plant Soil* **292**: 305-315.

Arkhipova, T.N., Veselov, S.U., Melentiev, A.I., Martynenko, E.V., and Kudoyarova, G.R. (2005) Ability of bacterium *Bacillus subtilis* to produce cytokinins and to control of plant diseases: The history and evolution of current concepts. *Plant Dis* **87**: 4-10.

Bailey, V.L., Smith, J.L., and Bolton, H. (2002) Fungal-to-bacterial ratios in soils investigated for enhanced C sequestration. *Soil Biol Biochem* **34**: 997-1007.

Barac, T., Taghavi, S., Borremans, B., Provoost, A., Oeyen, L., Colpaert, J.V. et al. (2004) Engineered endophytic bacteria improve phytoremediation of water-soluble, volatile, organic pollutants. *Nat Biotechnol* **22**: 583-588.

Bauer, W.D., and Mathesius, U. (2004) Plant responses to bacterial quorum sensing signals. *Curr Opin Plant Biol* **7**: 429-433.

Bazot, S., Ulff, L., Blum, H., Nguyen, C., and Robin, C. (2006) Effects of elevated CO₂ concentration on rhizodeposition from *Lolium perenne* grown on soil exposed to 9 years of CO₂ enrichment. *Soil Biol Biochem* **38**: 729-736.

Boddey, R.M., Urquiaga, S., Alves, B.J.R., and Reis, V. (2003) Endophytic nitrogen fixation in sugarcane: present knowledge and future applications. *Plant Soil* **252**: 139-149.

Boiero, L., Perrig, D., Masciarelli, O., Penna, C., Cassan, F., and Luna, V. (2007) Phytohormone production by three strains of *Bradyrhizobium japonicum* and possible physiological and technological implications. *Appl Microbiol Biotechnol* **74**: 874-880.

Böltner D, G.P., Muñoz-Rojas J, Duque E, Moreno-Morillas S, Sánchez L, Ramos JL (2008) Rhizoremediation of lindane by root-colonizing *Sphingomonas*. *Microb biotechnol* **1**: 87-93.

Brazil, G.M., Kenefick, L., Callanan, M., Haro, A., de Lorenzo, V., Dowling, D.N., and Ogara, F. (1995) Construction of a rhizosphere *Pseudomonad* with potential to degrade polychlorinated-biphenyls and detection of BPH gene expression in the rhizosphere. *Appl Environ Microbiol* **61**: 1946-1952.

Bremer, E., Janzen, H.H., Eliert, B.H., and McKenzie, R.H. (2008) Soil organic carbon after twelve years of various crop rotations in an Aridic Boroll. *Soil Sci Soc Am J.* **72**: 970-974.

Campbell, N. (1993) Biology 3rd Ed. Brady, E.B. (ed). Redwood City, CA: The Benjamin/Cummings Publishing Company Inc.

Catalán, A.I., Ferreira, F., Gill, P.R., and Batista, S. (2007) Production of polyhydroxyalkanoates by *Herbaspirillum seropedicae* grown with different sole carbon sources and on lactose when engineered to express the lacZlacY genes. *Enzyme Microb Technol* **40**: 1352-1357.

Compan, S., Duffy, B., Nowak, J., Clement, C., and Barka, E.A. (2005) Use of plant growth-promoting bacteria for biocontrol of plant diseases: Principles, mechanisms of action, and future prospects. *Appl Environ Microbiol* **71**: 4951-4959.

de Cárcer, D.A., Martin, M., Karlson, U., and Rivilla, R. (2007) Changes in bacterial populations

and in biphenyl dioxygenase gene diversity in a polychlorinated biphenyl-polluted soil after introduction of willow trees for rhizoremediation. *Appl Environ Microbiol* **73**: 6224-6232.

de Cárcer, D.A., Martin, M., Mackova, M., Macek, T., Karlson, U., and Rivilla, R. (2007) The introduction of genetically modified microorganisms designed for rhizoremediation induces changes on native bacteria in the rhizosphere but not in the surrounding soil. *Isme Journal* **1**: 215-223.

Degenhardt, J., Al-Masri, A.N., Kurkcuoglu, S., Szankowski, I., and Gau, A.E. (2005) Characterization by suppression subtractive hybridization of transcripts that are differentially expressed in leaves of apple scab-resistant and susceptible cultivars of *Malus domestica*. *Mol Genet Genomics* **273**: 326-335.

Del Giudice, L., Massardo, D.R., Pontieri, P., Berte, C.M., Mombello, D., Carata, E. et al. (2008) The microbial community of Vetiver root and its involvement into essential oil biogenesis. *Environ Microbiol* **10**: 2824-2841.

Dong, Y.H., Zhang, X.F., Xu, J.L., and Zhang, L.H. (2004) Insecticidal *Bacillus thuringiensis* silences *Erwinia carotovora* virulence by a new form of microbial antagonism, signal interference. *Appl Environ Microbiol* **70**: 954-960.

Douglas, A.E. (2008) Conflict, cheats and the persistence of symbioses. *New Phytol* **177**: 849-858.

Drissner, D., Blum, H., Tscherko, D., and Kandeler, E. (2007) Nine years of enriched CO₂ changes the function and structural diversity of soil microorganisms in a grassland. *Eur J Soil Sci* **58**: 260-269.

Elmer, P.A.G., and Reglinski, T. (2006) Biosuppression of *Botrytis cinerea* in grapes.

Plant Pathol **55**: 155-177.

Fiehn, O. (2002) Metabolomics - the link between genotypes and phenotypes. *Plant Mol Biol* **48**: 155-171.

Fillion, M. (2008) Do transgenic plants affect rhizobacteria? *Microb biotechnol* **1**: 463-475.

Gao, M.S., and Teplitski, M. (2008) RIVET - A tool for in vivo analysis of symbiotically relevant gene expression in *Sinorhizobium meliloti*. *Mol Plant-Microbe Interact* **21**: 162-170.

Gao, M.S., Teplitski, M., Robinson, J.B., and Bauer, W.D. (2003) Production of substances by *Medicago truncatula* that affect bacterial quorum sensing. *Mol Plant-Microbe Interact* **16**: 827-834.

Gau, A.E., Dietrich, C., and Kloppstech, K. (2002) Non-invasive determination of plant-associated bacteria in the phyllosphere of plants. *Environ Microbiol* **4**: 744-752.

Georgiev, M.I., Pavlov, A.I., and Bley, T. (2007) Hairy root type plant in vitro systems as sources of bioactive substances. *Appl Microbiol and Biotechnol* **74**: 1175-1185.

Gerhardt, K.E., Huang, X.D., Glick, B.R., and Greenberg, B.M. (2009) Phytoremediation and rhizoremediation of organic soil contaminants: Potential and challenges. *Plant Science* **176**: 20-30.

Guillon, S., Tremouillaux-Guiller, J., Pati, P.K., Rideau, M., and Gantet, P. (2006) Harnessing the potential of hairy roots: dawn of a new era. *Trends Biotechnol* **24**: 403-409.

Heath, K.D. (2008) The coevolutionary genetics of plant-microbe interactions. *New*

Phytol **180**: 268-270.

Heinemeyer, A., Hartley, I.P., Evans, S.P., De la Fuente, J.A.C., and Ineson, P. (2007)

Forest soil CO₂ flux: uncovering the contribution and environmental responses of ectomycorrhizas. *Glob Change Biol* **13**: 1786-1797.

Idris, E.E., Iglesias, D.J., Talon, M., and Borrius, R. (2007) Tryptophan-dependent production of indole-3-acetic acid (IAA) affects level of plant growth promotion by *Bacillus amyloliquefaciens* FZB42. *Mol Plant-Microbe Interact* **20**: 619-626.

influence the growth and endogenous hormone content of lettuce plants. *Plant Soil* **272**: 201-209.

Ingram, L.J., Stahl, P.D., Schuman, G.E., Buyer, J.S., Vance, G.F., Ganjegunte, G.K. et al. (2008) Grazing impacts on soil carbon and microbial communities in a mixed-grass ecosystem. *Soil Sci Soc Am J* **72**: 939-948.

Iniguez, A.L., Dong, Y.M., and Triplett, E.W. (2004) Nitrogen fixation in wheat provided by *Klebsiella pneumoniae* 342. *Mol Plant-Microbe Interact* **17**: 1078-1085.

James, E.K., Gyaneshwar, P., Mathan, N., Barraquio, Q.L., Reddy, P.M., Iannetta, P.P.M. et al. (2002) Infection and colonization of rice seedlings by the plant growth-promoting bacterium *Herbaspirillum seropedicae* Z67. *Mol Plant-Microbe Interact* **15**: 894-906.

Kanamori, N., Madsen, L.H., Radutoiu, S., Frantescu, M., Quistgaard, E.M.H., Miwa, H. et al. (2006) A nucleoporin is required for induction of Ca²⁺ spiking in legume nodule development and essential for rhizobial and fungal symbiosis. *Proc Natl Acad Sci U S A* **103**: 359-364.

Kamaludeen, S.P.B., and Ramasamy, K. (2008) Rhizoremediation of metals: harnessing

microbial communities. *Indian J Microbiol* **48**: 80-88.

Khan, M.S., Zaidi, A., Wani, P.A., and Oves, M. (2009) Role of plant growth promoting

rhizobacteria in the remediation of metal contaminated soils. *Environ Chem Lett* **7**: 1-19.

Kistner, C., Winzer, T., Pitzschke, A., Mulder, L., Sato, S., Kaneko, T. et al. (2005)

Seven *Lotus japonicus* genes required for transcriptional reprogramming of the root during fungal and bacterial symbiosis. *Plant Cell* **17**: 2217-2229.

Kuffner, M., Puschenreiter, M., Wieshammer, G., Gorfer, M., and Sessitsch, A. (2008)

Rhizosphere bacteria affect growth and metal uptake of heavy metal accumulating willows. *Plant Soil* **304**: 35-44.

Kumar, R., Pandey, S., and Pandey, A. (2006) Plant roots and carbon sequestration. *Curr*

Sci **91**: 885-890.

Kurkcuglu, S., Degenhardt, J., Lensing, J., Al-Masri, A.N., and Gau, A.E. (2007)

Identification of differentially expressed genes in *Malus domestica* after application of the non-pathogenic bacterium *Pseudomonas fluorescens* Bk3 to the phyllospere. *J Exp Bot* **58**: 733-741.

Kuzyakov, Y. (2006) Sources of CO₂ efflux from soil and review of partitioning

methods. *Soil Biol Biochem* **38**: 425-448.

Leake, J.R., Ostle, N.J., Rangel-Castro, J.I., and Johnson, D. (2006) Carbon fluxes from

plants through soil organisms determined by field (CO₂)-¹³C pulse-labelling in an upland grassland. *Appl Soil Ecol* **33**: 152-175.

Lee, W., Wood, T.K., and Chen, W. (2006) Engineering TCE-degrading rhizobacteria for heavy metal accumulation and enhanced TCE degradation. *Biotechnol Bioeng* **95**: 399-403.

Leigh, M.B., Prouzova, P., Mackova, M., Macek, T., Nagle, D.P., and Fletcher, J.S. (2006) Polychlorinated biphenyl (PCB)-degrading bacteria associated with trees in a PCB-contaminated site. *Appl Environ Microbiol* **72**: 2331-2342.

Li, P.J., Stagnitti, F., Xiong, X., and Peterson, J. (2009) Temporal and spatial distribution patterns of heavy metals in soil at a long-standing sewage farm. *Environ Monit Assess* **149**: 275-282.

Li, X.J., Fekete, A., Englmann, M., Gotz, C., Rothballer, M., Frommberger, M. et al. (2006) Development and application of a method for the analysis of N-acylhomoserine lactones by solid-phase extraction and ultra high pressure liquid chromatography. *J Chromatogr A* **1134**: 186-193.

Liu, L., Jiang, C.Y., Liu, X.Y., Wu, J.F., Han, J.G., and Liu, S.J. (2007) Plant-microbe association for rhizoremediation of chloronitroaromatic pollutants with *Comamonas* sp strain CNB-1. *Environ Microbiol* **9**: 465-473.

Marra, R., Ambrosino, P., Carbone, V., Vinale, F., Woo, S.L., Ruocco, M. et al. (2006) Study of the three-way interaction between *Trichoderma atroviride*, plant and fungal pathogens by using a proteomic approach. *Curr Genet* **50**: 307-321.

Mark, G.L., Morrissey, J.P., Higgins, P., and O'Gara, F. (2006) Molecular-based strategies to exploit *Pseudomonas* biocontrol strains for environmental biotechnology applications. *Fems Microbiol Ecol* **56**: 167-177.

Mayak, S., Tirosh, T., and Glick, B.R. (2004a) Plant growth-promoting bacteria that confer resistance to water stress in tomatoes and peppers. *Plant Sci* **166**: 525-530.

Mayak, S., Tirosh, T., and Glick, B.R. (2004b) Plant growth-promoting bacteria confer resistance in tomato plants to salt stress. *Plant Physiol Biochem* **42**: 565-572.

Molina, L., Ramos, C., Duque, E., Ronchel, M.C., Garcia, J.M., Wyke, L., and Ramos, J.L. (2000) Survival of *Pseudomonas putida* KT2440 in soil and in the rhizosphere of plants under greenhouse and environmental conditions. *Soil Biology & Biochemistry* **32**: 315-321.

Moore, F.P., Barac, T., Borrernans, B., Oeyen, L., Vangronsveld, J., van der Lelie, D. et al. (2006) Endophytic bacterial diversity in poplar trees growing on a BTEX-contaminated site: The characterisation of isolates with potential to enhance phytoremediation. *Syst Appl Microbiol* **29**: 539-556.

Narasimhan, K., Basheer, C., Bajic, V.B., and Swarup, S. (2003) Enhancement of plant-microbe interactions using a rhizosphere metabolomics-driven approach and its application in the removal of polychlorinated biphenyls. *Plant Phys* **132**: 146-153.

Newman, K.L., Chatterjee, S., Ho, K.A., and Lindow, S.E. (2008) Virulence of plant pathogenic bacteria attenuated by degradation of fatty acid cell-to-cell signaling factors. *Mol Plant-Microbe Interact* **21**: 326-334.

Niklaus, P.A., and Falloon, P. (2006) Estimating soil carbon sequestration under elevated CO₂ by combining carbon isotope labelling with soil carbon cycle modelling. *Glob Change Biol* **12**: 1909-1921.

Norby, R.J., and Cotrufo, M.F. (1998) Global change - A question of litter quality. *Nature* **396**: 17-18.

Olson, P.E., Wong, T., Leigh, M.B., and Fletcher, J.S. (2003) Allometric modeling of plant root growth and its application in rhizosphere remediation of soil contaminants. *Environ Sci Technol* **37**: 638-643.

Park, K., Kloepper, J.W., and Ryu, C.M. (2008) Rhizobacterial exopolysaccharides elicit induced resistance on cucumber. *J Microbiol Biotechnol* **18**: 1095-1100.

Partida-Martinez, L.P., and Hertweck, C. (2005) Pathogenic fungus harbours endosymbiotic bacteria for toxin production. *Nature* **437**: 884-888.

Partida-Martinez, L.P., de Looss, C.F., Ishida, K., Ishida, M., Roth, M., Buder, K., and Hertweck, C. (2007) Rhizonin, the first mycotoxin isolated from the zygomycota, is not a fungal metabolite but is produced by bacterial endosymbionts. *Appl Environ Microbiol* **73**: 793-797.

Patten, C.L., and Glick, B.R. (2002) Regulation of indoleacetic acid production in *Pseudomonas putida* GR12-2 by tryptophan and the stationary-phase sigma factor RpoS. *Can J Microbiol* **48**: 635-642.

Pendall, E., Bridgman, S., Hanson, P.J., Hungate, B., Kicklighter, D.W., Johnson, D.W. et al. (2004) Below-ground process responses to elevated CO₂ and temperature: a discussion of observations, measurement methods, and models. *New Phytol* **162**: 311-322.

Preininger, E. and Gyurjan, I. (2001) Trials to create artificial nitrogen-fixing symbioses and associations using in vitro methods: An outlook. *In Vitro Cell Dev Biol Plant* **37**: 139-148.

Pritchard, S.G., Strand, A.E., McCormack, M.L., Davis, M.A., and Oren, R. (2008) Mycorrhizal and rhizomorph dynamics in a loblolly pine forest during 5 years of free-air-CO₂-

enrichment. *Glob Change Biol* **14**: 1252-1264.

Rasse, D.P., Rumpel, C., and Dignac, M.F. (2005) Is soil carbon mostly root carbon? Mechanisms for a specific stabilisation. *Plant Soil* **269**: 341-356.

Ryan, R.P., Germaine, K., Franks, A., Ryan, D.J., and Dowling, D.N. (2008) Bacterial endophytes: recent developments and applications. *FEMS Microbiol Lett* **278**: 1-9.

Salvi, S., and Tuberosa, R. (2005) To clone or not to clone plant QTLs: present and future challenges. *Trends Plant Sci* **10**: 297-304.

Saravanan, V.S., Madhaiyan, M., Osborne, J., Thangaraju, M., and Sa, T.M. (2008) Ecological occurrence of *Gluconacetobacter diazotrophicus* and nitrogen-fixing Acetobacteraceae members: Their possible role in plant growth promotion. *Microb Ecol* **55**: 130-140.

Scherlach, K., Partida-Martinez, L.P., Dahse, H.M., and Hertweck, C. (2006) Antimitotic rhizoxin derivatives from a cultured bacterial endosymbiont of the rice pathogenic fungus *Rhizopus microsporus*. *J Am Chem Soc* **128**: 11529-11536.

Scott, R.A., Weil, J., Le, P.T., Williams, P., Fray, R.G., von Bodman, S.B., and Savka, M.A. (2006) Long- and short-chain plant-produced bacterial N-acyl-homoserine lactones become components of phyllosphere, rhizosphere, and soil. *Mol Plant-Microbe Interact* **19**: 227-239.

Shahroona, B., Jamro, G.M., Zahir, Z.A., Arshad, M., and Memon, K.S. (2007) Effectiveness of various *Pseudomonas* spp. and *Burkholderia caryophylli* containing ACC-Deaminase for improving growth and yield of wheat (*Triticum aestivum* L.). *J Microbiol Biotechnol* **17**: 1300-1307.

Sindhój, E., Andren, O., Katterer, T., Gunnarsson, S., and Pettersson, R. (2006) Projections of

30-year soil carbon balances for a semi-natural grassland under elevated CO₂ based on measured root decomposability. *Agric Ecosyst Environ* **114**: 360-368.

Singh, B.K., Millard, P., Whiteley, A.S., and Murrell, J.C. (2004a) Unravelling rhizosphere-microbial interactions: opportunities and limitations. *Trends Microbiol* **12**: 386-393.

Singh, P., Piotrowski, M., Kloppstech, K., and Gau, A.E. (2004b) Investigations on epiphytic living *Pseudomonas* species from *Malus domestica* with an antagonistic effect to *Venturia inaequalis* on isolated plant cuticle membranes. *Environ Microbiol* **6**: 1149-1158.

Sorensen, J. and Sessitsch, A. (2007) Plant-associated bacteria - lifestyle and molecular interactions. In Modern Soil Microbiology 2nd Edition. van Elsas, J.D., Jansson, J. D., and Trevors, J. T. (eds). Boca Raton: CRC Press, pp. 211-236.

Spaepen, S., Vanderleyden, J., and Remans, R. (2007) Indole-3-acetic acid in microbial and microorganism-plant signaling. *FEMS Microbiol Rev* **31**: 425-448.

Srivastava, S., and Srivastava, A.K. (2007) Hairy root culture for mass-production of high-value secondary metabolites. *Crit Rev Biotechnol* **27**: 29-43.

Stougaard, J. (2001) Genetics and genomics of root symbiosis. *Curr Opin Plant Biol* **4**: 328-335.

Taghavi, S., Barac, T., Greenberg, B., Borremans, B., Vangronsveld, J., and van der Lelie, D. (2005) Horizontal gene transfer to endogenous endophytic bacteria from poplar improves phytoremediation of toluene. *Appl Environ Microbiol* **71**: 8500-8505.

Uroz, S., Oger, P.M., Chapelle, E., Adeline, M.T., Faure, D., and Dessaix, Y. (2008) A *Rhodococcus qsdA*-encoded enzyme defines a novel class of large-spectrum quorum-

quenching lactonases. *Appl Environ Microbiol* **74**: 1357-1366.

Van Dillewijn, P., Caballero, A., Paz, J.A., Gonzalez-Perez, M.M., Oliva, J.M., and Ramos, J.L. (2007) Bioremediation of 2,4,6-trinitrotoluene under field conditions. *Environ Sci Technol* **41**: 1378-1383.

Verburg, P.S.J., Arnone, J.A., Obrist, D., Schorran, D.E., Evans, R.D., Leroux-Swarthout, D. et al. (2004) Net ecosystem carbon exchange in two experimental grassland ecosystems. *Glob Change Biol* **10**: 498-508.

Villacieros, M., Whelan, C., Mackova, M., Molgaard, J., Sanchez-Contreras, M., Lloret, J. et al. (2005) Polychlorinated biphenyl rhizoremediation by *Pseudomonas fluorescens* F113 derivatives, using a *Sinorhizobium meliloti* nod system to drive *bph* gene expression. *Appl Environ Microbiol* **71**: 2687-2694.

Watt, M., Hugenholtz, P., White, R., and Vinall, K. (2006) Numbers and locations of native bacteria on field-grown wheat roots quantified by fluorescence in situ hybridization (FISH). *Environ Microbiol* **8**: 871-884.

Whipps, J.M. (2004) Prospects and limitations for mycorrhizas in biocontrol of root pathogens. *Can J Bot* **82**: 1198-1227.

Whipps, J.M. and Gerhardson B. (2007) Biological Pesticides for control of seed- and soil-borne plant pathogens. In Modern Soil Microbiology 2nd Edition. van Elsas, J.D., Jansson, J. D., and Trevors, J. T. (eds). Boca Raton: CRC Press, pp. 479-501.

Wu, C.H., Hwang, Y.C., Lee, W., Mulchandani, A., Wood, T.K., Yates, M.V., and Chen, W. (2008) Detection of recombinant *Pseudomonas putida* in the wheat

rhizosphere by fluorescence in situ hybridization targeting mRNA and rRNA. *Appl Microbiol Biotechnol* **79**: 511-518.

Wu, C.H., Wood, T.K., Mulchandani, A., and Chen, W. (2006) Engineering plant-microbe symbiosis for rhizoremediation of heavy metals. *Appl Environ Microbiol* **72**: 1129-1134.

Wu, J.Y., Ng, J., Shi, M., and Wu, S.J. (2007) Enhanced secondary metabolite (tanshinone) production of *Salvia miltiorrhiza* hairy roots in a novel root-bacteria coculture process. *Appl Microbiol Biotechnol* **77**: 543-550.

You, M., Nishiguchi, T., Saito, A., Isawa, T., Mitsui, H., and Minamisawa, K. (2005) Expression of the *nifH* gene of a *Herbaspirillum* endophyte in wild rice species: Daily rhythm during the light-dark cycle. *Appl Environ Microbiol* **71**: 8183-8190.

Zhang, L.H. (2003) Quorum quenching and proactive host defense. *Trends Plant Sci* **8**: 238-244.

Table 1. Glossary of techniques used to study plant-microbe interactions.

| Techniques | Descriptions |
|---|---|
| <i>Nucleic acid amplification and fingerprinting</i> | DNA is amplified by polymerase chain reaction (PCR) while cDNA is synthesized from the extracted RNA. The amplicons can be analyzed with fingerprinting techniques such as denaturing gradient gel electrophoresis (DGGE), terminal restriction fragment analysis (T-RFLP), amplified rDNA restriction analysis (ARDRA) and BOX-PCR. |
| <i>Real-time PCR (RT-PCR)</i> | RT-PCR is based on the monitoring of PCR reaction using fluorescent reporter molecules. Particularly, analysis of the exponential curve of the PCR reaction allows the determination of the amount of starting material. Different detections methods have been developed such as SYBR green I, a fluorescent dye binding double-stranded DNA, and Taqman probes. |
| <i>Sequencing</i> | Determination of nucleotides sequence in single stranded DNA or cDNA using Sanger or Pyrosequencing methods. |
| <i>Chromatography</i> | Chromatography allows for separation and isolation of chemical mixture by suspending compounds in solvent and separating mobile fraction from stationary fraction through a column. The proportions of the mixture provide quantitative information, and the separated fractions can be used for further analysis. Several types of chromatography have been described: thin layer chromatography (TLC), liquid chromatography (LC), gas chromatography (GC), high performance liquid chromatography (HPLC), ultra high pressure liquid chromatography (UPLC) |
| <i>Mass spectrometry (MS)</i> | MS identifies the chemical composition of molecules based on the mass-charge ratio of charged particles. Stable Isotope Probing (SIP) with MS is being used to track a stable isotope atom from a particular plant substrate into microbial cells <i>in situ</i> . Electron-spray-ionization (ESI)-MS is a technique for measuring chemical speciation in root exudates metabolomics. Quadrupole time-of-flight (Q-TOF) hybrid MS consists of both analyzers resulting in high sensitivity in product ion scanning. |
| <i>Phospholipid fatty acid (PLFA)</i> | PLFA identifies phospholipids in cellular membranes that are esterified to fatty acids. GC is used to identify the fatty acids, and information regarding microbial biomass, metabolic status and community composition can be obtained. |
| <i>Microscopy</i> | The use of microscopy for visualization of plant-associated microbes provides useful spatial information that can be used to determine the functional relationships. Confocal laser scanning microscopy (CLSM) is used to obtain high-resolution optical sectioning images of fluorescent samples. Transmission electron microscopy (TEM) bombards ultra thin specimens with electrons in order to obtain images at higher resolution than light microscopes. |
| <i>Fourier Transform Infrared Spectroscopy (FTIR)</i> | FTIR generates an infrared absorption spectrum specific for a type of chemical bond thus allowing for the identification of molecules. |
| <i>Genetic modifications</i> | Manipulating specific genes by both genetic engineering and mutation is used to validate previous observations regarding gene functions. Transgenic plants and recombinant microbes have been used. |
| <i>Nuclear magnetic resonance (NMR)</i> | The technique measures physical resonance and is used to determine the structure of the chemical. |



| Applications | Techniques | References |
|--|--|---|
| Plant Growth Promotion – N-fixing endophytes | Isotope analysis using MS, GFP labeled <i>Klebsiella pneumoniae</i> 342 (Kp342), genetic manipulation of Kp342, RT-PCR | Iniguez et al., 2004; James et al., 2002; You et al., 2005. |
| Biocontrol - <i>Pseudomonas</i> suppression of <i>Venturia inaequalis</i> | CCD camera, CSLM, RT-PCR, cDNA libraries, QTOF-MS | Gau et al., 2002; Singh et al., 2004; Degenhardt et al., 2005; Kurkuoglu et al., 2007 |
| Biocontrol- resistance to <i>Xanthomonas</i> sp. and <i>Xylella fastidiosa</i> | Recombinant <i>E. coli</i> and <i>Pseudomonas</i> spp. | Newman et al., 2008 |
| Remediation – Toluene and BTEX | Recombinant <i>Burkholderia cepacia</i> , BOX-PCR genomic DNA fingerprinting, sequencing, GC-MS | Barac et al., 2004; Taghavi et al., 2005; Moore et al., 2005 |
| Plant Growth Promotion- phytohormone | Ethylene measurement using GC | Mayak et al., 2004a; 2004b |
| Bioactive compounds production – Tashinone | Hairy roots tissue culture, HPLC | Wu et al., 2007 |
| Bioactive compounds production – Rhizoxin | PCR, HPLC-MS, laser microscopy, ESI-MS, NMR, transmission electron microscopy | Partida-Martinez and Hertweck, 2005 |
| Bioactive compounds production – Essential oil | Box-PCR genomic DNA fingerprinting, GC-MS, RT-PCR, CSLM and TEM | Del Giudice et al., 2008 |
| Remediation – Lindane | PCR, sequencing | Böltner et al., 2008 |
| Remediation – PCB | Transgenic Arabidopsis and <i>P. putida</i> mutant, reverse-phase HPLC, ESI-MS, confocal microscopy | Narasimhan et al., 2003 |
| Remediation – 4CNB | GFP-labeled <i>Comamonas</i> sp., confocal microscopy, quantitative-competitive PCR, PCR-DGGE | Liu et al., 2007 |
| Remediation – Cadmium | Recombinant <i>P. putida</i> | Wu et al., 2006 |
| Remediation – Cadmium and TCE | Recombinant <i>Pseudomonas</i> and <i>Rhizobium</i> strains, GC | Lee et al., 2006 |
| Remediation – Cadmium and zinc | DNA extraction and partial 16S DNA sequencing | Kuffner et al., 2008 |
| Remediation – PCB and arsenic | Recombinant <i>P. fluorescens</i> , PCR | Ryan et al., 2007 |
| Remediation – PCB | Recombinant <i>P. fluorescens</i> , CSLM, HPLC, PCR, sequencing | Villacieros et al., 2005; Leigh et al., 2006 |
| Carbon sequestration | PLFA, Stable isotope probing | Bailey et al., 2002; Heinemeyer et al., 2007; Verburg et al., 2004; Bazot et al., 2006; Leake et al., 2006; Bremer et al., 2008; Pascal et al., 2006; Ingram et al., 2008 |

Figure 1. Examples of developing applications with plant-microbe interactions. The line separates interactions located in the shoot and root (279×215 mm, 600×600 dpi).