

LA-UR 95-2066

Conf 951006 --13

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36.

TITLE: BENCHMARKING THE LAHETTM FISSION MODELS

AUTHOR(S): R.E. Prael

SUBMITTED TO: ANS Winter Meeting, San Francisco, CA, 10/30 - 11/2/95

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

Los Alamos Los Alamos National Laboratory
Los Alamos, New Mexico 87545
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

**Portions of this document may be illegible
in electronic image products. Images are
produced from the best available original
document.**

Benchmarking the LAHETTM Fission Models

R. E. Prael (LANL)

Introduction

There has been considerable interest in improving the fission models in the LAHET¹ Monte Carlo code for the transport and interaction of nucleons, pions, muons, light ions, and antinucleons. Although subactinide fission contributes little to neutron production in lead or tungsten targets, it can be significant for simulation of target activation and fission product contamination. The availability of new data permits new comparisons to be made between experiment and calculation.

Benchmarking the Fission Models

Previous efforts² at benchmarking the models have shown that subactinide fission is not well estimated and that there is considerable sensitivity to the physics options employed in conjunction with the fission model. The question of model sensitivity is not addressed here. The examples of LAHET calculations shown all used the default LAHET physics options, e. g., Bertini intranuclear cascade, the RAL fission models, and the Gilbert-Cameron-Cook-Ignatyuk energy dependent level density; the multistage preequilibrium model was not used.

Since the absolute ^{235}U fission cross section is used as a standard to extract the fission cross section from fission cross section ratio measurements, it plays a central

role in testing the calculational capabilities. The "final" LANL³ experimental results have been made available for ²³⁵U fission. The comparison with the LAHET calculation shows the results are within 10% above 80 MeV, but there is considerable divergence below. However, the general features of the cross section are followed by the model to low energies. In addition, the real test of the model is a comparison of the fission yields (as a fraction of the nonelastic cross section), since that is exactly what the fission models predict. In LAHET, the prediction of the nonelastic reaction rate, and therefore the nonelastic cross section, is a function only of the intranuclear cascade model used.

Subactinide fission is much less well represented by the present model, but new experimental results^{4,5} show that the discrepancies may not be as large as previously indicated. Comparisons are shown in figures 1 and 2 for natural Pb and ²⁰⁹Bi. The comparison for Pb shows that although calculated results are everywhere low, the calculated cross sections are within a factor of 2 above 100 MeV and in good agreement near 150 MeV. For ²⁰⁹Bi, the agreement is considerably better. The behavior of the difference suggests that modification of the fission barrier within the model may well lead to improvement.

Summary

One must certainly conclude that the extension of the use of evaluated data libraries for proton and neutrons to 100 MeV or 150 MeV would eliminate most of the difficulties associated with developing good fission models for use with intranu-

clear cascade-evaporation codes! In any case, the proposed use in LAHET of a global nucleon-nucleus optical potential to determine the elastic and nonelastic reaction rates would allow study of the fission models to focus on the calculated fission yield rather than on the calculated fission rate.

However, there appears to be a growing number of consistent experimental measurements of Pb and Bi fission that may well be a basis for improvement of the subactinide fission model in LAHET. The latter is not an easy task, since, to be done properly, the fission model must be consistent with the intranuclear cascade model and the succeeding preequilibrium and evaporation models. The subactinide fission model is a good candidate for modification, since it is a true model with well-defined physical quantities employed; the actinide fission model is less treatable, since it only employs tabulations of observed ratios of fission- to neutron-widths.

1. Richard E. Prael and Henry Lichtenstein, "User Guide to LCS: The LAHET Code System", LA-UR-89-3014, Los Alamos National Laboratory (September 1989).
2. R. E. Prael, "The LAHET Code System: Introduction, Development, and Benchmarking," Transactions of the Workshop on Simulating Accelerator Radiation Environments, Santa Fe, Jan. 1993.
3. P. Lisowski, LANL, personal communication.
4. P. Staples and K.B. Morley, LANL, personal communication.
5. V. P. Eismont *et al.*, "Relative and Absolute Neutron-induced Fission Cross Sections of ^{208}Pb , ^{209}Bi , and ^{238}U in the Intermediate Energy Region", to be published.

LAHET™ is a trademark of the Regents of the University of California
and the Los Alamos National Laboratory.




Figure 1: Neutron induced natural Pb fission cross section. LAHET calculations with Bertini INC and default physics options (line).




Figure 2: Neutron induced ^{209}Bi fission cross section. LAHET calculations with Bertini INC and default physics options (line).