

LA-UR-03-2189

Approved for public release;
distribution is unlimited.

| | |
|----------------------|---|
| <i>Title:</i> | Quantum Simulations of Physics Problems |
| <i>Author(s):</i> | Rolando Somma, Gerardo Ortiz, Emanuel Knill, and James Gubernatis |
| <i>Submitted to:</i> | http://lib-www.lanl.gov/cgi-bin/getfile?00852379.pdf |

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-36. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Quantum Simulations of Physics Problems

Rolando Somma, Gerardo Ortiz, Emanuel Knill, and James Gubernatis

Los Alamos National Laboratory, Los Alamos, USA

(Dated: June 6, 2003)

Abstract

If a large Quantum Computer (QC) existed today, what type of physical problems could we efficiently simulate on it that we could not efficiently simulate on a classical Turing machine? In this paper we argue that a QC could solve some relevant physical “questions” more efficiently. The existence of one-to-one mappings between different algebras of observables or between different Hilbert spaces allow us to represent and imitate any physical system by any other one (e.g., a bosonic system by a spin-1/2 system). We explain how these mappings can be performed, and we show quantum networks useful for the efficient evaluation of some physical properties, such as correlation functions and energy spectra.

PACS numbers: Keywords: quantum mechanics, quantum computing, identical particles, spin systems, generalized Jordan-Wigner transformations

I. INTRODUCTION

Quantum simulation of physical systems on a QC has acquired importance during the last years since it is believed that QCs can simulate quantum physics problems more efficiently than their classical analogues [1–4]: The number of operations needed for deterministically solving a quantum many-body problem on a classical computer (CC) increases exponentially with the number of degrees of freedom of the system.

In quantum mechanics, each physical system has associated with it a language of operators and an algebra realizing this language, and can be considered as a possible model of quantum computation [3]. As we discussed in a previous paper [5], the existence of one-to-one mappings between different languages (e.g., the Jordan-Wigner transformation [6, 7] that maps fermionic operators onto spin-1/2 operators) and between quantum states of different Hilbert spaces allows the quantum simulation of one physical system by any other one. For example, a liquid nuclear magnetic resonance QC (NMR) can simulate a system of ^4He atoms (hard-core bosons) because an isomorphic mapping between the two algebras of observables exists.

The existence of mappings between operators allows us to construct quantum networks from sets of elementary gates, to which we map the operators of our physical system. An important remark is that these mappings can be performed efficiently: we need a number of steps that scales polynomially with the system size. However, this fact alone is not sufficient to establish that any quantum problem can be solved efficiently. One needs to show that all steps involved in the simulation (i.e., preparation of the initial state, evolution, measurement, and measurement control) can be performed with polynomial complexity. For example, the number of different eigenvalues in the two-dimensional Hubbard model scales exponentially with the system size, so QC algorithms for obtaining its energy spectrum will also require a number of operations that scales exponentially with the system size [5].

Generically, given a physical system over which we have quantum control, its operator algebra can be associated with a possible model of computation [5, 8]. In this paper, we consider the simulation of any physical system by the standard model of quantum computation (spin-1/2 system), since this might be the language needed for the practical implementation of the quantum algorithms (e.g., NMR). Therefore, the complexity of the quantum algorithms is analyzed from the point of view of the number of resources (elementary gates) needed for their implementation in the language of the standard model. If another model of computation were used, one would follow

the same qualitative steps (presented in this paper) although the mappings and network structure would be different.

The main purpose of this manuscript is to present a pedagogical review of previous works ([3, 5]) on the simulation of physical phenomena using a QC, including a new section explaining the simulation of systems of particles obeying canonical bosons’ commutation relations. We organized the paper in the following way: In section II we describe the standard model of quantum computation (spin-1/2 system). Section III shows the mappings between physical systems governed by a generalized Pauli’s exclusion principle (fermions, etc.) and the standard model, giving examples of algorithms for the first two steps (preparation of the initial state and evolution) of the quantum simulation. In section IV we develop similar steps for the simulation of quantum systems whose language has an infinite-dimensional representation, thus, there is no exclusion principle (e.g., canonical bosons). In section V we explain the measurement process used to extract information of some relevant and generic physical properties, such as correlation functions and energy spectra. We conclude with a discussion about efficiency and quantum errors (section VI) and a summary about the general statements (section VII).

II. STANDARD MODEL OF QUANTUM COMPUTATION

In the standard model of quantum computation, the fundamental unit is the *qubit*, represented by a two level quantum system with states of the form $|a\rangle = a|0\rangle + b|1\rangle$. For a spin-1/2 particle, for example, the two “levels” are the two different orientations of the spin, $|\uparrow\rangle = |0\rangle$ and $|\downarrow\rangle = |1\rangle$. In this model, the algebra assigned to a system of N -qubits is built upon the Pauli spin-1/2 operators σ_x^j , σ_y^j and σ_z^j acting on the j -th qubit (individual qubit). The commutation relations for these operators satisfy an $\bigoplus_{i=1}^N su(2)_i$ algebra defined by $(\mu, \nu, \lambda = x, y, z)$

$$[\sigma_\mu^j, \sigma_\nu^k] = 2i\delta_{jk}\epsilon_{\mu\nu\lambda}\sigma_\lambda^j, \quad (1)$$

where $[A, B] = AB - BA$ and $\epsilon_{\mu\nu\lambda}$ is the totally anti-symmetric Levi-Civita symbol. Sometimes it is useful to write the commutation relations in terms of the raising and lowering spin-1/2 operators

$$\sigma_\pm^j = \frac{\sigma_x^j \pm i\sigma_y^j}{2}. \quad (2)$$

Any computation on a QC is represented by a unitary operator U that evolves some initial state in a way that satisfies the time-dependent Schrödinger equation for some Hamiltonian $H(t)$. Any

unitary operation (evolution) U applied to a system of N qubits can be implemented on a QC by decomposing it into single qubit rotations $R_\mu^j(\vartheta) = e^{-i\frac{\vartheta}{2}\sigma_\mu^j}$ by an angle ϑ about the μ axis, and two qubits Ising interactions $R_{z^j,z^k} = e^{i\omega\sigma_z^j\sigma_z^k}$. (These elementary operations constitute a set of elementary gates). That this decomposition is possible is an important result of quantum information. It implies that with the given elementary gates one can perform universal quantum computation. Other, equivalent sets of elementary gates can be chosen. For example, we could also perform universal quantum computation with single qubit rotations and C-NOT gates [9] or even with different control Hamiltonians. The crucial point is that we need to have quantum control over those elementary operations in the real physical system. The complexity of a computation is given by the number of (elementary) gates used. Which universal gate set is used does not significantly affect the complexity of a computation, provided that the gates act on at most two qubits at a time.

In the following, we will write down our algorithms in terms of single qubit rotations and two qubits Ising interactions, since this is the language needed for the implementation of the algorithms, for example, in a liquid NMR QC. If we used a different set of local, universal elementary gates, our main results still hold but with modified quantum networks.

As an example of such decompositions, we consider the unitary operator $U(t) = e^{iHt}$, where $H = \alpha\sigma_x^1\sigma_z^2\sigma_x^3$ represents a time-independent Hamiltonian. After some simple calculations [3, 5] we decompose U into elementary gates (one qubit rotations and two qubits interactions) in the following way

$$U(t) = e^{i\alpha\sigma_x^1\sigma_z^2\sigma_x^3t} = e^{-i\frac{\pi}{4}\sigma_y^3}e^{i\frac{\pi}{4}\sigma_z^1\sigma_z^3}e^{i\frac{\pi}{4}\sigma_x^1}e^{i\alpha\sigma_z^1\sigma_z^2t}e^{-i\frac{\pi}{4}\sigma_x^1}e^{-i\frac{\pi}{4}\sigma_z^1\sigma_z^3}e^{i\frac{\pi}{4}\sigma_y^3}. \quad (3)$$

This decomposition is shown in Fig. 1, where the quantum network representation is displayed. In the same way, we could also decompose an operator $U'(t) = e^{-i\alpha\sigma_y^1\sigma_z^2\sigma_y^3t}$ using similar steps but replacing $\sigma_x^j \leftrightarrow \sigma_y^j$ in the right hand side of Eq. 3.

III. SIMULATION OF FERMIONIC SYSTEMS

As discussed in the Introduction, quantum simulations require simulations of systems with diverse degrees of freedom and particle statistics. Fermionic systems are governed by Pauli's exclusion principle, which implies that no more than one fermion can occupy the same quantum state at the same time. In this way, the Hilbert space of quantum states that represent a system of




FIG. 1: Decomposition of the unitary operator $U(t) = e^{i\alpha\sigma_x^1\sigma_z^2\sigma_x^3 t}$ into elementary single qubit rotations and two qubits interactions. Time t increases from left to right.

fermions in a solid is finite-dimensional (2^N for spinless fermions, where N is the number of sites or modes in the solid), and one could imagine in the existence of one-to-one mappings between the fermionic and Pauli's spin-1/2 algebras. Similarly, any language which involves operators with a finite-dimensional representation (e.g., hard-core bosons, higher irreps of $su(2)$, etc.) can be isomorphically mapped onto the standard model language [6–8].

In the second quantization representation, the (spinless) fermionic operators c_j^\dagger (c_j) are defined as the creation (annihilation) operators of a fermion in the j -th mode ($j = 1, \dots, N$). Due to the Pauli's exclusion principle and the antisymmetric nature of the fermionic wave function under the permutation of two fermions, the fermionic algebra is given by the following commutation

relations

$$\{c_i, c_j\} = 0, \{c_i^\dagger, c_j\} = \delta_{ij} \quad (4)$$

where $\{, \}$ denotes the anticommutator (i.e., $\{A, B\} = AB + BA$).

The Jordan-Wigner transformation [6, 7] is the isomorphic mapping that allows the description of a fermionic system by the standard model

$$c_j \rightarrow \left(\prod_{l=1}^{j-1} -\sigma_z^l \right) \sigma_-^j \quad (5)$$

$$c_j^\dagger \rightarrow \left(\prod_{l=1}^{j-1} -\sigma_z^l \right) \sigma_+^j, \quad (6)$$

where σ_μ^i are the Pauli operators defined in section II. One can easily verify that if the operators σ_μ^i satisfy the $su(2)$ commutation relations (Eq. 1), the operators c_j^\dagger and c_j obey Eqs. 4.

It is important to remark that the Jordan-Wigner transformation is an isomorphic mapping between operator algebras and is independent of the Hamiltonian of the fermionic system we want to simulate. Different Hamiltonians establish different connections (connectivity) between fermionic modes. Historically, Eqs. 5 and 6 correspond to lattices in one space dimension. Nevertheless, it is also valid for lattice systems in any dimension, since the set of modes j is countable. In particular, the set of all ordered p -tuples of integers can be placed in one-to-one correspondence with the set of integers. For example, the simulation of a two dimensional fermionic lattice system can be done by re-mapping each mode (l, m) into a new set of modes as $j = m + (l - 1)N_x$, where $[l = 1 \dots N_y]$ and $[m = 1 \dots N_x]$ are integer numbers that refer to the position of a site in the lattice, and N_x and N_y are the number of sites (modes) in the x and y direction respectively. The point is, however, that the Hamiltonian in the new mode-label connects *sites* separated by distances of order $O(N_x)$, but the complexity of the simulation remains polynomial. In particular, the simulation of the two dimensional Hubbard model is shown in Ref. [5].

We now need to show how to simulate a fermionic system by a QC. Just as for a simulation on a CC, the quantum simulation has three basic steps: the preparation of an initial state, the evolution of this state, and the measurement of a relevant physical property of the evolved state. We will now explain the first two steps, postponing the third until section V.

A. Preparation of the initial state

In the most general case, any quantum state $|\psi\rangle$ of N_e fermions can be written as a linear combination of Slater determinants $|\phi_\alpha\rangle$

$$|\psi\rangle = \sum_{\alpha=1}^L g_\alpha |\phi_\alpha\rangle, \quad (7)$$

where

$$|\phi_\alpha\rangle = \prod_{j=1}^{N_e} c_j^\dagger |\text{vac}\rangle \quad (8)$$

with the vacuum state $|\text{vac}\rangle$ defined as the state with no fermions: $c_j|\text{vac}\rangle = 0 \forall j$. In the spin language, $|\text{vac}\rangle = |\downarrow\downarrow\cdots\downarrow\rangle$.

The commutation relations of fermionic operators shown in Eqs. 4, where $c_i^\dagger c_j^\dagger = -c_j^\dagger c_i^\dagger$ if $i \neq j$, imply that the Slater determinants $|\phi_\alpha\rangle$ are antisymmetric under the permutation of any two fermions.

We can easily prepare the states $|\phi_\alpha\rangle$ by noticing that the quantum gate, represented by the unitary operator

$$U_m = e^{i\frac{\pi}{2}(c_m + c_m^\dagger)} \quad (9)$$

when acting on the vacuum state, creates a particle in the m -th mode. In other words, $U_m|\text{vac}\rangle = e^{i\frac{\pi}{2}c_m^\dagger}|\text{vac}\rangle$. Making use of the Jordan-Wigner transformation (Eqs. 5 and 6), we can write the operators U_m in the spin language

$$U_m = e^{i\frac{\pi}{2}\sigma_x^m \prod_{j=1}^{m-1} -\sigma_z^j}. \quad (10)$$

The successive application of N_e similar unitary operators will generate the state $|\phi_\alpha\rangle$ up to an irrelevant global phase.

A detailed preparation of the fermionic state $|\psi\rangle = \sum_{\alpha=1}^L g_\alpha |\phi_\alpha\rangle$ can be found in a previous work [3]. The basic idea is to use L extra (ancilla) qubits, then perform unitary evolutions controlled in the state of the ancillas, and finally perform a measurement of the z -component of the spin of the ancillas. In this way, the probability of successful preparation of $|\psi\rangle$ is $1/L$. (We need of the order of L trials before a successful preparation.)

Another important case is the preparation of a Slater determinant in a different basis than the one given before

$$|\phi_\beta\rangle = \prod_{j=1}^{N_e} d_j^\dagger |\text{vac}\rangle, \quad (11)$$

where the fermionic operators d_j^\dagger 's are related to the operators c_j^\dagger through the following canonical transformation

$$\vec{d}^\dagger = e^{iM} \vec{c}^\dagger \quad (12)$$

with $\vec{d}^\dagger = (d_1^\dagger, d_2^\dagger, \dots, d_N^\dagger)$, $\vec{c}^\dagger = (c_1^\dagger, c_2^\dagger, \dots, c_N^\dagger)$, and M is an $N \times N$ Hermitian matrix. Making use of Thouless's theorem [10], we observe that one Slater determinant evolves into the other, $|\phi_\beta\rangle = U|\phi_\alpha\rangle$, where the unitary operator $U = e^{-i\vec{c}^\dagger M \vec{c}}$ can be written in spin operators using the Jordan-Wigner transformation and can be decomposed into elementary gates [5], as described in section II. Since the number of gates scales polynomially with the system size (the operator $\vec{c}^\dagger M \vec{c}$ is a sum of a number of terms that scales polynomially with N), the state $|\phi_\beta\rangle$ can be efficiently prepared from the state $|\phi_\alpha\rangle$, as explained in the next section.

B. Evolution of the initial state

The second step in the quantum simulation is the evolution of the initial state. The unitary evolution operator of a time-independent Hamiltonian H is $U(t) = e^{iHt}$. In general, $H = K + V$ with K representing the kinetic energy and V the potential energy. Since we usually have $[K, V] \neq 0$, the decomposition of $U(t)$, written in the spin language through the Jordan-Wigner transformation (Eqs. 5 and 6), in terms of elementary gates (one qubit rotations and two qubits interactions), becomes complicated. To avoid this problem, we instead use a Trotter decomposition, so the evolution during a short period of time ($\Delta t = t/\mathcal{N}$ with $\Delta t \rightarrow 0$) is approximated. To order $O(\Delta t)$ (first order Trotter breakup)

$$U(t) = \prod_{g=1}^{\mathcal{N}} U(\Delta t), \quad (13)$$

$$U(\Delta t) = e^{iH\Delta t} = e^{i(K+V)\Delta t} \sim e^{iK\Delta t} e^{iV\Delta t}. \quad (14)$$

The potential energy V is usually a sum of commuting diagonal terms, and the decomposition of $e^{iV\Delta t}$ into elementary gates is straightforward. However, the kinetic energy K is usually a sum of noncommuting terms of the form $c_j^\dagger c_k + c_k^\dagger c_j$ (bilinear fermionic operators), so we need again to perform a Trotter approximation of the operator $e^{iK\Delta t}$. As an example of such a decomposition, we consider a typical term $e^{i(c_j^\dagger c_k + c_k^\dagger c_j)\Delta t}$ ($j < k$), when mapped onto the spin language gives

$$e^{-\frac{i}{2}(\sigma_x^j \sigma_x^k + \sigma_y^j \sigma_y^k) \prod_{l=j+1}^{k-1} (-\sigma_z^l)} = e^{-\frac{i}{2}\sigma_x^j \sigma_x^k \prod_{l=j+1}^{k-1} (-\sigma_z^l)} e^{-\frac{i}{2}\sigma_y^j \sigma_y^k \prod_{l=j+1}^{k-1} (-\sigma_z^l)}. \quad (15)$$

The decomposition of each term on the right hand side of Eq. 15 into elementary gates was already described in previous work [5]. In section II and Fig. 1, we also showed an example of such a decomposition for $j = 1$ and $k = 3$. It is important to mention that the required number of elementary gates scales polynomially with the length $|k - j|$. Notice that this step is not necessary for bosonic systems since no string of σ_z^l operators is involved (see section IV).

The accuracy of this method increases as Δt decreases, so we might require a large number of gates to perform the evolution with small errors. To overcome this problem, one could use a Trotter approximation of higher order in Δt [11]. Physical Hamiltonians with higher order products of creation and annihilation operators can also be efficiently simulated.

C. Generalization: simulation of anyonic systems

The concepts described in sections III A and III B can be easily generalized to other more general particle statistics, namely hard-core anyons. By “hard-core”, we mean that only zero or one particle can occupy a single mode (Pauli’s exclusion principle).

The commutation relations between the anyonic creation and annihilation operators a_i^\dagger and a_i , are given by

$$\begin{aligned} [a_i, a_j]_\theta &= [a_i^\dagger, a_j^\dagger]_\theta = 0, \\ [a_i, a_j^\dagger]_{-\theta} &= \delta_{ij}(1 - (e^{-i\theta} + 1)n_j), \\ [n_i, a_j^\dagger] &= \delta_{ij}a_j^\dagger, \end{aligned} \tag{16}$$

($i \leq j$) where $n_j = a_j^\dagger a_j$, $[\hat{A}, \hat{B}]_\theta = \hat{A}\hat{B} - e^{i\theta}\hat{B}\hat{A}$, with $0 \leq \theta < 2\pi$ defining the statistical angle. In particular, $\theta = \pi \bmod(2\pi)$ corresponds to canonical spinless fermions, while $\theta = 0 \bmod(2\pi)$ represents hard-core bosons.

In order to simulate this problem with a QC made of qubits, we need to apply the following isomorphic and efficient mapping between algebras

$$\begin{aligned} a_j^\dagger &= \prod_{l < j} \left[\frac{e^{-i\theta} + 1}{2} + \frac{e^{-i\theta} - 1}{2} \sigma_z^l \right] \sigma_+^j, \\ a_j &= \prod_{l < j} \left[\frac{e^{i\theta} + 1}{2} + \frac{e^{i\theta} - 1}{2} \sigma_z^l \right] \sigma_-^j, \\ n_j &= \frac{1}{2}(1 + \sigma_z^j), \end{aligned} \tag{17}$$

where the Pauli operators σ_μ^i were defined in section II, and since they satisfy Eq. 1, the corresponding commutation relations for the anyonic operators (Eqs. 16) are satisfied, too.

We can now proceed in the same way as in the fermionic case, writing our anyonic evolution operator in terms of single qubit rotations and two qubits interactions in the spin-1/2 language. As we already mentioned, anyon statistics have fermion and hard-core boson statistics as limiting cases. In the next section we relax the hard-core condition on the bosons and consider the important case of canonical bosons.

IV. SIMULATION OF BOSONIC SYSTEMS

Quantum computation is based on the manipulation of quantum systems that possess a finite number of degrees of freedom (e.g., qubits). From this point of view, the simulation of bosonic systems appears to be impossible, since the non-existence of an exclusion principle implies that the Hilbert space used to represent bosonic quantum states is infinite-dimensional; that is, there is no limit to the number of bosons that can occupy a given mode j . However, sometimes we might be interested in simulating and studying properties such that the use of the complete Hilbert space is unnecessary, and only a finite sub-basis of states is sufficient. This is the case for physical systems with interactions given by the Hamiltonian

$$H = \sum_{i,j=1}^N \alpha_{ij} b_i^\dagger b_j + \beta_{ij} \hat{n}_i \hat{n}_j, \quad (18)$$

where the operators b_i^\dagger (b_i) create (destroy) a boson at site i , and $\hat{n}_i = b_i^\dagger b_i$ is the number operator; that is

$$b_i^\dagger |n_1, n_2, \dots, n_i, \dots, n_N\rangle = \sqrt{n_i + 1} |n_1, n_2, \dots, n_i + 1, \dots, n_N\rangle, \quad (19)$$

$$b_i |n_1, n_2, \dots, n_i, \dots, n_N\rangle = \sqrt{n_i} |n_1, n_2, \dots, n_i - 1, \dots, n_N\rangle, \quad (20)$$

$$\hat{n}_i |n_1, n_2, \dots, n_i, \dots, n_N\rangle = n_i |n_1, n_2, \dots, n_i, \dots, n_N\rangle, \quad (21)$$

where the bosonic state $|n_1, n_2, \dots, n_i, \dots, n_N\rangle$ represents a quantum state with n_i bosons in the i -th site.

The space dimension of the lattice is encoded in the parameters α_{ij} and β_{ij} of the Hamiltonian. Obviously, the total number of bosons N_P in the system is conserved, and we restrict ourselves to work with a finite sub-basis of states, where the dimension depends on the value of N_P .

The corresponding bosonic commutation relations (in an infinite-dimensional Hilbert space) are

$$[b_i, b_j] = 0, [b_i, b_j^\dagger] = \delta_{ij}. \quad (22)$$

However, if the operators b_i^\dagger are restricted to the finite basis of states represented by $\{|n_1, n_2, \dots, n_N\rangle$ with $n_i = 0, \dots, N_P\}$, where N_P is the maximum number of bosons per site, then they have the following matrix representation (see Eq. 19)

$$\bar{b}_i^\dagger = \mathbb{1} \otimes \cdots \otimes \mathbb{1} \otimes \underbrace{\hat{b}^\dagger}_{i^{\text{th}} \text{ factor}} \otimes \mathbb{1} \otimes \cdots \otimes \mathbb{1} \quad (23)$$

where \otimes indicates the usual tensorial product between matrices, and the $(N_P + 1) \times (N_P + 1)$ dimensional matrices $\mathbb{1}$ and \hat{b}^\dagger are given by

$$\mathbb{1} = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}, \quad \hat{b}^\dagger = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & \sqrt{2} & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \sqrt{N_P} & 0 \end{pmatrix}. \quad (24)$$

It is important to note that in this finite basis, the commutation relations of the \bar{b}_i^\dagger differ from the standard bosonic ones (Eq. 22) [8]

$$[\bar{b}_i, \bar{b}_j] = 0, \quad [\bar{b}_i, \bar{b}_j^\dagger] = \delta_{ij} \left[1 - \frac{N_P + 1}{N_P!} (\bar{b}_i^\dagger)^{N_P} (\bar{b}_i)^{N_P} \right], \quad (25)$$

and clearly $(\bar{b}_i^\dagger)^{N_P+1} = 0$.

As we mentioned in the Introduction, our goal is to simulate any physical system in a QC made of qubits. For this purpose, we need to map the bosonic algebra into the spin-1/2 language. However, since Eqs. 25 imply that the linear span of the operators \bar{b}_i^\dagger and \bar{b}_i is not closed under the bracket (commutator), a direct mapping between the bosonic algebra and the spin-1/2 algebra (such as the case of the Jordan-Wigner transformation between the fermionic and spin-1/2 algebra) is not possible. Therefore, we could imagine in a one-to-one mapping between the bosonic and spin-1/2 quantum states, instead of an isomorphic mapping between algebras. We will now show one possible such mapping of quantum states.

We start by considering only the i -th site in the chain. Since this site can be occupied with at most N_P bosons, it is possible to associate an $N_P + 1$ qubits quantum state to each particle number

state, in the following way

$$\begin{aligned}
|0\rangle_i &\leftrightarrow |\uparrow_0\downarrow_1\downarrow_2 \cdots \downarrow_{N_P}\rangle_i \\
|1\rangle_i &\leftrightarrow |\downarrow_0\uparrow_1\downarrow_2 \cdots \downarrow_{N_P}\rangle_i \\
|2\rangle_i &\leftrightarrow |\downarrow_0\downarrow_1\uparrow_2 \cdots \downarrow_{N_P}\rangle_i \\
&\vdots \quad \vdots \\
|N_P\rangle_i &\leftrightarrow |\downarrow_0\downarrow_1\downarrow_2 \cdots \uparrow_{N_P}\rangle_i
\end{aligned} \tag{26}$$

where $|n\rangle_i$ denotes a quantum state with n bosons in site i . Therefore, we need $N(N_P + 1)$ qubits for the simulation (where N is the number of sites). In Fig. 2 we show an example of this mapping for a quantum state with 7 bosons in a chain of 5 sites, where the maximum number of bosons per site is $N_P = 3$.

By definition (see Eqs. 19, 23, and 24) $\bar{b}_i^\dagger |n\rangle_i = \sqrt{n+1} |n+1\rangle_i$, so the operator

$$\bar{b}_i^\dagger = \sum_{n=0}^{N_P-1} \sqrt{n+1} \sigma_-^{n,i} \sigma_+^{n+1,i}, \tag{27}$$

where the pair (n, i) indicates the qubit n that represents the i -th site, acts in the N_P+1 qubits states of Eqs. 26 as $\bar{b}_i^\dagger |\downarrow_0 \cdots \downarrow_{n-1} \uparrow_n \downarrow_{n+1} \cdots \downarrow_{N_P}\rangle_i = \sqrt{n+1} |\downarrow_0 \cdots \downarrow_n \uparrow_{n+1} \downarrow_{n+2} \cdots \downarrow_{N_P}\rangle_i$. Then, its matrix representation in this basis is the same matrix representation of b_i^\dagger in the basis of bosonic states. Similarly, the number operator can be written

$$\bar{n}_i = \sum_{n=0}^{N_P} n \frac{\sigma_z^{n,i} + 1}{2}, \tag{28}$$

and act as $\bar{n}_i |\downarrow_0 \cdots \downarrow_{n-1} \uparrow_n \downarrow_{n+1} \cdots \downarrow_{N_P}\rangle_i = n |\downarrow_0 \cdots \downarrow_n \uparrow_{n+1} \downarrow_{n+2} \cdots \downarrow_{N_P}\rangle_i$. Notice that $[\bar{b}_i^\dagger, \sum_{n=0}^{N_P} \sigma_z^{n,i}] = 0$, which means that these operators conserve the total z -component of the spin and, thus, always keep states within the same subspace.

We can now write down the Hamiltonian in Eq. 18 in the spin-1/2 algebra as

$$H = \sum_{i,j=1}^N \alpha_{ij} \bar{b}_i^\dagger \bar{b}_j + \beta_{ij} \bar{n}_i \bar{n}_j, \tag{29}$$

where the operators \bar{b}_i^\dagger (\bar{b}_i) are given by Eq. 27 and \bar{n}_i by Eq. 28. In this way, we obtain physical properties of the bosonic system (such as the mean value of an observable, the mean value of the evolution operator, etc.) in a QC made of qubits. It is important to note that the type of Hamiltonian given by Eq. 18 is not the only one simulatable using the described method. The only constraint is a fixed maximum number of bosons per site (or mode).




FIG. 2: Mapping of the bosonic state $|\phi_\alpha\rangle$, of a chain of 5 sites and 7 bosons ($N_P = 3$), into a spin-1/2 state (Eq. 26).

A. Preparation of the initial state

As in the fermionic case, the most general bosonic state of an N sites quantum system with N_P bosons can be written as a linear combination of product states like

$$|\phi_\alpha\rangle = K(b_1^\dagger)^{n_1}(b_2^\dagger)^{n_2} \cdots (b_N^\dagger)^{n_N} |\text{vac}\rangle, \quad (30)$$

where K is a normalization factor, n_i is the number of bosons at site i ($\sum_{i=1}^N n_i = N_P$), and $|\text{vac}\rangle$ is the boson vacuum state (no particle state). Using the mapping described in Eq. 26, we can write the vacuum state in the spin language as $|\text{vac}\rangle = |\uparrow_0 \downarrow_1 \cdots \downarrow_{N_P}\rangle_1 \otimes \cdots \otimes |\uparrow_0 \downarrow_1 \cdots \downarrow_{N_P}\rangle_N$ and $|\phi_\alpha\rangle = |\downarrow_0 \cdots \uparrow_{n_1} \cdots \downarrow_{N_P}\rangle_1 \otimes \cdots \otimes |\downarrow_0 \cdots \uparrow_{n_N} \cdots \downarrow_{N_P}\rangle_N$ (see Fig. 2 for an example). Therefore, the preparation of $|\phi_\alpha\rangle$ in a QC made of qubits is an easy process: only N spins are flipped from the fully polarized state, where all spins are pointing down.

The preparation of a bosonic initial state of the form $|\psi\rangle = \sum_{\alpha=1}^L g_\alpha |\phi_\alpha\rangle$ is realized as in the fermionic case. Again, we need to add L ancillas (extra qubits), perform controlled evolutions on

their states, and finally perform a measurement of an spin component [3].

B. Evolution of the initial state

The basic idea is to use the first order Trotter approximation (see the fermionic case) to separate those terms of the Hamiltonian that belong to the kinetic energy K , from the ones that belong to the potential energy V ($H = K + V$, $[K, V] \neq 0$), i.e.,

$$e^{iH\Delta t} \sim e^{iK\Delta t} e^{iV\Delta t}. \quad (31)$$




FIG. 3: Decomposition of the unitary operator $U(t) = e^{\frac{i}{8}\sigma_x^1\sigma_y^2\sigma_y^3\sigma_x^4t}$ into single qubit rotations and two qubits interactions. Time t increases from left to right.

In general, K is a sum of non commuting terms of the form $b_k^\dagger b_l + b_l^\dagger b_k$, and we need to perform another first order Trotter approximation to decompose it into elementary gates (in the

spin language). Then, a typical term $e^{i(b_k^\dagger b_l + b_l^\dagger b_k)t}$ when mapped onto the spin language (Eq. 27) gives

$$\exp\left[\frac{it}{8} \sum_{n,n'=0}^{N_P-1} \sqrt{(n+1)(n'+1)} \left[(\sigma_x^{n,k} \sigma_x^{n+1,k} + \sigma_y^{n,k} \sigma_y^{n+1,k}) (\sigma_x^{n',l} \sigma_x^{n'+1,l} + \sigma_y^{n',l} \sigma_y^{n'+1,l}) \right. \right. \\ \left. \left. + (\sigma_x^{n,k} \sigma_y^{n+1,k} - \sigma_y^{n,k} \sigma_x^{n+1,k}) (\sigma_x^{n',l} \sigma_y^{n'+1,l} - \sigma_y^{n',l} \sigma_x^{n'+1,l}) \right] \right], \quad (32)$$

where N_P is the number of bosons. The terms in the exponent of Eq. 32 commute with each other, so the decomposition into elementary gates becomes straightforward. As an example (see Fig.3), we consider a system of two sites with maximal one boson per site ($N_P = 1$). We need then $2(1+1) = 4$ qubits for the simulation, and Eq. 27 implies that $\bar{b}_1^\dagger = \sigma_-^{0,1} \sigma_+^{1,1}$ and $\bar{b}_2^\dagger = \sigma_-^{0,2} \sigma_+^{1,2}$. Then, $e^{i(b_k^\dagger b_l + b_l^\dagger b_k)t}$ becomes

$$\exp\left(\frac{it}{8} \sigma_x^{0,1} \sigma_x^{1,1} \sigma_x^{0,2} \sigma_x^{1,2}\right) \times \exp\left(\frac{it}{8} \sigma_x^{0,1} \sigma_x^{1,1} \sigma_y^{0,2} \sigma_y^{1,2}\right) \times \exp\left(\frac{it}{8} \sigma_y^{0,1} \sigma_y^{1,1} \sigma_x^{0,2} \sigma_x^{1,2}\right) \quad (33) \\ \times \exp\left(\frac{it}{8} \sigma_y^{0,1} \sigma_y^{1,1} \sigma_y^{0,2} \sigma_y^{1,2}\right) \times \exp\left(\frac{it}{8} \sigma_y^{0,1} \sigma_x^{1,1} \sigma_y^{0,2} \sigma_x^{1,2}\right) \times \exp\left(-\frac{it}{8} \sigma_y^{0,1} \sigma_x^{1,1} \sigma_x^{0,2} \sigma_y^{1,2}\right) \\ \times \exp\left(-\frac{it}{8} \sigma_x^{0,1} \sigma_y^{1,1} \sigma_y^{0,2} \sigma_x^{1,2}\right) \times \exp\left(\frac{it}{8} \sigma_x^{0,1} \sigma_y^{1,1} \sigma_x^{0,2} \sigma_y^{1,2}\right),$$

where the decomposition of each of the terms in Eq. 33 in elementary gates can be done using the methods described in previous works [3, 5]. In particular, in Fig. 3 we show the decomposition of the term $\exp\left(\frac{it}{8} \sigma_x^{0,1} \sigma_y^{1,1} \sigma_y^{0,2} \sigma_x^{1,2}\right)$, where the qubits were relabeled as $(n, j) \equiv n + 2j - 1$ (e.g., $(0, 1) \rightarrow 1$).

On the other hand, it is important to mention that the number of operations involved in the decomposition is not related to the distance between the sites k and l , as in the fermionic case (see section III B). Since typical bosonic Hamiltonians (like the one in Eq. 18) involve a number of terms which scales polynomially in N , their simulation complexity will scale polynomially in N and N_P .

V. MEASUREMENT: CORRELATION FUNCTIONS AND ENERGY SPECTRA

In previous work [3, 5] we introduced an efficient algorithm for the measurement of correlation functions in quantum systems. The idea is to make an indirect measurement; that is, we prepare an ancilla qubit (extra qubit) in a given initial state, then interact with the system whose properties one wants to measure, and finally we measure some observable of the ancilla to obtain information about the system. In particular, we might be interested in the measurement of dynamical

correlation functions of the form

$$G(t) = \langle \psi | T^\dagger A_i^\dagger T B_j \psi \rangle \quad (34)$$

where A_i and B_j are unitary operators (any operator can be decomposed in a unitary operator basis as $A = \sum_i \alpha_i A_i$, $B = \sum_j \beta_j B_j$), $T = e^{-iHt}$ is the time evolution operator of a time-independent Hamiltonian H , and $|\psi\rangle$ is the state of the system whose correlations one wants to determine. If we were interested in the evaluation of spatial correlation functions, we would replace the evolution operator T by the space translation operator. In Fig. 4 we show the quantum network for the evaluation of $G(t)$. As explained before [3, 5], the initial state (ancilla plus system) has to be prepared in the quantum state $|+\rangle_a \otimes |\psi\rangle$ (where a denotes the ancilla qubit and $|+\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}$). Additionally, we have to perform an evolution (unitary operation) in the following three steps: i) a controlled evolution in the state $|1\rangle$ of the ancilla $C-B = |0\rangle\langle 0| \otimes I + |1\rangle\langle 1| \otimes B_j$, ii) a time evolution T , and iii) a controlled evolution in the state $|0\rangle$ of the ancilla $C-A = |0\rangle\langle 0| \otimes A_i + |1\rangle\langle 1| \otimes I$. Finally we measure the observable $\langle 2\sigma_+^a \rangle = \langle \sigma_x^a + i\sigma_y^a \rangle = G(t)$.




FIG. 4: Quantum network for the evaluation of $G(t) = \langle \psi | T^\dagger A_i^\dagger T B_j \psi \rangle$.

On the other hand, sometimes we might be interested in obtaining the spectrum (eigenvalues) of a given observable \hat{Q} (i.e., an hermitian operator). A quantum algorithm (network) for this purpose was also given in previous work [5]. Again, the basic idea is to perform an indirect measurement

using an extra qubit (see Fig. 5). Basically, we prepare the initial state (ancilla plus system) $|+\rangle_a \otimes |\phi\rangle$, then apply the evolution $e^{i\hat{Q}\sigma_z^a \frac{t}{2}}$, and finally measure the observable $\langle 2\sigma_+^a(t) \rangle = \langle \phi | e^{-i\hat{Q}t} \phi \rangle$. Since the initial state of the system can be written as a linear combination of eigenstates of \hat{Q} , $|\phi\rangle = \sum_{n=0}^L \gamma_n |\psi_n\rangle$, where γ_n are complex coefficients and $|\psi_n\rangle$ are eigenstates of \hat{Q} with eigenvalue λ_n , the classical Fourier transform applied to the function of time $\langle 2\sigma_+^a(t) \rangle$ gives us λ_n

$$\hat{F}(\lambda) = \sum_{n=0}^L 2\pi |\gamma_n|^2 \delta(\lambda - \lambda_n). \quad (35)$$

Without loss of generality, we can choose $\hat{Q} = H$, with H some particular Hamiltonian.




FIG. 5: Quantum network for the evaluation of the spectrum of an observable \hat{Q} .

It is important to note that in order to obtain the different eigenvalues of \hat{Q} , the overlap between the initial state and the eigenstates of \hat{Q} must be different from zero. One can use different mean-field solutions of \hat{Q} as initial states $|\phi\rangle$ depending on the part of the spectrum one wants to determine with higher accuracy.

VI. ALGORITHM EFFICIENCY AND ERRORS

An algorithm is considered efficient if the number of operations involved scales polynomially with the system size and $1/\epsilon$, where ϵ is the maximal tolerable error in the measurement of a relevant property.

While the evolution step involves a number of unitary operations that scales polynomially with the system size (such is the case for the Trotter approximation) whenever the Hamiltonian H is physical (i.e., is a sum of a number of terms that scales polynomially with the system size), the preparation of the initial state could be inefficient. Such inefficiency would arise, for example, if the state $|\psi\rangle$ defined in Eq. 8 or Eq. 30 is a linear combination of an exponential number of elementary states ($L \sim x^N$, with N the number of sites in the system and $x > 1$). However, if we assume that $|\psi\rangle$ is a polynomial (in N) combination of states, its preparation can be done efficiently. This construction generalizes to more general coherent states [12]. On the other hand, the measurement process described in section V is always an efficient step, since it only involves the measurement of the spin of one qubit, despite the number of qubits or sites N of the quantum system.

Errors ϵ come from gate imperfections, the use of the Trotter approximation in the evolution operator, and the statistics in measuring the spin of the ancilla qubit (sections III B, IV B, and V). A precise description and study of the error sources can be found in previous work [3]. The result is that the algorithms described here, for the simulation of physical systems and processes, are efficient if the preparation of the initial state is efficient, too.

VII. CONCLUSIONS

We studied the implementation of quantum algorithms for the simulation of an arbitrary quantum physical system or phenomenon on a QC made of qubits, making a distinction between systems that are governed by Pauli's exclusion principle (fermions, hard-core bosons, anyons, spins, etc.), and systems that are not (e.g, canonical bosons). For the first class of quantum systems, we showed that a mapping between the corresponding algebra of operators and the spin-1/2 algebra exists, since both have a finite-dimensional representation. On the other hand, the operator representation of quantum systems that are not governed by an exclusion principle is infinite-dimensional, and an isomorphic mapping to the spin-1/2 algebra is not possible. However, one can work with a finite set of quantum states, setting a constraint, such as fixing the number of bosons in the system. Then, the representation of bosonic operators becomes finite-dimensional, and we showed that we can write down bosonic operators in the spin-1/2 language (Eq. 27), mapping bosonic states to spin-1/2 states (Eq. 26) .

We also showed how to perform quantum simulations in a QC made of qubits (quantum net-

works), giving algorithms for the preparation of the initial state, the evolution, and the measurement of a relevant physical property, where in the most general case the unitary operations have to be approximated (sections III B, IV B).

The mappings explained are efficient in the sense that we can perform them in a number of operations that scales polynomially with the system size. This implies that the evaluation of some correlation functions in quantum states that can be prepared efficiently is also efficient, showing an exponential speed-up of these algorithms with respect to their classical simulation. However, these mappings are insufficient to establish that quantum networks can simulate any physical problem efficiently. As we mentioned in the Introduction, this is the case for the determination of the spectrum of the Hamiltonian in the two-dimensional Hubbard model [5], where the signal-to-noise ratio decays exponentially with the system size.

Finally, in Fig. 6 a table displays the advantages of simulating some known algorithms with a QC than with a CC, affirming that QCs behave as efficient devices for some quantum simulations.

| Algorithm | Speed-Up |
|-----------------------|-----------------|
| Correlation functions | Exponential |
| Factoring (Shor) | Superpolynomial |
| Search (Grover) | Quadratic |

FIG. 6: Quantum vs. classical simulations. Speed-up refers to the gain in speed of the quantum algorithms compared to the known classical ones.

[1] R. P. Feynman, *Int. J. Theor. Phys.* **21**, 467 (1982).

[2] S. Lloyd, *Science* **273**, 1073 (1996); D. A. Meyer, *J. Stat. Phys.* **85**, 551 (1996). S. Wiesner, quant-ph/9603028; B. M. Boghosian and W. Taylor IV, *Physica D* **120**, 30 (1998); C. Zalka, *Proc. R. Soc. Lond. A* **454**, 313 (1998).

[3] G. Ortiz, J. E. Gubernatis, E. Knill, and R. Laflamme, *Phys. Rev. A* **64**, 042323 (2001).

[4] D. A. Meyer, *Comp. Phys. Comm.* **146**, 295 (2002).

- [5] R. Somma, G. Ortiz, J. E. Gubernatis, E. Knill, and R. Laflamme, Phys. Rev. A **65**, 22319 (2002).
- [6] P. Jordan and E. Wigner, Z. Phys. **47**, 631 (1928).
- [7] C. Batista and G. Ortiz, Phys. Rev. Lett. **86**, 1082 (2001).
- [8] C. D. Batista and G. Ortiz, cond-mat/0207106.
- [9] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, Cambridge, 2000.
- [10] J. Blaizot and G. Ripka, *Quantum Theory of Finite Systems*, MIT Press, Cambridge, 1986.
- [11] M. Suzuki, *Quantum Monte Carlo Methods in Condensed Matter Physics*, World Scientific, Singapore, 1993.
- [12] A. Perelomov, *Generalized Coherent States and their Applications* (Springer, Berlin, 1986).