

LA-UR-07-**** 0949

Approved for public release;
distribution is unlimited.

Title: Bison Antiquus Occurrence and Pleistocene-Holocene Stratigraphy, Cañada del Buey, Pajarito Plateau, New Mexico

Author(s): Paul G. Drakos
Steven L. Renneau
Gary S. Morgan

Submitted to: New Mexico Geological Society 2007 Field Conference Guidebook



Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-36. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

BISON ANTIQUUS OCCURRENCE AND PLEISTOCENE- HOLOCENE STRATIGRAPHY,
CAÑADA DEL BUEY, PAJARITO PLATEAU, NEW MEXICO

PAUL G. DRAKOS¹, STEVEN L. RENEAU², AND GARY S. MORGAN³

¹Glorieta Geoscience, Inc., 1723 Second St, Santa Fe, NM 87505, drakos@glorietageo.com

²Los Alamos National Laboratory, EES-9, MS D452, Los Alamos, NM 87545, sreneau@lanl.gov

³New Mexico Museum of Natural History and Science, 1801 Mountain Rd NW, Albuquerque, NM 87104, gary.morgan@state.nm.us

ABSTRACT

A *Bison* (probable *Bison antiquus*) distal humerus fragment was found within a Pleistocene colluvial deposit on a hillslope above Cañada del Buey near White Rock, New Mexico. The *Bison* fossil is preserved within a buried soil with an inferred age of ca. 50-100 ka, based on soil properties and on stratigraphic position below a deposit of ca. 50-60 ka El Cajete pumice. This represents the second oldest dated *Bison* in New Mexico, and one of the few occurrences of this genus in the northern mountains of the state. It is also only the second record of a Pleistocene vertebrate from Los Alamos County, and is a rare occurrence of a pre-25 ka *Bison* fossil in good stratigraphic context. Hillslopes in the study area are underlain by a sequence of truncated Pleistocene and Holocene soils that are inferred to represent colluvial deposition and soil formation followed by erosion in the mid Pleistocene (buried soil "b3"), the late Pleistocene (buried soil "b2"), and the mid-to-late Holocene (buried soil "b1"). The surface soil is developed in deposits that overlie 600-800 year-old Ancestral Puebloan sites. Colluvium is dominated by relatively fine-grained (fine to very fine sand) slopewash colluvium deposited by overland flow, but also includes rocky colluvium on hillslopes below mesas. The fine-grained colluvium is likely derived mainly from reworking of eolian deposits. Episodic colluvial deposition appears to, at least in part, accompany and follow episodic eolian events, with intervening periods dominated by erosion and the development of truncated soils.

INTRODUCTION

A recently discovered distal humerus fragment of a *Bison sp.* (probable *Bison antiquus*) specimen near White Rock, New Mexico, provides an opportunity to examine a rare bison fossil in good stratigraphic context, in a buried soil horizon below a deposit of ca. 50-60 ka El Cajete pumice. The fossil was found during archaeological investigations conducted at Los Alamos National Laboratory (LANL) in 2002 by Dr. Bradley Vierra, the archaeologist in charge of the project. This is the second recorded Pleistocene fossil from Los Alamos County, and is also one of only two bison records in New Mexico with dates older than about 25 ka. The *Bison sp.* site was examined as part of geomorphic studies conducted for LANL in support of archaeological investigations within the White Rock land transfer parcel, located within the Cañada del Buey watershed (Fig. 1). The fossil locality is situated on an east-facing colluvial slope approximately 0.5 km west of the town of White Rock. The fossil was preserved in late Pleistocene colluvium overlying older Pleistocene colluvium and Pliocene Cerros del Rio basalt. The preservation of this specimen in a colluvial deposit is a somewhat unusual occurrence. In this paper we discuss the Pleistocene and Holocene stratigraphy of the White Rock parcel and the associated setting of the *Bison sp.* fossil.

METHODS

A surficial geologic map of the study area was prepared at a scale of 1:1200. The mapping was focused on units with potential archaeological significance. Soil descriptions were made following methods discussed in Birkeland (1999). Soil horizon nomenclature is from Birkeland (1999) and Soil Survey Staff (1999). Buried soil horizons were numbered based on the overall stratigraphy for the study area, rather than for individual profiles (e.g. some profiles may have the surface soil profile sitting directly on buried soil b2 or b3 horizons, or the b1 soil may sit directly on the b3 soil). Carbonate stage for soils follows nomenclature developed by Gile et al. (1966). Preliminary age estimates for deposits were made based on soil descriptions and comparison of the general degree of soil development to previously dated sites on the

Pajarito Plateau, and to soils described during the present investigation where radiocarbon dates were obtained. A colluvial deposit in Fence Canyon with a calibrated (cal) radiocarbon age of ca. 5.0 ka (Stop 1-4c, Reneau and McDonald 1996, pp. 62-64), at the same general elevation as the White Rock parcel, was used as a key reference for the degree of soil development in a mid-Holocene unit on that part of the plateau. The presence of the ca. 50-60 ka El Cajete pumice interbedded with colluvial sediments provided an upper age constraint for the *Bison sp.* fossil (tephra age from Toyoda et al. 1995; Reneau et al. 1996b).

SETTING

Cañada del Buey is located on the Pajarito Plateau, an area of gently east-sloping mesas and intervening narrow canyons (Fig. 1). The study area is in the eastern part of the plateau at an elevation of 1950 to 2000 m, in an area of relatively low relief. The modern climate is semiarid, with mean annual precipitation of about 350 mm (Bowen, 1990). Vegetation is dominated by piñon-juniper woodlands (McKown et al., 2003).

SURFICIAL GEOLOGIC UNITS AND STRATIGRAPHY OF UNDERLYING DEPOSITS

The study area includes part of the Cañada del Buey stream channel and adjacent floodplains, colluvial slopes, and alluvial fans (Fig. 2). Bedrock beneath most of the area is Pliocene basalt of the Cerros del Rio volcanic field (unit Tb). The early Pleistocene Tshirege Member of the Bandelier Tuff (unit Qbt), which overlies the Cerros del Rio basalt, is also present along the northern margin, and as an isolated mesa in the western part of the study area (Fig. 2). Large parts of the study area are covered by locally derived colluvial, alluvial fan, or slopewash deposits of a variety of ages. Geologic maps of this area have been prepared by Griggs (1964), Rogers (1995), and Dethier (1997).

Unit Qal consists of young alluvium in the main stream channel of Cañada del Buey and tributary drainages, and adjoining floodplains and stream terraces. Sediment ranges in size from silt to coarse sand and gravel, and is dominated by coarse sand in the main channels and very fine sand on the floodplains (Drakos et al., 2000). The upper sediment layers along the main

channel and floodplains (approximately 0.5 to 2.0 m thick) are largely historic in age, although older sediment may be locally present at depth. Higher stream terraces (unit Qt) along Cañada del Buey are generally above the level of historic flooding, and are inferred to be late Holocene to Pleistocene in age. The stream terraces are in part overlain by colluvium (unit Qc).

Unit Qf consists of young alluvial fans that emanate from side drainages, typically below eroding areas of colluvium. Qf is dominated by stratified fine to very fine sand, and also includes coarse sand and fine gravel layers. The upper parts of these deposits are historic in age, and older deposits are commonly present at depth. Greater than 1 m of late Holocene sediment is present in some Qf units.

Unit Qc is dominated by relatively fine-grained (fine to very fine sand) slopewash colluvium deposited by overland flow, and also includes rocky colluvium on hillslopes below mesas. Qc likely includes alluvial fan surfaces and underlying deposits, and eolian deposits and/or locally reworked eolian sediment. Qc deposits have a wide range in age, and typically have buried soils that indicate pauses in deposition, in part accompanied by local erosion. Several soil profiles include surficial and buried deposits that indicate at least two episodes of colluvial deposition since mid Holocene time, with a lower colluvial layer likely deposited around 2 to 4 ka, and an upper colluvial layer that was likely deposited within the past 1000 years (Drakos and Reneau, 2003). However, in many locations, the upper colluvial layer overlies late Pleistocene or early Holocene to latest Pleistocene deposits.

Although unit Qc is characterized by spatial complexity in its depositional history, it can be subdivided into two units, Qc1 and Qc2, based on the presence or absence of middle Holocene colluvium beneath the pervasive latest Holocene colluvium. Unit Qc1 is characterized by latest Holocene (<1 ka?) Qc overlying Pleistocene or early Holocene Qc (Fig. 3). In the area east of the Bandelier Tuff mesa, the late Holocene Qc thins downslope from approximately 0.7 m thick at the base of the mesa to less than 0.1 m thick at SP15, the *Bison sp.* fossil locality (Fig. 3).

Unit Qc2 is characterized by latest Holocene (<1 ka?) Qc overlying inferred middle Holocene Qc. Middle Holocene deposits in unit Qc2 are approximately 1 m thick at SP6, and are overlain by approximately 0.2 to 0.7 m of late Holocene deposits (Fig. 4). In general, Qc1 underlies east- and southeast-facing slopes in areas of relatively thin colluvial deposits overlying bedrock units Tb and Qbt (Figs. 2 and 3). Unit Qc2 underlies aggrading toe slopes below embayments in the Qbt mesa north of the study area and the north-facing slope between the small Qbt mesa and Cañada del Buey within the western part of the study area (Fig. 2).

Sediment in unit Qc with estimated ages younger than ca. 5 ka, based on comparison with the Fence Canyon reference section, ranges in thickness from 6 cm to >1 m. Its soils lack Stage I carbonate or Bt horizons. The thickest deposit was recorded in the eastern part of the study area, where greater than 1.1 m of late Holocene colluvium is present at location SP22, within site LA 12765 (Drakos and Reneau, 2003). Farther west, 70-80 cm of colluvium younger than ca. 4 ka is present on the south side of an isolated mesa of Bandelier Tuff (SP17 and SP18, Fig. 2). The total thickness of Holocene or possibly latest Pleistocene sediment (< ~10-15 ka) reaches about 1.7 m in a gullied area in the northwestern part of the parcel (SP3a, Fig. 2).

Bedrock and Pumice Units

Unit Qec is the ca. 50-60 ka El Cajete pumice (Toyoda et al. 1995; Reneau et al. 1996b). It is present in a relatively thick (≥ 50 cm) layer within Qc on the north side of the isolated Bandelier Tuff mesa in the western part of the study area (locations SP1 and SP1a, Fig. 2), but has been largely eroded from the rest of the parcel. Thin Qec remnants were observed within Qc farther east, in the vicinity of the *Bison* sp. site (site SP15b, Fig. 3 and Table 1).

Unit Qbt is the early Pleistocene Tshirege Member of the Bandelier Tuff. There are no soils or only thin soils present in much of this unit, particularly along the edges of mesas. Where present, mesa top soils are formed in thin, discontinuous, fine-grained deposits dominated by very fine sand (locations SP21 and SP 21a, Fig. 4), and represent either eolian or locally reworked eolian sediment. Thin deposits overlying Qbt are in part late Holocene in age (likely

less than 1 ka) based on the degree of soil development (Drakos and Reneau, 2003, and this volume).

Unit Tb is basalt of the Cerros del Rio volcanic field. There are no soils or only thin soils present throughout the area of exposure of this unit. In other areas discontinuous colluvial or eolian sediments overlie unit Tb.

Soil Stratigraphy of Hillslope Deposits

Hillslopes in the study area are underlain by a sequence of truncated Pleistocene and Holocene soils that are inferred to represent colluvial deposition and soil formation followed by erosion in the mid Pleistocene (buried soil "b3"), the late Pleistocene (buried soil "b2"), and the mid-to-late Holocene (buried soil "b1") (Figs. 3 and 4). The b3 soil exhibits 5YR color, moderately thick clay films, maximum Stage III carbonate morphology and, based on comparison with previous soils investigations on the Pajarito Plateau, has an estimated age of at least 100-200 ka (McFadden et al., 1996). The b2 soil exhibits 7.5YR color, thin to moderately thick clay films, and Stage I to I+ carbonate morphology. In a study of calcic soils of the southwestern U.S., soils formed in gravelly alluvial deposits develop stage II morphology over a time period of 10 ka to 100 ka (Machette, 1985). The b2 soil is older than the ca. 50-60 ka Qec and, based on carbonate morphology and stratigraphic position overlying the b3 soil, is younger than 100 ka. The b1 soil exhibits 7.5YR-10YR color, lacks clay films or has rare, thin films (e.g. Btjb1 horizon in SP6, Fig. 4), and maximum Stage I carbonate morphology. Well-developed Bt horizons, with many moderately thick clay films and 7.5 YR color, overlying stage I carbonate horizons, were observed in ca. 6.7-7.4 ka colluvial deposits in Rendija canyon at a site where older upslope soils were a possible clay source (Drakos and Reneau, 2004). Incipient Bt horizon development observed in Cañada del Buey b1 soils is therefore consistent with a middle Holocene age, likely less than 7 ka. Based on correlation with the ca. 5.0 ka Fence Canyon soil profile with stage I carbonate, the b1 soil has an estimated age of 4 to 6 ka. The surface soil typically exhibits an A-Bw profile, and is developed in deposits that overlie 600-800 year-old

Ancestral Puebloan sites. At some locations, the surface soil consists of an A, AC, or C horizon only, and is interpreted to be historic in age (Figs. 3 and 4).

The predominance of fine-grained colluvial deposits is likely due to reworking of eolian deposits as a significant source of colluvial sediment. In between eolian events, erosional processes dominate, during which time much of the sediment is stripped from hillslopes and soils are truncated. Deposition of fine-grained colluvium likely enhances the probability of fossil preservation within a colluvial deposit.

Stratigraphy of *Bison* Locality

The *Bison* fossil was located on an east-facing colluvial slope in unit Qc1 overlying basalt bedrock, in late Pleistocene colluvium that overlies a remnant older Pleistocene buried soil developed in a thin colluvial deposit (Fig. 3). The late Pleistocene colluvial deposit is overlain by thin, inferred historic-age colluvium that is reddened (note AC horizon with 7.5YR hue, location 15, Table 1), suggesting that it is derived from reworking of older soils. The piece of fossilized bone was found at a depth of about 20-30 cm eroding out of a gully wall stratigraphically below the ca. 50-60 ka El Cajete pumice (Fig. 3; Table 1, note Qec pumice at location SP 15b). The bone was situated in a reddened (7.5YR) Btb2 horizon with a soil profile that exhibited a Stage I+ carbonate in the underlying Btcb2 horizon (Table 1). Based on the stratigraphic position of the bone horizon below the El Cajete pumice, the fossil is > ca.50-60 ka. As discussed above, the Bt and Btk horizon development, with maximum Stage I+ carbonate, moderately thick clay films and 7.5YR hue, is consistent with an age of <100 ka. The *Bison* fossil therefore has an estimated age of ca. 50-100 ka.

PALEONTOLOGY

***Bison* sp.**

A partial distal end of a left humerus of an extinct species of *Bison* (New Mexico Museum of Natural History-NMMNH catalogue number 37623; Fig. 5B) was found in the Cañada del Buey site (NMMNH locality L-5214). In a previous reference to this same fossil, the

site was called White Rock for its proximity to the town of White Rock in Los Alamos County (Morgan and Lucas, 2005a). The specimen is very similar in morphology to a distal humerus from an extant American bison (*Bison bison*) skeleton from Fort Wingate Military Reservation in McKinley County, northwestern New Mexico (Fig. 5A). The distal articular surface of the fossil humerus compares closely to the humerus of *B. bison* in having the medial condyle noticeably deeper (proximo-distally) than the lateral condyle, a deeply excavated (concave) coronoid fossa just proximal to the articular surface, and the presence of a distinct medial ridge on the lateral condyle that extends as a low ridge proximally for a short distance into the coronoid fossa before curving laterally toward the edge of the shaft. The size and overall features of the fossil are superficially similar to the distal humerus of a large extinct horse (*Equus*), but the horse differs in having the medial and lateral condyles about the same depth, a much shallower coronoid fossa, and a less distinct medial ridge on the lateral condyle that does not extend into the coronoid fossa as a low ridge.

The species of *Bison* are identified by characters of their bony horn cores (McDonald, 1981), and thus the partial humerus from Cañada del Buey can only be referred to *Bison* sp. However, its occurrence in a late Pleistocene site (50-100 ka; late Rancholabrean land mammal age) and smaller size compared to *B. latifrons* suggest that NMMNH 37623 is more likely to be *B. antiquus*, an extinct species typical of late Pleistocene and early Holocene sites in New Mexico and elsewhere in western North America (McDonald, 1981). The much larger *B. latifrons* is more characteristic of medial Pleistocene sites (100-300 ka; early Rancholabrean land mammal age). Although the fossil appears to be similar in size to, or even somewhat smaller than, the recent male *B. bison* humerus in Fig. 5A, this is misleading because the fossil is incomplete and weathered. The medial edges of the shaft and the articular surface are both broken in the fossil (Fig. 5B). In features that are comparable (e.g., the breadth of the humeral shaft just proximal to the articular surface), the fossil is actually somewhat larger than the recent *B. bison*, which places it in the size range of *B. antiquus*.

The Cañada del Buey *Bison* represents the second known record of a Pleistocene vertebrate fossil from Los Alamos County. The first is a lower jaw of a small horse (*Equus*), found in 1966 by collectors from the Frick Laboratory of the American Museum of Natural History on the mesa top east of Portrillo Canyon, west of White Rock Canyon, 1 to 2 km west of Pajarito Springs (Morgan and Lucas, 2005a). According to the original notes written on the fossil by Ted Galusha, this horse jaw was collected from the early Pleistocene Bandelier Tuff, and thus would be considerably older than the bison humerus which has an inferred age of 50-100 ka. The Cañada del Buey humerus is the second oldest dated *Bison* known from New Mexico. The oldest record is a partial skull and horn core of *B. latifrons* from the Los Duranes Formation in a gravel pit in Bernalillo, Sandoval County (Smartt et al., 1991). According to Connell and Love (2001) and Connell (2004), the Los Duranes Formation is bracketed by dates of 156 ka on the Albuquerque volcanoes basalt (Peate et al., 1996) and 98 ka on the Cat Hills basalt above the Los Duranes Formation (Maldonado et al., 1999). Other dated records of *Bison* in New Mexico are younger than 25 ka, of which the identifiable specimens are all referable to *B. antiquus*.

Late Pleistocene records of *Bison* are uncommon in the mostly mountainous terrain of northern New Mexico (Morgan and Lucas, 2005b). The Cañada del Buey site is one of the higher elevation records of *Bison* in the state at 1975 m (6480 ft). Other late Pleistocene *Bison* occurrences in this region from elevations above 1830 m (6000 ft) include Navajo Lake in San Juan County, Abiquiu in Rio Arriba County, Mesa Vibora in Taos County, Folsom in Colfax County, and Snow Ranch in Santa Fe County. *Bison* are grazing ungulates, and during the late Pleistocene were more common in the extensive grasslands and savannas of the Great Plains in eastern New Mexico (Morgan and Lucas, 2005a).

CONCLUSIONS

Hillslopes in the study area are underlain by a sequence of truncated Pleistocene and Holocene soils that are inferred to represent colluvial deposition and soil formation followed by

erosion in the mid Pleistocene (buried soil "b3"), the late Pleistocene (buried soil "b2"), and the mid-to-late Holocene (buried soil "b1") (Figs. 3 and 4). Two episodes of widespread Pleistocene colluvial deposition are recorded as buried soils b2 and b3 (Fig. 3). The b3 soil is discontinuously preserved, often as remnant stripped soils in bedrock depressions, and has an estimated age of at least 100-200 ka; it may actually be developed in deposits with a wide range in age, although this cannot be determined at present. Based on stratigraphic relationships and soil characteristics, the b2 soil is inferred to be greater than 50 ka but less than 100 ka.

An examination of colluvial stratigraphy at sites throughout the study area indicates that there have been two general periods of relatively widespread colluvial deposition since the middle Holocene. An episode or several episodes of colluvial deposition occurred during the middle to late Holocene, likely around 2 to 5 ka (e.g. buried soil b1, Fig. 4), and a second period of colluvial deposition occurred within the past 800 years, likely contemporaneous with and/or post-dating Puebloan occupation (A-Bw surficial soil profiles, Figs. 3 and 4). A thin (typically less than 10 cm thick), very young colluvial layer, likely deposited within the past 100 years, is observed at several locations. Areas where middle to late Holocene colluvial deposits are preserved are mapped as Qc2 (Fig. 2). Areas where late Holocene colluvium overlies Pleistocene colluvium and middle Holocene deposits are not preserved are mapped as Qc1.

The presence of older early-to-middle Holocene colluvial deposits in other areas on the eastern Pajarito Plateau indicates that the record in the study area is incomplete, with likely loss of deposits of this age by erosion. For example, colluvial deposits dated at 6-8 ka have been found in Fence Canyon and in several other areas (Reneau and McDonald, 1996; Reneau et al., 1996a), but have not been clearly identified in the study area.

The small mesa top in the western part of the study area is characterized by Bandelier Tuff bedrock overlain by thin, discontinuous remnant Pleistocene soils and recent eolian or reworked eolian deposits typically less than 20 to 30 cm thick (Figs. 3 and 4). Similar thin, discontinuous deposits not greater than 20 to 30 cm thick were noted during archaeological

excavations on the Mesita del Buey mesa top west of the study area (Steen, 1982). Prior to approximately 800 years BP, many mesa top surfaces were apparently characterized by stripped surfaces with remnant eroded Pleistocene (b3) soils and exposed bedrock, with subsequent burial by eolian sediment.

The Bison fossil is located on an east-facing colluvial slope within buried soil b2 in unit Qc1, in late Pleistocene colluvium stratigraphically below the ca. 50-60 ka El Cajete pumice. Although the Bison fossil, a partial distal humerus, is not identifiable to the species level, its size and late Pleistocene age (inferred age of 50-100 ka) suggest it is probably B. antiquus. This is one of the very few bison records in New Mexico with a date older than about 25 ka. It is also only the second recorded Pleistocene fossil from Los Alamos County, and one of about a half dozen records of Bison from the mountains of northern New Mexico at elevations above 1830 m (6000 ft).

ACKNOWLEDGMENTS

Funding for this work was provided by the LANL Land Conveyance and Transfer Project, and we thank Brad Vierra for his support throughout this project. Scott Eberhardt provided GIS support, and Jim Riesterer collaborated on figure preparation. John Hawley, Emily Schultz and Dave Love provided helpful review comments.

REFERENCES

Birkeland, P. W., 1999, Soils and Geomorphology, Third Edition: Oxford, Oxford University Press, 430 p.

Bowen, B.M., 1990, Los Alamos climatology: Los Alamos National Laboratory Report LA-11735-MS, Los Alamos, New Mexico, 254 p.

Connell, S. D., 2004, Geology of the Albuquerque basin and tectonic development of the Rio Grande rift in north-central New Mexico; in Mack, G. H. and Giles, K. A., eds., The geology of New Mexico, a geologic history: New Mexico Geological Society, Special Publication 11, p. 359-388.

Connell, S. D. and Love, D. W., 2001, Stratigraphy of middle and upper Pleistocene fluvial deposits of the Rio Grande (post-Santa Fe Group) and the geomorphic development of the Rio Grande valley, northern Albuquerque basin, central New Mexico: New Mexico Bureau of Mines and Mineral Resources, Open-File Report 454A, p. J-67-78.

Dethier, D. P., 1997, Geology of the White Rock Quadrangle, Santa Fe and Los Alamos Counties, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Geological Map 73, Socorro, New Mexico.

Drakos, P., Rytí, R., Reneau, S., and Greene, K., 2000, Evaluation of possible sediment contamination in the White Rock land transfer parcel: Reach CDB-4: Los Alamos National Laboratory report LA-UR-00-5071, Los Alamos, New Mexico.

Drakos, P. G. and Reneau, S. L., 2003, Surficial units and processes associated with archaeological sites in selected land conveyance parcels, Los Alamos National Laboratory: 2002 through February 2003 field season investigations: Los Alamos National Laboratory report LA-UR-03-2630, Los Alamos, New Mexico.

Drakos, P. G. and Reneau, S. L., 2004, Surficial units and processes associated with archaeological sites in selected land conveyance parcels, Los Alamos National Laboratory, Volume II, 2003 field season investigations: Rendija Canyon and Airport Tract sites: Los Alamos National Laboratory report LA-UR-04-1842, Los Alamos, New Mexico.

Drakos, P. G. and Reneau, S. L., this volume, Episodic eolian events and preservation of mesa top archaeological sites on the Pajarito Plateau: New Mexico Geological Society 58th Field Conference, Guidebook.

Gile, L., Peterson, F. F., and Grossman, R. B., 1966, Morphologic and genetic sequences of carbonate accumulation in desert soils: Soil Science, v. 101, p. 347-360.

Griggs, R. L., 1964, Geology and groundwater resources of the Los Alamos area, New Mexico: U.S. Geological Survey Water-Supply Paper 1735, 107 p.

Machette, N. N., 1985, Calcic soils of the southwestern United States: Geological Society of America Special Paper 203, p. 1-21.

Maldonado, F., Connell, S. D., Love, D. W., Grauch, V. J. S., Slate, J. L., McIntosh, W. C., Jackson, P. B., and Byers, F. M. Jr., 1999, Neogene geology of the Isleta Reservation and vicinity, Albuquerque basin, central New Mexico: New Mexico Geological Society, Guidebook 50, p. 175-188.

McDonald, J. N., 1981, North American *Bison*: Their classification and evolution: Berkeley, University of California Press, 316 p.

McFadden, L.D., Watt, P.M., Reneau, S. L., and McDonald, E. V., 1996, General soil-landscape relationships and soil-forming processes in the Pajarito Plateau, Los Alamos National Laboratory area, New Mexico: New Mexico Geological Society 47th Field Conference, Guidebook, p. 357-366.

McKown, B., Koch, S.W. Balice, R.G. and Neville, P. 2003. Land cover map for the eastern Jemez region: Los Alamos National Laboratory Report LA-14029, Los Alamos, New Mexico, 84 p.

Morgan, G.S., and Lucas, S.G., 2005a, Pleistocene vertebrate faunas in New Mexico from alluvial, fluvial, and lacustrine deposits, *in* Lucas, S.G., Morgan, G.S., and Zeigler, K.E., eds., New Mexico's Ice Ages: New Mexico Museum of Natural History and Science, Bulletin 28, p. 185-248.

Morgan, G.S., and Lucas, S.G., 2005b, Pleistocene vertebrates from Rio Arriba and Taos Counties, northernmost New Mexico: New Mexico Geological Society, Guidebook 56, p. 416-424.

Peate, D. W., Chen, J. H., Wasserburg, G. J., and Papanastassiou, D. A., 1996, $^{238}\text{U}/^{230}\text{Th}$ dating of a geomagnetic excursion in Quaternary basalts of the Albuquerque volcanoes field, New Mexico (USA): Geophysical Research Letters, v. 23, n. 17, p. 2271-2274

Reneau, S. L., and McDonald, E. V., 1996, Landscape history and processes on the Pajarito Plateau, northern New Mexico: Rocky Mountain Cell, Friends of the Pleistocene, Field Trip Guidebook, Los Alamos National Laboratory report LA-UR-96-3035, Los Alamos, New Mexico, 195 p.

Reneau, S. L., McDonald, E. V., Gardner, J. N., Kolbe, T. R., Carney, J. S., Watt, P. M., and Longmire, P. A., 1996a, Erosion and deposition on the Pajarito Plateau, New Mexico, and implications for

geomorphic responses to late Quaternary climatic changes: New Mexico Geological Society, 47th Field Conference, Guidebook, p. 391-397.

Reneau, S. L., Gardner, J. N., and Forman, S. L., 1996b, New evidence for the age of the youngest eruptions in the Valles caldera, New Mexico: Geology, v. 24, p. 7-10.

Rogers, M. A., 1995, Geologic map of the Los Alamos National Laboratory Reservation: New Mexico Environment Department, Santa Fe.

Smartt, R. A., Lucas, S. G., and Hafner, D. J., 1991, The giant bison (*Bison latifrons*) from the middle Rio Grande Valley of New Mexico: Southwestern Naturalist, v. 36, p. 136-137.

Soil Survey Staff, 1999, Soil Taxonomy: A basic system of soil classification for making and interpreting soil surveys: United State Department of Agriculture, Natural Resources Conservation Service, Agriculture Handbook No. 436, 869 p.

Steen, C.R., 1982, Pajarito Plateau archaeological surveys and excavations, II: Los Alamos National Laboratory report LA-8860-NERP, Los Alamos, New Mexico, 60 p.

Stuiver, M., Reimer, P. J., and Reimer, R. W. 2005. CALIB 5.0. [WWW program and documentation].

Toyoda, S., Goff, F., Ikeda, S., and Ikeya, M., 1995, ESR dating of quartz phenocrysts in the El Cajete and Battleship Rock members of the Valles Rhyolite, Valles caldera, New Mexico: Journal of Volcanology and Geothermal Research, v. 67, p. 29-40.

APPENDIX A. KEY TO SYMBOLS USED IN DESCRIPTIONS OF SOIL MORPHOLOGY

From Birkeland (1999)

Structure

Grade	Size	Type
1 = weak	vc = very coarse	sbk = subangular blocky
2 = moderate	c = coarse	abk = angular blocky
3 = strong	m = medium	pr = prismatic
	f = fine	pl = platy
		sg = single grain
		m = massive

Consistence

Dry	Wet - Stickiness	Wet - Plasticity
lo = loose	so = non sticky	po = non-plastic
so = soft	vss = very slightly sticky	vps = very slightly plastic

sh = slightly hard
h = hard
vh = very hard

ss = slightly sticky
s = sticky

ps = slightly plastic
p = plastic

Cutans (clay films)

Abundance

n.o. = none observed
v1 = very few (< 5%)
1 = few (2 - 25%)
2 = common (25 - 50%)
3 = many (50 - 75%)
4 = continuous (75+%)

Thickness/(Distinctness)
n = thin (faint)
mk = moderately thick
k = thick

Location/Type
po = along pores

co = coating gravel, ped faces
br = bridging grains
pf = along ped faces (as co + br)
pr:pf along prismatic ped faces
bk:pf along blocky ped faces
Lam = lamellae

Horizon Boundary

Thickness

a = abrupt (< 2.5cm)
c = clear (2.5 - 6cm)
g = gradual (6-12.5cm)
d = diffuse (> 12.5 cm)

Topography
s = smooth
w = wavy
i = irregular
b = broken

Carbonate effervescence in HCl
none = non-effervescent
e = slightly effervescent
es = strongly effervescent
ev = violently effervescent

Texture

s = sand
ls = loamy sand
sl = sandy loam
l = loam

sil = silt loam
scl = sandy clay loam
sicl = silty clay loam
cl = clay loam

LIST OF FIGURES

FIGURE 1. Digital Elevation Map (DEM) of Pajarito Plateau showing study area.

FIGURE 2. Geomorphology, cross section and soil pit locations, Cañada del Buey study area. See Figs. 1 and 3 for cross sections A-A' and B-B'.

FIGURE 3. Cross section (bottom), soil profiles and correlations (top) showing *Bison sp.* occurrence and stratigraphic relationships in Unit Qc1. See Fig. 1 for cross section and soil description locations.

FIGURE 4. Cross section (bottom), soil profiles and correlations (top) showing stratigraphic relationships between alluvial fan (Unit Qf), hillslope (Unit Qc2), and mesa top (Unit Qe+Qc) deposits, Cañada del Buey. See Fig. 1 for cross section and soil description locations.

FIGURE 5. Photograph of left distal humerus of the extinct species of bison, *Bison sp.* (*B. antiquus*?) (lower image of bone fragment). Humerus of a modern bison, *Bison bison* (upper image of intact bone) shown for comparison. New Mexico Museum of Natural History catalogue number NMMNH 37623 and NMMNH locality number L-5214.

Table 1. Summary of Soil Morphology for *Bison sp.* site.

Table 1.

Horizon	Depth (cm)	Gravel (%)	Dry Color (Matrix)	Moist Color (Matrix)	Texture	Structure	Dry Consistency	Wet Consistency	Argillans	CaCO3 Stage	Lower Horizon Boundary	Age Estimate	Notes
Location 15, North facing gully wall at Bison antiquus bone site													
AC	0-6	<2	7.5YR3/3	7.5YR2.5/3	sicl	m	lo	s,p	no	non	-	cs	historic
ABwb2	6-17	<2	7.5YR4/3	7.5YR3/2	sicl	sfsbk	sh	ss,ps	no	non	-	cs	correlative to pre-El Cajete soil?
Btb2	17-30	<2	7.5YR4/3	7.5YR3/2	sicl	2-3msbk	sh-h	s,ps	1nbrpopf	non	-	cs	<i>Bison sp.</i> bone horizon, est age 50-100 ka
Btgb2	30-55	<2	7.5YR5/3	7.5YR4/3	sicl	2-3msbk	sh-h	s,p	2nbrpfpo	ev	l+	gs	50-100 ka abrupt increase in carbonate suggests second buried soil?
BCb2	55-71	<2	10YR5/4	10YR4/4	l	1-2msbk	so, sh-h	ss,ps	no	e	-	gs	abundant cicada burrows, sh-h, main structure soft dry
Coxb2	71-88+	<2	10YR5/4	10YR4/4	ls	1msbk	so-lo	so,po	no	es	-		fewer cicada burrows
Location 15a, Flat surface 6 m south of gully near bison bone locale, south-central Parcel													
AC	0-6	<2	7.5YR 5/4	7.5YR 4/3	sl	1mgr	lo-so	ss, ps	no	none		cs	<1 ka (historic?)
Ab2	6-17	<2	7.5YR 4/3	7.5YR 3/3	scl	1fsbk-2mgr	sh	s, ps	no	none		cs	fs-cs; young slopewash colluvium
Btb2	17-37	<2	7.5YR 5/4	7.5YR 4/3	sicl	3msbk	h	s, p	2mkpobrpf	none		cs	50-100 ka
Btgb2	37-50+	<2	7.5YR 5/4	7.5YR 4/3	sicl	3f-msbk	h	s, p	2npobr	es-ev	1+	-	vfs
Location 15b, South gully wall, 5 m west of bison bone locale, south-central Parcel													
AC	0-9	10-20	8.75YR 6/3	8.75YR 4/4	ls	1f-mpl	so-lo	so, po	no	e-		vas	ca. 50-60 ka
Ab2	9-22	<2	7.5YR 4/3	7.5YR 3/3	scl	2f-msbk	sh-h	s,p	vnpobr	none		cs	Qec pumice + fines (fs) very few thin bridges and pore fillings
Bk1b2	22-52	<2	7.5YR 5/3	7.5YR 4/3	sl	2msbk	h	ss, ps	no	es-ev	1+	as	filaments and coatings on ped faces
Bk2b2	52-104	<2	7.5YR 5/3	7.5YR 4/3	scl	2msbk	sh-h	ss, ps	vnpobr	es-ev	1-	vas	few CaCO3 coatings on ped faces; vfs, eolian?
Btgb3	104-114+	<5	7.5YR 8/2	7.5YR 5/4	sl	3m-cabk	vh	so, ps	2n-mkbrpo	ev	3-	-	> 100-200 ka
													7.5YR 6/6 mottles, clay films remnant from Bt horizon, largely impregnated with CaCO ₃





Figure 1. Digital Elevation Map (DEM) of Pajarito Plateau showing study area

Figure 2. Geomorphology, cross section and soil pit locations, Cañada del Buey study area.
See Figures 2 and 3 for cross sections A-A' and B-B'.






Figure 3. Cross section (bottom), soil profiles and correlations (top) showing *Bison sp.* occurrence and stratigraphic relationships in Unit Qc1. See Figure 1 for cross section and soil description locations.




Figure 4. Cross section (bottom), soil profiles and correlations (top) showing stratigraphic relationships between alluvial fan (Unit Qf), hillslope (Unit Qc2), and mesa top (Unit Qe+Qc) deposits, Cañada del Buey.




Figure 5. Photographs of the left distal humerus of *Bison*. A. *Bison bison*, recent male specimen from Fort Wingate Military Reservation, McKinley County, New Mexico (catalogue number NMMNH 2364-mammalogy); B. *Bison* sp. fossil, extinct species (most likely *B. antiquus*), from Canada del Buey (catalogue number NMMNH 37623-vertebrate paleontology; NMMNH locality number L-5214).